

# On the total restrained domination number of a graph

JOANNA CYMAN      JOANNA RACZEK

*Department of Discrete Mathematics  
Faculty of Applied Physics and Mathematics  
Gdańsk University of Technology  
Narutowicza 11/12, 80-952 Gdańsk  
Poland  
{joanna, gardenia}@mif.pg.gda.pl*

## Abstract

For a graph  $G = (V, E)$ , a set  $D \subseteq V(G)$  is a *total restrained dominating set* if it is dominating and both  $\langle D \rangle$  and  $\langle V(G) - D \rangle$  are isolate free. The cardinality of a minimum total restrained dominating set in  $G$  is the *total restrained domination number* and is denoted by  $\gamma_r^t(G)$ . We investigate several properties of total restrained dominating sets and give some bounds on the total restrained domination number.

## 1 Introduction

Let  $G = (V, E)$  be a simple graph with  $|V(G)| = n(G)$  and  $|E(G)| = m(G)$ . The *neighbourhood*  $N_G(v)$  of a vertex  $v$  is the set of all vertices adjacent to  $v$  in  $G$ . The *degree*  $d_G(v)$  of a vertex  $v$  is the number of edges incident to  $v$  in  $G$ , that is  $d_G(v) = |N_G(v)|$ . The *minimum* and *maximum degree* among vertices of  $V(G)$  are denoted by  $\delta(G)$  and  $\Delta(G)$ , respectively. If  $d_G(x) = 0$ , then  $x$  is *isolate* in  $G$ . Let  $\Omega(G)$  be the set of all leaves of  $G$ , that is the set of vertices degree 1, and let  $n_1(G)$  be the cardinality of  $\Omega(G)$ . A vertex which is a neighbour of a leaf is called a *support*. Let  $S(G)$  be the set of supports in  $G$  and let  $n_S(G)$  be the cardinality of  $S(G)$ . For notational convenience, we denote  $\Omega(G) \cup S(G)$  by  $J(G)$ .

A set  $D \subseteq V(G)$  is a *dominating set* of  $G$  if for every vertex  $v \in V(G) - D$ , there exists a vertex  $u \in D$  such that  $v$  and  $u$  are adjacent. The minimum cardinality of a dominating set in  $G$  is the *domination number* denoted  $\gamma(G)$ .

A set  $D \subseteq V(G)$  is a *total dominating set* if each vertex of  $V(G)$  has a neighbour in  $D$ . The cardinality of a minimum total dominating set in  $G$  is the *total domination number* and is denoted by  $\gamma_t(G)$ . Total domination in graphs is now well studied in graph theory (see for example [1, 6]).

A set  $D \subseteq V(G)$  is a *restrained dominating set* of  $G$  if each vertex of  $V(G) - D$  has a neighbour in  $D$  as well as another neighbour in  $V(G) - D$ . The cardinality of

a minimum restrained dominating set in  $G$  is the *restrained domination number* and is denoted by  $\gamma_r(G)$ . The concept of restrained domination was introduced by Telle and Proskurowski [8], albeit indirectly, as a vertex partitioning problem. Restrained domination was studied further for example by Domke et al. [3, 4].

In this paper we study the total restrained domination number of a graph defined by De-Xiang Ma, Xue-Gang Chen and Liang Sun in [7]. A set  $D \subseteq V(G)$  is a *total restrained dominating set* if it is dominating and the induced subgraphs  $\langle D \rangle$  and  $\langle V(G) - D \rangle$  do not contain isolate vertices. The cardinality of a minimum total restrained dominating set in  $G$  is the *total restrained domination number* and is denoted by  $\gamma_r^t(G)$ . We note that every graph without isolates has a total restrained dominating set, since  $D = V(G)$  is such a set.

For unexplained terms and symbols see [5].

## 2 Bounds on the $\gamma_r^t$

In this section we first illustrate the total restrained domination number by presenting the value of  $\gamma_r^t(G)$  for some classes of graphs. We begin with some basic properties of total restrained dominating sets.

**Proposition 1** *Let  $D$  be a total restrained dominating set of a connected graph  $G$  on  $n \geq 3$  vertices. Then*

- (i) *every leaf is in  $D$ ;*
- (ii) *every support is in  $D$ ;*
- (iii)  $\gamma_r^t(G) \geq n_1(G) + n_S(G) \geq 2n_S(G)$ .

**Proof.**

- (i) Suppose  $x$  is a leaf such that  $x \notin D$  and let  $y \in N_G(x)$ . Then  $y \in D$ , because  $D$  is dominating. But then  $x$  is isolate in  $\langle V(G) - D \rangle$ , a contradiction.
- (ii) Suppose  $x$  is a leaf and let  $y \in N_G(x)$  be a support such that  $y \notin D$ . Then  $x \in D$ , because  $D$  is dominating. But then  $x$  is isolate in  $\langle D \rangle$ , a contradiction.
- (iii) By (i) and (ii) and simply observation that  $n_1(G) \geq n_S(G)$  the inequalities follows. ■

In [7] we find the following result.

**Proposition 2** *Let  $G$  be a connected graph on  $n \geq 2$  vertices. Then*

- (i)  $\gamma_r^t(K_n) = \begin{cases} 3, & n = 3, \\ 2, & \text{otherwise;} \end{cases}$



$$(ii) \gamma_r^t(K_{p,q}) = \begin{cases} p+q, & \min\{p, q\} = 1, \\ 2, & \text{otherwise;} \end{cases}$$

$$(iii) \gamma_r^t(P_n) = n - 2 \lfloor \frac{n-2}{4} \rfloor;$$

$$(iv) \gamma_r^t(C_n) = n - 2 \lfloor \frac{n}{4} \rfloor.$$

A *caterpillar* is a tree with the property that the removal of its leaves results in a path, called the *spine* of the caterpillar. The *code*  $C$  of a caterpillar  $T$  with spine  $v_1, v_2, \dots, v_s$  is the sequence of nonnegative integers  $(t_1, t_2, \dots, t_s)$  where  $t_i$  is the number of leaves adjacent to  $v_i$  in  $T$ . The substrings of consecutive zeros in  $C$  are called the *zero strings* of  $C$  and are labeled from 1 to  $k$ . For  $i = 1, 2, \dots, k$  the number of zeros in string  $i$  is denoted by  $z_i$ . For example, the caterpillar with code  $(2, 0, 0, 1, 0, 4, 3, 0, 0, 0, 1)$  has  $z_1 = 2$ ,  $z_2 = 1$  and  $z_3 = 3$ .

**Proposition 3** For any caterpillar  $T$  with code  $C$  is  $\gamma_r^t(T) = n(T) - 2 \sum_{i=1}^k \lfloor \frac{z_i+2}{4} \rfloor$ .

**Proof.** Of course each leaf of  $T$  belongs to the total restrained dominating set  $D$ . Moreover, if  $t_i \neq 0$ , then  $v_i$  is a support and thus  $v_i \in D$ . Let  $v_j, v_{j+1}, \dots, v_{j+z_i-1}$  be the vertices of zero string and let  $x$  and  $y$  be leaves adjacent to  $v_{j-1}$  and  $v_{j+z_i}$ , respectively. Then  $x, v_{j-1}, v_j, v_j + 1, \dots, v_{j+z_i-1}, v_{j+z_i}, y$  induces a path  $P$  on  $z_i + 4$  vertices in  $T$  and by Proposition 2,  $\gamma_r^t(P) = z_i + 4 - 2 \lfloor \frac{z_i+2}{4} \rfloor$ . As  $\{x, v_{j-1}, v_{j+z_i}, y\} \subseteq D$ , we conclude that exactly  $2 \lfloor \frac{z_i+2}{4} \rfloor$  vertices of the zero string  $j$  do not belong to  $D$ . Consequently,  $|D| = n(T) - 2 \sum_{i=1}^k \lfloor \frac{z_i+2}{4} \rfloor$ . ■

Now we find some bounds on the total restrained domination number. For this purpose, denote by  $\mathcal{A}$  the family of connected graphs such that  $G$  belongs to  $\mathcal{A}$  if and only if every edge of  $G$  is incident to a support or if  $G$  is a cycle on three vertices.

**Theorem 4** If a graph  $G$  is connected, then  $\gamma_r^t(G) = n(G)$  if and only if  $G$  belongs to the family  $\mathcal{A}$ .

**Proof.** For  $n(G) = 2$  and  $n(G) = 3$  the result is obvious. Thus assume  $n(G) \geq 4$ . Let  $G \in \mathcal{A}$ . By Proposition 1,  $J(G)$  is a subset of every total restrained dominating set  $D$ . Moreover, no two vertices of  $V(G) - J(G)$  are adjacent, which implies that  $D = V(G)$  is a minimum total restrained dominating set of  $G$ .

Now we prove that if  $G \notin \mathcal{A}$ , then  $\gamma_r^t(G) \leq n(G) - 2$ . If  $G \notin \mathcal{A}$ , then there exists an edge  $uv$  where neither  $u$  nor  $v$  is a support. Of course  $d_G(u) > 1$  and  $d_G(v) > 1$ . Let  $D = V(G) - \{u, v\}$ . Obviously  $D$  is a dominating set and  $\langle V(G) - D \rangle$  is connected. If  $\langle D \rangle$  is without isolates, then  $D$  is a total restrained dominating set of  $G$  of cardinality  $n(G) - 2$ . If it is not the case, then there exists a vertex  $x \in V(G) - J(G)$  such that  $N_G(x) = \{u, v\}$ . As  $G$  is not a cycle on 3 vertices, we conclude that  $u$  or  $v$  has degree 3 or more. Without loss of generality we assume that  $d_G(u) \geq 3$  and let  $D' = V(G) - \{v, x\}$ . This time  $\langle V(G) - D' \rangle$  is connected,  $D'$  is dominating and  $\langle D' \rangle$  is without isolates. As  $|D'| = n(G) - 2$ , we obtain that  $\gamma_r^t(G) \leq n(G) - 2$ . ■



**Corollary 5** *If  $G$  is a connected graph with  $n(G) \geq 4$  and  $\delta(G) \geq 2$ , then  $\gamma_r^t(G) \leq n(G) - 2$ .*

**Theorem 6** *If  $G$  is a connected graph on  $n \geq 4$  vertices and  $2 \leq \delta(G) \leq n - 2$ , then  $\gamma_r^t(G) \leq n - \delta(G)$ .*

**Proof.** Let  $x$  be a vertex with  $d_G(x) = \delta(G)$ . We consider two cases:

*Case 1.* There exists a vertex  $y \in N_G(x)$  such that  $N_G[y] \neq N_G[x]$ . Let  $A = N_G[x] - \{y\}$  and  $D = V(G) - A$ . As  $|A| = \delta(G)$ , each vertex of  $A$  has a neighbour in  $D$ , so  $D$  is dominating. Moreover,  $\langle A \rangle$  is isolate-free. Suppose now that  $u$  is isolate in  $\langle D \rangle$ . Then obviously  $u \neq y$  and  $N_G(u) = A$ . Hence  $u$  is a neighbour of  $x$ . But then  $d_G(x) > \delta(G)$ , a contradiction. We conclude that  $D$  is a total restrained dominating set of  $G$  and thus  $\gamma_r^t(G) \leq |D| = n - \delta(G)$ .

*Case 2.*  $N_G[y] = N_G[x]$  for every vertex  $y \in N_G(x)$ . Then the connectivity of  $G$  implies that  $G$  is complete. But then  $\delta(G) = n - 1$ , a contradiction. ■

**Proposition 7** *Let  $G$  be a connected graph on  $n(G) \geq 3$ . Then  $\gamma_r^t(G) = n_1(G) + n_S(G)$  if and only if  $\langle V(G) - J(G) \rangle$  has no isolates and each vertex of  $V(G) - J(G)$  has a neighbour in  $J(G)$ .*

**Proof.** Assume first that  $G$  is connected, each vertex of  $V(G) - J(G)$  has a neighbour in  $J(G)$  and  $\langle V(G) - J(G) \rangle$  has no isolates. Then by Proposition 1,  $\gamma_r^t(G) \geq n_1(G) + n_S(G)$ . We claim that  $J(G)$  is a total restrained dominating set of  $G$ . It is clear that  $\langle J(G) \rangle$  does not contain isolates. Moreover, we know that  $\langle V(G) - J(G) \rangle$  has no isolates and each vertex of  $V(G) - J(G)$  has a neighbour in  $J(G)$ . Hence,  $\gamma_r^t(G) = |J(G)| = n_1(G) + n_S(G)$ .

Conversely, if  $\gamma_r^t(G) = n_1(G) + n_S(G)$ , then  $J(G)$  is a minimum total restrained dominating set of  $G$ . Thus,  $\langle V(G) - J(G) \rangle$  has no isolates and each vertex of  $V(G) - J(G)$  is dominated by a vertex of  $J(G)$ , so each vertex of  $V(G) - J(G)$  has a neighbour in  $J(G)$ . ■

In order to present our next bound on the total restrained domination number, we denote by  $\mathcal{B}$  the family of all connected graphs  $G$  in which there exists a set  $D \subseteq V(G)$  with property that

- the induced subgraphs  $\langle D \rangle$  and  $\langle V(G) - D \rangle$  are isolate free;
- each vertex of  $D$  has exactly  $\Delta(G) - 1$  neighbours in  $V(G) - D$ ;
- each vertex of  $V(G)$  has exactly one neighbour in  $D$ .

**Theorem 8** *If  $G$  is a connected graph without isolates on  $n$  vertices and with  $m$  edges, then*

$$\frac{n}{\Delta(G)} \leq \gamma_r^t(G) \leq 2m - n + 2,$$

*with equality for the lower bound if and only if  $G$  belongs to the family  $\mathcal{B}$  and equality for the upper bound if and only if  $G$  is a tree belonging to the family  $\mathcal{A}$ .*



**Proof.** Since  $\frac{n}{\Delta(G)} \leq \gamma_t(G) \leq \gamma_r^t(G)$  the lower bound follows. If  $G \in \mathcal{B}$ , then  $D$  is a total restrained dominating set of  $G$  and hence  $|D| \geq \gamma_r^t(G) \geq \frac{n}{\Delta(G)}$ . On the other hand,  $n(G) = |D| + (\Delta(G) - 1) \cdot |D| = \Delta(G) \cdot |D|$ , so  $|D| = \frac{n}{\Delta(G)}$ . Thus  $\gamma_r^t(G) = \frac{n}{\Delta(G)}$ .

Conversely, assume that  $\gamma_r^t(G) = \frac{n}{\Delta(G)}$  and let  $D$  be a minimum total restrained dominating set of  $G$ . Then  $\langle D \rangle$  and  $\langle V(G) - D \rangle$  are isolate free and each vertex of  $V(G)$  has at least one neighbour in  $D$ . We prove that each vertex of  $D$  has exactly  $\Delta(G) - 1$  neighbours in  $V(G) - D$ . If not, then there exists a vertex  $u \in D$  such that  $u$  has at most  $\Delta(G) - 2$  neighbours in  $V(G) - D$ . For this reason

$$n \leq \gamma_r^t(G) + (\Delta(G) - 1) \cdot (\gamma_r^t(G) - 1) + \Delta(G) - 2 < \Delta(G) \cdot \gamma_r^t(G)$$

and hence  $\gamma_r^t(G) > \frac{n}{\Delta(G)}$ . Further, since  $D$  is a total restrained dominating set of  $G$ , it follows that each vertex of  $D$  has exactly one neighbour in  $D$ .

If there exists a vertex  $u \in V(G)$  such that  $u$  has at least two neighbors in  $D$ , then  $u \notin D$  and similarly

$$n \leq \gamma_r^t(G) + (\Delta(G) - 1) \cdot (\gamma_r^t(G) - 1) + \Delta(G) - 2 < \Delta(G) \cdot \gamma_r^t(G)$$

and hence  $\gamma_r^t(G) > \frac{n}{\Delta(G)}$ .

Now we prove the upper bound. Obviously,  $\gamma_r^t(G) \leq n = 2(n - 1) - n + 2$  and since  $G$  is connected,  $m \geq n - 1$ . Thus,  $\gamma_r^t(G) \leq 2m - n + 2$ . Now we show that  $\gamma_r^t(G) = 2m - n + 2$  if and only if  $G$  is a tree belonging to the family  $\mathcal{A}$ . If  $G$  is a tree belonging to the family  $\mathcal{A}$ , then  $m = n - 1$  and  $\gamma_r^t(G) = n = 2m - n + 2$ . Conversely, let  $\gamma_r^t(G) = 2m - n + 2$ . Then  $2m - n + 2 \leq n$ , which implies that  $m \leq n - 1$ . Since  $G$  is connected,  $m \geq n - 1$ . We conclude that  $m = n - 1$ , so  $G$  is a tree with  $\gamma_r^t(G) = n$  and by Theorem 4,  $G$  belongs to the family  $\mathcal{A}$ . ■

In order to present our next bound on the total restrained domination number, we denote by  $\mathcal{C}$  a family of all connected graphs such that  $G$  belongs to  $\mathcal{C}$  if and only if there exists a set  $A \subseteq V(G)$  such that  $\langle A \rangle = pK_2$  and  $\langle V(G) - A \rangle = qK_2$  for some non-negative integers  $p$  and  $q$ , and each vertex of  $V(G) - A$  has exactly one neighbour in  $A$ .

**Theorem 9** *If  $G$  is a graph without isolates on  $n$  vertices and with  $m$  edges, then  $\gamma_r^t(G) \geq \frac{3}{2}n - m$  with equality for the bound if and only if  $G$  belongs to the family  $\mathcal{C}$ .*

**Proof.** Let  $D$  be a minimum total restrained dominating set in  $G$ . Since  $\langle D \rangle$  and  $\langle V - D \rangle$  are isolate free and  $D$  is dominating, we have the following inequalities:

$$\begin{aligned} m(\langle D \rangle) &\geq \frac{\gamma_r^t(G)}{2}, \\ m(\langle V(G) - D \rangle) &\geq \frac{n - \gamma_r^t(G)}{2}, \\ m_{\gamma_r^t} &\geq n - \gamma_r^t(G), \end{aligned} \tag{1}$$

where  $m_{\gamma_r^t}$  is the number of the edges connecting vertices of  $V(G) - D$  to vertices of  $D$ . By summing the inequalities we obtain

$$m = m(\langle D \rangle) + m(\langle V - D \rangle) + m_{\gamma_r^t} \geq \frac{3}{2}n - \gamma_r^t(G),$$



and thus  $\gamma_r^t(G) \geq \frac{3}{2}n - m$ .

If  $G$  belongs to the family  $\mathcal{C}$ , then  $n = 2p + 2q$ ,  $m = p + 3q$  and  $A$  is a total restrained dominating set of  $G$ . As  $|A| \geq \gamma_r^t(T) \geq \frac{3}{2}n - m$  we conclude that  $|A| = 2p \geq \frac{3}{2}n - m = 2p$ . This implies that  $A$  is a minimum total restrained dominating set and  $\gamma_r^t(G) = \frac{3}{2}n - m$ .

Conversely, if  $\gamma_r^t(G) = \frac{3}{2}n - m$  and  $D$  is a minimum total restrained dominating set of  $G$ , then we have equalities in (1). This implies that  $\langle D \rangle = pK_2$ ,  $\langle V(G) - D \rangle = qK_2$  and each vertex of  $V(G) - D$  has exactly one neighbour in  $D$ . Hence  $G$  belongs to the family  $\mathcal{C}$ . ■

**Corollary 10** *If  $G$  is a graph without isolates and does not contain a perfect matching, then  $\gamma_r^t(G) \geq \frac{3}{2}n - m + 1$ .*

**Corollary 11** *If  $G$  is a graph without isolates and with odd number of vertices, then  $\gamma_r^t(G) \geq \frac{3(n+1)}{2} - m$ .*

**Proposition 12** *If  $G$  is hamiltonian, then  $\gamma_r^t(G) \leq n - 2\lfloor \frac{n}{4} \rfloor$ .*

**Proof.** Let  $G$  be hamiltonian and let  $C$  be a hamiltonian cycle of  $G$ . By Proposition 2,  $\gamma_r^t(C) = n - 2\lfloor \frac{n}{4} \rfloor$ . As adding an edge does not increase the total restrained domination number of  $G$ , we obtain the result. ■

### 3 Edge subdivision and vertex removal

In this section we study the influence of edge subdivision and vertex removal on the total restrained domination number of a graph. An *edge subdivision* in a nonempty graph  $G$  is an operation of removal of an edge  $e = uv$  and the addition of a new vertex  $w$  and edges  $uw$  and  $vw$ . A graph obtained from  $G$  by subdividing the edge  $e = uv$  is denoted by  $G \oplus w_{uv}$ .

**Theorem 13** *For every integer  $k$ ,  $k \geq 2$ , there exists a graph  $G$  such that  $\gamma_r^t(G \oplus w_{uv}) - \gamma_r^t(G) = k$ .*

**Proof.** We construct graphs  $G$  and  $G \oplus w_{uv}$  as follows. We begin with two stars  $K_{1,k-1}$  and denote its centers by  $u$  and  $v$ . Next we add a vertex  $x$  and edges joining  $x$  with all vertices of the stars. Finally we add an edge  $uv$  and a pendant edge  $xx'$  (see Fig. 1). It is easy to observe that  $\{x, x'\}$  is a minimum total restrained dominating set of  $G$ .

For the graph  $G \oplus w_{uv}$  notice, that the set  $N_G[v] \cup \{x, x'\}$  is a minimum total restrained dominating set of cardinality  $k + 2$ . Thus  $\gamma_r^t(G \oplus w_{uv}) - \gamma_r^t(G) = k$ . ■

**Theorem 14** *For every integer  $k$ ,  $k \geq 2$ , there exists a graph  $H$  such that  $\gamma_r^t(H) - \gamma_r^t(H \oplus z_{uv}) = k - 2$ .*






Figure 1: Graphs  $G$  and  $G \oplus w_{uv}$  for  $k=5$




Figure 2: Graphs  $H$  and  $H \oplus z_{uw}$  for  $k=5$

**Proof.** Let  $H$  be the graph  $G \oplus w_{uv}$  constructed as in the proof of Theorem 13. Then  $\gamma_r^t(H) = k + 2$  (see Fig. 2).

For the graph  $H \oplus z_{uw}$  notice, that the set  $\{x, x', w, z\}$  is a minimum total restrained dominating set of cardinality 4. Thus  $\gamma_r^t(H) - \gamma_r^t(H \oplus z_{uw}) = k + 2 - 4 = k - 2$ . ■

Now we investigate how removal of a vertex influences the total restrained domination number. Let us denote by  $G - x$  the graph received from  $G$  by removing the vertex  $x \in V(G)$  with all edges incident to  $x$ .

**Theorem 15** For every integer  $k, k \geq 1$ , there exists a graph  $G$  such that  $\gamma_r^t(G) - \gamma_r^t(G - x) = k + 2$ .

**Proof.** Let  $G$  be the graph obtained from the join  $K_{1,k} + K_1$  by adding to each of the two vertices of degree  $k + 1$  a leaf (see Fig. 3). It is easy to observe that  $V(G)$  is a minimum total restrained dominating set of  $G$ . Thus,  $\gamma_r^t(G) = k + 4$ .

Denote by  $x$  a leaf in  $G$  and consider the graph  $G - x$ . Obviously  $\gamma_r^t(G - x) = 2$  and finally  $\gamma_r^t(G) - \gamma_r^t(G - x) = k + 2$ . ■






Figure 3: Graphs  $G$  and  $G - x$  for  $k = 4$

**Theorem 16** For every integer  $k, k \geq 2$ , there exists a graph  $G$  such that  $\gamma_r^t(G - x) - \gamma_r^t(G) = k - 1$ .

**Proof.** Let  $G$  be a join  $K_{1,k} + K_1$ . Then  $\gamma_r^t(G) = 2$ . Let  $x$  be a vertex of degree  $k + 1$ . As  $G - x$  is a star,  $\gamma_r^t(G - x) = k + 1$ .




Figure 4: Graphs  $G$  and  $G - x$  for  $k = 4$

Thus  $\gamma_r^t(G - x) - \gamma_r^t(G) = k - 1$ . ■

#### 4 Comparison $\gamma_r^t$ to other types of domination numbers

In this section we investigate relations between total restrained domination number and other types of domination numbers.

A set  $D \subseteq V(G)$  is a *connected dominating set* if it is dominating and the induced subgraph  $\langle D \rangle$  is connected. The cardinality of a minimum connected dominating set in  $G$  is the *connected domination number* and is denoted by  $\gamma_c(G)$ .

Numbers  $\gamma_r^t(G)$  and  $\gamma_c(G)$  are incomparable. Consider, for example, any star and a path on more than 10 vertices.

A set  $D \subseteq V(G)$  is a *doubly connected dominating set* if it is dominating and the induced subgraphs  $\langle D \rangle$  and  $\langle V(G) - D \rangle$  are connected. The cardinality of a minimum doubly connected dominating set in  $G$  is the *doubly connected domination number* and is denoted by  $\gamma_{cc}(G)$ . Properties of the doubly connected domination number are





studied in [2]. Here we note that for any connected graph  $G$ ,  $1 \leq \gamma_{cc}(G) \leq n(G) - 1$  and  $\gamma_{cc}(G) = 1$  if and only if  $G$  has a non-cut vertex of degree  $n(G) - 1$ .

**Theorem 17** *For an arbitrary connected graph  $G$  with  $n(G) > 1$ ,*

- (i)  $\gamma_r^t(G) \leq \gamma_{cc}(G) + 2$ ;
- (ii)  $\gamma_r^t(G) = \gamma_{cc}(G) + 2$  if and only if  $G$  is the cycle  $C_3$ ;
- (iii) For any integer  $k$ , where  $k \geq -2$ , there exists a graph  $G$  such that  $\gamma_{cc}(G) - \gamma_r^t(G) = k$ .

**Proof.**

- (i) If  $2 \leq \gamma_{cc}(G) \leq n - 2$ , then every doubly connected dominating set is a total restrained dominating set. That is, if  $2 \leq \gamma_{cc}(G) \leq n - 2$ , then  $\gamma_r^t(G) \leq \gamma_{cc}(G)$ . Hence,  $\gamma_r^t(G) \leq \gamma_{cc}(G) + 2$ .
- (ii) If  $G$  is the cycle  $C_3$ , the result is obvious. Thus assume  $\gamma_r^t(G) = \gamma_{cc}(G) + 2$ . By the proof of (i),  $\gamma_{cc}(G) = 1$  and hence  $\gamma_r^t(G) = 3$ . This implies that there exists a vertex  $u \in V(G)$  of degree  $n(G) - 1$  such that  $\langle V(G) - \{u\} \rangle$  is connected, and so  $\delta(G) \geq 2$ . Moreover, for each  $v \in V(G) - \{u\}$  the set  $\{u, v\}$  is not a total restrained dominating set, although  $\{u, v\}$  is dominating and connected. Hence we conclude that for each  $v \in V(G) - \{u\}$  the induced subgraph  $\langle V(G) - \{u, v\} \rangle$  has isolates. Denote by  $x$  an isolate vertex of  $\langle V(G) - \{u, v\} \rangle$ . Of course  $N_G(x) = \{u, v\}$ . Similarly,  $v$  must be the only isolate in  $\langle V(G) - \{u, x\} \rangle$  and the connectivity of  $\langle V(G) - \{u\} \rangle$  implies that  $n(G) = 3$ . Consequently,  $G = C_3$ .
- (iii) For  $k = -2$  let  $G = C_3$ , for  $k = -1$  let  $G = P_4$  and for  $k = 0$  let  $G = C_5$ . For  $k \geq 1$  let  $G$  be the graph as in Fig. 5.




Figure 5: Graph  $G$  with  $\gamma_{cc}(G) - \gamma_r^t(G) = k$  for  $k \geq 1$ .

It is left to the reader as an easy exercise to check that  $\gamma_{cc}(G) - \gamma_r^t(G) = k$ . ■

For an arbitrary graph  $G$  which is isolate free, we have the following inequalities:  $\gamma(G) \leq \gamma_r^t(G)$ ,  $\gamma_t(G) \leq \gamma_r^t(G)$  and  $\gamma_r(G) \leq \gamma_r^t(G)$ . However, each of these differences may be arbitrarily large.



**Proposition 18** For each positive integer  $k$  there exists graphs  $G_1, G_2, G_3$  such that  $\gamma_r^t(G_1) - \gamma(G_1) = k$ ,  $\gamma_r^t(G_2) - \gamma_t(G_2) = k$  and  $\gamma_r^t(G_3) - \gamma_r(G_3) = k$ .

**Proof.** Let  $G_1$  be the star  $K_{1,k}$ , let  $G_2$  be the star  $K_{1,k+1}$  and let  $G_3$  be the subdivided star  $K_{1,k}^*$  for  $k \geq 2$  and let  $G_3$  be a graph obtained from the cycle  $C_4$  by adding a pendant edge for  $k = 1$ . ■

## References

- [1] E. J. Cockayne, R. M. Dawes and S. T. Hedetniemi, Total domination in graphs, *Networks* **10** (1980), 211–219.
- [2] J. Cyman, M. Lemańska and J. Raczek, On the doubly connected domination number of a graph, *Central European J. Math.* **4** (2006), 34–45.
- [3] G. S. Domke, J. H. Hattingh, S. T. Hedetniemi, R. C. Laskar and L. R. Markus, Restrained domination in graphs, *Discrete Math.* **203** (1999), 61–69.
- [4] G. S. Domke, J. H. Hattingh, S. T. Hedetniemi and L. R. Markus, Restrained domination in trees, *Discrete Math.* **211** (2000), 1–9.
- [5] T. W. Haynes, S. T. Hedetniemi and P. J. Slater, *Fundamentals of domination in graphs*, Marcel Dekker, New York, 1998.
- [6] T. W. Haynes, S. T. Hedetniemi and P. J. Slater (eds), *Domination in Graphs: Advanced Topics*, Marcel Dekker, New York, 1998.
- [7] D. Ma, X. Chen and L. Sun, On total restrained domination in graphs, *Czechoslovak Math. J.* **55** (2005), 165–173.
- [8] J. A. Telle and A. Proskurowski, Algorithms for vertex partitioning problems on partial  $k$ -trees, *SIAM J. Discrete Math.* **10** (1997), 529–550.

(Received 20 Apr 2005; revised 17 Oct 2005)

