

MINIMIZATION OF THE NUMBER OF PERIODIC POINTS FOR SMOOTH SELF-MAPS OF CLOSED SIMPLY-CONNECTED 4-MANIFOLDS

GRZEGORZ GRAFF

Faculty of Applied Physics and Mathematics
Gdansk University of Technology
Narutowicza 11/12, 80-233 Gdansk, Poland

JERZY JEZIERSKI

Institute of Applications of Mathematics
Warsaw University of Life Sciences (SGGW)
Nowoursynowska 159, 00-757 Warsaw, Poland

ABSTRACT. Let M be a smooth closed simply-connected m -dimensional manifold, f be a smooth self-map of M and r be a given natural number. The invariant $D_r^m[f]$ defined by the authors in [Forum Math. 21 (2009)] is equal to the minimum of $\#\text{Fix}(g^r)$ over all maps g smoothly homotopic to f . In this paper we calculate the invariant $D_r^4[f]$ for the class of smooth self-maps of 4-manifolds with fast growth of Lefschetz numbers and for r being a product of different primes.

1. Introduction. One of the fundamental problems in periodic point theory is to find minimal number of periodic points in the homotopy class of a given map. Let f be a self-map of a compact manifold M . B. Jiang introduced in 1983 the invariant $NF_r(f)$ which estimates from above $\#\text{Fix}(g^r)$ for all g homotopic to f [14]. J. Jezierski proved in 2006 that the invariant is the best estimation if the dimension of M is at least 3 [12]. This means that $NF_r(f)$ is equal to the minimal number of elements in $\text{Fix}(g^r)$ over all g homotopic to f . In the last years the invariant was computed in many special cases, see for example: [10], [13], [16], [18].

In the recent papers [4], [6] the authors developed the theory for the smooth (i.e. C^1) category, searching for the minimum in smooth homotopy class. As a result, two counterparts of $NF_r(f)$ were found: $D_r^m[f]$ for simply-connected manifolds [4] and its generalization $NJD_r^m[f]$ for non simply-connected ones [6]. The crucial demanding for effective computation of the invariants is the knowledge of all sequences of local fixed point indices of iterations at a periodic p -orbit for smooth maps in the given dimension m , called $DD^m(p)$ sequences. This information was provided in dimension 3 in the paper [9], which made it possible to compute the value of $D_r^3[f]$ for $S^2 \times I$ [4], S^3 [5], a two-holed 3-dimensional closed ball [3] and $NJD_r^3[f]$ for $\mathbb{R}P^3$ [7]. Recently, in [8] we provided the list of all possible sequences of local indices of iterations in arbitrary dimension, which allows one to calculate the invariants for self-maps of higher dimensional manifolds. In this paper we partially realize this

2000 *Mathematics Subject Classification.* Primary: 37C25, 55M20; Secondary: 37C05.

Key words and phrases. Indices of iterations, smooth maps, Nielsen number.

Research supported by Polish National Research Grant No. N N201 373236.

programme for simply-connected manifolds and $m = 4$. We calculate $D_r^4[f]$ under the assumption that the so-called *periodic expansion* of $\{L(f^n)\}_{n=1}^\infty$, the sequence of the Lefschetz numbers of iterations, has only non-zero coefficients. This property holds for example for maps with fast grow of the sequence of Lefschetz numbers, such as self-maps of S^4 with degree d satisfying $|d| > 1$.

The paper is organized in the following way. First, in Section 2 we give the definition of $D_r^m[f]$ which is expressed in terms of $DD^m(p)$ sequences. Next, in Section 3 we provide the list of all $DD^m(1)$ sequences and prove that in order to calculate $D_r^4[f]$ it is enough to use only $DD^4(1)$ sequences. Finally, in Section 4 we calculate $D_r^4[f]$ for r being a product of different primes (Theorem 4.8).

2. The invariant $D_r^m[f]$. The notion of *Differential Dold* sequences (DD sequences in short) introduced in [4] is used in the definition of the invariant $D_r^m[f]$. A $DD^m(p)$ sequence is a sequence of integers that can be locally realized as a sequence of indices on an isolated p -orbit for some smooth map.

Definition 2.1. A sequence of integers $\{c_n\}_{n=1}^\infty$ is called a $DD^m(p)$ sequence if there is a C^1 map $\phi : U \rightarrow \mathbb{R}^m$ ($U \subset \mathbb{R}^m$) and its isolated p -orbit P such that $c_n = \text{ind}(\phi^n, P)$. If this equality holds for $n|r$, where r is fixed, then the finite sequence $\{c_n\}_{n|r}$ will be called a $DD^m(p|r)$ sequence.

Let r be fixed. The minimal decomposition of the sequence of Lefschetz numbers of iterations into $DD^m(p|r)$ sequences gives the value of $D_r^m[f]$.

Definition 2.2. Let $\{L(f^n)\}_{n|r}$ be a finite sequence of Lefschetz numbers. We decompose $\{L(f^n)\}_{n|r}$ into the sum:

$$L(f^n) = c_1(n) + \dots + c_s(n), \quad (1)$$

where c_i is a $DD^m(l_i|r)$ sequence for $i = 1, \dots, s$. Each such decomposition determines the number $l = l_1 + \dots + l_s$. We define the number $D_r^m[f]$ as the smallest l which can be obtained in this way.

The invariant $D_r^m[f]$ is equal to the minimal number of r -periodic points in smooth homotopy class of f .

Theorem 2.3. ([4]) *Let M be a smooth closed connected and simply-connected manifold of dimension $m \geq 3$ and $r \in \mathbb{N}$ a fixed number. Then,*

$$D_r^m[f] = \min\{\#\text{Fix}(g^r) : g \text{ is smoothly homotopic to } f\}.$$

Periodic expansion is a convenient method of storing the data connected with the sequence of indices of iterations. Each such sequence can be expanded as a combination of some basic periodic sequences $\{\text{reg}_k\}_n$ taken with integral coefficients.

Definition 2.4. For a given k we define the basic sequence:

$$\text{reg}_k(n) = \begin{cases} k & \text{if } k|n, \\ 0 & \text{if } k \nmid n. \end{cases}$$

A sequence of indices of iterations (as well as a sequence of Lefschetz numbers of iterations) may be written down in the form of *periodic expansion* (cf. [15]), namely:

$$\text{ind}(f^n, x_0) = \sum_{k=1}^{\infty} a_k \text{reg}_k(n), \quad (2)$$

where $a_n = \frac{1}{n} \sum_{k|n} \mu(k) \text{ind}(f^{(n/k)}, x_0)$, a_n are integers, μ is the classical Möbius function, i.e. $\mu : \mathbb{N} \rightarrow \mathbb{Z}$ is defined by the following three properties: $\mu(1) = 1$, $\mu(k) = (-1)^s$ if k is a product of s different primes, $\mu(k) = 0$ otherwise.

The fact that the coefficients a_n are integers follows from the result of Dold [2].

The invariant $D_r^m[f]$ is defined in terms of $DD^m(p)$ sequences. On the other hand, it is enough to know only the forms of $DD^m(1)$ sequences, because every $DD^m(p)$ sequence can be obtained from some $DD^m(1)$ one.

Definition 2.5. We will say that the $DD^m(p)$ sequence $\{\tilde{c}_n\}_n$ comes from the given $DD^m(1)$ sequence $\{c_n\}_n$ with the periodic expansion $c_n = \sum_{d=1}^{\infty} a_d \text{reg}_d(n)$ if the periodic expansion of $\{\tilde{c}_n\}_n$ has the form:

$$\tilde{c}_n = \sum_{d=1}^{\infty} a_d \text{reg}_{pd}(n).$$

Theorem 2.6 ([4]). *Every $DD^m(p)$ sequence comes from some $DD^m(1)$ sequence.*

3. Local indices of iterations in dimension 4. In this section we give the complete list of all sequences of local indices of iterations of a smooth map in dimension 4 i.e. the list of all $DD^4(1)$ sequences. Let us mention here that the forms of indices of iterations for continuous maps are known since 1991 [1], and recently indices of iterations have been found also for other important classes of maps, such as holomorphic maps [22] and planar homeomorphisms [17], [21].

Definition 3.1. Let H be a finite subset of natural numbers, we introduce the following notation.

By $\text{LCM}(H)$ we mean the least common multiple of all elements in H with the convention that $\text{LCM}(\emptyset) = 1$. We define the set \bar{H} by: $\bar{H} = \{\text{LCM}(Q) : Q \subset H\}$.

For natural s we denote by $L(s)$ any set of natural numbers of the form \bar{L} with $\#L = s$ and $1, 2 \notin L$.

By $L_2(s)$ we denote any set of natural numbers of the form \bar{L} with $\#L = s + 1$ and $1 \notin L, 2 \in L$.

Theorem 3.2 (Main Theorem I in [8]). *Let f be a C^1 self-map of \mathbb{R}^m , where m is even. Then the sequence of local indices of iterations $\{\text{ind}(f^n, 0)\}_{n=1}^{\infty}$ has one of the following forms:*

$$(A^e) \quad \text{ind}(f^n, 0) = \sum_{k \in L_2(\frac{m-4}{2})} a_k \text{reg}_k(n).$$

$$(B^e) \quad \text{ind}(f^n, 0) = \sum_{k \in L(\frac{m-2}{2})} a_k \text{reg}_k(n).$$

$$(C^e), (D^e), (E^e) \quad \text{ind}(f^n, 0) = \sum_{k \in L_2(\frac{m-2}{2})} a_k \text{reg}_k(n),$$

where

$$a_1 = \begin{cases} 1 & \text{in the case } (C^e), \\ -1 & \text{in the case } (D^e), \\ 0 & \text{in the case } (E^e). \end{cases}$$

$$(F^e) \operatorname{ind}(f^n, 0) = \sum_{k \in L(\frac{m}{2})} a_k \operatorname{reg}_k(n),$$

where $a_1 = 1$.

By $[d, l]$ we denote the least common multiple of d and l .

Theorem 3.3. *The list of all $DD^4(1)$ sequences is the following:*

$$(A) \quad c_A(n) = a_1 \operatorname{reg}_1(n) + a_2 \operatorname{reg}_2(n);$$

$$(B) \quad c_B(n) = a_1 \operatorname{reg}_1(n) + a_d \operatorname{reg}_d(n);$$

$$(C - E)_{\text{odd}}$$

$$c_X(n) = \varepsilon_X \operatorname{reg}_1(n) + a_2 \operatorname{reg}_2(n) + a_d \operatorname{reg}_d(n) + a_{2d} \operatorname{reg}_{2d}(n),$$

where $\varepsilon_X \in \{-1, 0, 1\}$, $X \in \{C, D, E\}$, d is odd.

$$(C - E)_{\text{even}}$$

$$c_X(n) = \varepsilon_X \operatorname{reg}_1(n) + a_2 \operatorname{reg}_2(n) + a_d \operatorname{reg}_d(n),$$

where $\varepsilon_X \in \{-1, 0, 1\}$, $X \in \{C, D, E\}$, d is even.

$$(F) \quad c_F(n) = \operatorname{reg}_1(n) + a_d \operatorname{reg}_d(n) + a_l \operatorname{reg}_l(n) + a_{[d,l]} \operatorname{reg}_{[d,l]}(n),$$

In all cases $d, l \geq 3$ and $a_i \in \mathbb{Z}$.

Proof. We apply Theorem 3.2 for $m = 4$, obtaining the corresponding parts of the thesis. For example, to obtain the case (F), we use (F^e) and get:

$$L\left(\frac{m}{2}\right) = L(2) = \overline{\{d, l\}} = \operatorname{LCM}\{Q \subset \{d, l\}\} = \{1, d, l, [d, l]\}.$$

□

Corollary 1. *Let us notice that any $DD^4(1)$ sequence has one of the following forms:*

$$1. \quad a_1 \operatorname{reg}_1(n) + a_d \operatorname{reg}_d(n);$$

for $a_1, a_d \in \mathbb{Z}$.

$$2. \quad \varepsilon \operatorname{reg}_1(n) + a_2 \operatorname{reg}_2(n) + a_d \operatorname{reg}_d(n) + \gamma_d a_{2d} \operatorname{reg}_{2d}(n);$$

for $a_2, a_d \in \mathbb{Z}$, $\varepsilon = 0, \pm 1$, $\gamma_d = 0$ if d is even and $\gamma_d = 1$ if d is odd.

$$3. \quad \operatorname{reg}_1(n) + a_d \operatorname{reg}_d(n) + a_l \operatorname{reg}_l(n) + a_{[d,l]} \operatorname{reg}_{[d,l]}(n);$$

for $a_d, a_l \in \mathbb{Z}$, $d, l \geq 3$.

The next two lemmas show that during the calculation of $D_r^4[f]$ we may consider only $DD^4(1)$ sequences, which makes the computation much easier.

Lemma 3.4 (Remark 4.6 in [4]). *For $m \geq 3$ in Definition 2.2 of $D_r^m[f]$ we can equivalently use only $DD^m(p|r)$ sequences such that $p < 2^{\lceil \frac{m+1}{2} \rceil}$.*

Lemma 3.5. *To calculate $D_r^4[f]$ it is enough to consider only $DD^4(1)$ sequences.*

Proof. By Lemma 3.4 it is enough to consider only such $DD^4(p|r)$ sequences for which $p \leq 3$.

We show that

(1) every $DD^4(2|r)$ sequence is a sum of at most two $DD^4(1|r)$ sequences.

(2) every $DD^4(3|r)$ sequence is a sum of at most three $DD^4(1|r)$ sequences.

Proof of (1). Using Theorem 2.6 we find the forms of all $DD^4(2|r)$ sequences, each of which comes from some $DD^4(1|r)$ sequences of one of the types (A)-(F). Next, we represent each $DD^4(2|r)$ sequence as a sum of at most two $DD^4(1|r)$ sequences.



(A) $a_2\text{reg}_2(n) + a_4\text{reg}_4(n)$ is in fact the $DD^4(1|r)$ sequence of the type $(D)_{\text{even}}$.

(B) $a_2\text{reg}_2(n) + a_{2d}\text{reg}_{2d}(n)$ the same argument as above is true.

(C-E) For d odd we have that $[4, 2d] = 4d$, then

$$\varepsilon_X \text{reg}_2(n) + a_4 \text{reg}_4(n) + a_{2d} \text{reg}_{2d}(n) + a_{4d} \text{reg}_{4d}(n) = \text{reg}_1(n) + a_4 \text{reg}_4(n) + a_{2d} \text{reg}_{2d}(n) + a_{4d} \text{reg}_{4d}(n) \tag{F}$$

$$-\text{reg}_1(n) + \varepsilon_X \text{reg}_2(n) \tag{A}$$

where on the right-hand side of the above formula we indicated that the first sum is realized by a sequence of the type (F) and the second by (A).

In the same way we deal with the case of d even (every sequence is a sum of a sequence of the type (F) and (A)).

(F) Notice that $[2d, 2l] = 2[d, l]$, thus we get

$$\text{reg}_2(n) + a_{2d} \text{reg}_{2d}(n) + a_{2l} \text{reg}_{2l}(n) + a_{2[d,l]} \text{reg}_{2[d,l]}(n) = -\text{reg}_1(n) + \text{reg}_2(n) + \tag{A}$$

$$+\text{reg}_1(n) + a_{2d} \text{reg}_{2d}(n) + a_{2l} \text{reg}_{2l}(n) + a_{[2d,2l]} \text{reg}_{[2d,2l]}(n). \tag{F}$$

Proof of (2). Let us now consider a $DD^4(3|r)$ sequence.

Notice that by Theorem 2.6 and Corollary 1 it has always the form with no more than four basic sequences reg_i , i.e.

$$a_p \text{reg}_p(n) + a_q \text{reg}_q(n) + a_r \text{reg}_r(n) + a_s \text{reg}_s(n),$$

where $p, q, r, s \geq 3$. Then we may represent this sequence as a sum of three $DD^4(1|r)$ sequences in the following way:

$$-\text{reg}_1(n) + a_p \text{reg}_p(n) + \tag{B}$$

$$+ a_q \text{reg}_q(n) + \tag{D}$$

$$+\text{reg}_1(n) + a_r \text{reg}_r(n) + a_s \text{reg}_s(n) \tag{F}$$

This completes the proof. □

4. Calculation of the invariant. We work under the following standing assumptions

Standing Assumptions

1. $f : M^4 \rightarrow M^4$ is a smooth self-map of a smooth closed connected and simply-connected 4-manifold,
2. $r = p_1 \dots p_s$ is a product of different prime numbers,
3. in the periodic expansion of Lefschetz numbers

$$L(f^k) = \sum_{i=1}^{\infty} a_i \text{reg}_i(k)$$

$a_i \neq 0$ for all $i \neq 1$ dividing r .

Remark 1. The assumption (3) is satisfied for a self-map $f : S^4 \rightarrow S^4$ with $|\text{deg}(f)| > 1$ [20]. In general, it often takes place if the growth of $\{L(f^k)\}_k$ is quick.

We will find the formula for $D_r^4[f]$, under the above assumptions.

It turns out that first it is convenient to find the minimal decomposition of the sum

$$\sum_{i|r} a_i \text{reg}_i$$

into $DD^4(1|r)$ sequences modulo reg_1 i.e. we require that the equality holds only for all divisors $i|r$ different than 1. In other words, we will temporarily ignore the coefficient at reg_1 .

Lemma 4.1. *The two following numbers are equal:*

1. *the minimal number of summands in the decomposition of the sum*

$$\sum_{i|r} a_i \text{reg}_i$$

modulo reg_1 into $DD^4(1|r)$ sequences,

2. *the minimal number $h(s)$ determining the family of pairs of subsets of $I_s = \{1, \dots, s\}$:*

$$\{A_1, B_1\}, \{A_2, B_2\}, \dots, \{A_{h(s)}, B_{h(s)}\}$$

such that

$$\bigcup_{i=1}^{h(s)} \{A_i, B_i, A_i \cup B_i\} = 2^{I_s} \setminus \{\emptyset\}$$

i.e. for each nonempty subset $X \subset I_s$ there is an i such that either $X = A_i$ or $X = B_i$ or $X = A_i \cup B_i$.

Proof. Let us notice that to get the minimal decomposition of

$$\sum_{i|r} a_i \text{reg}_i \quad \text{modulo } \text{reg}_1,$$

we should use as much as possible the most “greedy” $DD^4(1|r)$ sequences, with the greatest number of basic expressions reg_i i.e. of the type (2) or (3) of Corollary 1. In both of these cases we have the sequences of the form:

$$\varepsilon \text{reg}_1 + a_d \text{reg}_d + a_l \text{reg}_l + \gamma a_{[d,l]} \text{reg}_{[d,l]}, \quad (3)$$

where d, l are divisors of r different than 1, $\gamma \in \{0, 1\}$.

Since $r = p_1 \cdots p_s$ is a product of different primes, there is a bijection $G : 2^{I_s} \rightarrow \text{Div}(r)$ between $\text{Div}(r)$, the set of all divisors of r , and the family of all subsets of $I_s = \{1, \dots, s\}$:

$$\{1, \dots, s\} \supset A \rightarrow \prod_{i \in A} p_i \in \text{Div}(r),$$

with the convention that $\prod_{i \in \emptyset} p_i = 1$. Moreover

$$G(A \cup B) = [G(A), G(B)].$$

As a result, every triple of divisors $d, l, [d, l]$ determining the sequence (3) is associated with a triple of subsets of I_s : $A_j, B_j, A_j \cup B_j$.

Now, a decomposition of the sum $\sum_{1 \neq i|r} a_i \text{reg}_i(k)$ into $h(s)$ $DD^4(1|r)$ sequences of the form (3) is equivalent to the existence of $h(s)$ families of subsets of I_s

$$\{A_1, B_1\}, \{A_2, B_2\}, \dots, \{A_{h(s)}, B_{h(s)}\}$$

such that

$$\bigcup_{i=1}^{h(s)} \{A_i, B_i, A_i \cup B_i\} = 2^{I_s} \setminus \{\emptyset\}$$

i.e. for each nonempty subset $X \subset I_s$ there is an i such $X = A_i$, $X = B_i$ or $X = A_i \cup B_i$. \square

Now our problem reduces to the following combinatorial question:

Problem 4.1. *Let s be a natural number. Find the minimal number $h(s)$ such that there exist $h(s)$ families of subsets $\mathcal{A}_1, \dots, \mathcal{A}_{h(s)} \subset 2^{I_s}$ satisfying*

1. *$\#\mathcal{A}_i \leq 2$ i.e. each family consists of at most two subsets,*



2. for each nonempty subset $X \subset \{1, \dots, s\}$ there exists $i \in \{1, \dots, s\}$ such that X is one of the sets A_i, B_i or $A_i \cup B_i$, where $\mathcal{A}_i = \{A_i, B_i\}$.

Theorem 4.2. *The minimal number searched in Problem 4.1 is given by the formula*

$$h(s) = \frac{2^s + (-1)^{s+1}}{3}. \tag{4}$$

The proof of Theorem 4.2 is a consequence of the following three lemmas.

Lemma 4.3. *The formula (4) for $h(s)$ can be given inductively as follows:*

$$h(2) = 1, \quad h(s + 1) = 2 \cdot h(s) + (-1)^s.$$

Proof.

$$\begin{aligned} 2 \cdot h(s) + (-1)^s &= 2 \cdot \frac{2^s + (-1)^{s+1}}{3} + (-1)^s \\ &= \frac{2^{s+1} + 2 \cdot (-1)^{s+1} + 3 \cdot (-1)^s}{3} = \frac{2^{s+1} + (-1)^s}{3} = h(s + 1). \end{aligned}$$

□

Lemma 4.4. *$h(s)$ given by the formula (4) is less or equal to the minimal number satisfying the conditions in Problem 4.1.*

Proof. We notice that each family containing two subsets $\{A, B\} \subset 2^{I_s}$ determines at most three nonempty subsets $A, B, A \cup B \subset I_s$. Thus, to realize all nonempty subsets in I_s we need at least $(2^s - 1)/3$ pairs. The last means that the minimal number in Problem 4.1 is greater or equal to $(2^s - 1)/3$. On the other hand, the least natural number $\geq (2^s - 1)/3$ is equal to $(2^s - 1)/3$ when s is even and $(2^s + 1)/3$ when s is odd. It remains to notice that in both cases we get $h(s)$. □

Lemma 4.5. (I) *For each $s \geq 2$ there exist $h(s) = \frac{2^s + (-1)^{s+1}}{3}$ families satisfying the conditions in Problem 4.1.*

(II) *Moreover, if s is even then each family must contain two different subsets, while if s is odd then $h(s) - 1$ families must contain two different subsets and the last family can contain only one subset consisting of a single, arbitrarily chosen, element.*

Proof. We will show inductively that (for $s \geq 2$): there exists a family $\mathcal{A}_s = \{\{A_i, B_i\} : i = 1, \dots, h(s)\}$ whose elements are nonempty subsets $A_i, B_i \subset I_s$ realizing all nonempty subsets in I_s and moreover

1. $A_i \neq B_i$ if $i = 1, \dots, h(s)$ and s is even,
2. $A_i \neq B_i$ if $i = 1, \dots, h(s) - 1$ and s is odd.
3. $A_{h(s)} = B_{h(s)} = \{s\}$ for s odd.

For $s = 2$ all nonempty subsets of $I_2 = \{1, 2\}$ can be obtained from the family $\{\{1\}, \{2\}\}$ which implies $h(2) = 1$.

Now we assume that for even s a family $\mathcal{A}_s = \{\{A_i, B_i\} : i = 1, \dots, h(s)\}$ where $A_i \neq B_i$ realizes all nonempty subsets in $I_s = \{1, \dots, s\}$. Then the family

$$\mathcal{A}_{s+1} = \{\{A_i, B_i\}, \{A_i \cup \{s + 1\}, B_i \cup \{s + 1\}\}, \{\{s + 1\}\} : i = 1, \dots, h(s)\}$$

realizes all nonempty subsets in $I_{s+1} = \{1, \dots, s, s + 1\}$. Moreover,

$$\#\mathcal{A}_{s+1} = 2 \cdot \#\mathcal{A}_s + 1 = 2 \cdot h(s) + 1 = 2 \cdot h(s) + (-1)^s = h(s + 1)$$

since s is even.

Now, the family

$$\mathcal{A}_{s+2} = \{\{A'_i, B'_i\}, \{A'_i \cup \{s+2\}, B'_i \cup \{s+2\}\}, \{\{s+1\}, \{s+2\}\}\}$$

where $\{A'_i, B'_i\} \in \mathcal{A}_{s+1} \setminus \{\{s+1\}\}$

realizes all subsets in I_{s+2} and moreover

$$\#\mathcal{A}_{s+2} = 2 \cdot (\#\mathcal{A}_{s+1} - 1) + 1 = 2 \cdot h(s+1) - 1 = 2 \cdot h(s+1) + (-1)^{s+1} = h(s+2)$$

since $s+1$ is odd.

This ends the proof of part (I). Part (II) of Lemma 4.5 follows from Lemma 4.4 and the observation that for $s+1$ odd in the above inductive construction, the family $\{\{s+1\}\}$, i.e. the last element in \mathcal{A}_{s+1} , consists of one subset containing a single element. It is evident that after a permutation $\{\{s+1\}\}$ can be replaced with $\{\{i\}\}$ for an arbitrarily prescribed $i \in I_{s+1}$. \square

Proof of Theorem 4.2

Lemma 4.4 gives

$$h(s) \leq \text{minimal number in Problem 4.1}$$

while Lemma 4.5 proves the opposite inequality. \square

By Theorem 4.2 we obtain

Corollary 2. *The minimal decomposition of the sum*

$$\sum_{i|r} a_i \text{reg}_i$$

modulo reg_1 into $DD^4(1|r)$ sequences contains exactly

$$h(s) = \frac{2^s + (-1)^{s+1}}{3}$$

sequences.

Moreover, by Lemma 4.5 (II) we get:

(A) *if s is even then the minimal decomposition must contain $h(s)$ sequences of the type*

$$\varepsilon \cdot \text{reg}_1 + a_d \text{reg}_d + a_l \text{reg}_l + \gamma a_{[d,l]} \text{reg}_{[d,l]}, \quad (5)$$

i.e. of the form (2) or (3) of Corollary 1 ($\gamma \in \{0, 1\}$);

(B) *if s is odd then the minimal decomposition must contain $h(s) - 1$ sequences of the type (5) while the remaining sequence may be $a_1 \text{reg}_1(n) + a_d \text{reg}_d(n)$ (i.e. of the type (1) of Corollary 1), where $d \neq 1$ is an arbitrarily prescribed divisor of r .*

Remark 2. Let us notice that in all sequences (5), appearing in the minimal decomposition modulo 1 described in Corollary 2, the divisors d, l must be different as they correspond to different subsets in Lemma 4.5, so both $\text{reg}_d(n)$ and $\text{reg}_l(n)$ appear with nonzero coefficients.

Now we are in a position to find the formula for $D_r^4[f]$, i.e. we take into account also the coefficient at reg_1 .

Let us remark that $D_r^4[f] \geq h(s)$. In fact, in the minimal realization modulo reg_1 we need $h(s)$ of $DD^4(1|r)$ sequences. The following lemmas make it precise when

these sequences are sufficient to obtain the decomposition with $a_1 \text{reg}_1$ and when one additional sequence, to realize $a_1 \text{reg}_1$, is necessary.

Lemma 4.6. *Assume our Standing Assumptions are satisfied and s is even, then*

$$D_r^4[f] = \begin{cases} h(s) & \text{if } (r \text{ is odd and } L(f) = h(s)) \\ & \text{or } (r \text{ is even and } h(s) - 2 \leq L(f) \leq h(s)), \\ h(s) + 1 & \text{otherwise.} \end{cases}$$

Proof. By Corollary 2 (A) to realize

$$\sum_{1 \neq i|r} a_i \text{reg}_i$$

we need at least $h(s) DD^4(1|r)$ sequences of the type (2) or (3) of Corollary 1.

If we assume that r is odd then they all must be of the type (3). Then the contribution of each of them to the coefficient at reg_1 is 1. If moreover $L(f) = h(s)$ then $D_r^4[f] = h(s)$, since $a_1 = L(f)$. Otherwise, we need one sequence of the type (1) more to realize the difference $(a_1 - h(s)) \cdot \text{reg}_1(n)$.

Now we consider the case of even r . Then exactly one sequence in the minimal decomposition must be of the type (2) and the remaining $h(s) - 1$ sequences are of the type (3). Their contribution to the coefficient at reg_1 is $(h(s) - 1) + \varepsilon$ where $\varepsilon = 0, +1, -1$. Now, if $h(s) - 2 \leq L(f) \leq h(s)$, then a_1 can be realized by these sequences. Otherwise we need one more sequence of the type (1). \square

Lemma 4.7. *Assume our Standing Assumptions are satisfied and s is odd, then*

$$D_r^4[f] = h(s).$$

Proof. It is enough to show that $\sum_{i|r} a_i \text{reg}_i(n)$ is the sum of exactly $h(s) DD^4(1|r)$ sequences.

Since s is odd, by Corollary 2 (B), $h(s) - 1$ sequences of the types (2) or (3) of Corollary 1 realize

$$\sum_i a_i \text{reg}_i,$$

where the summation runs through the set $\text{Div}(r) \setminus \{1, d\}$, for some $d|r$. Again by Corollary 2 (B), it remains to add one expression of the type (1) realizing the sum $a_1 \text{reg}_1 + a_d \text{reg}_d$. \square

We sum up our considerations in the following

Theorem 4.8. *Let $f : M^4 \rightarrow M^4$ be a smooth self-map of a smooth closed connected and simply-connected 4-manifold, $r = p_1 \dots p_s$ be a product of different prime numbers. We assume that the coefficients a_i in the periodic expansion of $L(f^k) = \sum_{i=1}^{\infty} a_i \text{reg}_i(k)$, are nonzero for all $i|r, i \neq 1$. Then*

$$D_r^4[f] = \begin{cases} h(s) & \text{if } (s \text{ is odd}) \text{ or } (r \text{ is odd and } L(f) = h(s)) \\ & \text{or } (r \text{ is even and } h(s) - 2 \leq L(f) \leq h(s)), \\ h(s) + 1 & \text{otherwise.} \end{cases}$$

where $h(s) = (2^s + (-1)^{s+1})/3$.

Remark 3. If in Theorem 4.8 we drop the part (3) of the Standing Assumption according which $a_i \neq 0$ for all $i \neq 1$ dividing r then the equality becomes the inequality and we get the estimation for $D_r^4[f]$ from above.

REFERENCES

- [1] I. K. Babenko and S. A. Bogatyĭ, *The behavior of the index of periodic points under iterations of a mapping*, Math. USSR Izv. **38** (1992), 1–26.
- [2] A. Dold, *Fixed point indices of iterated maps*, Invent. Math. **74** (1983), 419–435.
- [3] G. Graff, *Minimal number of periodic points for smooth self-maps of two-holed 3-dimensional closed ball*, Topol. Methods Nonlinear Anal. **33** (2009), no. 1, 121–130.
- [4] G. Graff and J. Jezierski, *Minimal number of periodic points for C^1 self-maps of compact simply-connected manifolds*, Forum Math. **21** (2009), no. 3, 491–509.
- [5] G. Graff and J. Jezierski, *Minimal number of periodic points for smooth self-maps of S^3* , Fund. Math. **204** (2009), 127–144.
- [6] G. Graff and J. Jezierski, *Minimizing the number of periodic points for smooth maps. Non-simply connected case*, Topology Appl. **158** (2011), no. 3, 276–290.
- [7] G. Graff, J. Jezierski and M. Nowak-Przygodzki, *Minimal number of periodic points for smooth self-maps of $\mathbb{R}P^3$* , Topology Appl. **157** (2010), no. 10–11, 1784–1803.
- [8] G. Graff, J. Jezierski and P. Nowak-Przygodzki, *Fixed point indices of iterated smooth maps in arbitrary dimension*, J. Differential Equations (2011), doi:10.1016/j.jde.2011.05.024
- [9] G. Graff and P. Nowak-Przygodzki, *Fixed point indices of iterations of C^1 maps in \mathbb{R}^3* , Discrete Cont. Dyn. Systems **16** (2006), no. 4, 843–856.
- [10] E. Hart, P. Heath and E. Keppelmann, *Algorithms for Nielsen type periodic numbers of maps with remnant on surfaces with boundary and on bouquets of circles. I.*, Fund. Math. **200** (2008), no. 2, 101–132.
- [11] E. Hart and E. Keppelmann, *Nielsen periodic point theory for periodic maps on orientable surfaces*, Topology Appl. **153** (2006), no. 9, 1399–1420.
- [12] J. Jezierski, *Wecken’s theorem for periodic points in dimension at least 3*, Topology Appl. **153** (2006), no. 11, 1825–1837.
- [13] J. Jezierski, *Homotopy periodic sets of selfmaps of real projective spaces*, Bol. Soc. Mat. Mexicana (3) **11** (2005), no. 2, 294–302.
- [14] B. J. Jiang, “*Lectures on the Nielsen Fixed Point Theory*,” Contemp. Math. **14**, Amer. Math. Soc., Providence 1983.
- [15] J. Jezierski and W. Marzantowicz, “*Homotopy Methods in Topological Fixed and Periodic Points Theory*,” Topological Fixed Point Theory and Its Applications, 3. Springer, Dordrecht, 2005.
- [16] H. J. Kim, J. B. Lee and W. S. Yoo, *Computation of the Nielsen type numbers for maps on the Klein bottle*, J. Korean Math. Soc. **45** (2008), no. 5, 1483–1503.
- [17] P. Le Calvez, *Dynamique des homéomorphismes du plan au voisinage d’un point fixe*, Ann. Sci. École Norm. Sup. (4) **36**, (2003), no. 1, 139–171.
- [18] J. B. Lee and X. Zhao, *Nielsen type numbers and homotopy minimal periods for maps on the 3-nilmanifolds*, Sci. China Ser. A **51** (2008), no. 3, 351–360.
- [19] J. B. Lee and X. Zhao, *Nielsen type numbers and homotopy minimal periods for maps on 3-solvmanifolds*, Algebr. Geom. Topol. **8** (2008), no. 1, 563–580.
- [20] J. Llibre, J. Paranõs and J. A. Rodriguez, *Periods for transversal maps on compact manifolds with a given homology*, Houston J. Math. **24** (1998), no. 3, 397–407.
- [21] F. Ruiz del Portal and J.M. Salazar, *A Poincaré formula for the fixed point indices of the iterates of arbitrary planar homeomorphisms*, Fixed Point Theory Appl., Volume 2010 (2010), Article ID 323069, 31 pages.
- [22] G.Y. Zhang, *Fixed point indices and periodic points of holomorphic mappings*, Math. Ann. **337** (2007), no. 2, 401–433.

Received July 2010; revised February 2011.

E-mail address: graff@mif.pg.gda.pl

E-mail address: jezierski@acn.waw.pl

