

# Hat problem on a graph

Marcin Krzywkowski

e-mail: marcin.krzywkowski@gmail.com

*Faculty of Applied Physics and Mathematics  
Gdańsk University of Technology  
Narutowicza 11/12, 80-952 Gdańsk, Poland*

## Abstract

The topic of our paper is the hat problem. In that problem, each of  $n$  people is randomly fitted with a blue or red hat. Then everybody can try to guess simultaneously his own hat color looking at the hat colors of the other people. The team wins if at least one person guesses his hat color correctly and no one guesses his hat color wrong, otherwise the team loses. The aim is to maximize the probability of win. In this version every person can see everybody excluding him. In this paper we consider such problem on a graph, where vertices are people and a person can see these people, to which he is connected by an edge. We prove some general theorems about the hat problem on a graph and solve the problem on trees. We also consider the hat problem on a graph with given degrees of vertices. We give an upper bound that is based only on the degrees of vertices on the chance of success of any strategy for the graph  $G$ . We show that this upper bound together with integrality constraints is tight on some toy examples.

**Keywords:** hat problem, graph, path, tree, vertex degree.

**AMS Subject Classification:** 05C05, 05C38, 05C57, 91A43.

## 1 Introduction

In the hat problem, a team of  $n$  people enters a room and a blue or red hat is randomly placed on the head of each person. Each person can see the hats of all of the other people but not his own. No communication of any sort is allowed, except for an initial strategy session before the game begins. Once they have had

a chance to look at the other hats, each person must simultaneously guess the color of his own hat or pass. The team wins if at least one person guesses his hat color correctly and no one guesses his hat color wrong, otherwise the team loses. The aim is to maximize the probability of win.

The hat problem with seven people called "seven prisoners puzzle" was formulated by T. Ebert in his Ph.D. Thesis [10]. The hat problem was also the subject of articles in The New York Times [20], Die Zeit [5], and abcNews [19]. The hat problem with  $n$  people and two colors of hat was investigated in [6]. It was solved for  $2^k - 1$  people in [12]. The hat problem and Hamming codes were also the subject of an article in Polish math–physics–informatic magazine [9].

There are also known some variants and generalizations of hat problem. The authors of [18] investigate the generalized hat problem with  $q \geq 2$  colors, they also consider variants in which there are arbitrary input distributions, randomized playing strategies, and symmetric strategies. In the papers [1], [8], and [17] there is considered another variant of hat problem in which passing is not allowed, thus everybody has to try to guess his hat color. The aim is to maximize the number of correct guesses. In [14] the authors investigate several variants of hat problem in which the aim is to design a strategy such that the number of correct guesses is greater than or equal to the given positive integer. In the paper [15] there is considered the hat problem, and also a variant in which the probabilities of getting hats of each colors do not have to be equal. The authors of [2] investigate a problem similar to the hat problem. There are  $n$  people which have random bits on foreheads, and they have to vote on the parity of the  $n$  bits.

The hat problem and its variants have many applications and connections to other areas of science, for example: information technology [4], linear programming [14, 16], genetic programming [7], economy [1, 17], biology [15], approximating Boolean functions [2], and autoreducibility of random sequences [3,10–13]. Therefore, it is hoped that the hat problem on a graph considered in this paper, as a natural generalization, is worth exploring, and may also have many applications.

We consider the hat problem on a graph, where vertices are people and a person can see these people, to which he is connected by an edge. We prove some general theorems about the hat problem on a graph and solve the problem on trees. We also consider the hat problem on a graph with given degrees of vertices. We give an upper bound that is based only on the degrees of vertices on the chance of success of any strategy for the graph  $G$ . We show that this upper

bound together with integrality constraints is tight on some toy examples.

The paper is organized as follows. In Section 2 we give the notation and terminology used. In Section 3 first we make some general observations about the hat problem on a graph. In Theorem 4 we solve that problem on paths, and in Theorem 5 we solve the hat problem on trees. Then we consider the hat problem on a graph with given degrees of vertices.

## 2 Preliminaries

For a graph  $G$ , by  $V(G)$  and  $E(G)$  we denote the set of vertices and the set of edges of this graph, respectively. If  $H$  is a subgraph of  $G$ , then we write  $H \subseteq G$ . Let  $v \in V(G)$ . By  $N_G(v)$  we denote the open neighbourhood of  $v$ , that is  $N_G(v) = \{x \in V(G) : vx \in E(G)\}$ . By  $N_G[v]$  we denote the closed neighbourhood of  $v$ , that is  $N_G[v] = N_G(v) \cup \{v\}$ . By  $d_G(v)$  we denote the degree of the vertex  $v$ , that is the number of its neighbours, thus  $d_G(v) = |N_G(v)|$ . By  $P_n$  we denote the path with  $n$  vertices. By  $C_n$  we denote the cycle with  $n$  vertices. By  $K_n$  we denote the complete graph with  $n$  vertices. Let  $f: X \rightarrow Y$  be a function. If  $Z \subseteq X$ , then by  $f|_Z$  we denote the restriction of  $f$  to  $Z$ . If  $y \in Y$ , then by  $f \equiv y$  we denote that for every  $x \in X$  we have  $f(x) = y$ .

Without loss of generality we may assume an ordering of the vertices of a graph  $G$ , that is  $V(G) = \{v_1, v_2, \dots, v_n\}$ .

Let  $\{b, r\}$  be the set of colors ( $b$  means blue and  $r$  means red). If  $v_i \in V(G)$ , then  $c(v_i)$  is the color of  $v_i$ , so  $c: V(G) \rightarrow \{b, r\}$  is a function. By a case for the graph  $G$  we mean a sequence  $(c(v_1), c(v_2), \dots, c(v_n))$ . The set of all cases for the graph  $G$  we denote by  $C(G)$ , of course  $|C(G)| = 2^{|V(G)|}$ .

If  $v_i \in V(G)$ , then by  $s_i$  we denote a function  $s_i: V(G) \rightarrow \{b, r, *\}$ , where  $s_i(v_j) \in \{b, r\}$  is the color of  $v_j$  if  $v_i$  sees  $v_j$ , and mark  $*$  otherwise, that is,  $s_i(v_j) = c(v_j)$  if  $v_j \in N_G(v_i)$ , while  $s_i(v_j) = *$  if  $v_j \in V(G) \setminus N_G(v_i)$ . By a situation of the vertex  $v_i$  in the graph  $G$  we mean the sequence  $(s_i(v_1), s_i(v_2), \dots, s_i(v_n))$ . The set of all possible situations of  $v_i$  in the graph  $G$  we denote by  $St_i(G)$ . Of course,  $|St_i(G)| = 2^{|N_G(v_i)|}$ .

Let  $v_i \in V(G)$ . We say that a case  $(c_1, c_2, \dots, c_n)$  for the graph  $G$  corresponds to a situation  $(t_1, t_2, \dots, t_n)$  of the vertex  $v_i$  in the graph  $G$  if it is created from this situation only by changing every mark  $*$  to  $b$  or  $r$ . So, a case corresponds to a situation of  $v_i$  if every vertex adjacent to  $v_i$ , in that case has the same color as

in that situation. To every situation of the vertex  $v_i$  in the graph  $G$  correspond  $2^{|V(G)| - |N_G(v_i)|}$  cases, because every situation of  $v_i$  has  $|V(G)| - |N_G(v_i)|$  marks  $*$ .

Let  $G$  and  $H$  be graphs such that  $V(H) = \{v_1, v_2, \dots, v_m\}$ ,  $V(G) = \{v_1, v_2, \dots, v_m, \dots, v_n\}$ , and  $E(H) \subseteq E(G)$ . We say that a case  $(a_1, a_2, \dots, a_m, \dots, a_n)$  for the graph  $G$  corresponds to a case  $(b_1, b_2, \dots, b_m)$  for the graph  $H$  if  $(a_1, a_2, \dots, a_m) = (b_1, b_2, \dots, b_m)$ , that is, every vertex from the graph  $H$  in both cases has the same color. Of course, to every case for the graph  $H$  correspond  $2^{n-m}$  cases for the graph  $G$ .

Let  $G$  and  $H$  be graphs such that  $V(H) = \{v_1, v_2, \dots, v_m\}$ ,  $V(G) = \{v_1, v_2, \dots, v_m, \dots, v_n\}$ , and  $E(H) \subseteq E(G)$ . Let  $i \in \{1, 2, \dots, m\}$ . We say that a situation  $(t_1, t_2, \dots, t_m, \dots, t_n)$  of the vertex  $v_i$  in the graph  $G$  corresponds to a situation  $(u_1, u_2, \dots, u_m)$  of the vertex  $v_i$  in the graph  $H$  if  $(t_1, t_2, \dots, t_m) = (u_1, u_2, \dots, u_m)$ , that is, every vertex adjacent to  $v_i$  in the graph  $H$ , in both of these situations has the same color.

By a statement of a vertex we mean its declaration about the color it guesses it is. By the result of a case we mean a win or a loss. According to the definition of the hat problem, the result of a case is a win if at least one vertex states its color correctly and no vertex states its color wrong. The result of a case is a loss if no vertex states its color or some vertex states its color wrong.

By a guessing instruction for the vertex  $v_i \in V(G)$  (denoted by  $g_i$ ) we mean a function  $g_i: St_i(G) \rightarrow \{b, r, p\}$  which, for a given situation, gives  $b$  or  $r$  meaning the color  $v_i$  guesses it is, or the letter  $p$  if  $v_i$  passes. Thus a guessing instruction is a rule which determines the behavior of the vertex  $v_i$  in every situation. By a strategy for the graph  $G$  we mean a sequence  $(g_1, g_2, \dots, g_n)$ . By  $\mathcal{F}(G)$  we denote the family of all strategies for the graph  $G$ .

Let  $v_i \in V(G)$  and  $S \in \mathcal{F}(G)$ . We say that  $v_i$  never states its color in the strategy  $S$  if  $v_i$  passes in every situation, that is  $g_i \equiv p$ . We say that  $v_i$  always states its color in the strategy  $S$  if  $v_i$  states its color in every situation, that is, for every  $T \in St_i(G)$  we have  $g_i(T) \in \{b, r\}$  ( $g_i(T) \neq p$ , equivalently).

If  $S \in \mathcal{F}(G)$ , then by  $Cw(S)$  and  $Cl(S)$  we denote the sets of cases for the graph  $G$  in which the team wins or loses, respectively. Of course,  $|Cw(S)| + |Cl(S)| = |C(G)|$ . Consequently, by the chance of success of the strategy  $S$  we mean the number  $p(S) = \frac{|Cw(S)|}{|C(G)|}$ . By the hat number of the graph  $G$  we mean the number  $h(G) = \max\{p(S) : S \in \mathcal{F}(G)\}$ . Certainly  $p(S) \leq h(G)$ . We say that the strategy  $S$  is optimal for the graph  $G$  if  $p(S) = h(G)$ . By  $\mathcal{F}^0(G)$  we denote the family of all optimal strategies for the graph  $G$ .

Let  $t, m_1, m_2, \dots, m_t \in \{1, 2, \dots, n\}$  be such that  $m_j \neq m_k$  and  $c_{m_j} \in \{b, r\}$ , for every  $j, k \in \{1, 2, \dots, t\}$ .

By  $C(G, v_{m_1}^{c_{m_1}}, v_{m_2}^{c_{m_2}}, \dots, v_{m_t}^{c_{m_t}})$  we denote the set of cases for the graph  $G$  such that the color of  $v_{m_j}$  is  $c_{m_j}$ .

Let  $S \in \mathcal{F}(G)$ . By  $Cw(S, v_{m_1}^{c_{m_1}}, v_{m_2}^{c_{m_2}}, \dots, v_{m_t}^{c_{m_t}})$  ( $Cl(S, v_{m_1}^{c_{m_1}}, v_{m_2}^{c_{m_2}}, \dots, v_{m_t}^{c_{m_t}})$ , respectively) we denote the set of cases for  $G$  which belong to the set  $C(G, v_{m_1}^{c_{m_1}}, v_{m_2}^{c_{m_2}}, \dots, v_{m_t}^{c_{m_t}})$ , and in which the team wins (loses, respectively).

Let  $v_i \in V(G)$ . If for every  $j \in \{1, 2, \dots, t\}$  we have  $v_{m_j} \in N_G(v_i)$ , then by  $St_i(G, v_{m_1}^{c_{m_1}}, v_{m_2}^{c_{m_2}}, \dots, v_{m_t}^{c_{m_t}})$  we denote the set of possible situations of  $v_i$  in the graph  $G$  such that the color of  $v_{m_j}$  is  $c_{m_j}$ .

### 3 Results

First let us observe that indeed we can confine to deterministic strategies (that is strategies such that the decision of each person is determined uniquely by the hat colors of other people). We can do this since for any randomized strategy there exists a not worse deterministic one. It is true, because every randomized strategy is a convex combination of some deterministic strategies. The probability of winning is a linear function on the convex polyhedron corresponding to the set of all randomized strategies which can be achieved combining those deterministic strategies. It is well known that this function achieves its maximum on a vertex of the polyhedron which corresponds to a deterministic strategy.

Let  $G$  and  $H$  be graphs. Assume that  $H \subseteq G$ . Since every vertex from the set  $V(G) \setminus V(H)$  can always pass, and every vertex  $v_i \in V(H)$  can ignore the colors of vertices from the set  $N_G(v_i) \setminus N_H(v_i)$ , it is easy to see that the hat number of the graph  $G$  is greater than or equal to the hat number of the graph  $H$ . It is that if  $H \subseteq G$ , then  $h(H) \leq h(G)$ .

Since  $K_1$  is a subgraph of every graph, we get  $h(G) \geq \frac{1}{2}$ .

Let  $S$  be an optimal strategy for the graph  $G$ . By definition we have  $p(S) = h(G)$ . Since  $h(G) \geq \frac{1}{2}$ , we get  $p(S) \geq \frac{1}{2}$ .

Now we prove a fact characterizing the number of cases in which the loss of the team is caused by a statement of a vertex.

**Fact 1** *Let  $G$  be a graph and let  $v_i$  be a vertex of  $G$ . Let  $S \in \mathcal{F}(G)$ . If  $v_i$  states its color in a situation, then the team loses in at least half of all cases corresponding to this situation.*

**Proof.** Assume that  $v_i$  states its color in a situation  $T$ . Without loss of generality we assume that in this situation  $v_i$  states it is blue, that is  $g_i(T) = b$ . In half of all cases corresponding to  $T$  we have  $c(v_i) = r$ , it means that  $v_i$  is red. Thus, the team loses in every one of these cases, because  $v_i$  states its color wrong, as  $g_i(T) = b \neq r = c(v_i)$ . ■

**Corollary 2** *Let  $G$  be a graph and let  $v$  be a vertex of  $G$ . If  $S \in \mathcal{F}^0(G)$  is a strategy such that  $v$  always states its color, then  $h(G) = \frac{1}{2}$ .*

**Proof.** Assumption indicates that in every case  $v$  states its color, so by Fact 1 we have  $|Cl(S)| \geq \frac{|C(G)|}{2}$ . Consequently,

$$p(S) = \frac{|Cw(S)|}{|C(G)|} = \frac{|C(G)| - |Cl(S)|}{|C(G)|} \leq \frac{|C(G)| - \frac{|C(G)|}{2}}{|C(G)|} = \frac{1}{2}.$$

Since  $p(S) \leq \frac{1}{2}$  and  $S \in \mathcal{F}^0(G)$ , we have  $h(G) \leq \frac{1}{2}$  (by definition). On the other hand we have  $h(G) \geq \frac{1}{2}$ . ■

In the following theorem we give a sufficient condition for deleting a vertex of a graph without changing its hat number.

**Theorem 3** *Let  $G$  be a graph and let  $v$  be a vertex of  $G$ . If  $S \in \mathcal{F}^0(G)$  is a strategy such that  $v$  never states its color, then  $h(G) = h(G - v)$ .*

**Proof.** Let  $S' \in \mathcal{F}(G - v)$  be the strategy as follows: Every vertex not adjacent to  $v$  in  $G$  behaves in the same way as in  $S$ , that is, if  $v_i \notin N_G(v)$ , then  $g'_i = g_i$ , where  $g'_i$  and  $g_i$  are the guessing instructions for the vertex  $v_i$  in the strategies  $S'$  and  $S$ , respectively. First assume that  $|Cw(S, v^b)| \geq |Cw(S, v^r)|$ . Let every vertex adjacent to  $v$  in  $G$  behave in the same way as in  $S$  when  $v$  is blue, that is, if  $v_i \in N_G(v)$ , then  $g'_i = g_i|_{St_i(G, v^b)}$ . The result of any case  $C'$  in the strategy  $S'$  is the same as the result of the case  $C$  in the strategy  $S$ , where  $C$  is the corresponding case in which  $v$  is blue, because in both strategies  $S'$  and  $S$  the vertex  $v$  never states its color and every vertex in the strategy  $S'$  behaves in the same way as in  $S$  when  $v$  is blue. This implies that  $|Cw(S')| = |Cw(S, v^b)|$ . Now we get

$$p(S') = \frac{|Cw(S')|}{2^{|V(G-v)|}} = \frac{|Cw(S, v^b)|}{2^{|V(G)|-1}} = \frac{2|Cw(S, v^b)|}{2^{|V(G)|}}$$

$$\geq \frac{|Cw(S, v^b)| + |Cw(S, v^r)|}{2^{|V(G)|}} = p(S).$$

If  $|Cw(S, v^b)| < |Cw(S, v^r)|$ , then similarly we get a strategy  $S'$  such that  $p(S') > p(S)$ . Since  $S \in \mathcal{F}^0(G)$  and  $S' \in \mathcal{F}(G - v)$ , we have  $h(G) = p(S) \leq p(S') \leq h(G - v)$ . On the other hand we have  $h(G) \geq h(G - v)$ . ■

Let  $S$  be a strategy for the graph  $G$ . Let  $C$  be a case in which some vertex states its color. Since the rules of the hat problem are such that one correct statement suffices to win, and one wrong statement causes the loss, it is easy to see that a statement of any other vertex cannot improve the result of the case  $C$ .

Now we solve the hat problem on paths.

**Theorem 4** *For every path  $P_n$  we have  $h(P_n) = \frac{1}{2}$ .*

**Proof.** Let  $E(P_n) = \{v_1v_2, v_2v_3, \dots, v_{n-1}v_n\}$ . We distinguish six possibilities:  $n = 1$ ,  $n = 2$ ,  $n = 3$ ,  $n = 4$ ,  $n = 5$ , and  $n \geq 6$ .

First, we assume that  $n = 1$ . Since  $P_1 = K_1$ , we have  $h(P_1) = h(K_1) = \frac{1}{2}$ .

Now assume that  $n = 2$ . Let  $S$  be an optimal strategy for  $P_2$ . If some vertex, say  $v_i$ , never states its color, then by Theorem 3 we have  $h(P_2) = h(P_2 - v_i)$ . Since  $P_2 - v_i = P_1$ , we have  $h(P_2) = h(P_1) = \frac{1}{2}$ . Now assume that  $v_1$  and  $v_2$  state their colors. If one of them always states its color, then by Corollary 2 we have  $h(P_2) = \frac{1}{2}$ . If, neither  $v_1$  nor  $v_2$  always states its color, then without loss of generality we assume that  $v_1$  states its color when  $v_2$  is blue, and in this situation it states it is blue. We consider the following four possibilities:  $g_2(b, *) = b$  (Table 1);  $g_2(b, *) = r$  (Table 2);  $g_2(r, *) = b$  (Table 3);  $g_2(r, *) = r$  (Table 4). In the next tables b means blue, r means red, + means correct statement (success), - means wrong statement (loss), and blank square means passing.

In Tables 1, 2, and 3 we have  $|Cw(S)| = 1$ ,  $|C(P_2)| = 4$ , so  $p(S) = \frac{1}{4} < \frac{1}{2}$ , a contradiction.

Table 1

No	The color of		The statement of		Result
	$v_1$	$v_2$	$v_1$	$v_2$	
1	b	b	+	+	+
2	b	r		-	-
3	r	b	-		-
4	r	r			-

Table 2

No	The color of		The statement of		Result
	$v_1$	$v_2$	$v_1$	$v_2$	
1	b	b	+	-	-
2	b	r		+	+
3	r	b	-		-
4	r	r			-

Table 3

No	The color of		The statement of		Result
	$v_1$	$v_2$	$v_1$	$v_2$	
1	b	b	+		+
2	b	r			-
3	r	b	-	+	-
4	r	r		-	-

Table 4

No	The color of		The statement of		Result
	$v_1$	$v_2$	$v_1$	$v_2$	
1	b	b	+		+
2	b	r			-
3	r	b	-	-	-
4	r	r		+	+

In Table 4 we have  $|Cw(S)| = 2$ ,  $|C(P_2)| = 4$ , so  $p(S) = \frac{2}{4} = \frac{1}{2}$ . Since  $S \in \mathcal{F}^0(P_2)$ , we have  $h(P_2) = \frac{1}{2}$ .

Now assume that  $n = 3$ . Let  $S$  be an optimal strategy for  $P_3$ . If  $v_1$  or  $v_3$  never states its color, then without loss of generality we assume that it is  $v_1$ . By Theorem 3 we have  $h(P_3) = h(P_3 - v_1)$ . Since  $P_3 - v_1 = P_2$ , we have  $h(P_3) = h(P_2) = \frac{1}{2}$ . Now assume that  $v_1$  and  $v_3$  state their colors. If  $v_1$  or  $v_3$  always states its color, then by Corollary 2 we have  $h(P_3) = \frac{1}{2}$ . If neither  $v_1$  nor  $v_3$  always states its color, then without loss of generality we assume that  $v_1$  states its color when  $v_2$  is blue, and in this situation it states it is blue. We have the following two possibilities: (1)  $v_3$  states its color when  $v_2$  is blue; (2)  $v_3$  does not state its color when  $v_2$  is blue.

(1) Let the strategy  $S'$  differ from  $S$  only in that  $v_3$  does not state its color when  $v_2$  is blue. Since in every case in which  $v_2$  is blue  $v_1$  states its color,



the statement of  $v_3$  cannot improve the result of any of these cases. Therefore,  $p(S) \leq p(S')$ . Since  $S \in \mathcal{F}^0(P_3)$ , the strategy  $S'$  is also optimal for  $P_3$ . If  $v_3$  never states its color in the strategy  $S'$ , then we have the possibility already considered. The other possibility when  $v_3$  states its color we consider in the next paragraph.

(2) Certainly,  $v_3$  states its color when  $v_2$  is red. Since  $v_1$  ( $v_3$ , respectively) states its color when  $v_2$  is blue (red, respectively), by Fact 1 we have

$$|Cl(S, v_2^b)| \geq \frac{|C(P_3, v_2^b)|}{2} \quad \left( |Cl(S, v_2^r)| \geq \frac{|C(P_3, v_2^r)|}{2}, \text{ respectively} \right).$$

This implies that

$$|Cl(S)| = |Cl(S, v_2^b)| + |Cl(S, v_2^r)| \geq \frac{|C(P_3, v_2^b)|}{2} + \frac{|C(P_3, v_2^r)|}{2} = \frac{|C(P_3)|}{2}.$$

Consequently,

$$p(S) = \frac{|Cw(S)|}{|C(P_3)|} = \frac{|C(P_3)| - |Cl(S)|}{|C(P_3)|} \leq \frac{|C(P_3)| - \frac{|C(P_3)|}{2}}{|C(P_3)|} = \frac{1}{2}.$$

Since  $S \in \mathcal{F}^0(P_3)$ , we have  $h(P_3) \leq \frac{1}{2}$ . Since  $h(P_3) \geq \frac{1}{2}$ , we get  $h(P_3) = \frac{1}{2}$ .

Now assume that  $n = 4$ . Let  $S$  be an optimal strategy for  $P_4$ . If some vertex, say  $v_i$ , never states its color, then by Theorem 3 we have  $h(P_4) = h(P_4 - v_i)$ . If  $i \in \{1, 4\}$ , then  $P_4 - v_i = P_3$ , so  $h(P_4) = h(P_3) = \frac{1}{2}$ . If  $i \in \{2, 3\}$ , then  $P_4 - v_i = P_1 \cup P_2$ . Since  $P_1 \cup P_2 \subseteq P_3$ , we have  $h(P_1 \cup P_2) \leq h(P_3) = \frac{1}{2}$ . Therefore,  $h(P_4) = h(P_1 \cup P_2) \leq \frac{1}{2}$ . Since  $h(P_4) \geq \frac{1}{2}$ , we get  $h(P_4) = \frac{1}{2}$ . Now assume that every vertex states its color. If some vertex always states its color, then by Corollary 2 we have  $h(P_4) = \frac{1}{2}$ . If no vertex always states its color, then without loss of generality we assume that  $v_1$  states its color when  $v_2$  is blue, and in this situation it states it is blue. Similarly, since  $N_{P_4}[v_1] \cap N_{P_4}[v_4] = \emptyset$ , we may assume that  $v_4$  states its color when  $v_3$  is blue, and in this situation it states it is blue. We consider the following two possibilities: (1)  $v_2$  states its color when  $v_3$  is blue, or  $v_3$  states its color when  $v_2$  is blue; (2)  $v_2$  does not state its color when  $v_3$  is blue, and  $v_3$  does not state its color when  $v_2$  is blue.

(1) Let the strategy  $S'$  differ from  $S$  only in that  $v_2$  does not state its color when  $v_3$  is blue, and  $v_3$  does not state its color when  $v_2$  is blue. Since in every case in which  $v_3$  ( $v_2$ , respectively) is blue  $v_4$  ( $v_1$ , respectively) states its color, the statement of  $v_2$  ( $v_3$ , respectively) cannot improve the result of any of these cases. Therefore,  $p(S) \leq p(S')$ . Since  $S \in \mathcal{F}^0(P_4)$ , the strategy  $S'$  is also optimal for  $P_4$ . If  $v_2$  or  $v_3$  never states its color in the strategy  $S'$ , then we have the possibility

already considered. The other possibility when  $v_2$  and  $v_3$  state their colors we consider in the next paragraph.

(2) If  $c(v_1) = r$  and  $c(v_2) = b$ , or  $c(v_3) = b$  and  $c(v_4) = r$ , then in each of the 7 cases, the team loses. Certainly,  $v_2$  can state its color only when  $v_3$  is red. Thus there are the following four possibilities: (2.1)  $g_2(b, *, r, *) = b$ ; (2.2)  $g_2(b, *, r, *) = r$ ; (2.3)  $g_2(r, *, r, *) = b$ ; (2.4)  $g_2(r, *, r, *) = r$ .

(2.1) Since  $|Cl(S, v_1^b, v_2^r, v_3^r)| = |C(P_4, v_1^b, v_2^r, v_3^r)| = 2$  and  $C(P_4, v_1^b, v_2^r, v_3^r) \cap (C(P_4, v_1^r, v_2^b) \cup C(P_4, v_3^b, v_4^r)) = \emptyset$ , the team loses in at least  $7 + 2 = 9$  cases, and wins in at most 7 cases. It means that  $p(S) \leq \frac{7}{16} < \frac{1}{2}$ , a contradiction.

Possibilities (2.2) and (2.3) are similar to (2.1).

(2.4) Certainly,  $v_3$  can state its color only when  $v_2$  is red. Thus we have the following four possibilities: (2.4.1)  $g_3(*, r, *, b) = b$ ; (2.4.2)  $g_3(*, r, *, b) = r$ ; (2.4.3)  $g_3(*, r, *, r) = b$ ; (2.4.4)  $g_3(*, r, *, r) = r$ .

In possibilities (2.4.1), (2.4.2), and (2.4.3), without considering the consequences of statements of  $v_2$ , we get a similar contradiction as in (2.1), (2.2), and (2.3).

(2.4.4) In this possibility, analyzed in Table 5, we have  $|Cw(S)| = 8$ ,  $|C(P_4)| = 16$ , so  $p(S) = \frac{8}{16} = \frac{1}{2}$ . Since  $S \in \mathcal{F}^0(P_4)$ , we have  $h(P_4) = \frac{1}{2}$ .

Now assume that  $n = 5$ . Let  $S$  be an optimal strategy for  $P_5$ . If for some  $i \in \{1, 3, 5\}$  the vertex  $v_i$  never states its color, then by Theorem 3 we have  $h(P_5) = h(P_5 - v_i)$ . If  $i \in \{1, 5\}$ , then  $P_5 - v_i = P_4$ , so  $h(P_5) = h(P_4) = \frac{1}{2}$ . If  $i = 3$ , then  $P_5 - v_3 = P_2 \cup P_2$ . Since  $P_2 \cup P_2 \subseteq P_4$ , we have  $h(P_2 \cup P_2) \leq h(P_4) = \frac{1}{2}$ , so  $h(P_5) = h(P_2 \cup P_2) \leq \frac{1}{2}$ . Since  $h(P_5) \geq \frac{1}{2}$ , we get  $h(P_5) = \frac{1}{2}$ . Now assume that every vertex from the set  $\{v_1, v_3, v_5\}$  states its color. If some of these vertices always states its color, then by Corollary 2 we have  $h(P_5) = \frac{1}{2}$ . If no vertex from the set  $\{v_1, v_3, v_5\}$  always states its color, then without loss of generality we assume that  $v_1$  states its color when  $v_2$  is blue, and in this situation it states it is blue. Similarly, since  $N_{P_5}[v_1] \cap N_{P_5}[v_5] = \emptyset$ , we may assume that  $v_5$  states its color when  $v_4$  is blue, and in this situation it states it is blue. We consider the following two possibilities: (1)  $v_3$  states its color when  $v_2$  or  $v_4$  is blue; (2)  $v_3$  does not state its color when  $v_2$  or  $v_4$  is blue.

(1) Let the strategy  $S'$  differ from  $S$  only in that  $v_3$  does not state its color when  $v_2$  or  $v_4$  is blue. Since in every case in which  $v_2$  ( $v_4$ , respectively) is blue,  $v_1$  ( $v_5$ , respectively) states its color, the statement of  $v_3$  cannot improve the result of any of these cases. Therefore,  $p(S) \leq p(S')$ . Since  $S \in \mathcal{F}^0(P_5)$ , the strategy  $S'$  is also optimal for  $P_5$ . If  $v_3$  never states its color in the strategy  $S'$ , then we



Table 5

No	The color of				The statement of				Result
	$v_1$	$v_2$	$v_3$	$v_4$	$v_1$	$v_2$	$v_3$	$v_4$	
1	b	b	b	b	+			+	+
2	b	b	b	r	+			-	-
3	b	b	r	b	+				+
4	b	b	r	r	+				+
5	b	r	b	b				+	+
6	b	r	b	r			-	-	-
7	b	r	r	b					-
8	b	r	r	r			+		+
9	r	b	b	b	-			+	-
10	r	b	b	r	-			-	-
11	r	b	r	b	-	-			-
12	r	b	r	r	-	-			-
13	r	r	b	b				+	+
14	r	r	b	r			-	-	-
15	r	r	r	b		+			+
16	r	r	r	r		+	+		+

have the possibility already considered. The other possibility when  $v_3$  states its color we consider in the next paragraph.

(2) If  $c(v_1) = r$  and  $c(v_2) = b$ , or  $c(v_4) = b$  and  $c(v_5) = r$ , then in each of the  $2^3 + 2^3 - 2 = 14$  cases the team loses. Certainly,  $v_3$  states its color only when  $v_2$  and  $v_4$  are red. Without loss of generality we assume that in this situation  $v_3$  states it is blue. If  $c(v_2) = c(v_3) = c(v_4) = r$ , then in each of the 4 cases, the team loses. Since  $(C(P_5, v_1^r, v_2^b) \cup C(P_5, v_4^b, v_5^r)) \cap C(P_5, v_2^r, v_3^r, v_4^r) = \emptyset$ , the team loses in at least  $14 + 4 = 18$  cases, and wins in at most 14 cases. This implies that  $p(S) \leq \frac{14}{32} < \frac{1}{2}$ , a contradiction.

The result for  $n \geq 6$  we prove by the induction on the number of vertices of a path. Let us assume that  $n$  is an integer such that  $n \geq 6$ , and  $h(P_{n-1}) = \frac{1}{2}$ . We will prove that  $h(P_n) = \frac{1}{2}$ . Let  $S$  be an optimal strategy for  $P_n$ . If for some  $i \in \{1, 3, n\}$  the vertex  $v_i$  never states its color, then by Theorem 3 we have  $h(P_n) = h(P_n - v_i)$ . If  $i \in \{1, n\}$ , then  $P_n - v_i = P_{n-1}$ , so  $h(P_n) = h(P_{n-1}) = \frac{1}{2}$ . If  $i = 3$ , then  $P_n - v_3 = P_2 \cup P_{n-3}$ . Since  $P_2 \cup P_{n-3} \subseteq P_{n-1}$ , we have  $h(P_2 \cup P_{n-3}) \leq h(P_{n-1}) = \frac{1}{2}$ , so  $h(P_n) = h(P_2 \cup P_{n-3}) \leq \frac{1}{2}$ . Since

$h(P_n) \geq \frac{1}{2}$ , we get  $h(P_n) = \frac{1}{2}$ . Now assume that every vertex from the set  $\{v_1, v_3, v_n\}$  states its color. If some from these vertices always states its color, then by Corollary 2 we have  $h(P_n) = \frac{1}{2}$ . If no vertex from the set  $\{v_1, v_3, v_n\}$  always states its color, then without loss of generality we assume that  $v_1$  states its color when  $v_2$  is blue, and in this situation it states it is blue. Similarly, since  $N_{P_n}[v_1] \cap N_{P_n}[v_n] = \emptyset$ , we may assume that  $v_n$  states its color when  $v_{n-1}$  is blue, and in this situation it states it is blue. We consider the following two possibilities: (1)  $v_3$  states its color when  $v_2$  is blue; (2)  $v_3$  does not state its color when  $v_2$  is blue.

(1) Let the strategy  $S'$  differ from  $S$  only in that  $v_3$  does not state its color when  $v_2$  is blue. Since in every case in which  $v_2$  is blue,  $v_1$  states its color, the statement of  $v_3$  cannot improve the result of any of these cases. Therefore,  $p(S) \leq p(S')$ . Since  $S \in \mathcal{F}^0(P_n)$ , the strategy  $S'$  is also optimal for  $P_n$ . If  $v_3$  never states its color in the strategy  $S'$ , then we have the possibility already considered. The other possibility when  $v_3$  states its color we consider in the next paragraph.

(2) If  $c(v_1) = r$  and  $c(v_2) = b$ , or  $c(v_{n-1}) = b$  and  $c(v_n) = r$ , then in each of the  $(\frac{1}{4} + \frac{1}{4} - \frac{1}{4^2})|C(P_n)| = \frac{7}{16}|C(P_n)|$  cases the team loses. Certainly,  $v_3$  can state its color only when  $v_2$  is red. Without loss of generality we assume that  $v_3$  states its color when  $v_2$  is red and  $v_4$  is blue, and in this situation it states it is blue. If  $c(v_2) = c(v_3) = r$  and  $c(v_4) = b$ , then the team loses. All the cases in which  $c(v_{n-1}) = b$  and  $c(v_n) = r$  have been counted, so it remains to count the such ones that  $c(v_2) = c(v_3) = r$ ,  $c(v_4) = b$ , and  $(c(v_{n-1}) = r$  or  $c(v_n) = b)$ . There are  $\frac{1}{2^3} \cdot \frac{3}{4} \cdot |C(P_n)| = \frac{3}{32}|C(P_n)|$  such cases. This implies that the team loses in at least  $(\frac{7}{16} + \frac{3}{32})|C(P_n)| = \frac{17}{32}|C(P_n)|$  cases, and wins in at most  $\frac{15}{32}|C(P_n)|$  cases. It means that  $p(S) \leq \frac{15}{32} < \frac{1}{2}$ , a contradiction. ■

Now we solve the hat problem on trees.

**Theorem 5** *For every tree  $T$  we have  $h(T) = \frac{1}{2}$ .*

**Proof.** The result we prove by induction on the number of vertices of a tree. If  $T$  has one vertex, that is  $T = K_1$ , it is obvious that the theorem is true. Let  $T$  be any tree with  $n \geq 2$  vertices, and let us assume that  $h(T') = \frac{1}{2}$  for every tree  $T'$  with  $n - 1$  vertices. Every tree has at least two leaves (that is vertices of a tree having exactly one neighbour). If  $T$  has exactly two leaves, then  $T$  is a path, and by Theorem 4 we have  $h(T) = \frac{1}{2}$ . If  $T$  has at least three leaves, then

let  $v_1, v_2$ , and  $v_3$  be any three of them. Let  $S$  be an optimal strategy for  $T$ . Since  $v_1, v_2$ , and  $v_3$  are leaves, there are exactly two possible situations of each of them. If for some  $i \in \{1, 2, 3\}$  the vertex  $v_i$  never states its color, then by Theorem 3 we have  $h(T) = h(T - v_i)$ . Since  $T - v_i$  is a tree with  $n - 1$  vertices, by the inductive assumption we have  $h(T - v_i) = \frac{1}{2}$ , and therefore  $h(T) = \frac{1}{2}$ . Now assume that every vertex from the set  $\{v_1, v_2, v_3\}$  states its color. If one of them always states its color, then by Corollary 2 we have  $h(T) = \frac{1}{2}$ . Now assume that every vertex from the set  $\{v_1, v_2, v_3\}$  states its color in exactly one situation. We consider the following two possibilities: (1) at least two leaves from the set  $\{v_1, v_2, v_3\}$  have the same neighbour, that is,  $N_T(v_i) = N_T(v_j)$  for certain  $i, j \in \{1, 2, 3\}, i \neq j$ ; (2) every leaf from the set  $\{v_1, v_2, v_3\}$  has another neighbour, that is,  $N_T(v_1) \neq N_T(v_2) \neq N_T(v_3)$  and  $N_T(v_1) \neq N_T(v_3)$ .

(1) Let us denote  $\{x\} = N_T(v_i) = N_T(v_j)$ . We consider the following two possibilities: (1.1)  $v_i$  and  $v_j$  state their colors in the same situation; (1.2)  $v_i$  and  $v_j$  state their colors in different situations.

(1.1) Without loss of generality we assume that  $v_i$  and  $v_j$  state their colors when  $x$  is blue. Let the strategy  $S'$  differ from  $S$  only in that  $v_j$  does not state its color when  $x$  is blue, that is,  $v_j$  never states its color. Since in every case in which  $x$  is blue  $v_i$  states its color, the statement of  $v_j$  cannot improve the result of any of these cases. Therefore,  $p(S) \leq p(S')$ . Since  $S \in \mathcal{F}^0(T)$ , the strategy  $S'$  is also optimal for  $T$ . Since  $v_j$  never states its color in the strategy  $S'$ , we have the possibility already considered.

(1.2) Without loss of generality we assume that  $v_i$  states its color when  $x$  is blue and  $v_j$  states its color when  $x$  is red. By Fact 1 we have

$$|Cl(S, x^b)| \geq \frac{|C(T, x^b)|}{2} \text{ and } |Cl(S, x^r)| \geq \frac{|C(T, x^r)|}{2}.$$

This implies that

$$|Cl(S)| = |Cl(S, x^b)| + |Cl(S, x^r)| \geq \frac{|C(T, x^b)|}{2} + \frac{|C(T, x^r)|}{2} = \frac{|C(T)|}{2}.$$

Consequently,

$$p(S) = \frac{|Cw(S)|}{|C(T)|} = \frac{|C(T)| - |Cl(S)|}{|C(T)|} \leq \frac{|C(T)| - \frac{|C(T)|}{2}}{|C(T)|} = \frac{1}{2}.$$

Since  $S \in \mathcal{F}^0(T)$ , we have  $h(T) \leq \frac{1}{2}$ . Since  $h(T) \geq \frac{1}{2}$ , we get  $h(T) = \frac{1}{2}$ .

(2) If  $i \in \{1, 2, 3\}$ , then let us denote  $N_T(v_i) = \{v'_i\}$ . Without loss of generality we assume that  $v_1$  states its color when  $v'_1$  is blue, and in this situation it states

it is blue. Similarly, since  $v'_1 \neq v'_2 \neq v'_3$  and  $v'_1 \neq v'_3$ , we may assume that  $v_2$  states its color when  $v'_2$  is blue and in this situation it states it is blue, and  $v_3$  states its color when  $v'_3$  is blue and in this situation it states it is blue. No vertex from the set  $\{v_1, v_2, v_3\}$  states its color if and only if  $c(v'_1) = c(v'_2) = c(v'_3) = r$ . If  $(c(v_1) = r \text{ and } c(v'_1) = b)$  or  $(c(v_2) = r \text{ and } c(v'_2) = b)$ , or  $(c(v_3) = r \text{ and } c(v'_3) = b)$ , then in each of the  $(1 - (1 - \frac{1}{4})^3)|C(T)| = \frac{37}{64}|C(T)|$  cases the team loses. This implies that the team wins in at most  $\frac{27}{64}|C(T)|$  cases. Consequently,

$$p(S) = \frac{|Cw(S)|}{|C(T)|} \leq \frac{\frac{27}{64}|C(T)|}{|C(T)|} = \frac{27}{64} < \frac{1}{2},$$

a contradiction. ■

Now we consider the hat problem on a graph such that the only information we know about are the degrees of vertices. In the following theorem we give an upper bound on the chance of success of any strategy for the hat problem on a graph with given degrees of vertices.

**Theorem 6** *Let  $G$  be a graph and let  $S$  be any strategy for this graph. Then*

$$|Cw(S)| \leq \sum_{v \in V(G)} \left[ 2^{d_G(v)+1} - \frac{|Cw(S)|}{2^{|V(G)|-d_G(v)-1}} \right] \cdot 2^{|V(G)|-d_G(v)-1}.$$

**Proof.** Let  $v_i$  be a vertex of  $G$ . Every statement of the color in any situation done by  $v_i$  is wrong in exactly  $2^{|V(G)|-d_G(v_i)-1}$  cases, because to every situation of  $v_i$  correspond  $2^{|V(G)|-d_G(v_i)}$  cases, and in the half of them  $v_i$  has another color than it states it has. The vertex  $v_i$  cannot state its color in at least

$$\left[ 2^{d_G(v_i)+1} - \frac{|Cw(S)|}{2^{|V(G)|-d_G(v_i)-1}} \right] + 1$$

situations, otherwise its statements are wrong in at least

$$\begin{aligned} & 2^{|V(G)|-d_G(v_i)-1} \left( \left[ 2^{d_G(v_i)+1} - \frac{|Cw(S)|}{2^{|V(G)|-d_G(v_i)-1}} \right] + 1 \right) \\ & > 2^{|V(G)|-d_G(v_i)-1} \left( 2^{d_G(v_i)+1} - \frac{|Cw(S)|}{2^{|V(G)|-d_G(v_i)-1}} \right) = 2^{|V(G)|} - |Cw(S)| \end{aligned}$$

cases. This implies that the team loses in more than  $2^{|V(G)|} - |Cw(S)|$  cases, and wins in less than

$$|C(G)| - (2^{|V(G)|} - |Cw(S)|) = 2^{|V(G)|} - 2^{|V(G)|} + |Cw(S)| = |Cw(S)|$$

cases, but  $|Cw(S)|$  is the number of cases in which the team wins, a contradiction. Since the vertex  $v_i$  does not state its color in at least

$$\left[ 2^{d_G(v_i)+1} - \frac{|Cw(S)|}{2^{|V(G)|-d_G(v_i)-1}} \right] + 1$$

situations, it states its color in at most

$$\left[ 2^{d_G(v_i)+1} - \frac{|Cw(S)|}{2^{|V(G)|-d_G(v_i)-1}} \right]$$

situations. Every statement of the color in any situation done by  $v_i$  is correct in exactly  $2^{|V(G)|-d_G(v_i)-1}$  cases, because to every situation of  $v_i$  correspond  $2^{|V(G)|-d_G(v_i)-1}$  cases, and in the half of them  $v_i$  has the color it states it has. Therefore, the statements of  $v_i$  are correct in at most

$$\left[ 2^{d_G(v_i)+1} - \frac{|Cw(S)|}{2^{|V(G)|-d_G(v_i)-1}} \right] \cdot 2^{|V(G)|-d_G(v_i)-1}$$

cases. This implies that the team wins in at most

$$\sum_{v \in V(G)} \left[ 2^{d_G(v)+1} - \frac{|Cw(S)|}{2^{|V(G)|-d_G(v)-1}} \right] \cdot 2^{|V(G)|-d_G(v)-1}$$

cases. ■

In the following three facts we show that the upper bound from the previous theorem together with integrality constraints is tight on complete graphs with two, three, and four vertices, respectively.

**Fact 7**  $h(K_2) = \frac{1}{2}$ .

**Proof.** Let  $S$  be any strategy for  $K_2$ . By Theorem 6 we have

$$|Cw(S)| \leq \sum_{v \in V(K_2)} \left[ 2^{d_{K_2}(v)+1} - \frac{|Cw(S)|}{2^{|V(K_2)|-d_{K_2}(v)-1}} \right] \cdot 2^{|V(K_2)|-d_{K_2}(v)-1}.$$

Since  $|V(K_2)| = 2$  and every vertex in  $K_2$  has exactly one neighbour, we get

$$|Cw(S)| \leq 2 \cdot [2^2 - |Cw(S)|] \Leftrightarrow |Cw(S)| \leq 8 - 2|Cw(S)| \Leftrightarrow |Cw(S)| \leq \frac{2}{3}.$$

This implies that  $|Cw(S)| \leq 2$ , as  $n \in \mathbb{N}$ . Consequently,

$$p(S) = \frac{|Cw(S)|}{|C(K_2)|} \leq \frac{2}{2^2} = \frac{1}{2}.$$

Since  $S$  is any strategy for  $K_2$ , we have  $h(K_2) \leq \frac{1}{2}$ . Since  $h(K_2) \geq \frac{1}{2}$ , we get  $h(K_2) = \frac{1}{2}$ . ■



**Fact 8**  $h(K_3) = \frac{3}{4}$ .

**Proof.** Let  $S$  be any strategy for  $K_3$ . By Theorem 6 we have

$$|Cw(S)| \leq \sum_{v \in V(K_3)} \left[ 2^{d_{K_3}(v)+1} - \frac{|Cw(S)|}{2^{|V(K_3)|-d_{K_3}(v)-1}} \right] \cdot 2^{|V(K_3)|-d_{K_3}(v)-1}.$$

Since  $|V(K_3)| = 3$  and every vertex in  $K_3$  has exactly two neighbours, we get

$$|Cw(S)| \leq 3 \cdot [2^3 - |Cw(S)|] \Leftrightarrow |Cw(S)| \leq 24 - 3|Cw(S)| \Leftrightarrow |Cw(S)| \leq 6.$$

Consequently,

$$p(S) = \frac{|Cw(S)|}{|C(K_3)|} \leq \frac{6}{2^3} = \frac{3}{4}.$$

Since  $S$  is any strategy for  $K_3$ , we have  $h(K_3) \leq \frac{3}{4}$ . Let  $S_1 \in \mathcal{F}(K_3)$  be the strategy such that every vertex considers colors of its two neighbours, and if they are the same, it states it has the opposite color. If they are different, it passes. It is easy to verify that  $|Cw(S_1)| = 6$ . Since  $|C(K_3)| = 2^3 = 8$ , we have  $p(S_1) = \frac{|Cw(S_1)|}{|C(K_3)|} = \frac{6}{8} = \frac{3}{4}$ . Since  $p(S_1) \leq h(K_3)$ , we have  $h(K_3) \geq \frac{3}{4}$ . Since  $h(K_3) \geq \frac{3}{4}$  and  $h(K_3) \leq \frac{3}{4}$ , we get  $h(K_3) = \frac{3}{4}$ . ■

**Fact 9**  $h(K_4) = \frac{3}{4}$ .

**Proof.** Let  $S$  be any strategy for  $K_4$ . By Theorem 6 we have

$$|Cw(S)| \leq \sum_{v \in V(K_4)} \left[ 2^{d_{K_4}(v)+1} - \frac{|Cw(S)|}{2^{|V(K_4)|-d_{K_4}(v)-1}} \right] \cdot 2^{|V(K_4)|-d_{K_4}(v)-1}.$$

Since  $|V(K_4)| = 4$  and every vertex in  $K_4$  has three neighbours, we get

$$|Cw(S)| \leq 4 \cdot [2^4 - |Cw(S)|] \Leftrightarrow |Cw(S)| \leq 64 - 4|Cw(S)| \Leftrightarrow |Cw(S)| \leq 12\frac{4}{5}.$$

This implies that  $|Cw(S)| \leq 12$ , as  $|Cw(S)| \in N$ . Consequently,

$$p(S) = \frac{|Cw(S)|}{|C(K_4)|} \leq \frac{12}{2^4} = \frac{3}{4}.$$

Since  $S$  is any strategy for  $K_4$ , we have  $h(K_4) \leq \frac{3}{4}$ . Since  $K_3 \subseteq K_4$  and  $h(K_3) = \frac{3}{4}$ , we get  $h(K_3) \leq h(K_4)$ . Since  $h(K_3) = \frac{3}{4}$ , we have  $h(K_4) \geq \frac{3}{4}$ . This implies that  $h(K_4) = \frac{3}{4}$ . ■

In the next fact we solve the hat problem on the graph  $K_3 \cup K_2$ .



**Fact 10**  $h(K_3 \cup K_2) = \frac{3}{4}$ .

**Proof.** Let  $E(K_3 \cup K_2) = \{v_1v_2, v_2v_3, v_3v_1, v_4v_5\}$ . Let  $S$  be any strategy for the graph  $K_3 \cup K_2$ . By Theorem 6 we have

$$|Cw(S)| \leq \sum_{v \in V(K_3 \cup K_2)} \left[ 2^{d_{K_3 \cup K_2}(v)+1} - \frac{|Cw(S)|}{2^{|V(K_3 \cup K_2)| - d_{K_3 \cup K_2}(v) - 1}} \right] \cdot 2^{|V(K_3 \cup K_2)| - d_{K_3 \cup K_2}(v) - 1}.$$

Since  $d_{K_3 \cup K_2}(v_1) = d_{K_3 \cup K_2}(v_2) = d_{K_3 \cup K_2}(v_3) = 2$  and  $d_{K_3 \cup K_2}(v_4) = d_{K_3 \cup K_2}(v_5) = 1$ , we get

$$\begin{aligned} |Cw(S)| &\leq 3 \cdot 2^2 \cdot \left\lfloor 2^3 - \frac{|Cw(S)|}{2^2} \right\rfloor + 2 \cdot 2^3 \cdot \left\lfloor 2^2 - \frac{|Cw(S)|}{2^3} \right\rfloor \\ &= 12 \left\lfloor 8 - \frac{|Cw(S)|}{4} \right\rfloor + 16 \left\lfloor 4 - \frac{|Cw(S)|}{8} \right\rfloor. \end{aligned}$$

This implies that

$$\begin{aligned} |Cw(S)| &\leq 12 \left( 8 - \frac{|Cw(S)|}{4} \right) + 16 \left( 4 - \frac{|Cw(S)|}{8} \right) \\ &= 96 - 3|Cw(S)| + 64 - 2|Cw(S)| = 160 - 5|Cw(S)|. \end{aligned}$$

Now we easily get  $|Cw(S)| \leq \frac{160}{6} = 26\frac{2}{3}$ . Since  $|Cw(S)|$  is an integer, we have  $|Cw(S)| \leq 26$ . Assume that  $|Cw(S)| = 26$ . We have  $26 \leq 12\lfloor 8 - \frac{26}{4} \rfloor + 16\lfloor 4 - \frac{26}{8} \rfloor = 12 \cdot 1 + 16 \cdot 0 = 12$ , a contradiction. Now assume that  $|Cw(S)| = 25$ . We have  $25 \leq 12\lfloor 8 - \frac{25}{4} \rfloor + 16\lfloor 4 - \frac{25}{8} \rfloor = 12 \cdot 1 + 16 \cdot 0 = 12$ , a contradiction. This implies that  $|Cw(S)| \leq 24$ , and consequently,  $p(S) = \frac{|Cw(S)|}{|C(K_3 \cup K_2)|} \leq \frac{24}{32}$ . Since  $S$  is any strategy for  $K_3 \cup K_2$ , we have  $h(K_3 \cup K_2) \leq \frac{3}{4}$ . Since  $K_3 \subseteq K_3 \cup K_2$  and  $h(K_3) = \frac{3}{4}$ , we get  $h(K_3 \cup K_2) \geq h(K_3) = \frac{3}{4}$ . This implies that  $h(K_3 \cup K_2) = \frac{3}{4}$ . ■

## References

- [1] G. Aggarwal, A. Fiat, A. Goldberg, J. Hartline, N. Immorlica, and M. Sudan, *Derandomization of auctions*, in Proceedings of the 37th ACM Symposium on Theory of Computing, pages 619–625, 2005.
- [2] J. Aspnes, R. Beigel, M. Furst, and S. Rudich, *The expressive power of voting polynomials*, *Combinatorica* 14 (1994) 135-148.

- [3] R. Beigel, L. Fortnow, and F. Stephan, *Infinitely–Often Autoreducible Sets*, SIAM Journal on Computing 36 (2006), 595–608.
- [4] M. Bernstein, *The Hat Problem and Hamming Codes*, MAA Focus, November, 2001, 4–6.
- [5] W. Blum, *Denksport für Hutträger*, Die Zeit, May 3, 2001.
- [6] M. Breit, D. Deshommes, and A. Falden, *Hats required: Perfect and Imperfect strategies for the Hat Problem*, preprint.
- [7] E. Burke, S. Gustafson, and G. Kendall, *A Puzzle to Challenge Genetic Programming*, in Genetic Programming, 136–147, Lecture Notes in Computer Science, Springer, 2002.
- [8] S. Butler, M. T. Hajiaghayi, R. D. Kleinberg, and T. Leighton, *Hat Guessing Games*, SIAM Journal on Discrete Mathematics 22 (2008), 592–605.
- [9] A. Dąbrowski, *Kolorowe czapeczki*, Delta 7/2005, 1–5, in Polish.
- [10] T. Ebert, *Applications of recursive operators to randomness and complexity*, Ph.D. Thesis, University of California at Santa Barbara, 1998.
- [11] T. Ebert and W. Merkle, *Autoreducibility of Random Sets: a Sharp Bound on the Density of Guessed Bits*, in Mathematical Foundations of Computer Science 2002, Lecture Notes in Computer Science, 2420: 221–233, Springer, 2002.
- [12] T. Ebert, W. Merkle, and H. Vollmer, *On the Autoreducibility of Random Sequences*, SIAM Journal on Computing 32 (2003), 1542–1569.
- [13] T. Ebert and H. Vollmer, *On the Autoreducibility of Random Sequences*, in M. Nielsen and B. Rovan (editors) Mathematical Foundations of Computer Science 2000, Lecture Notes in Computer Science, 1893: 333–342, Springer, 2000.
- [14] U. Feige, *You can leave Your hat on (if you guess its color)*, Technical Report MCS04-03, Computer Science and Applied Mathematics, The Weizmann Institute of Science, pp. 10, 2004.

- [15] W. Guo, S. Kasala, M.B. Rao, and B. Tucker, *The Hat Problem and Some Variations*, in N. Balakrishnan, J. M. Sarabia, and E. Castillo (editors) *Advances in Distribution Theory, Order Statistics, and Interference*, Statistics for Industry and Technology, Birkhäuser Boston, 2007, 459–479.
- [16] A. Hoffman and J. Kruskal, *Integral boundary points of convex polyhedra*, in H. Kuhn and A. Tucker (editors) *Linear Inequalities and Related Systems*, Princeton University Press, 1956.
- [17] N. Immorlica, *Computing with Strategic Agents*, Ph.D. Thesis, Massachusetts Institute of Technology, 2005.
- [18] H. W. Lenstra and G. Seroussi, *On Hats and other Covers*, IEEE International Symposium on Information Theory, Lausanne, 2002.
- [19] J. A. Poulos, *Could You Solve This \$1 Million Hat Trick?*, abcNews, November 29, 2001.
- [20] S. Robinson, *Why mathematicians now care about their hat color*, The New York Times, Science Times Section, page D5, April 10, 2001.