

catena-Poly[[[bis(thiourea- κ S)cadmium]-di- μ -thiocyanato- κ^2 N:S; κ^2 S:N]dihydrate]

Anna Mietlarek-Kropidłowska* and Jaroslaw Chojnacki

 Department of Inorganic Chemistry, Chemical Faculty, Gdansk University of Technology, 11/12 G. Narutowicza Str., 80-233 Gdańsk, Poland
 Correspondence e-mail: anna.mietlarek-kropidłowska@pg.gda.pl


Received 18 June 2012; accepted 3 July 2012

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{N}-\text{C}) = 0.003$ Å; R factor = 0.028; wR factor = 0.062; data-to-parameter ratio = 26.7.

The title compound, $[\{\text{Cd}(\text{NCS})_2(\text{CH}_4\text{N}_2\text{S})_2\} \cdot 2\text{H}_2\text{O}]_n$, forms a one-dimensional chain parallel to the a axis, caused by the presence of the bridging thiocyanate groups. Two solvent molecules per complex are present in the lattice. The Cd^{II} ion is situated on an inversion centre and is coordinated in a distorted octahedral fashion by two N and two S atoms from four thiourea ligands and by two S atoms from two thiocyanate ligands. Weak $\text{O}-\text{H} \cdots \text{S}$, $\text{N}-\text{H} \cdots \text{O}$ and $\text{N}-\text{H} \cdots \text{N}$ interactions reinforce the structure.

Related literature

For a general introduction to thiocyanato complexes, see: Nardelli *et al.* (1957). For the syntheses and structures of a series of cadmium complexes with thiourea derivatives and thiocyanato ligands, see: Wang *et al.* (2002); Cavalca *et al.* (1960); Zhu *et al.* (2000); Yang *et al.* (2001); Ahmad *et al.* (2008); Williams *et al.* (1992). For information on the properties of complexes incorporating these ligands, see: Yuan *et al.* (1997); Krunko *et al.* (1997); Amutha *et al.* (2011); Machura *et al.* (2011). For the use of Cd^{II} complexes with mixed S-donor ligands as precursors to CdS, see: Kropidłowska *et al.* (2008).



Experimental

Crystal data

$[\text{Cd}(\text{NCS})_2(\text{CH}_4\text{N}_2\text{S})_2] \cdot 2\text{H}_2\text{O}$
 $M_r = 416.84$
 Triclinic, $P\bar{1}$
 $a = 5.8533$ (3) Å
 $b = 7.3527$ (3) Å
 $c = 8.8630$ (4) Å
 $\alpha = 73.413$ (4)°
 $\beta = 76.926$ (4)°

$\gamma = 88.856$ (4)°
 $V = 355.69$ (3) Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 2.12$ mm⁻¹
 $T = 293$ K
 $0.53 \times 0.42 \times 0.23$ mm

Data collection

Oxford Diffraction KM-4-CCD diffractometer
 Absorption correction: analytical [*CrysAlis PRO* (Oxford Diffraction, 2008), based on expressions

derived by Clark & Reid (1995)]
 $T_{\text{min}} = 0.558$, $T_{\text{max}} = 0.725$
 7642 measured reflections
 2268 independent reflections
 1985 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.062$
 $S = 1.05$
 2268 reflections
 85 parameters
 3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.47$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.51$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1A} \cdots \text{O1}^{\text{i}}$	0.86	2.26	3.049 (3)	153
$\text{N1}-\text{H1B} \cdots \text{N3}^{\text{ii}}$	0.86	2.3	3.147 (3)	167
$\text{N2}-\text{H2A} \cdots \text{O1}^{\text{i}}$	0.86	2.4	3.159 (3)	147
$\text{N2}-\text{H2B} \cdots \text{O1}^{\text{iii}}$	0.86	2.19	3.050 (3)	175
$\text{O1}-\text{H1C} \cdots \text{S2}$	0.80 (2)	2.54 (2)	3.340 (2)	177 (4)
$\text{O1}-\text{H1D} \cdots \text{S1}^{\text{iv}}$	0.81 (2)	2.59 (2)	3.377 (2)	165 (3)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x + 1, y, z$; (iii) $-x, -y + 1, -z + 1$; (iv) $x, y + 1, z$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2008); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *publCIF* (Westrip, 2010).

The research was supported by grants from the Polish Ministry of Education and Science (grant Nos. N N204 543339 and N N204 150237).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2575).

References

- Ahmad, S., Altaf, M., Malik, M. R., Ali, S. & Stoeckli-Evans, H. (2008). Private communication (CSD ID: XOMSUN01). CCDC, Cambridge, England.
 Amutha, R., Muruganandham, M., Lee, G. J. & Wu, J. J. (2011). *J. Nanosci. Nanotechnol.* **11**, 7940–7944.
 Cavalca, L., Nardelli, M. & Fava, G. (1960). *Acta Cryst.* **13**, 125–130.
 Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.

- Kropidłowska, A., Chojnacki, J., Fahmi, A. & Becker, B. (2008). *Dalton Trans.* pp. 6825–6831.
- Krunks, M., Madarasz, J., Hiltunen, L., Mannonen, R., Mellikov, E. & Niinistö, L. (1997). *Acta Chem. Scand.* **51**, 294–301.
- Machura, B., Nawrot, I. & Michalik, K. (2011). *Polyhedron*, **30**, 2619–2626.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Nardelli, M., Braibanti, A. & Fava, G. (1957). *Gazz. Chim. Ital.* **87**, 1209–1231.
- Oxford Diffraction (2008). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, X. Q., Yu, W. T., Xu, D., Lu, M. K. & Yuan, D. R. (2002). *Acta Cryst.* **C58**, m336–m337.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Williams, D. J., VanDerveer, D., Lipscomb, L. A. & Jones, R. L. (1992). *Inorg. Chim. Acta*, **192**, 51–57.
- Yang, G., Liu, G.-F., Zheng, S.-L. & Chen, X.-M. (2001). *J. Coord. Chem.* **53**, 269–279.
- Yuan, D. R., Xu, D., Fang, Q., Yu, W. T. & Jiang, M. H. (1997). *Appl. Phys. Lett.* **70**, 544–546.
- Zhu, H.-G., Yang, G., Chen, X.-M. & Ng, S. W. (2000). *Acta Cryst.* **C56**, e430–e431.

supporting information

Acta Cryst. (2012). E68, m1051–m1052 [https://doi.org/10.1107/S1600536812030267]

***catena*-Poly[[[bis(thiourea- κ S)cadmium]-di- μ -thiocyanato- κ^2 N:S; κ^2 S:N] dihydrate]**

Anna Mietlarek-Kropidłowska and Jaroslaw Chojnacki

S1. Comment

The interest in the coordination compounds possessing simultaneously thiourea and thiocyanato ligands dates back to the 1950s (*e.g.* Nardelli *et al.*, 1957) when the nature of coordination compounds formed by divalent cations ($M = \text{Mn, Co, Ni, Cd, Pb}$) and organic molecules containing sulfur was extensively studied. Besides the polymeric *catena*-poly[bis(thiocyanato- κ N)bis(μ -thiourea- κ^2 S:S)cadmium(II)] (Wang *et al.*, 2002), also structures of four other complexes with thiourea derivatives of $[\text{Cd}(\text{SCN})_2(\text{TU})_2]_n$ type, where TU = ethylenethiourea (Cavalca *et al.*, 1960), N,N' -diphenylthiourea (Zhu *et al.*, 2000), N -phenylthiourea (Yang *et al.*, 2001; Ahmad *et al.*, 2008) or 1,3-dimethyl-2(3*H*)-imidazolethione (Williams *et al.*, 1992), can be found. The interest in these compounds is related either with their non-linear optical properties (Yuan *et al.*, 1997) or with the possibility to use them as single-source precursors of semiconducting materials based on CdS (Krunks *et al.*, 1997; Amutha *et al.*, 2011). Moreover, the use of SCN ligands, with bridging abilities, may lead to intriguing architectures and topologies, often generating one-dimensional chains (Machura *et al.*, 2011). It was the reason why, during our studies on the new molecular precursors (Kropidłowska *et al.*, 2008), we have turned our attention to systems of this type now and we have obtained a new complex containing thiourea and thiocyanato ligands connected to cadmium center.

The cadmium atom in *catena*-Poly[[[bis(thiourea- κ S)cadmium]-di- μ -thiocyanato- κ^2 N:S; κ^2 S:N] dihydrate] is located at the inversion center and is octahedrally coordinated by two S atoms and two N atoms from four thiocyanate groups as well as by two S atoms from thiourea molecules. The neighbouring Cd^{II} ions are bridged by two μ -SCN- κ^2 N:S ligands, thus forming eight-membered ring of $[\text{Cd-SCN}]_2$ type with the Cd...Cd distance of 5.853 Å, which is close to the values observed in other bridged systems (Machura *et al.*, 2011). These units form one-dimensional chains of slightly distorted edge-shared Cd-centered octahedra along the [100] crystallographic direction. The Cd—S i Cd—N distances are typical for cadmium(II) thiocyanate complexes. The IR spectra clearly show the presence of the thiocyanato groups (with the maxima of ν_{CN} absorption at 2078 cm^{-1}).

In the structure of $[[\text{Cd}\{\text{SC}(\text{NH}_2)_2\}_2(\text{SCN})_2] \cdot 2\text{H}_2\text{O}]_n$, several weak interactions may be assumed, leading to the alternating arrangement of water and complex molecules. Each water molecule interacts with S or N atoms from the three neighboring polymeric chains. Thus, it can serve as a donor of a weak hydrogen bond to the sulfur atom from one of the thiourea moieties (O1(H1D)—S1^{viii}) in one chain, as well as to sulfur from one thiocyanato ligand (O1(H1C)—S2) in the other. The oxygen lone pairs act as acceptors towards NH_2 groups from thiourea moieties located within the third chain (N2(H2B)—O1^{vii}, N1(H1A)—O1^v, N2(H2A)—O1^v). Finally, one "interchain" interaction, N1—H1B...N3ⁱ, operates between NH_2 and SCN groups.



S2. Experimental

The reaction was carried out between 0.50 g cadmium(II) thiocyanate, $\text{Cd}(\text{SCN})_2$, and 1.34 g thiourea (molar ratio 1:8) which were dissolved in a small amount of water. The mixture was heated to 70°C and stirred using magnetic stirrer for 50 minutes and then left for crystallization at room temperature. After a few days two types of crystals appeared in the flask: needles (0.2 g) and blocks (0.1 g), which were mechanically separated under the microscope. The structure of needle-like crystals $[\text{Cd}(\text{SCN})_2\{\mu\text{-SC}(\text{NH}_2)_2\}]_n$ (m.p. 189°C) has been already described (Wang *et al.*, 2002), while the block-like crystals appeared to be a new compound, crystallizing as diaqua solvate $[[\text{Cd}\{\text{SC}(\text{NH}_2)_2\}_2(\text{SCN})_2]\cdot 2\text{H}_2\text{O}]_n$ (m.p. 187°C). The product, when taken from the mother liquor and dried using the filter paper, changes - becomes opaque and finally takes the form of a powder (most probably because of the removal of the solvent molecules). IR spectra were recorded using Mattson Genesis II Gold spectrometer equipped with Momentum Microscope as detector.

S3. Refinement

All N—H atoms were placed in calculated positions and refined as riding on their carrier atoms with N—H = 0.86 \AA (NH_2) and $U_{\text{iso}}(\text{H}) = 1.2$ times $U_{\text{eq}}(\text{N})$. Solvent O—H hydrogen atoms were found in the Fourier map and refined as constrained to: O—H bond length of 0.80 \AA , H1C - H1D distance of 1.30 \AA and $U_{\text{iso}}(\text{H}) = 1.5$ times $U_{\text{eq}}(\text{O})$ with the default uncertainties.




Figure 1

Molecular structure and atom-numbering scheme for $[[\text{Cd}\{\text{SC}(\text{NH}_2)_2\}_2(\text{SCN})_2]\cdot 2\text{H}_2\text{O}]_n$ with displacement ellipsoids drawn at 50% probability level. H atoms are represented as arbitrary circles. Symmetry codes: (i) $x+1, y, z$; (ii) $-x, -y+1, -z$; (iii) $-x+1, -y+1, -z$.




Figure 2

Weak interaction present in the crystal structure of $[[\text{Cd}\{\text{SC}(\text{NH}_2)_2\}_2(\text{SCN})_2]\cdot 2\text{H}_2\text{O}]_n$ between the water and complex molecules (on the left). The arrangement of water and complex molecules in the crystal (on the right). Dashed lines denote the possible weak interactions.

catena-Poly[[[bis(thiourea- κ S)cadmium]-di- μ -thiocyanato- κ^2 N:S; κ^2 S:N] dihydrate]

Crystal data



$M_r = 416.84$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 5.8533$ (3) Å

$b = 7.3527$ (3) Å

$c = 8.8630$ (4) Å

$\alpha = 73.413$ (4)°

$\beta = 76.926$ (4)°

$\gamma = 88.856$ (4)°

$V = 355.69$ (3) Å³

$Z = 1$

$F(000) = 206$

$D_x = 1.946$ Mg m⁻³

Melting point: 460 K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4676 reflections

$\theta = 2.9\text{--}33.8^\circ$

$\mu = 2.12$ mm⁻¹

$T = 293$ K

Block, colourless

$0.53 \times 0.42 \times 0.23$ mm

Data collection

Oxford Diffraction KM-4-CCD
diffractometer

Graphite monochromator

ω -scan

Absorption correction: analytical

[*CrysAlis PRO* (Oxford Diffraction, 2008),
based on expressions derived by Clark & Reid
(1995)]

$T_{\min} = 0.558$, $T_{\max} = 0.725$

7642 measured reflections

2268 independent reflections

1985 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 31^\circ$, $\theta_{\min} = 2.9^\circ$

$h = -8 \rightarrow 8$

$k = -10 \rightarrow 10$

$l = -12 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.028$

$wR(F^2) = 0.062$

$S = 1.05$

2268 reflections

85 parameters

3 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0266P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.47 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.51 \text{ e } \text{\AA}^{-3}$$

Special details

Experimental. CrysAlisPro, (Oxford Diffraction, 2008). Analytical numeric absorption correction using a multifaceted crystal model based on expressions derived by (Clark & Reid, 1995).

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.5	0.5	0	0.03082 (8)
S1	0.24899 (10)	0.38566 (9)	0.30122 (7)	0.03890 (15)
S2	0.25651 (10)	0.82898 (8)	-0.03961 (7)	0.03377 (13)
N1	0.6502 (3)	0.3430 (3)	0.3988 (3)	0.0424 (5)
H1A	0.7326	0.2965	0.4686	0.051*
H1B	0.7155	0.4193	0.3061	0.051*
N2	0.3288 (4)	0.1803 (3)	0.5771 (3)	0.0481 (6)
H2A	0.4141	0.1352	0.6452	0.058*
H2B	0.1813	0.1488	0.6023	0.058*
N3	-0.1946 (3)	0.6528 (3)	0.0619 (3)	0.0391 (5)
C1	0.4240 (4)	0.2975 (3)	0.4340 (3)	0.0321 (5)
C2	-0.0091 (4)	0.7257 (3)	0.0194 (3)	0.0281 (4)
O1	0.1962 (4)	0.9249 (3)	0.3133 (2)	0.0524 (5)
H1C	0.216 (7)	0.902 (5)	0.229 (3)	0.079*
H1D	0.193 (7)	1.038 (3)	0.296 (4)	0.079*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.02632 (12)	0.03241 (13)	0.03125 (13)	-0.00184 (9)	-0.00529 (9)	-0.00613 (10)
S1	0.0255 (3)	0.0545 (4)	0.0282 (3)	0.0010 (3)	-0.0048 (2)	0.0002 (3)
S2	0.0298 (3)	0.0265 (3)	0.0402 (3)	-0.0021 (2)	-0.0038 (2)	-0.0050 (2)
N1	0.0333 (10)	0.0557 (13)	0.0361 (11)	-0.0005 (10)	-0.0120 (9)	-0.0065 (10)
N2	0.0454 (12)	0.0560 (13)	0.0333 (11)	-0.0053 (11)	-0.0110 (10)	0.0044 (10)
N3	0.0330 (10)	0.0371 (11)	0.0500 (13)	-0.0009 (9)	-0.0106 (9)	-0.0158 (10)
C1	0.0357 (12)	0.0316 (11)	0.0295 (11)	0.0009 (9)	-0.0074 (9)	-0.0100 (9)
C2	0.0335 (11)	0.0264 (10)	0.0266 (10)	0.0050 (9)	-0.0101 (9)	-0.0088 (8)
O1	0.0662 (13)	0.0522 (11)	0.0412 (11)	-0.0028 (11)	-0.0197 (10)	-0.0113 (10)

Geometric parameters (\AA , $^\circ$)

Cd1—N3 ⁱ	2.3734 (19)	N1—H1A	0.86
Cd1—N3 ⁱⁱ	2.3734 (19)	N1—H1B	0.86
Cd1—S1	2.6431 (6)	N2—C1	1.318 (3)
Cd1—S1 ⁱⁱⁱ	2.6431 (6)	N2—H2A	0.86
Cd1—S2 ⁱⁱⁱ	2.7585 (6)	N2—H2B	0.86
Cd1—S2	2.7585 (6)	N3—C2	1.154 (3)
S1—C1	1.714 (2)	N3—Cd1 ^{iv}	2.3734 (19)
S2—C2	1.649 (2)	O1—H1C	0.797 (17)
N1—C1	1.317 (3)	O1—H1D	0.805 (17)
N3 ⁱ —Cd1—N3 ⁱⁱ	180	C1—S1—Cd1	111.15 (8)
N3 ⁱ —Cd1—S1	95.01 (6)	C2—S2—Cd1	96.75 (7)
N3 ⁱⁱ —Cd1—S1	84.99 (6)	C1—N1—H1A	120
N3 ⁱ —Cd1—S1 ⁱⁱⁱ	84.99 (6)	C1—N1—H1B	120
N3 ⁱⁱ —Cd1—S1 ⁱⁱⁱ	95.01 (6)	H1A—N1—H1B	120
S1—Cd1—S1 ⁱⁱⁱ	180	C1—N2—H2A	120
N3 ⁱ —Cd1—S2 ⁱⁱⁱ	89.81 (5)	C1—N2—H2B	120
N3 ⁱⁱ —Cd1—S2 ⁱⁱⁱ	90.19 (5)	H2A—N2—H2B	120
S1—Cd1—S2 ⁱⁱⁱ	92.043 (19)	C2—N3—Cd1 ^{iv}	148.15 (19)
S1 ⁱⁱⁱ —Cd1—S2 ⁱⁱⁱ	87.957 (19)	N1—C1—N2	118.6 (2)
N3 ⁱ —Cd1—S2	90.19 (5)	N1—C1—S1	122.51 (18)
N3 ⁱⁱ —Cd1—S2	89.81 (5)	N2—C1—S1	118.87 (18)
S1—Cd1—S2	87.957 (19)	N3—C2—S2	179.5 (2)
S1 ⁱⁱⁱ —Cd1—S2	92.043 (19)	H1C—O1—H1D	108 (3)
S2 ⁱⁱⁱ —Cd1—S2	180.00 (3)		
N3 ⁱ —Cd1—S1—C1	-43.20 (10)	N3 ⁱⁱ —Cd1—S2—C2	30.81 (10)
N3 ⁱⁱ —Cd1—S1—C1	136.80 (10)	S1—Cd1—S2—C2	-54.18 (8)
S2 ⁱⁱⁱ —Cd1—S1—C1	46.79 (9)	S1 ⁱⁱⁱ —Cd1—S2—C2	125.82 (8)
S2—Cd1—S1—C1	-133.21 (9)	Cd1—S1—C1—N1	22.3 (2)
N3 ⁱ —Cd1—S2—C2	-149.19 (10)	Cd1—S1—C1—N2	-157.98 (17)

Symmetry codes: (i) $x+1, y, z$; (ii) $-x, -y+1, -z$; (iii) $-x+1, -y+1, -z$; (iv) $x-1, y, z$.Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots O1 ^v	0.86	2.26	3.049 (3)	153
N1—H1B \cdots N3 ⁱ	0.86	2.3	3.147 (3)	167
N2—H2A \cdots O1 ^v	0.86	2.4	3.159 (3)	147
N2—H2B \cdots O1 ^{vi}	0.86	2.19	3.050 (3)	175
O1—H1C \cdots S2	0.80 (2)	2.54 (2)	3.340 (2)	177 (4)
O1—H1D \cdots S1 ^{vii}	0.81 (2)	2.59 (2)	3.377 (2)	165 (3)

Symmetry codes: (i) $x+1, y, z$; (v) $-x+1, -y+1, -z+1$; (vi) $-x, -y+1, -z+1$; (vii) $x, y+1, z$.