

RESEARCH ARTICLE | SEPTEMBER 10 2014

## Valence and ionic lowest-lying electronic states of ethyl formate as studied by high-resolution vacuum ultraviolet photoabsorption, He(I) photoelectron spectroscopy, and *ab initio* calculations

M. A. Śmiałek; M. Łabuda; J. Guthmuller; M.-J. Hubin-Franskin; J. Delwiche; D. Dufлот; N. J. Mason; S. V. Hoffmann; N. C. Jones; P. Limão-Vieira




*J. Chem. Phys.* 141, 104311 (2014)  
<https://doi.org/10.1063/1.4894762>



CrossMark

This article may be downloaded for personal use only. Any other use requires prior permission of the author and AIP Publishing. This article appeared in (citation of published article) and may be found at <https://doi.org/10.1063/1.4894762>

26 February 2024 10:38:35



Chemical Physics Reviews

Special Topic: AI and Machine Learning  
in Chemical and Materials Science

Submit Today



# Valence and ionic lowest-lying electronic states of ethyl formate as studied by high-resolution vacuum ultraviolet photoabsorption, He(I) photoelectron spectroscopy, and *ab initio* calculations

M. A. Śmiałek,<sup>1,a)</sup> M. Łabuda,<sup>2</sup> J. Guthmuller,<sup>2</sup> M.-J. Hubin-Franskin,<sup>3</sup> J. Delwiche,<sup>3</sup> D. Duflot,<sup>4</sup> N. J. Mason,<sup>5</sup> S. V. Hoffmann,<sup>6</sup> N. C. Jones,<sup>6</sup> and P. Limão-Vieira<sup>7,b)</sup>

<sup>1</sup>*Department of Control and Energy Engineering, Faculty of Ocean Engineering and Ship Technology, Gdańsk University of Technology, Gabriela Narutowicza 11/12, 80-233 Gdańsk, Poland*

<sup>2</sup>*Department of Theoretical Physic and Quantum Information, Faculty of Applied Physics and Mathematics, Gdańsk University of Technology, Gabriela Narutowicza 11/12, 80-233 Gdańsk, Poland*

<sup>3</sup>*Département de Chimie, Université de Liège, Institut de Chimie-Bât. B6C, B-4000 Liège, Belgium*

<sup>4</sup>*Laboratoire de Physique des Lasers, Atomes et Molécules (PhLAM), UMR CNRS 8523, Université Lille 1 Sciences et Technologies, F-59655 Villeneuve d'Ascq Cedex, France*

<sup>5</sup>*Department of Physical Sciences, The Open University, Walton Hall, Milton Keynes MK7 6AA, United Kingdom*

<sup>6</sup>*ISA, Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, Building 1520, DK-8000 Aarhus C, Denmark*

<sup>7</sup>*Laboratório de Colisões Atômicas e Moleculares, CEFITEC, Departamento de Física, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal*

(Received 17 July 2014; accepted 25 August 2014; published online 10 September 2014)

The highest resolution vacuum ultraviolet photoabsorption spectrum of ethyl formate, C<sub>2</sub>H<sub>5</sub>OCHO, yet reported is presented over the wavelength range 115.0–275.5 nm (10.75–4.5 eV) revealing several new spectral features. Valence and Rydberg transitions and their associated vibronic series, observed in the photoabsorption spectrum, have been assigned in accordance with new *ab initio* calculations of the vertical excitation energies and oscillator strengths. Calculations have also been carried out to determine the ionization energies and fine structure of the lowest ionic state of ethyl formate and are compared with a newly recorded He(I) photoelectron spectrum (from 10.1 to 16.1 eV). New vibrational structure is observed in the first photoelectron band. The photoabsorption cross sections have been used to calculate the photolysis lifetime of ethyl formate in the upper stratosphere (20–50 km). © 2014 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4894762>]

## I. INTRODUCTION

Ethyl formate, also known as ethyl methanoate, is a volatile compound formerly used as a fumigant for grain, now restricted to use on dried fruit and processed cereal products.<sup>1</sup> Together with the simplest ester, methyl formate,<sup>2</sup> ethyl formate has been detected in the interstellar space in the dust cloud, Sagittarius B2(N).<sup>3</sup> Ethyl formate is also used as a model compound for combustion chemistry studies of ethyl esters, which are typical constituents of biodiesel fuels, showing their contribution in benzene and soot formation upon fuel combustion.<sup>4</sup> In a recent publication, we reported the electronic state spectroscopy of methyl formate using high resolution vacuum ultraviolet (VUV) photoabsorption, He(I) photoelectron spectroscopy, and *ab initio* calculations, where a detailed analysis of the vibrational progressions and several Rydberg series were proposed for the first time.<sup>5</sup>

Ethyl formate has been the subject of many theoretical and experimental studies resulting in a description of the geometry of the molecule and assignment of the observed frequencies to the vibrational modes of the molecule.<sup>6–9</sup> Ionization and dissociation of ethyl formate by electron impact has also been investigated.<sup>10</sup> Photoionization energies together with ion mass spectrometry were obtained through a He(I) ionization source,<sup>11,12</sup> synchrotron radiation, and a femtosecond laser.<sup>13</sup> The values obtained by other investigators are in a good agreement with the present data despite the higher resolution achieved in this study.

In this paper, we present high resolution VUV photoabsorption spectra with absolute cross sections, together with theoretical calculations of the vertical excitation energies and oscillator strengths of the electronic transitions of ethyl formate. To our knowledge, no high resolution studies have been reported for this molecule yet. Additionally, the He(I) photoelectron spectrum was measured and analyzed in order to clarify Rydberg assignments in the VUV spectrum. These data also provide, for the first time, a vibrational resolution of the first ionic band of ethyl formate. Absolute photoabsorption cross sections are needed in modeling studies of the Earth atmosphere and radiation chemistry of aliphatic esters.

<sup>a)</sup>Electronic mail: smialek@pg.gda.pl. Tel.: +48583472614. Fax: +48583486372. Also at Department of Physical Sciences, The Open University, Walton Hall, Milton Keynes MK7 6AA, United Kingdom.

<sup>b)</sup>Electronic mail: plimaovieira@fct.unl.pt. Tel.: +351212947859. Fax: +351212948549. Also at Department of Physical Sciences, The Open University, Walton Hall, Milton Keynes MK7 6AA, United Kingdom.




FIG. 1. High resolution photoabsorption spectrum of ethyl formate.

## II. EXPERIMENTAL PROCEDURE

### A. Ethyl formate sample

The liquid sample used both in the VUV measurements and the PES experiment was purchased from Sigma-Aldrich, with a purity of 97%. The sample was degassed by repeated freeze–pump–thaw cycles in both sets of experiments.

### B. VUV photoabsorption

The high-resolution VUV photoabsorption spectrum of ethyl formate (Fig. 1) was measured at the UV1 beam line, using the ASTRID synchrotron facility in Aarhus University, Denmark. The experimental apparatus has been described in detail previously.<sup>14</sup> Synchrotron radiation passes through a static gas sample and a photomultiplier is used to measure the transmitted light intensity. The incident wavelength is selected using a toroidal dispersion grating with 2000 lines/mm, providing a resolution of 0.1 nm, corresponding to 3 meV at the midpoint of the energy range studied. The minimum wavelength to which scans are performed, 115 nm (10.8 eV), is determined by the windows of the gas cell (LiF entrance and MgF<sub>2</sub> exit).

The sample pressure is measured using a capacitance manometer (Baratron). To ensure that the data were free of any saturation effects,<sup>15,16</sup> the cross sections were measured over the pressure range 0.07–1.27 mbar with typical attenuations <40%. A background scan is recorded with the cell evacuated. Absolute photoabsorption cross sections are then obtained using the Beer–Lambert attenuation law

$$I_t = I_0 \times \exp(-n\sigma x),$$

where  $I_t$  is the radiation intensity transmitted through the gas sample,  $I_0$  is that through the evacuated cell,  $n$  is the molecular number density of the sample gas,  $\sigma$  is the absolute photoabsorption cross section, and  $x$  is the absorption path length (25 cm). The accuracy of the cross section is estimated to be better than  $\pm 5\%$ . Only when absorption by the sample is very weak ( $I_0 \approx I_t$ ), does the uncertainty increase as a percentage of the measured cross section.




FIG. 2. He(I) photoelectron spectrum of ethyl formate, C<sub>3</sub>H<sub>6</sub>O<sub>2</sub>, in the 10.1–16.1 eV region (\*N<sub>2</sub><sup>+</sup> X<sup>2</sup>Σ<sub>g</sub><sup>+</sup>, v' = 0 produced by the He<sub>β</sub> line; \*\*N<sub>2</sub><sup>+</sup> X<sup>2</sup>Σ<sub>g</sub><sup>+</sup>, v' = 0 produced by the He<sub>α</sub> line; \*\*\*N<sub>2</sub><sup>+</sup> A<sup>2</sup>Π<sub>u</sub>, v' = 1 produced by the He<sub>α</sub> line) and assignment of the IEs.

### C. Photoelectron spectroscopy

The He(I) (21.22 eV) photoelectron spectrum of ethyl formate (Fig. 2) was measured at the Université de Liège, Belgium. The apparatus has been described in detail previously.<sup>17</sup> Briefly, it consists of a 180° hemispherical electrostatic analyzer with a mean radius of 5 cm. The analyzer is used in the constant pass energy mode. The incident photons are produced by a D.C. discharge in a two-stage differentially pumped lamp. The energy scale is calibrated using the X<sup>2</sup>Σ<sub>g</sub><sup>+</sup>, v' = 0 and A<sup>2</sup>Π<sub>u</sub>, v' = 0 peaks of N<sub>2</sub><sup>+</sup>, rounded to three decimal places.<sup>18,19</sup> The resolution of the present spectrum is 55 meV and the accuracy of the energy scale is estimated to be  $\pm 2$  meV. The photoelectron spectrum presented in this paper is the sum of 70 individual spectra. This procedure allows us to obtain a good signal-to-noise ratio while keeping the pressure in the spectrometer low ( $< 5 \times 10^{-6}$  mbar), thus minimizing the occurrence of dimers.

## III. COMPUTATIONAL METHODS

The ground state geometry, harmonic vibrational frequencies, and normal coordinates of the neutral singlet state (S<sub>0</sub>) and ionic doublet state (D<sub>0</sub>) of ethyl formate were obtained with the Gaussian 09 program<sup>20</sup> by means of second-order Møller-Plesset (MP2) calculations in association with the aug-cc-pVTZ basis set.<sup>21</sup> The ionic state was described by open-shell unrestricted calculations. The first ionization energies (IE) were computed from the energy difference between the neutral and ionic ground states. The vertical IE was calculated at the ground state geometry of the neutral compound and the adiabatic IE was evaluated using the optimized geometries of the neutral and ionic ground states. Additionally, the first and second ionization energies were calculated with the coupled-cluster singles and doubles (CCSD), and the coupled-cluster singles, doubles and perturbative triples (CCSD(T)) methods, employing the optimized geometries at the MP2/aug-cc-pVTZ level of approximation. The first IE of the *s-cis*, *trans* conformer of ethyl formate was also

calculated at the MP2, CCSD, and CCSD(T) levels using the aug-cc-pVQZ basis set in order to estimate the basis set effect. Furthermore, the zero point vibrational energy (ZPVE) corrections to the relative energies of the conformers and to the adiabatic IEs were determined from the MP2/aug-cc-pVTZ harmonic vibrational frequencies. These corrections were also applied to the values calculated with the CCSD and CCSD(T) methods. Higher IEs were obtained with the Partial Third Order (P3) propagator method using the aug-cc-pVTZ basis set and the MP2/aug-cc-pVTZ geometry. The Franck-Condon (FC) factors, associated to the first and second photoelectron bands of ethyl formate, were calculated using recursive relations and include Duschinsky rotation effects.<sup>22–24</sup> These calculations made use of the ground state geometries, harmonic frequencies, and normal coordinates calculated with the MP2/aug-cc-pVTZ method for the neutral ( $S_0$  state) and ionic ( $D_0$  state) forms of ethyl formate. Next, the vertical excitation energies, oscillator strengths ( $f_L$ , in the length gauge), and electronic radial spatial extents ( $\langle r^2 \rangle$ ) of *s-cis, trans* ethyl formate were calculated with the equation of motion coupled cluster method restricted to single and double excitations (EOM-CCSD) using the MP2/aug-cc-pVTZ geometry. Although the calculations have been made only for the more abundant conformer, the authors expect the changes in excited state energies between the conformers to be smaller than the intrinsic errors in calculations. To provide a better description of the Rydberg states, a set of diffuse functions (6s, 6p, 4d), taken from Kaufmann *et al.*<sup>25</sup> and localized on the central oxygen atom (O1), was added to the aug-cc-pVDZ basis set (named aug-cc-pVDZ + R). The electronic radial spatial extents of the states were calculated in order to help in the assignment of Rydberg states, which show comparable  $r^2$  value for a given  $nl$  configuration. EOM-CCSD calculations were performed using the MOLPRO program.<sup>26</sup>

#### IV. STRUCTURE AND PROPERTIES OF ETHYL FORMATE

In ethyl formate,  $C_3H_6O_2$ , there are two internal rotations that give rise to conformational isomerism.<sup>8</sup> The rotation around the (O=C–O) bond generates the *s-cis* and *s-trans* conformers, whereas the rotation about the O–R (R is the ethyl group) gives another two conformers: *trans* and *gauche*. From there, four stable conformations of ethyl formate can be obtained, *s-cis, trans, s-cis, gauche, s-trans, trans, and s-trans, gauche*.

It has been shown that at room temperature (RT) both in the liquid and gas phase only two conformers are present, *s-cis, trans, s-cis, gauche*,<sup>7</sup> with *s-cis, trans* being more stable.<sup>8</sup> Both configurations are shown in Fig. 3 and the relative energies of both conformers, calculated using MP2 geometries, are summarized in Table I. From the relative energies it can be seen that both conformers are present at RT with estimated populations of about 60% for *s-cis, trans* and about 40% for *s-cis, gauche*. The calculated spectroscopic properties of both conformers show many similarities. Therefore, in order to clarify the results presented here, only the results of




FIG. 3. Structure of the two most stable conformers of ethyl formate: *s-cis, trans*, and *s-cis, gauche*.

*s-cis,trans* are described in the paper, whereas for the results on *s-cis,gauche* conformer see the supplementary material.<sup>27</sup>

The *s-cis,trans* ethyl formate has  $C_s$  symmetry and its 27 normal coordinates separate in 17 vibrations of  $A'$  symmetry and 10 modes of  $A''$  symmetry. Furthermore, the calculated Hartree-Fock electron configuration of the  $\tilde{X}^1A'$  ground state is as follows: (a) core orbitals  $(1a')^2 (2a')^2 (3a')^2 (4a')^2 (5a')^2$ , (b) valence orbitals  $(6a')^2 (7a')^2 (8a')^2 (9a')^2 (10a')^2 (11a')^2 (1a'')^2 (12a'')^2 (2a'')^2 (13a'')^2 (14a'')^2 (15a'')^2 (3a'')^2 (4a'')^2 (16a'')^2$ . The highest occupied molecular orbital (HOMO,  $16a'$ ) in the neutral ground state is localized predominantly on the oxygen in-plane lone pair ( $n_o$ ). The second highest occupied molecular orbital (HOMO-1,  $4a''$ ) corresponds mostly to the bonding  $\pi(C=O)$ . The orbital  $15a'$  (HOMO-3) shows a main contribution on the oxygen (O1) in-plane lone pair. The lowest unoccupied molecular orbital is mainly of  $\pi^*$  antibonding character and it is localized on the C=O bond.

## V. RESULTS AND DISCUSSION

### A. He(I) photoelectron

The He(I) photoelectron spectrum of ethyl formate, measured over the energy range of 10.0–16.1 eV, is shown in Fig. 2. The calculated vertical and adiabatic IEs, obtained through various methods are shown in Table II for *s-cis, trans* conformer. See the supplementary material for the *s-cis, gauche* calculated IEs.<sup>27</sup> In agreement with previous work,<sup>5</sup> the IEs calculated with the MP2 method are overestimated in comparison to the experimental results. However, the lowest vertical (10.771 eV) and adiabatic (10.588 eV) IEs, observed in the photoelectron spectrum, agree reasonably well with the values obtained from calculations with the CCSD (10.903 eV and 10.491 eV, respectively) and CCSD(T) (10.988 eV and 10.605 eV, respectively). In particular, the CCSD(T) adiabatic value (10.605 eV), obtained with the large aug-cc-pVQZ basis

TABLE I. Relative energies of the two most stable conformers of ethyl formate calculated using aug-cc-pVTZ basis set without and with inclusion of zero-point vibrational energies. The Boltzmann factors (in %), calculated at RT, are given in brackets.

$\Delta E$ (kcal mol <sup>-1</sup> )	<i>s-cis, trans</i>	<i>s-cis, gauche</i>
MP2	0.0 (52)	0.05 (48)
MP2+ZPVE	0.0 (58)	0.20 (42)
CCSD+ZPVE	0.0 (65)	0.36 (35)
CCSD(T)+ZPVE	0.0 (59)	0.21 (41)



TABLE II. Calculated vertical/adiabatic ionization energies (in eV) of ethyl formate (*s-cis*, *trans*) at the MP2/aug-cc-pVTZ geometry, compared with experimental values.

Configuration	Calculated				Experimental		
	P3	MP2	CCSD	CCSD(T)	This work	Benoit and Harrison <sup>11</sup>	Sweigart and Turner <sup>12</sup>
${}^2A'$ ( $16a'^{-1}$ )	11.162	11.327/10.890	10.847/10.432	10.927/10.540	10.771/10.588	10.61	10.96/10.62
...	...	11.396/10.962 <sup>a</sup>	10.903/10.491 <sup>a</sup>	10.988/10.605 <sup>a</sup>	...	...	...
${}^2A''$ ( $4a''^{-1}$ )	11.640	12.445/11.992	11.435	11.500	11.227	...	11.28
${}^2A'$ ( $15a'^{-1}$ )	13.082	...	...	...	12.582	...	...
${}^2A''$ ( $3a''^{-1}$ )	13.273	...	...	...	13.255	...	...
${}^2A'$ ( $14a'^{-1}$ )	14.099	...	...	...	13.947	...	...
${}^2A'$ ( $13a'^{-1}$ )	14.529	...	...	...	14.535	...	...
${}^2A''$ ( $2a''^{-1}$ )	14.954	...	...	...	...	...	...
${}^2A'$ ( $12a'^{-1}$ )	16.735	...	...	...	...	...	...
${}^2A''$ ( $1a''^{-1}$ )	17.050	...	...	...	...	...	...
${}^2A'$ ( $11a'^{-1}$ )	18.124	...	...	...	...	...	...

<sup>a</sup>Ionization energies calculated with the aug-cc-pVQZ basis set using the MP2/aug-cc-pVTZ geometry.

set and including ZPVE correction, is in very good agreement with the experimental value (10.588 eV). Additionally, the calculated vertical IEs for  $4a''^{-1}$  (11.435 eV with CCSD and 11.500 eV with CCSD(T)) are in reasonable agreement with the experimental values (11.227 eV and 11.28 eV<sup>12</sup>). The higher IEs calculated with the P3 methods allowed an assignment of the bands observed in the photoelectron spectrum (Fig. 2). Similar to what was found in our previous work on methyl formate,<sup>5</sup> the P3 method has a tendency of overestimating the IEs of ethyl formate. This is particularly the case for the IEs  $4a''^{-1}$  and  $15a'^{-1}$ , which present deviation with respect to experiment as large as 0.5 eV. Finally, the experimental values obtained in our study are within reasonable agreement with the results of Benoit and Harrison<sup>11</sup> and Sweigart and Turner.<sup>12</sup>

The calculations presented in Table II show that the low energy part of the photoelectron spectrum with the resolved vibronic structure corresponds to the  $16a'^{-1}$  ionic state (Fig. 4). The observed progressions can be classified into two groups, corresponding to separations of approximately 0.180 eV and 0.090 eV (Table III). In order to determine, which vibrational transitions are observed in the spectrum, calculations of frequencies together with Franck-Condon fac-

tors were performed, and are presented in Table IV and Fig. 5. According to the calculations, the first group of the more intense peaks, separated by  $\sim 0.180$  eV, is mainly attributed to mode  $\nu_{21}$ , which designates C–O and C=O stretches. The vibration  $\nu_{16}$ , involving mostly CH<sub>2</sub> wagging, also presents a non-negligible contribution to this first group. The second group, separated by about 0.090 eV, is mostly ascribed to mode  $\nu_6$  that is O=C–O deformation with C–O stretch. In particular, the normal mode configuration may lead to Fermi resonances. Moreover, mode  $\nu_7$  provides an additional contribution to this second group of bands and involves C–O stretch and CH<sub>3</sub> rocking. Next, the band, centered at 11.227 eV, is assigned to  $4a''^{-1}$  state (see Fig. 2 and Table II). This band does not exhibit vibrational structure and overlaps with the  $16a'^{-1}$  band, although the nature of the features suggests that further modes and combinations may also contribute to the observed structure. Calculations of the FC factors for this state show that the lack of vibrational structure is due to overlapping peaks resulting from various combinations of vibrations (see Fig. 5).

## B. Valence states and transitions

The results of calculations presented in Table V allowed the assignment of absorption bands that are centered at 5.753, 7.465, and 8.292 eV to ( $\pi^*(C=O) \leftarrow n_O$ ,  $16a'$ ), ( $3s\sigma/\sigma^*(CH) \leftarrow n_O$ ,  $16a'$ ), and ( $\pi^*(C=O) \leftarrow \pi(C=O)$ ,  $4a''$ ) transitions, respectively (Figs. 6–8). Pure Rydberg transitions of high oscillator strengths in this energy range are discussed later (see




FIG. 4. He(I) photoelectron spectrum of first two ionic states of ethyl formate with vibronic excitation assignment.

TABLE III. Energy positions and vibrational analysis of features observed in the first photoelectron band ( $16a'^{-1}$ ) of ethyl formate.

Peak energy (eV)	Assignment	$\Delta E(\nu_{21})$ (eV)	$\Delta E(\nu_6)$ (eV)
10.588	Adiabatic IE	...	...
10.672	$1\nu_6$	...	0.089
10.771	$1\nu_{21}$	0.183	...
10.860	$1\nu_{21} + \nu_6$	...	0.089
10.950	$2\nu_{21}$	0.179	...
11.043	$2\nu_{21} + \nu_6$	...	0.093
11.129	$3\nu_{21}$	0.179	...

TABLE IV. Main vibrational transitions and Franck-Condon factors participating to the vibrational structure of the first photoelectron band for the *s-cis*, *trans* conformer of ethyl formate (MP2/aug-cc-pVTZ).

Vibrational state	Frequency <sup>a</sup> (cm <sup>-1</sup> )	FC factor
0-0 (adiabatic)	0	0.0742
$\nu_6$ (O=C-O deformation, C-O stretch)	596.1	0.0307
$\nu_7$ (C-O stretch, CH <sub>3</sub> rocking)	713.4	0.0163
$\nu_{16}$ (CH <sub>2</sub> wagging)	1344.6	0.0225
$\nu_{21}$ (C-O stretch, C=O stretch)	1567.5	0.0991
$\nu_{21} + \nu_6$	2163.6	0.0418
$\nu_{21} + \nu_7$	2280.9	0.02
$\nu_{21} + \nu_{16}$	2912.1	0.0286
$2\nu_{21}$	3135	0.0715
$2\nu_{21} + \nu_6$	3731	0.0306
$2\nu_{21} + \nu_7$	3848.4	0.0133
$2\nu_{21} + \nu_{16}$	4479.6	0.0196
$3\nu_{21}$	4702.5	0.0368
$3\nu_{21} + \nu_6$	5298.5	0.016
$3\nu_{21} + \nu_7$	5415.8	0.0063
$3\nu_{21} + \nu_{16}$	6047	0.0096
$4\nu_{21}$	6270	0.0152

<sup>a</sup>The MP2 vibrational frequencies were corrected by a scaling factor of 0.96.<sup>28</sup>

Sec. V C). Detailed assignment of the vibronic progressions that were found in these three bands is presented in Table VI.

The first band (Fig. 6) with a maximum absolute cross section of 0.217 Mb has been identified as the transition from the oxygen in-plane lone pair to the first  $\pi$  antibonding molecular orbital ( $\pi^*(\text{C}=\text{O}) \leftarrow n_{\text{O}}, 16a'$ ) ( $1^1A'' \leftarrow 1^1A'$ ). The calculated oscillator strength for this transition yields  $1.43 \times 10^{-3}$  and is comparable to the value obtained for methyl formate.<sup>5</sup> The associated vibronic series in this band, seen between 5.0 and 6.5 eV were attributed to CC stretching ( $\nu_{13}$ ), and assigned based on data of Maes *et al.*<sup>8</sup> The broad nature of this band suggests that further modes, not resolved here, may be involved. The feature at 5.276 eV was tentatively assigned as the origin of this band.

A second absorption band (Fig. 7), centered at 7.465 eV with a maximum cross section of 11.603 Mb, was assigned




FIG. 5. Comparison between experimental (dashed) and theoretical Franck-Condon (solid) vibrational structures of the first (upper, ion  $2A'(16a'^{-1})$ ) and second (lower, ion  $2A''(4a''^{-1})$ ) photoelectron bands of ethyl formate.

to the ( $3s\sigma/\sigma^*(\text{CH}) \leftarrow n_{\text{O}}, 16a'$ ) ( $2^1A' \leftarrow 1^1A'$ ) transition. The calculations indicate a mixed valence/Rydberg character ( $3s\sigma$ ) for this state, nevertheless the high value of calculated oscillator strength ( $3.70 \times 10^{-3}$ ) is primarily due to valence ( $\sigma^*(\text{CH})$ ) character of this molecular orbital. The vibronic structure at 7.464 eV is proposed to be due mixed Rydberg/valence character, combined with two quanta of CH<sub>3</sub> rocking ( $\nu_{12}$ ),<sup>8</sup> one quantum of combined CO stretch ( $\nu_{21}$ ), and one quantum of O=C-O deformation, coupled with C-O stretch ( $\nu_6$ ).

The third band, centered at 8.292 eV with a cross section of 21.469 Mb, was assigned to the ( $\pi^*(\text{C}=\text{O}) \leftarrow \pi(\text{C}=\text{O}), 4a''$ ) ( $3^1A' \leftarrow 1^1A'$ ) transition. The calculated oscillator strength for this transition has the highest value among all transitions identified in the spectrum, yielding 0.172, compared to methyl formate: 0.153.

Continued overlap of vibronic structures associated both with Rydberg and valence transitions is proposed for features above 7.9 eV (Figs. 7 and 8 together with Table VI). Due to a large number of bands (see Table V) in this region it was very difficult to assign transitions unambiguously.

### C. Rydberg series

The photoabsorption spectrum above 7.4 eV shows a number of structures superimposed on a diffuse absorption feature extending to the lowest ionization energy. The proposed Rydberg transitions, that are labeled in Fig. 7 and Fig. 8 and presented in Table VII and Table VIII, originate mainly from the HOMO ( $16a'$ ) and HOMO-1 ( $4a''$ ) according to computations. The positions of the peaks,  $E_n$ , have been tested using the Rydberg formula:  $E_n = E_i - R/(n - \delta)^2$ , where  $E_i$  is the ionization energy with adiabatic value of 10.588 eV,  $n$  is the principal quantum number of the Rydberg orbital of energy  $E_n$ , and  $\delta$  is the quantum defect. Quantum defects are expected to be in range of 1.0–0.9, 0.6–0.3, and <0.25 for  $ns$ ,  $np$ , and  $nd$  series, respectively.

The feature at 7.465 eV was assigned to ( $3s\sigma \leftarrow n_{\text{O}}, 16a'$ ) Rydberg transition with a quantum defect of 0.91 (Table VII) and is accompanied by vibronic structure for valence transition, discussed in Sec. V B. The higher members of this Rydberg series are proposed to extend to  $n = 6$ . State  $n = 7$  was assigned purely on the value of the quantum defect.

The first members of  $np\sigma$ ,  $np\sigma'$ , and  $np\pi$  series were found at 8.046 eV, 8.222 eV, and 8.382 eV with quantum defects of 0.69, 0.60, and 0.52, respectively (Table VII). Calculated energies for these transitions (Table V), yielding 8.333 eV, 8.524 eV, and 8.675 eV agree with experimental values reasonably well, when the valence character of these transitions is accounted for. They are also associated with vibronic structure, as shown in Fig. 7. The higher members of this Rydberg series are proposed to extend to  $n = 6$  for the  $\sigma$  series,  $n = 4$  for the  $\pi$  series, and  $n = 5$  for the  $\sigma'$  series. The feature at 8.774 eV was assigned to first transition of the  $nd\sigma$ , with  $\delta = 0.26$  and is also in a good agreement with the calculated value of 9.083 eV. For the  $n = 3$ , the  $nd\sigma$  and  $nd\sigma'$  series show vibrational excitation with one quantum of combined CO stretch ( $\nu_{21}$ ) and one quantum of O=C-O deformation, coupled with C-O stretch ( $\nu_6$ ) each. The higher members of

TABLE V. Calculated vertical excitation energies (EOM-CCSD/aug-cc-pVDZ+R) (in eV) and oscillator strengths compared with the present experimental vertical energies and VUV absorption cross sections of *s* – *cis*, *trans* ethyl formate; numbers in brackets indicate shoulder peak.

State	$E$ (eV)	$f_L$	$\langle r^2 \rangle$	Main character	$E_{exp}$ (eV)	Cross section (Mb)
$^1A''$	5.875	0.00143	69	$16 a' \rightarrow \text{LUMO}$	5.753	0.22
$^1A'$	7.755	0.00370	110	$16 a' \rightarrow 3s \sigma/\sigma^*(\text{CH})$	7.465	11.60
$^1A'$	8.129	0.17179	82	$4 a'' \rightarrow \text{LUMO}$	8.292	21.47
$^1A''$	8.221	0.00451	105	$4 a'' \rightarrow 3s \sigma/\sigma^*(\text{CH})$	8.114	20.02
$^1A'$	8.333	0.06113	119	$16 a' \rightarrow 3p \sigma/\sigma^*(\text{CH}_3)$	8.046	19.76
$^1A''$	8.524	0.00246	128	$16 a' \rightarrow 3p \pi$	8.222	21.28
$^1A'$	8.675	0.00693	137	$16 a' \rightarrow 3p \sigma'/\sigma'^*(\text{CH})$	8.383	19.72
$^1A''$	8.790	0.01273	123	$4 a'' \rightarrow 3p \sigma/\sigma^*(\text{CH}_3)$	8.707	21.45
$^1A'$	8.958	0.05008	126	$4 a'' \rightarrow 3p \pi$	8.913	25.69
$^1A'$	9.083	0.02393	148	$16 a' \rightarrow 3d \sigma/\sigma^*$	8.77(4)	23.26
$^1A''$	9.120	0.02713	141	$4 a'' \rightarrow 3p \sigma'/\sigma'^*(\text{CH})$	9.030	24.18
$^1A''$	9.149	0.00042	76	$15 a' \rightarrow \text{LUMO}$		
$^1A''$	9.326	0.00005	184	$16 a' \rightarrow 3d \pi$		
$^1A''$	9.351	0.00011	132	$4 a'' \rightarrow 3d \sigma$	9.386	18.97
$^1A'$	9.354	0.00415	224	$16 a' \rightarrow 3d \sigma'$	9.037	23.99
$^1A'$	9.382	0.00229	201	$16 a' \rightarrow 3d \sigma''$		
$^1A''$	9.458	0.00015	214	$16 a' \rightarrow 3d \pi'$		
$^1A'$	9.512	0.00628	297	$16 a' \rightarrow 4s \sigma$	9.16(3)	19.67
$^1A'$	9.639	0.01165	359	$16 a' \rightarrow 4p \sigma$	9.336	18.75
$^1A''$	9.678	0.00070	384	$16 a' \rightarrow 4p \pi$	9.407	19.62
$^1A'$	9.712	0.01146	274	$16 a' \rightarrow 4p \sigma'$	9.443	21.59
$^1A'$	9.732	0.00265	229	$4 a'' \rightarrow 3d \pi$		
$^1A''$	9.749	0.00034	197	$4 a'' \rightarrow 3d \sigma'$	9.694	30.14
$^1A'$	9.772	0.00101	196	$15 a' \rightarrow 3s \sigma$		
$^1A''$	9.819	0.00063	193	$4 a'' \rightarrow 3d \sigma''$		
$^1A'$	9.844	0.01363	400	$16 a' \rightarrow 4d \sigma$		
$^1A'$	9.895	0.00043	213	$4 a'' \rightarrow 3d \pi'$		
$^1A''$	9.968	0.00319	221	$4 a'' \rightarrow 4s \sigma$	9.793	33.05
$^1A''$	9.986	<0.00001	566	$16 a' \rightarrow 4d \pi$		
$^1A'$	9.991	0.00244	626	$16 a' \rightarrow 4d \sigma'$	9.709	30.13
$^1A''$	10.005	0.00160	269	$4 a'' \rightarrow 4p \sigma$	9.967	36.54
$^1A'$	10.006	0.00112	609	$16 a' \rightarrow 4d \sigma''$		
$^1A''$	10.036	0.00027	631	$16 a' \rightarrow 4d \pi'$		
$^1A'$	10.063	0.00279	783	$16 a' \rightarrow 5s \sigma$	9.755	30.59
$^1A''$	10.109	0.02326	281	$4 a'' \rightarrow 4p \sigma'$	10.07(1)	38.71
$^1A'$	10.113	0.00131	909	$16 a' \rightarrow 5p \sigma$	9.861	34.23
$^1A''$	10.134	0.00021	993	$16 a' \rightarrow 5p \pi$		
$^1A'$	10.135	0.01303	385	$4 a'' \rightarrow 4p \pi$	10.047	38.89
$^1A'$	10.158	0.00457	939	$16 a' \rightarrow 5p \sigma'$	9.91(1)	35.153
$^1A'$	10.188	0.01538	811	$16 a' \rightarrow 5d \sigma$		
$^1A''$	10.201	0.00001	340	$4 a'' \rightarrow 4d \sigma$		
$^1A''$	10.242	<0.00001	398	$4 a'' \rightarrow 4d \sigma'$	10.341	40.82
$^1A'$	10.258	0.01479	185	$15 a' \rightarrow 3p \sigma$		
$^1A'$	10.290	0.00325	1463	$16 a' \rightarrow 5d \sigma'$		
$^1A'$	10.342	0.00014	1347	$16 a' \rightarrow 5d \sigma''$		
$^1A''$	10.348	0.00003	939	$16 a' \rightarrow 5d \pi$		
$^1A'$	10.360	0.00296	1412	$16 a' \rightarrow 6s \sigma$	10.055	38.77
$^1A''$	10.370	0.00047	1519	$16 a' \rightarrow 5d \pi'$		
$^1A'$	10.372	0.00318	1253	$16 a' \rightarrow 6p \sigma$		
$^1A''$	10.393	0.02606	526	$15 a' \rightarrow 3p \pi$		
$^1A''$	10.413	0.00628	726	$16 a' \rightarrow 6p \pi$		
$^1A'$	10.414	0.00092	1616	$16 a' \rightarrow 6p \sigma'$		
$^1A'$	10.427	0.01369	520	$4 a'' \rightarrow 4d \pi$		
$^1A''$	10.452	0.00588	472	$4 a'' \rightarrow 4d \sigma''$		
$^1A'$	10.463	0.00467	1395	$16 a' \rightarrow 6d \sigma$		
$^1A''$	10.468	0.00005	532	$4 a'' \rightarrow 5s \sigma$	10.428	40.90
$^1A'$	10.480	0.00914	1635	$16 a' \rightarrow 6d \sigma'$		
$^1A''$	10.500	<0.00001	583	$4 a'' \rightarrow 5p \sigma'$	10.543	41.27
$^1A''$	10.538	0.00052	753	$4 a'' \rightarrow 5p \sigma$	10.480	40.99
$^1A''$	10.609	0.00370	924	$4 a'' \rightarrow 5p \pi$		




FIG. 6. Vibrational progressions in the 4.5–6.9 eV absorption band of ethyl formate,  $C_3H_6O_2$ .

these Rydberg series, of lowering relative intensity, are difficult to assign due to the overlap with other transitions and vibronic structures.

Some tentative assignment of features belonging to other Rydberg series, which converge to the ionic electronic first excited state  $\tilde{A}^2A''$  ( $4a''^{-1}$ ) was performed (Figures 7 and 8 together with Table VIII). For the high energy members, the assignment was based purely on the values of the quantum defects.

#### D. Absolute photoabsorption cross sections and atmospheric photolysis

The photoabsorption cross sections presented in this work were measured for the first time for this compound. These absolute cross sections can be used in combination with




FIG. 7. Vibrational progressions and Rydberg series assignment in the 6.5–9.3 eV absorption band of ethyl formate,  $C_3H_6O_2$ .

solar actinic flux<sup>29</sup> measurements from the literature to estimate the photolysis rate of ethyl formate in the atmosphere from an altitude close to the ground to the stratopause at 50 km. Details of the programme are presented in a previous publication.<sup>30</sup> The quantum yield for dissociation following absorption is assumed to be unity. The reciprocal of the photolysis rate at a given altitude corresponds to the local photolysis lifetime. Photolysis lifetimes of less than 72 sunlit hours (6 days) were calculated at altitudes above 20 km. This indicates that ethyl formate molecules can be broken up quite efficiently by UV absorption at these altitudes. While there is a fairly substantial photolysis rate at the higher altitudes leading to a correspondingly short lifetime of a few days or less, at lower altitudes the lack of solar actinic flux at the




FIG. 8. Vibrational progressions and Rydberg series assignment in the 9.25–10.8 eV absorption band of ethyl formate,  $C_3H_6O_2$ .



TABLE VI. Proposed vibrational assignments in the 5.0–9.2 eV absorption bands of ethyl formate, C<sub>3</sub>H<sub>6</sub>O<sub>2</sub>.

Energy (eV)	Assignment	$\Delta E(v'_0)$ (eV)	$\Delta E(v'_{12})$ (eV)	$\Delta E(v'_{13})$ (eV)	$\Delta E(v'_{21})$ (eV)
First band: $\pi^*(\text{C=O}) \leftarrow n_{\text{O}}(16a')$					
5.276	$\nu_{00}$	...	...	...	...
5.402	$1\nu_{13}$	...	...	0.126	...
5.523	$2\nu_{13}$	...	...	0.121	...
5.636	$3\nu_{13}$	...	...	0.113	...
5.753(v) <sup>a</sup>	$4\nu_{13}$	...	...	0.117	...
5.862	$5\nu_{13}$	...	...	0.109	...
5.975	$6\nu_{13}$	...	...	0.113	...
6.093	$7\nu_{13}$	...	...	0.118	...
Second band: $3s\sigma/\sigma^* \leftarrow n_{\text{O}}(16a')$					
7.464	$\nu_{00}$	...	...	...	...
7.602	$1\nu_{12}$	...	0.138	...	...
7.744	$2\nu_{12}$	...	0.142	...	...
7.815	$2\nu_{12}+1\nu_6$	0.071	...	...	...
7.920	$2\nu_{12}+1\nu_{21}$	...	...	...	0.176
Third band: $\pi^*(\text{C=O}) \leftarrow n_{\text{O}}(4a'')$					
8.046	$\nu_{00}$	...	...	...	...
8.114	$\nu_{00}+1\nu_6$	0.068	...	...	...
8.223	$1\nu_{21}$	...	...	...	0.177
8.288	$1\nu_{21}+1\nu_6$	0.065	...	...	...
8.389	$2\nu_{21}$	...	...	...	0.166
8.463	$2\nu_{21}+1\nu_6$	0.074	...	...	...
8.557	$3\nu_{21}$	...	...	...	0.168
8.640	$3\nu_{21}+1\nu_6$	0.083	...	...	...

<sup>a</sup>(v) is a vertical value.

absorption wavelengths of this molecule lead to an extremely long lifetime (extending to months).

Rate constants for gas-phase reactions of OH radicals with ethyl formate have been studied and a value of

TABLE VII. Energies (in eV), quantum defects and assignments of the ns, np, and nd Rydberg series converging to the  $\bar{X}^2A'(16a'^{-1})$  ionic electronic ground state of ethyl formate.

Vertical transition energy	Quantum defect, $\delta$	Assignment
7.464	0.91	$3s\sigma$
9.163	0.91	$4s\sigma$
9.755	0.95	$5s\sigma$
9.935	...	$5s\sigma+1\nu_{21}$
10.055	0.95	$6s\sigma$
10.221	0.91	$7s\sigma$
10.115	...	$5s\sigma+2\nu_{21}$
10.295	...	$5s\sigma+3\nu_{21}$
8.046	0.69	$3p\sigma$
9.336	0.70	$4p\sigma$
9.863	0.67	$5p\sigma$
10.113	0.65	$6p\sigma$
10.165	...	$4d\sigma+3\nu_{21}$
8.222	0.60	$3p\pi$
9.407	0.61	$4p\pi$
8.383	0.52	$3p\sigma'$
9.443	0.55	$4p\sigma'$
9.910	0.52	$5p\sigma$
8.774	0.26	$3d\sigma$
9.619	0.25	$4d\sigma$
9.967	0.32	$5d\sigma$
9.037	0.04	$3d\sigma'$
9.709	0.07	$4d\sigma'$

$(5.66 \pm 1.10) \times 10^{-13} \exp[(134 \pm 53)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  was measured in a temperature range of 233–372 K, such that these molecules may be transported further from their emission sources.<sup>31</sup> These results were confirmed later for ethyl formate oxidation in presence of chlorine atoms.<sup>32,33</sup> Calculated tropospheric lifetimes of ethyl formate reactions with OH have yielded a value of 13.6 days,<sup>31</sup> which is still shorter than the photolysis lifetimes at the altitude of 19 km (16.4 days) and below (51 days at 18 km). This means that

TABLE VIII. Energies (in eV), quantum defects, and assignments of the ns, np, and nd Rydberg series converging to the  $\bar{A}^2A''(4a''^{-1})$  ionic electronic first excited state of ethyl formate.

Vertical transition energy	Quantum defect, $\delta$	Assignment
8.114	0.91	$3s\sigma$
9.786	0.93	$4s\sigma$
10.428	0.87	$5s\sigma$
10.686	0.98	$6s\sigma$
8.707	0.68	$3p\sigma$
9.967	0.71	$4p\sigma$
10.480	0.73	$5p\sigma$
8.913	0.58	$3p\pi$
10.047	0.60	$4p\pi$
9.030	0.51	$3p\sigma'$
10.071	0.57	$4p\sigma'$
10.543	0.54	$5p\sigma'$
9.386	0.28	$3d\sigma$
10.298	0.17	$4d\sigma$
9.694	0.02	$3d\sigma'$
10.341	0.08	$4d\sigma'$
9.523	0.17	$3d\pi$
10.280	0.21	$4d\pi$

radical reactions must all be considered as the main loss processes for ethyl formate and UV photolysis is not expected to play a significant role in the tropospheric removal of these molecules.

## VI. CONCLUSIONS

In this work, we present for the first time complete electronic spectra of ethyl formate together with absolute photoabsorption cross sections from 4.5 to 10.9 eV. The structures that can be observed in the spectrum can be assigned to both valence and Rydberg transitions, based on *ab initio* calculations of vertical excitation energies and oscillator strengths of this molecule. Fine structure has been assigned to vibrational series involving predominantly excitations of  $\nu_{21}$  and  $\nu_6$  modes from C=O and C–O stretching. The high resolution He(I) photoelectron spectrum of ethyl formate has enabled vibrational excitations in the  $\tilde{X}^2A'(16a''^{-1})$  ionic electronic ground state of this molecule to be resolved and also assigned to  $\nu_{21}$  and  $\nu_6$  modes. The theoretical calculations presented here are in a good agreement with experimental data, as well as in predicting a significant mixing of Rydberg and  $\pi^*$  states. Based on the photoabsorption cross section, photolysis lifetimes of ethyl formate have been calculated for the Earth's troposphere and stratosphere.

## ACKNOWLEDGMENTS

M.A.Ś. acknowledges the visiting fellow position in the Molecular Physics group, Open University, UK. P.L.V. acknowledges his visiting Professor position at The Open University, UK and partial funding from the research Grant Nos. PESt-OE/FIS/UI0068/2014 and PTDC/FIS-ATO/1832/2012 through FCT-MEC. The authors wish to acknowledge the beam time at the ISA synchrotron at Aarhus University, Denmark, supported by the European Union (EU) I3 programme ELISA, Grant Agreement No. 226716. We also acknowledge the financial support provided by the European Commission through the Access to Research Infrastructure action of the Improving Human Potential Programme. J.G. is thankful to the 7th Framework Programme of the European Union. All calculations have been performed at the Academic Center (CI TASK) in Gdańsk and at Universität-rechenzentrum of the Friedrich-Schiller University in Jena. D.D. acknowledges support from the CaPPA project (Chemical and Physical Properties of the Atmosphere), funded by the French National Research Agency (ANR) through the PIA (Programme d'Investissement d'Avenir) under Contract No. ANR-10-LABX-005.

- <sup>1</sup>C. H. Bell, *Crop Protection* **19**, 563 (2000).
- <sup>2</sup>I. Medvedev, F. C. D. Lucia, and E. Herbst, *Astrophys. J. Suppl. Ser.* **181**, 433 (2009).
- <sup>3</sup>I. Medvedev, F. C. D. Lucia, and E. Herbst, *Astron. Astrophys.* **499**, 215 (2009).
- <sup>4</sup>P. Osswald, U. Struckmeier, T. Kasper, K. Kohse-Hoeinghaus, J. Wang, T. A. Cool, N. Hansen, and P. R. Westmoreland, *J. Phys. Chem. A* **111**, 4093 (2007).
- <sup>5</sup>Y. Nunes, G. Martins, N. J. Mason, D. Duflo, S. V. Hoffmann, J. Delwiche, M.-J. Hubin-Franskin, and P. Limão-Vieira, *Phys. Chem. Chem. Phys.* **12**, 15734 (2010).
- <sup>6</sup>J. M. Riveros and E. Bright Wilson, Jr., *J. Chem. Phys.* **46**, 4605 (1967).
- <sup>7</sup>S. W. Charles, G. I. L. Jones, N. L. Owen, S. J. Cyvin, and B. N. Cyvin, *J. Mol. Struct.* **16**, 225 (1973).
- <sup>8</sup>I. I. Maes, W. Herrebout, and J. van der Veken, *J. Raman Spectrosc.* **25**, 679 (1994).
- <sup>9</sup>M. Chhibba, F. Tristram, and G. Vergoten, *J. Mol. Struct.* **405**, 113 (1997).
- <sup>10</sup>D. van Raalte and A. G. Harris, *Can. J. Chem.* **41**, 2054 (1963).
- <sup>11</sup>F. M. Benoit and A. G. Harrison, *J. Am. Chem. Soc.* **99**, 3980 (1977).
- <sup>12</sup>D. A. Sweigart and D. W. Turner, *J. Am. Chem. Soc.* **94**, 5592 (1972).
- <sup>13</sup>M. Liu, C. Wu, Z. Wu, Q. Liang, Y. Deng, Q. Gong, Y. Sun, Y. Xu, Y. Zhao, X. Shan, F. Liu, and L. Sheng, *Chem. Phys. Lett.* **468**, 153 (2009).
- <sup>14</sup>S. Eden, P. Limão-Vieira, S. Hoffmann, and N. Mason, *Chem. Phys.* **323**, 313 (2006).
- <sup>15</sup>N. Mason, J. Gingell, J. Davies, H. Zhao, I. Walker, and M. Siggel, *J. Phys. B* **29**, 3075 (1996).
- <sup>16</sup>W. Chan, G. Cooper, and C. Brion, *Phys. Rev. A* **44**, 186 (1991).
- <sup>17</sup>J. Delwiche, P. Natalis, J. Momigny, and J. E. Collin, *J. Electron Spectrosc. Relat. Phenom.* **1**, 219 (1972).
- <sup>18</sup>K. Huber and C. Jungen, *J. Chem. Phys.* **92**, 850 (1990).
- <sup>19</sup>D. Shaw, D. Holland, M. MacDonald, A. Hopkirk, M. Hayes, and S. McSweeney, *Chem. Phys.* **166**, 379 (1992).
- <sup>20</sup>M. J. Frisch, G. W. Trucks, H. B. Schlegel *et al.*, Gaussian 09, Revision A.02, Gaussian, Inc., Wallingford, CT, 2009.
- <sup>21</sup>T. H. Dunning, Jr., *J. Chem. Phys.* **90**, 1007 (1989).
- <sup>22</sup>M. Łabuda and J. Guthmuller, *Eur. Phys. J. Spec. Top.* **222**, 2257 (2013).
- <sup>23</sup>J. Guthmuller, F. Zutterman, and B. Champagne, *J. Chem. Theory Comput.* **4**, 2094 (2008).
- <sup>24</sup>J. Guthmuller, F. Zutterman, and B. Champagne, *J. Chem. Phys.* **131**, 154302 (2009).
- <sup>25</sup>K. Kaufmann, W. Baumeister, and M. Jungen, *J. Phys. B* **22**, 2223 (1989).
- <sup>26</sup>H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby, M. Schütz *et al.*, MOLPRO, version 2010.1, a package of *ab initio* programs, 2012, see <http://www.molpro.net>.
- <sup>27</sup>See supplementary material at <http://dx.doi.org/10.1063/1.4894762> for additional information on *s-cis, gauche* conformer.
- <sup>28</sup>P. Sinha, S. E. Boesch, C. Gu, R. A. Wheeler, and A. K. Wilson, *J. Phys. Chem. A* **108**, 9213 (2004).
- <sup>29</sup>W. B. DeMore, S. P. Sander, D. M. Golden, R. F. Hampson, M. J. Kurylo, C. J. Howard, A. R. Ravishankara, C. E. Kolb, and M. J. Molina, "Chemical kinetics and photochemical data for use in stratospheric modeling," Evaluation No. 12, 15 January 1997.
- <sup>30</sup>P. Limão-Vieira, S. Eden, P. Kendall, N. Mason, and S. Hoffmann, *Chem. Phys. Lett.* **364**, 535 (2002).
- <sup>31</sup>S. LeCalve, G. LeBras, and A. Mellouki, *J. Phys. Chem. A* **101**, 5489 (1997).
- <sup>32</sup>T. J. Wallington, M. D. Hurley, and A. Haryanto, *Chem. Phys. Lett.* **432**, 57 (2006).
- <sup>33</sup>J. J. Orlando and G. S. Tyndall, *Int. J. Chem. Kinet.* **42**, 397 (2010).