

**EXISTENCE AND UNIQUENESS OF SOLUTIONS
FOR SINGLE-POPULATION
MCKENDRICK-VON FOERSTER MODELS
WITH RENEWAL**

AGNIESZKA BARTŁOMIEJCZYK, HENRYK LESZCZYŃSKI
AND PIOTR ZWIERKOWSKI

ABSTRACT. We study a McKendrick-von Foerster type equation with renewal. This model is represented by a single equation which describes one species which produces young individuals. The renewal condition is linear but takes into account some history of the population. This model addresses nonlocal interactions between individuals structured by age. The vast majority of size-structured models are also treatable. Our model generalizes a number of earlier models with delays and integrals. The existence and uniqueness is proved through a fixed-point approach to an equivalent integral problem in $L^\infty \cap L^1$.

1. Introduction. Von Foerster-McKendrick models (originated in [12]) describe populations with structure given by age [15], size [17] or level of maturation of individuals [21]. In the literature there are discrete models of that type with finite [22] or infinite matrices [28].

We consider one population with a structure given by the size of individual members or level of maturity and with the birth process expressed by a linear renewal equation. An elementary outline of such models, together with their biological interpretation, is provided in [3]. The best general reference here is the seminal work [26] and, for the case of age structure, the books [9, 31]. There are a number of existence and uniqueness proofs in literature for different versions of the McKendrick-von Foerster equations and which extend size and age-structured problems, e.g., [2, 6]. Closer to the techniques used in the paper are the papers [4, 17, 19]. In [11], a model concerning

2010 AMS *Mathematics subject classification.* Primary 35L45, 35L50, 35D05, 92D25.

Keywords and phrases. Existence, uniqueness, characteristics, renewal.

Received by the editors on October 28, 2012, and in revised form on March 24, 2013.

demographic and economics problems of ageing populations is studied. The model consists of two McKendrick type equations: for a population and for a capital stock.

The governing equation of a structured population is formulated either in the conservation law form

$$\frac{\partial u}{\partial t} + \frac{\partial(cu)}{\partial x} = \tilde{\lambda}u$$

or in the standard form of a hyperbolic equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = \lambda u.$$

These equations are closely related to each other; it suffices to put $\tilde{\lambda} = \lambda + \frac{\partial c}{\partial x}$. In our work $\tilde{\lambda}$ is denoted by W , and it is associated with the change of variables in the integral $\int u dx$. The occurrence of $\frac{\partial c}{\partial x}$ follows from the Liouville theorem. This change of variables shows the dynamics of the initial mass transport $\int \varphi \exp(\int_0^t W) dx$. The global Lipschitz condition for c and λ is not sufficient for the global existence and uniqueness, because the nonlinearity $u \lambda$ is strong, e.g.,

$$\frac{\partial u}{\partial t} + \frac{\partial u}{\partial x} = u^2$$

possesses local solutions. Global existence is due to boundedness of c and λ . This assumption is reasonable and commonly used for these terms.

We continue the sequence of results [7, 23, 24], which are focused on integral fixed-point equations, generated by the differential-functional problems. As a main tool, we construct integral fixed-point equations and a functional space, invariant with respect to these equations. The space consists of u and z describing densities and sizes, respectively. These densities are absolutely continuous in t and Lipschitz continuous in x , and the total sizes are continuous. The Banach contraction principle is applied in this functional space. The renewal condition causes serious problems for any fixed point theorem, not to mention a functional dependence. For instance, [27] deals with a simple McKendrick-von Foerster model without functionals, but the Banach fixed point theorem demands some sophisticated technicalities.

We formulate the differential problem. Let $a > 0$ and denote $E = [0, a] \times \mathbb{R}_+$ and $E_a = [-\tau, a] \times \mathbb{R}_+$, where $\mathbb{R}_+ = [0, +\infty)$. If



$t \in [0, a]$ and $z: [-\tau, a] \rightarrow \mathbb{R}_+$, then the Hale functional z_t is given by $z_t(s) = z(t+s)$ for $s \in [-\tau, 0]$, see [16]. If $u: E_a \rightarrow \mathbb{R}_+$, then we consider a natural family of Hale functionals $u_t(\cdot, x): [-\tau, 0] \rightarrow \mathbb{R}_+$ for $x \in \mathbb{R}_+$, defined by $u_t(s, x) = u(t+s, x)$ for $s \in [-\tau, 0]$ (this is the same Hale functional with the parameter x). Suppose that $c: E \times \mathcal{C}_+ \rightarrow \mathbb{R}$ and $\lambda: E \times \mathcal{C}_+ \times \mathcal{C}_+ \rightarrow \mathbb{R}$, where \mathcal{C}_+ is the positive cone of the space of continuous functions from $[-\tau, 0]$ into \mathbb{R}_+ . Let $\varphi: E_0 \rightarrow \mathbb{R}$, where $E_0 = [-\tau, 0] \times \mathbb{R}_+$. Consider the differential-functional equation

$$(1) \quad \partial_t u(t, x) + c(t, x, z_t) \partial_x u(t, x) = u(t, x) \lambda(t, x, u_t(\cdot, x), z_t)$$

with the initial condition

$$(2) \quad u(t, x) = \varphi(t, x) \quad \text{for } (t, x) \in E_0,$$

and the renewal condition

$$(3) \quad u(t, 0) = \int_0^\infty K(t, x) u_t(\cdot, x) dx \quad \text{for } t \in [0, a],$$

where $K: E \rightarrow \mathcal{C}_+^*$, \mathcal{C}_+^* is the cone of positive continuous functionals over \mathcal{C}_+ and

$$(4) \quad z(t) = \int_0^\infty u(t, x) dx \quad \text{for } t \in [-\tau, a].$$

Since $u = \varphi$ on E_0 , the well posedness of the problem requires the following consistency condition

$$\varphi(0, 0) = \int_0^\infty K(0, x) \varphi(\cdot, x) dx,$$

which is valid throughout the paper. We illustrate the functional dependence appearing in the right sides of equations (1) and (3) by several examples:

1. Classical age structured models without delays or integrals:

$$-u(t, x) \mu(t, x, u(t, x), z(t))$$

and

$$\int_0^\infty \tilde{K}(t, x) u(t, x) dx,$$

where $\mu: E \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$ and $\tilde{K}: E \rightarrow \mathbb{R}_+$, [14, 15],



2. delayed structures, where death and birth rates depend on certain past states of u and z :

$$-u(t, x) \mu(t, x, u(t - \tau, x), z(t - \tau))$$

and

$$\int_0^\infty \tilde{K}(t, x) u(t - \tau, x) dx$$

with the same functions \tilde{K} , μ , [25, 29];

3. moving averages for densities and total sizes, typical for cumulation effects in mathematical biology and medicine:

$$-u(t, x) \mu \left(t, x, \frac{1}{\tau} \int_{t-\tau}^t u(s, x) ds, \frac{1}{\tau} \int_{t-\tau}^t z(s) ds \right)$$

and

$$\int_0^\infty \tilde{K}(t, x) \frac{1}{\tau} \int_{t-\tau}^t u(s, x) ds dx$$

with the same functions \tilde{K} , μ , [8],

4. the size structured model:

$$-u(t, x) [\mu(x, z(t)) + \partial_x \gamma(x, z(t))],$$

where μ and γ denote the mortality and growth rates of individuals, respectively, and $K \equiv 0$, [10].

In the literature one can observe various differential-functional models: classical arguments [14, 15], delays [5, 13, 20], integrals [8], Hale functionals z_t [16, 18], mixed-types (e.g., z_t and its multidimensional generalization [32]). We have chosen a unified approach to both functional arguments by means of one-dimensional Hale functional, applied to u and z . Then the description becomes simple but sufficiently general. This approach, despite its simplicity, is surprisingly feasible in mathematical biology. It is worth mentioning that the results of our paper are new even for classical arguments. Our paper is the first work in which there are such unified Hale functionals for u and z . One can raise the question whether anything is missing when solutions u , z are considered in the subclass of continuous functions instead the whole space $L^\infty \cap L^1$, while it is more natural to consider integrable densities u in the biological modeling. The main reason of this restriction lies in the method of the proof where the class of continuous functions seems



to be unavoidable. On the other hand, having proved existence results for continuous functions, we can consider problem (1)–(4) with the initial function of the class L^1 . This initial function is approximated by continuous functions of the class L^1 . This gives a sequence of approximate solutions which converges weakly to the unique solution of our problem. Since our paper is very extensive and technical, we do not intend to provide any details of this corollary.

The aim of this paper is to look for Carathéodory's solutions to (1)–(4), i.e., continuous functions $u: E_a \rightarrow \mathbb{R}_+$ which satisfy (1) almost everywhere on E , their derivatives $\partial_t u$, $\partial_x u$ exist almost everywhere on E , and conditions (2), (4) hold. Condition (3) can be regarded as a definition of $z(t)$ by means of $u(t, \cdot)$. In the present paper we understand that the solutions to problem (1)–(4) consist of pairs (u, z) such that u is the Carathéodory solution and z is given by (4). We recall its biological interpretation: (u, z) means the density and total size of the population. We focus on equivalent integral equations.

The paper is organized as follows. In Section 2, we introduce bicharacteristics of the hyperbolic equation and give their basic properties. We also formulate main assumptions and define the space of admissible functions, where the solution of problem (1)–(4) will be found. In Section 3, we prove the main existence and uniqueness theorem by virtue of the Banach contraction principle. The space of admissible functions is mapped into the same space. The integral operator is a contraction with respect to a Bielecki type norm. Because of the number of technical details the proofs of auxiliary lemmas are collected in the last section.

2. Preliminaries. We start with the formulation of characteristic equations and an analysis of u along these characteristics. For a given continuous function $z: [-\tau, a] \rightarrow \mathbb{R}_+$, consider the characteristic equations for problem (1)–(2):

$$(5) \quad \frac{d}{ds} \eta(s) = c(s, \eta(s), z_s), \quad \eta(t) = x,$$

where $(t, x) \in E$. Let $\eta(\cdot) = \eta[z](\cdot; t, x)$ be the characteristic curve passing through the point $(t, x) \in E$, i.e., the solution of (5) in the Carathéodory sense, cf., [30]. We denote the maximal existence interval of $\eta[z](\cdot; t, x)$ by $[\alpha, a]$, where $\alpha = \alpha[z](t, x)$. It is clear that



either $\alpha = 0$ or $\alpha \in (0, a]$. If $\alpha = 0$, then the characteristic curve starts from $(0, \eta(0))$. If $\alpha > 0$, then it starts from $(\alpha, 0)$. Equation (1) along a characteristic $\eta(\cdot) = \eta[z](\cdot; t, x)$ is rewritten in the form

$$(6) \quad \frac{d}{ds} u(s, \eta(s)) = u(s, \eta(s)) \lambda(s, \eta(s), u_s(\cdot, \eta(s)), z_s)$$

with the initial and boundary condition

$$u(\alpha, \eta(\alpha)) = \begin{cases} \varphi(\alpha, \eta(\alpha)) & \text{for } \alpha = 0, \\ \int_0^\infty K(\alpha, y) u_t(\alpha, y) dy & \text{for } \alpha > 0. \end{cases}$$

If $\alpha = \alpha[z](t, x) = 0$, then equation (6) is accompanied by the initial condition (2). If $\alpha = \alpha[z](t, x) > 0$, then (6) is equipped with the boundary condition (3). Denote by η_0 the characteristic which starts from $(0, 0)$, i.e., $\eta_0(t) = \eta[z](t; 0, 0)$.

Remark 2.1. The differential equation (5) leads to the integral equation

$$(7) \quad \eta[z](s; t, x) = x - \int_s^t c(\zeta, \eta[z](\zeta; t, x), z_\zeta) d\zeta.$$

We denote by $C_b(X, Y)$ the space of all continuous and bounded functions. By $L^1(X, Y)$, we understand the space of all integrable functions with a natural L^1 -norm, denoted by $\|\cdot\|_1$. The symbol $\|\cdot\|$ stands for any supremum norm. We need the following assumptions.

Assumption $[\varphi]$. Suppose that:

1. $\varphi \in C_b(E_0, \mathbb{R}_+)$, $\varphi(t, \cdot) \in L^1(\mathbb{R}_+, \mathbb{R}_+)$ for $t \in [-\tau, 0]$, and the function $[-\tau, 0] \ni t \mapsto \int_0^\infty \varphi(t, x) dx$ is continuous and

$$\|\varphi\|_{\infty, 1} := \int_0^\infty \sup_{t' \in [-\tau, 0]} \varphi(t', x) dx < \infty.$$

2. There is a constant $L_\varphi > 0$ such that:

$$|\varphi(t, \bar{x}) - \varphi(t, x)| \leq L_\varphi |\bar{x} - x| \quad \text{on } E_0.$$

Assumption $[c]$. Suppose that:



1. the function $c: E \times \mathcal{C}_+ \rightarrow \mathbb{R}$ is bounded and measurable in $t \in [0, a]$ for every $(x, q) \in \mathbb{R}_+ \times \mathcal{C}_+$,
2. $|c(t, \bar{x}, \bar{q}) - c(t, x, q)| \leq L_c(t)(|\bar{x} - x| + \|\bar{q} - q\|)$ on $E \times \mathcal{C}_+$, where $L_c \in L^1([0, a], \mathbb{R}_+)$,
3. the function c satisfies the estimates

$$\widehat{c}(t) \geq \|c(t, x, q)\| \quad \text{on } E \times \mathcal{C}_+$$

and

$$c(t, x, q) \geq \varepsilon_0 \widehat{c}(t) \quad \text{for } 0 \leq t \leq a,$$

$$0 \leq x \leq \int_0^t \widehat{c}(s) ds,$$

with some $\varepsilon_0 \in (0, 1)$ and $\widehat{c} \in L^1([0, a], \mathbb{R}_+)$.

Concerning Assumption [c, 3], notice that the function c must be strictly positive near the lateral boundary, in reach of characteristic curves which start from this boundary. The condition $c \geq 0$ is natural in mathematical biology, when c describes ageing or maturation, see examples in [15, 21]. In size-structured population models, the renewal condition may occur not only at 0, but also at other points. All newborns have the same age 0 but not the same size. Our general model refers to the classical Kermack-McKendrick-von Foerster model with $c = \text{const} > 0$, [14]. Assumption [c, 3] is not satisfied in the Lasota model ([21]), where $c(t, x, q) = x$, hence $c(t, 0, q) = 0$. Our results can be generalized to the case of the nonlocal and nonlinear renewal condition

$$c(t, 0, z_t)u(t, 0) = \int K(t, x, z_t)u_t(\cdot, x) dx,$$

see [4]. Under our assumption $c(t, 0, q) > 0$, the coefficient $c(t, 0, z_t)$ can be incorporated in K ; hence, we can write it as follows

$$u(t, 0) = \int K(t, x, z_t)u_t(\cdot, x) dx.$$

The results of our paper carry over to equation (1) with such renewal conditions. Due to the large number of details we omit the generalization.

Assumption [λ]. Suppose that the function $\lambda: E \times \mathcal{C}_+ \times \mathcal{C}_+ \rightarrow \mathbb{R}$ satisfies the conditions:



1. λ is Lebesgue integrable in $t \in [0, a]$ for every $(x, w, q) \in \mathbb{R}_+ \times \mathcal{C}_+ \times \mathcal{C}_+$,
2. there is $L_\lambda \in L^1([0, a], \mathbb{R}_+)$ such that

$$|\lambda(t, \bar{x}, \bar{w}, \bar{q}) - \lambda(t, x, w, q)| \leq L_\lambda(t)(|\bar{x} - x| + \|\bar{w} - w\| + \|\bar{q} - q\|)$$

for $t \in [0, a]$, $x, \bar{x} \in \mathbb{R}_+$, $w, \bar{w}, q, \bar{q} \in \mathcal{C}_+$,

3. there exists $M_\lambda \in L^1([0, a], \mathbb{R}_+)$ such that

$$|\lambda(t, x, w, q)| \leq M_\lambda(t) \quad \text{for } (t, x) \in E, \quad w, q \in \mathcal{C}_+.$$

Denote

$$(8) \quad W(t, x, w, q) = \lambda(t, x, w, q) + \partial_x c(t, x, q)$$

for $(t, x) \in E$, $w, q \in \mathcal{C}_+$.

Assumption [W]. Suppose that $W: E \times \mathcal{C}_+ \times \mathcal{C}_+ \rightarrow \mathbb{R}$ satisfies the conditions:

1. W is Lebesgue integrable in $t \in [0, a]$ for every $(x, w, q) \in \mathbb{R}_+ \times \mathcal{C}_+ \times \mathcal{C}_+$,
2. there exists $L_W \in L^1([0, a], \mathbb{R}_+)$ such that

$$|W(t, \bar{x}, \bar{w}, \bar{q}) - W(t, x, w, q)| \leq L_W(t)(|\bar{x} - x| + \|\bar{w} - w\| + \|\bar{q} - q\|)$$

for $t \in [0, a]$, $x, \bar{x} \in \mathbb{R}_+$, $w, \bar{w}, q, \bar{q} \in \mathcal{C}_+$,

3. there is $M_W \in L^1([0, a], \mathbb{R}_+)$ such that

$$|W(t, x, w, q)| \leq M_W(t) \quad \text{for } (t, x) \in E, \quad w, q \in \mathcal{C}_+.$$

Assumption [K]. Suppose that:

- (1) $\|K(t, x)\|_{\mathcal{C}_+^*} \leq \kappa$ on E for some constant $\kappa \in \mathbb{R}_+$, where $\|\cdot\|_{\mathcal{C}_+^*}$ is the standard functional norm,
- (2) K is absolutely continuous on E in the following sense:

$$\|K(\bar{t}, \bar{x}) - K(t, x)\|_{\mathcal{C}_+^*} \leq L_K \left[\left| \int_t^{\bar{t}} \widehat{c}(s) ds \right| + |\bar{x} - x| \right]$$

with some $L_K \in \mathbb{R}_+$ and the function \widehat{c} from Assumption [c].

Assumption [L, M]. The functions L_c/\widehat{c} , L_λ/\widehat{c} , L_W/\widehat{c} , M_λ/\widehat{c} , M_W/\widehat{c} are bounded on $[0, a]$.



Remark 2.2. The most convenient realization of Assumption $[L, M]$ can be achieved by the substitution $L_c = \text{const.} \widehat{c}$, $L_\lambda = \text{const.} \widehat{c}$, $L_W = \text{const.} \widehat{c}$, $M_\lambda = \text{const.} \widehat{c}$, $M_W = \text{const.} \widehat{c}$. In particular, one can put $L_c = L_\lambda = L_W = M_\lambda = M_W = \widehat{c}$.

3. Main results. Now we are ready to define a space \mathcal{X} of admissible functions in terms of constants from the previous assumptions, where a priori estimates of solutions to (1)–(4) are fulfilled.

Definition 3.1. We say that a pair (u, z) belongs to \mathcal{X} if and only if $u: E_a \rightarrow \mathbb{R}_+$ and $z: [-\tau, a] \rightarrow \mathbb{R}_+$ are continuous and

1. (u, z) satisfies conditions (2) and (4),
2. $u(t, 0) \leq \kappa Z(t)$, $z(t) \leq Z(t)$, $u(t, x) \leq U(t)$ for $t \in [0, a]$ and $x \in \mathbb{R}_+$, where

$$Z(t) = \|\varphi\|_{\infty, 1} \exp \left(\int_0^t [\kappa \widehat{c}(s) + M_W(s)] ds \right),$$

$$U(t) = \exp \left(\int_0^t M_\lambda(s) ds \right) \max \{ \|\varphi\|, \kappa Z(t) \},$$

3. $|u(\bar{t}, 0) - u(t, 0)| \leq G_u \int_t^{\bar{t}} \widehat{c}(s) ds$ for $0 \leq t \leq \bar{t} \leq a$, where

$$G_u = \left\{ \kappa^2 Z(a) + \left[2L_K + \kappa^2 Z(a) \left\| \frac{M_W}{\widehat{c}} \right\| \right] \times \left[\int_0^a \widehat{c}(s) ds + \|\varphi\|_{\infty, 1} \right] \right\} \exp \left(\int_0^a M_W(s) ds \right),$$

4. $|u(t, \bar{x}) - u(t, x)| \leq L_u(t) |\bar{t} - x|$ for $t \in [0, a]$ and $x, \bar{t} \in \mathbb{R}_+$, where L_u is given by

$$L_u(t) = -1 + (1 + L_u(0))$$

$$\times \exp \left\{ [\kappa Z(a) + \|\varphi\|] \int_0^t L_\lambda(s) ds \right.$$

$$\left. \times \exp \left(\int_0^a (L_c(s) + M_\lambda(s)) ds \right) \right\},$$



where

$$L_u(0) = \left\{ L_\varphi + \frac{G_u + \kappa Z(a) \|M_\lambda / \hat{c}\|}{\varepsilon_0} \right\} \exp \left(\int_0^a (L_c(s) + M_\lambda(s)) ds \right).$$

Remark 3.2. If $u: E_a \rightarrow \mathbb{R}_+$ is a bounded, continuous and integrable function, then the corresponding function $z: [-\tau, a] \rightarrow \mathbb{R}_+$ is measurable and bounded. However, for $(u, z) \in \mathcal{X}$, the function z has an enhanced regularity, so that it becomes absolutely continuous on $[0, a]$. In fact, the function z is as regular as $u(\cdot, 0)$ on $[0, a]$.

Let us formulate the main result of this paper.

Theorem 3.3. *Suppose that Assumptions $[\varphi]$, $[c]$, $[\lambda]$, $[W]$, $[K]$ and $[L, M]$ are satisfied. Then there exists exactly one solution (u, z) of problem (1)–(4) in the class \mathcal{X} .*

Our main existence theorem will be proved by means of the Banach contraction principle in the space \mathcal{X} . Suppose that $(u, z) \in \mathcal{X}$. We construct a new pair of functions (\tilde{u}, \tilde{z}) via the renewal condition (3) as follows. Suppose that $\eta = \eta[z](\cdot; t, x)$ is a characteristic determined by the Cauchy problem (5). Denote

$$P_{t,x}^{u,z}(s) := (s, \eta[z](s; t, x), u_s(\cdot, \eta[z](s; t, x)), z_s).$$

Let $\tilde{u}(t, 0)$ for $t \in [0, a]$ be the solution of the following Volterra integral equation

$$\begin{aligned} \tilde{u}(t, 0) &= \int_0^t K(t, \eta[z](t; \xi, 0)) \tilde{u}_\xi(\cdot, 0) c(\xi, 0, z_\xi) \\ &\times \exp \left(\int_\xi^t W \left(P_{\xi,0}^{u,z}(s) \right) ds \right) d\xi \\ (9) \quad &+ \int_0^\infty K(t, \eta[z](t; 0, y)) \varphi(\cdot, y) \\ &\times \exp \left(\int_0^t W \left(P_{0,y}^{u,z}(s) \right) ds \right) dy, \end{aligned}$$

where W is defined by (8). The explanation of the changes of variables $x \mapsto \xi$ and $x \mapsto y$ is given in Remark 4.2 at the very end of the paper. Because the Volterra integral equation (9) provides a natural boundary



condition, the function \tilde{u} on the whole set E will be the only solution of the PDE:

$$\partial_t \tilde{u}(t, x) + c(t, x, z_t) \partial_x \tilde{u}(t, x) = \tilde{u}(t, x) \lambda(t, x, u_t(\cdot, x), z_t)$$

with the initial condition $\tilde{u} = \varphi$ on E_0 . Considering this problem along the characteristics satisfying (5), we get its solution by the explicit formula

$$(10) \quad \tilde{u}(t, x) = \tilde{u}(\alpha, \eta[z](\alpha; t, x)) \exp\left(\int_{\alpha}^t \lambda(P_{t,x}^{u,z}(s)) ds\right)$$

for $(t, x) \in E$, where $\alpha = \alpha[z](t, x)$. According to (4), we have

$$\tilde{z}(t) = \int_0^{\infty} \tilde{u}(t, x) dx \quad \text{for } t \in [-\tau, a].$$

This way we have constructed an integral operator \mathcal{T} which maps a pair of functions (u, z) to a pair $(\tilde{u}, \tilde{z}) = \mathcal{T}(u, z)$. By virtue of the Banach contraction principle we show that the operator \mathcal{T} has exactly one fixed point $(u, z) \in \mathcal{X}$. This fixed point satisfies the differential-functional problem (1)–(4). This goal will be achieved in three auxiliary lemmas:

- (1) If $(u, z) \in \mathcal{X}$, then $\mathcal{T}(u, z)$ satisfies conditions 1–2 of Definition 3.1,
- (2) $\mathcal{T}(u, z)$ satisfies conditions 3–4 of Definition 3.1,
- (3) the operator $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ is a contraction.

Because of multitudes of technical details we relegate the proofs of these lemmas to the next section.

Lemma 3.4. *Suppose that Assumptions [c, 3], [λ , 3], [W, 3] and [K, 1] are satisfied. If $(u, z) \in \mathcal{X}$, then \tilde{u} is bounded and continuous, the pair $(\tilde{u}, \tilde{z}) = \mathcal{T}(u, z)$ satisfies condition (4), and the following estimates hold true:*

$$\begin{aligned} 0 &\leq \tilde{u}(t, 0) \leq \kappa Z(t), \\ \tilde{z}(t) &\leq Z(t), \\ \tilde{u}(t, x) &\leq U(t) \quad \text{for } (t, x) \in E. \end{aligned}$$

Lemma 3.5. *Suppose that Assumptions [φ], [c], [λ], [W], [K] and [L, M] are satisfied. If $(u, z) \in \mathcal{X}$, then the pair $(\tilde{u}, \tilde{z}) = \mathcal{T}(u, z)$*



satisfies the conditions

$$|\tilde{u}(\bar{t}, 0) - \tilde{u}(t, 0)| \leq G_u \int_t^{\bar{t}} \hat{c}(s) ds \quad \text{for } 0 \leq t \leq \bar{t} \leq a,$$

$$|\tilde{u}(t, \bar{x}) - \tilde{u}(t, x)| \leq L_u(t) |\bar{x} - x| \quad \text{for } t \in [0, a], \bar{x}, x \in \mathbb{R}_+,$$

where G_u and $L_u(t)$ are the same as in Definition 3.1, 3–4.

Definition 3.6. The Bielecki norm is given by

$$\|(u, z)\|_B = \max\{\|u/B\|, \|z/B\|\},$$

where $B = B(t)$, and $B: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a positive, continuous, nondecreasing function. The meaning of the supremum norms $\|u/B\|$ and $\|z/B\|$ is obvious, see [1].

Lemma 3.7. *Suppose that Assumptions $[\varphi]$, $[c]$, $[\lambda]$, $[W]$, $[K]$ and $[L, M]$ are satisfied. Then the operator $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ is a contraction with respect to a Bielecki norm $\|\cdot\|_B$ for some $B: \mathbb{R}_+ \rightarrow \mathbb{R}_+$, that is: there is a $\Theta \in (0, 1)$ such that*

$$\|\mathcal{T}(\bar{u}, \bar{z}) - \mathcal{T}(u, z)\|_B \leq \Theta \|(\bar{u}, \bar{z}) - (u, z)\|_B \quad \text{on } \mathcal{X}.$$

In fact, for any $\Theta \in (0, 1)$, we can find a function B of the form $B(t) = \exp(C \int_0^t \hat{c}(s) ds)$ such that the above contraction inequality holds true.

Corollary 3.8. *If the functions λ , c and $\partial_x c$ are bounded and continuous; λ is Lipschitz continuous in x, p, q ; c and $\partial_x c$ are Lipschitz continuous in x, q ; K is nonnegative and Lipschitz continuous; c is nonnegative on E ; $c(t, x, q) \geq \varepsilon_1 > 0$ for all $t \in [0, a]$ and $x \in [0, tx_0/a]$ with some $x_0 > 0$; φ is Lipschitz continuous and satisfies Assumption $[\varphi]$, then there is a solution to (1)–(4) which is Lipschitz continuous with respect to both variables t, x .*

Proof. All assumptions of Theorem 3.3 are satisfied with the functions $M_\lambda, M_W, L_\lambda, L_W, \hat{c}, L_c$ that are constant. Observe that $\varepsilon_0 = \varepsilon_1/\hat{c}$, that is, Assumption $[c, 3]$ is also satisfied. Since the functions mentioned above are constant, the functions $\alpha[z], \eta[z]$ and $u(\cdot, 0)$ inherit the Lipschitz continuity with respect to t . Therefore, using (10), we obtain that u is Lipschitz continuous. □



Corollary 3.9. *Suppose that the assumptions of Corollary 3.8 are satisfied. If φ , c and K are C^1 functions, then the solution of (1)–(4) is a C^1 function in the whole domain E , except the characteristic curve which starts from $(0, 0)$.*

Proof. Since the initial function φ is of the class C^1 and $(\tilde{u}, \tilde{z}) = \mathcal{T}(u, z) \in \mathcal{X}$, it follows from (10) that \tilde{u} is of the class C^1 for (t, x) such that $x > \eta_0(t)$, $t \in [0, a]$. Due to the Volterra integral equation (9), C^1 -regularity of K results in the same regularity of \tilde{u} for $x < \eta_0(t)$, $t \in [0, a]$, provided that $\alpha[z]$ is C^1 , which follows from the smoothness property of the function c . \square

4. Proofs of lemmas.

Proof of Lemma 3.4. Let $t \in [0, a]$. It follows from the Volterra equation (9) that

$$\begin{aligned} \tilde{u}(t, 0) &\leq \kappa \int_0^t \hat{c}(\xi) \|\tilde{u}_\xi(\cdot, 0)\| \exp\left(\int_\xi^t M_W(s) ds\right) d\xi \\ &\quad + \kappa \|\varphi\|_{\infty, 1} \exp\left(\int_0^t M_W(s) ds\right). \end{aligned}$$

Since the right-hand side is increasing with respect to t , the left-hand side can be replaced by $\|\tilde{u}_t(\cdot, 0)\|$. Applying the Gronwall lemma, we get the inequality

$$\|\tilde{u}_t(\cdot, 0)\| \exp\left(-\int_0^t M_W(s) ds\right) \leq \kappa \|\varphi\|_{\infty, 1} \exp\left(\int_0^t \kappa \hat{c}(s) ds\right).$$

Therefore, we have $\tilde{u}(t, 0) \leq \kappa Z(t)$, where $Z(t)$ is described in Definition 3.1 2. Since \tilde{z} has the estimate

$$\begin{aligned} \tilde{z}(t) &\leq \int_0^t \kappa Z(\xi) \hat{c}(\xi) \\ &\quad \times \exp\left(\int_\xi^t M_W(s) ds\right) d\xi + \|\varphi\|_{\infty, 1} \exp\left(\int_0^t M_W(s) ds\right), \end{aligned}$$

it is easy to observe that $\tilde{z}(t) \leq Z(t)$.

Now we show the estimate for $\tilde{u}(t, x)$. From (10), we deduce that



1) if $x \geq \eta_0(t)$, then

$$\begin{aligned}\tilde{u}(t, x) &= \varphi(0, \eta[z](0; t, x)) \exp\left(\int_0^t \lambda(P_{t,x}^{u,z}(s)) ds\right) \\ &\leq \|\varphi\| \exp\left(\int_0^t M_\lambda(s) ds\right);\end{aligned}$$

2) if $x \leq \eta_0(t)$, then

$$\begin{aligned}\tilde{u}(t, x) &= u(\alpha, 0) \exp\left(\int_\alpha^t \lambda(P_{t,x}^{u,z}(s)) ds\right) \\ &\leq \kappa Z(\alpha) \exp\left(\int_\alpha^t M_\lambda(s) ds\right) \\ &= \kappa \|\varphi\|_{\infty,1} \exp\left(\int_0^\alpha [\kappa \hat{c}(s) + M_W(s)] ds\right) \\ &\quad \times \exp\left(\int_\alpha^t M_\lambda(s) ds\right) \\ &\leq \kappa \|\varphi\|_{\infty,1} \exp\left(\int_0^t \max\{\kappa \hat{c}(s) + M_W(s), M_\lambda(s)\} ds\right),\end{aligned}$$

where $\alpha = \alpha[z](t, x)$. Both estimates in cases 1) and 2) can be unified as follows

$$\begin{aligned}\tilde{u}(t, x) &\leq \max\{\|\varphi\|, \kappa \|\varphi\|_{\infty,1}\} \\ &\quad \times \exp\left(\int_0^t \max\{\kappa \hat{c}(s) + M_W(s), M_\lambda(s)\} ds\right),\end{aligned}$$

hence $\tilde{u}(t, x) \leq U(t)$ on E . Regularity assertions and condition (4) for \tilde{u} , \tilde{z} are trivial. This completes the proof. \square

Proof of Lemma 3.5. Let $(u, z) \in \mathcal{X}$. In order to demonstrate the Lipschitz condition of $\tilde{u}(t, \cdot)$ and the absolute continuity of $\tilde{u}(\cdot, 0)$, we analyze properties of $\eta[z]$ and $\alpha[z]$. Take arbitrary $(t, x) \in E$ and $\bar{x} \in \mathbb{R}_+$.

Step 1. Estimate of increments of η for \bar{x} and x . We derive from



(7) the integral inequality

$$|\eta[z](s; t, \bar{x}) - \eta[z](s; t, x)| \leq |\bar{x} - x| + \left| \int_s^t L_c(\zeta) |\eta[z](\zeta; t, \bar{x}) - \eta[z](\zeta; t, x)| d\zeta \right|.$$

Applying the Gronwall lemma, we get

$$|\eta[z](s; t, \bar{x}) - \eta[z](s; t, x)| \leq |\bar{x} - x| \exp \left(\left| \int_s^t L_c(\zeta) d\zeta \right| \right).$$

Step 2. Estimate of increments of η for \bar{t} and t . From (7), we get the inequality

$$|\eta[z](s; \bar{t}, x) - \eta[z](s; t, x)| \leq \int_t^{\bar{t}} \widehat{c}(\zeta) d\zeta + \left| \int_s^t L_c(\zeta) |\eta[z](\zeta; \bar{t}, x) - \eta[z](\zeta; t, x)| d\zeta \right|$$

for $0 \leq t \leq \bar{t} \leq a$. Applying the Gronwall lemma, we get

$$|\eta[z](s; \bar{t}, x) - \eta[z](s; t, x)| \leq \int_t^{\bar{t}} \widehat{c}(\zeta) d\zeta \exp \left(\left| \int_s^t L_c(\zeta) d\zeta \right| \right).$$

Similarly, we have

$$\begin{aligned} |\eta[z](\bar{s}; t, x) - \eta[z](s; t, x)| &= \left| \int_s^{\bar{s}} c(\zeta, \eta[z](\zeta; t, x), z_\zeta) d\zeta \right| \\ &\leq \left| \int_s^{\bar{s}} \widehat{c}(\zeta) d\zeta \right|. \end{aligned}$$

Step 3. Estimate of some integrals. By the definition of $\alpha = \alpha[z](t, x)$, we have the integral identity

$$0 = x - \int_\alpha^t c(\zeta, \eta[z](\zeta; t, x), z_\zeta) d\zeta \quad \text{for } \alpha > 0.$$

Denote $\bar{\alpha} = \alpha[z](t, \bar{x})$. Suppose that $\alpha \leq \bar{\alpha}$. Then we have

$$0 = \bar{x} - \int_{\bar{\alpha}}^t c(\zeta, \eta[z](\zeta; t, \bar{x}), z_\zeta) d\zeta.$$



If we subtract these identities, then

$$\begin{aligned} & \left| \int_{\alpha}^{\bar{\alpha}} c(\zeta, \eta[z](\zeta; t, \bar{x}), z_{\zeta}) d\zeta \right| \\ & \leq |\bar{x} - x| + \int_{\alpha}^t L_c(\zeta) |\eta[z](\zeta; t, \bar{x}) - \eta[z](\zeta; t, x)| d\zeta. \end{aligned}$$

Applying Assumption [c, 2, 3] and Step 1 we get

$$\varepsilon_0 \int_{\alpha}^{\bar{\alpha}} \widehat{c}(\zeta) d\zeta \leq |\bar{x} - x| + \int_{\alpha}^t L_c(\zeta) |\bar{x} - x| \exp\left(\int_{\zeta}^t L_c(s) ds\right) d\zeta.$$

Consequently, we have

$$\varepsilon_0 \int_{\alpha}^{\bar{\alpha}} \widehat{c}(\zeta) d\zeta \leq |\bar{x} - x| \exp\left(\int_{\alpha}^t L_c(\zeta) d\zeta\right).$$

Step 4. Estimate of increments of λ and W along characteristics. Since $u(s, \cdot)$ is Lipschitz continuous, we get

$$\begin{aligned} & |u(s, \eta[z](s; t, \bar{x})) - u(s, \eta[z](s; t, x))| \\ & \leq L_u(s) |\eta[z](s; t, \bar{x}) - \eta[z](s; t, x)| \\ & \leq L_u(s) |\bar{x} - x| \exp\left(\int_s^t L_c(\zeta) d\zeta\right) \end{aligned}$$

for $0 \leq s \leq t$. This inequality is applied to the estimates of increments of λ and W ; in particular, we have

$$\begin{aligned} & \left| \lambda(P_{t, \bar{x}}^{u, z}(s)) - \lambda(P_{t, x}^{u, z}(s)) \right| \\ & \leq L_{\lambda}(s) |\eta[z](s; t, \bar{x}) - \eta[z](s; t, x)| \\ & \quad + L_{\lambda}(s) |u(s, \eta[z](s; t, \bar{x})) - u(s, \eta[z](s; t, x))| \\ & \leq L_{\lambda}(s) |\bar{x} - x| (1 + L_u(s)) \exp\left(\int_s^t L_c(\zeta) d\zeta\right) \end{aligned}$$

for $0 \leq s \leq t$. The estimate of $|W(P_{t, \bar{x}}^{u, z}(s)) - W(P_{t, x}^{u, z}(s))|$ is similar.

Step 5. Estimate of $\tilde{u}(\bar{t}, 0) - \tilde{u}(t, 0)$. Take $t \leq \bar{t}$. From (9), we obtain the inequality

$$|\tilde{u}(\bar{t}, 0) - \tilde{u}(t, 0)| \leq \int_t^{\bar{t}} \|K(\bar{t}, \eta[z](\bar{t}; \xi, 0))\|_{C^*_+} \|\tilde{u}_{\xi}(\cdot, 0)\| c(\xi, 0, z_{\xi})$$



$$\begin{aligned}
& \times \exp \left(\int_{\xi}^{\bar{t}} W \left(P_{\xi,0}^{u,z}(s) \right) ds \right) d\xi \\
& + \int_0^t \|K(\bar{t}, \eta[z](\bar{t}; \xi, 0)) - K(t, \eta[z](t; \xi, 0))\|_{C_+^*} \\
& \quad \times \|\tilde{u}_{\xi}(\cdot, 0)\| c(\xi, 0, z_{\xi}) \\
& \times \exp \left(\int_{\xi}^{\bar{t}} W \left(P_{\xi,0}^{u,z}(s) \right) ds \right) d\xi \\
& + \int_0^t \|K(t, \eta[z](t; \xi, 0))\|_{C_+^*} \|\tilde{u}_{\xi}(\cdot, 0)\| c(\xi, 0, z_{\xi}) \\
& \times \left| \exp \left(\int_{\xi}^{\bar{t}} W \left(P_{\xi,0}^{u,z}(s) \right) ds \right) \right. \\
& \quad \left. - \exp \left(\int_{\xi}^t W \left(P_{\xi,0}^{u,z}(s) \right) ds \right) \right| d\xi \\
& + \int_0^{\infty} \left\| K(\bar{t}, \eta[z](\bar{t}; 0, y)) \right. \\
& \quad \left. - K(t, \eta[z](t; 0, y)) \right\|_{C_+^*} \|\varphi(\cdot, y)\| \\
& \times \exp \left(\int_0^{\bar{t}} W \left(P_{0,y}^{u,z}(s) \right) ds \right) dy \\
& + \int_0^{\infty} \|K(t, \eta[z](t; 0, y))\|_{C_+^*} \|\varphi(\cdot, y)\| \\
& \times \left| \exp \left(\int_0^{\bar{t}} W \left(P_{0,y}^{u,z}(s) \right) ds \right) \right. \\
& \quad \left. - \exp \left(\int_0^t W \left(P_{0,y}^{u,z}(s) \right) ds \right) \right| dy.
\end{aligned}$$

Applying Lemma 3.4, Step 4 and Assumptions $[K]$, $[c, 3]$ and $[W, 3]$, we obtain

$$\begin{aligned}
|\tilde{u}(\bar{t}, 0) - \tilde{u}(t, 0)| & \leq \kappa^2 \int_t^{\bar{t}} Z(\xi) \hat{c}(\xi) \exp \left(\int_{\xi}^{\bar{t}} M_W(s) ds \right) d\xi \\
& + \kappa L_K \int_0^t \left[\int_t^{\bar{t}} \hat{c}(s) ds \right.
\end{aligned}$$



$$\begin{aligned}
 & + |\eta[z](\bar{t}; \xi, 0) - \eta[z](t; \xi, 0)| \Big] Z(\xi) \widehat{c}(\xi) \\
 & \times \exp \left(\int_{\xi}^{\bar{t}} M_W(s) ds \right) d\xi \\
 & + \kappa^2 \int_0^t Z(\xi) \widehat{c}(\xi) \int_t^{\bar{t}} |W(P_{\xi,0}^{u,z}(s))| ds \\
 & \times \exp \left(\int_{\xi}^{\bar{t}} M_W(s) ds \right) d\xi \\
 & + L_K \int_0^{\infty} \left[\int_t^{\bar{t}} \widehat{c}(s) ds \right. \\
 & \left. + |\eta[z](\bar{t}; 0, y) - \eta[z](t; 0, y)| \right] \|\varphi(\cdot, y)\| \\
 & \times \exp \left(\int_0^{\bar{t}} M_W(s) ds \right) dy \\
 & + \kappa \int_0^{\infty} \|\varphi(\cdot, y)\| \int_t^{\bar{t}} |W(P_{0,y}^{u,z}(s))| ds \\
 & \times \exp \left(\int_0^{\bar{t}} M_W(s) ds \right) dy.
 \end{aligned}$$

By the last inequality from Step 2 we arrive at condition 4 for \tilde{u} :

$$|\tilde{u}(\bar{t}, 0) - \tilde{u}(t, 0)| \leq G_u \int_t^{\bar{t}} \widehat{c}(s) ds,$$

where G_u is the same as in Definition 3.1 3.

Step 6. Estimates of increments of \tilde{u} for \bar{x} and x . Denote $\alpha = \alpha[z](t, x)$ and $\bar{\alpha} = \alpha[z](t, \bar{x})$. If $\alpha = \bar{\alpha} = 0$, then Assumptions $[\varphi]$ and $[\lambda, 3]$, together with Steps 1 and 4, applied to equation (10), imply the estimates

$$\begin{aligned}
 |\tilde{u}(t, \bar{x}) - \tilde{u}(t, x)| & \leq L_{\varphi} |\eta[z](0; t, \bar{x}) - \eta[z](0; t, x)| \exp \left(\int_0^t M_{\lambda}(s) ds \right) \\
 & + \|\varphi\| \exp \left(\int_0^t M_{\lambda}(s) ds \right) \\
 & \times \int_0^t L_{\lambda}(s) |\bar{x} - x| (1 + L_u(s))
 \end{aligned}$$



$$\begin{aligned}
& \times \exp \left(\int_s^t L_c(\zeta) d\zeta \right) ds \\
& \leq |\bar{x} - x| \exp \left(\int_0^t (L_c(s) + M_\lambda(s)) ds \right) \\
& \times \left\{ L_\varphi + \|\varphi\| \int_0^t L_\lambda(s)(1 + L_u(s)) ds \right\}.
\end{aligned}$$

If $0 < \alpha < \bar{\alpha}$, then, utilizing Step 4, we obtain

$$\begin{aligned}
|\tilde{u}(t, \bar{x}) - \tilde{u}(t, x)| & \leq G_u \int_\alpha^{\bar{\alpha}} \hat{c}(s) ds \exp \left(\int_\alpha^t M_\lambda(s) ds \right) \\
& + \kappa Z(\alpha) \exp \left(\int_\alpha^t M_\lambda(s) ds \right) \\
& \times \left\{ \int_\alpha^{\bar{\alpha}} M_\lambda(s) ds + \int_\alpha^t L_\lambda(s) |\bar{x} - x| (1 + L_u(s)) \right. \\
& \left. \times \exp \left(\int_s^t L_c(\zeta) d\zeta \right) ds \right\}.
\end{aligned}$$

By Step 3, we have

$$\int_\alpha^{\bar{\alpha}} \hat{c}(s) ds \leq \frac{1}{\varepsilon_0} |\bar{x} - x| \exp \left(\int_\alpha^t L_c(s) ds \right).$$

Hence, we get the inequality

$$\begin{aligned}
|\tilde{u}(t, \bar{x}) - \tilde{u}(t, x)| & \leq |\bar{x} - x| \exp \left(\int_0^t (L_c(s) + M_\lambda(s)) ds \right) \\
& \times \left\{ \frac{G_u}{\varepsilon_0} + \kappa Z(\alpha) \left[\frac{\|M_\lambda/\hat{c}\|}{\varepsilon_0} + \int_0^t L_\lambda(s) (1 + L_u(s)) ds \right] \right\}.
\end{aligned}$$

For arbitrary $x, \bar{x} \in \mathbb{R}_+$, one can find an intermediate point x^* between them such that the differences $|\tilde{u}(t, \bar{x}) - \tilde{u}(t, x^*)|$ and $|\tilde{u}(t, x^*) - \tilde{u}(t, x)|$ have the upper bounds from the above two cases. Thus, we deduce the desired inequality

$$|\tilde{u}(t, \bar{x}) - \tilde{u}(t, x)| \leq L_u(t) |\bar{x} - x|$$

for all $x, \bar{x} \in \mathbb{R}_+$, where L_u is defined in Definition 3.1 4. \square



Remark 4.1. The function L_u from Definition 3.1 4 satisfies the following integral equation:

$$\begin{aligned}
 L_u(t) &= \exp \left(\int_0^a (L_c(s) + M_\lambda(s)) ds \right) \\
 &\times \left\{ L_\varphi + \frac{G_u + \kappa Z(a) \|M_\lambda/\widehat{c}\|}{\varepsilon_0} + [\kappa Z(a) + \|\varphi\|] \right. \\
 &\times \left. \int_0^t L_\lambda(s) (1 + L_u(s)) ds \right\}.
 \end{aligned}$$

Proof of Lemma 3.7. Take any $(\bar{u}, \bar{z}), (u, z) \in \mathcal{X}$ and $(t, x) \in E$. Let $B: [0, a] \rightarrow \mathbb{R}$ be a positive, continuous and nondecreasing function whose precise specification will be given later.

Step 1. Estimate of increments of η for \bar{z} and z . By the Gronwall lemma, we get

$$\begin{aligned}
 &|\eta[\bar{z}](s; t, x) - \eta[z](s; t, x)| \\
 &\leq \|(\bar{z} - z)/B\| \left| \int_s^t B(\zeta) L_c(\zeta) d\zeta \exp \left(\int_s^t L_c(\zeta) d\zeta \right) \right|
 \end{aligned}$$

whenever s belongs to the domains of both characteristics.

Step 2. Estimate of some integrals for $\alpha[\bar{z}]$ and $\alpha[z]$. Denote $\alpha = \alpha[z](t, x)$ and $\bar{\alpha} = \alpha[\bar{z}](t, x)$. Assume that $0 < \alpha < \bar{\alpha}$. Applying (7) to both characteristics and Assumption [c, 3], we obtain the estimate

$$\begin{aligned}
 &\varepsilon_0 \int_\alpha^{\bar{\alpha}} \widehat{c}(s) ds \\
 &\leq \int_\alpha^t L_c(s) \{ |\eta[\bar{z}](s; t, x) - \eta[z](s; t, x)| + \|\bar{z}_s - z_s\| \} ds \\
 &\leq \|(\bar{z} - z)/B\| \int_\alpha^t L_c(s) \left\{ \int_s^t B(\zeta) L_c(\zeta) d\zeta \right. \\
 &\times \left. \exp \left(\int_s^t L_c(\zeta) d\zeta \right) + B(s) \right\} ds.
 \end{aligned}$$

The second inequality is a simple consequence of Step 1.

Step 3. Estimate of increments of λ and W along characteristics. We start with the difference of \bar{u} and u taken along their characteristics $\eta[\bar{z}]$ and $\eta[z]$. Using the function L_u , defined by the formula in



Remark 4.1, we get

$$\begin{aligned} & |\bar{u}(s, \eta[\bar{z}](s; t, x)) - u(s, \eta[z](s; t, x))| \\ & \leq \|(\bar{u} - u)/B\| B(s) \\ & \quad + L_u(s) \|(\bar{z} - z)/B\| \left| \int_s^t B(\zeta) L_c(\zeta) d\zeta \exp \left(\int_s^t L_c(\zeta) d\zeta \right) \right|. \end{aligned}$$

By Assumption $[\lambda, 2]$ and the above inequality, we obtain

$$\begin{aligned} \left| \lambda \left(P_{t,x}^{\bar{u}, \bar{z}}(s) \right) - \lambda \left(P_{t,x}^{u,z}(s) \right) \right| & \leq L_\lambda(s) \left\{ (1 + L_u(s)) \|(\bar{z} - z)/B\| \right. \\ & \quad \times \left| \int_s^t B(\zeta) L_c(\zeta) d\zeta \exp \left(\int_s^t L_c(\zeta) d\zeta \right) \right| \\ & \quad \left. + \|(\bar{u} - u)/B\| B(s) + \|(\bar{z} - z)/B\| B(s) \right\}. \end{aligned}$$

A similar estimate can be derived for increments of W .

Step 4. Estimate of $\tilde{u}(t, 0) - \tilde{u}(t, 0)$. From (9), we have

$$\begin{aligned} & |\tilde{u}(t, 0) - \tilde{u}(t, 0)| \\ & \leq \kappa L_K \int_0^t |\eta[\bar{z}](t; \xi, 0) - \eta[z](t; \xi, 0)| Z(\xi) \hat{c}(\xi) \\ & \quad \times \exp \left(\int_\xi^t M_W(s) ds \right) d\xi \\ & \quad + \kappa \int_0^t \|\tilde{u}_\xi(\cdot, 0) - \tilde{u}_\xi(\cdot, 0)\| \hat{c}(\xi) \\ & \quad \times \exp \left(\int_\xi^t M_W(s) ds \right) d\xi \\ & \quad + \kappa^2 \int_0^t Z(\xi) L_c(\xi) \|\bar{z}_\xi - z_\xi\| \\ & \quad \times \exp \left(\int_\xi^t M_W(s) ds \right) d\xi \\ & \quad + \kappa^2 \int_0^t Z(\xi) \hat{c}(\xi) \left| \exp \left(\int_\xi^t W \left(P_{\xi,0}^{\bar{u}, \bar{z}}(s) \right) ds \right) \right. \\ & \quad \left. - \exp \left(\int_\xi^t W \left(P_{\xi,0}^{u,z}(s) \right) ds \right) \right| d\xi \end{aligned}$$



$$\begin{aligned}
& + L_K \int_0^\infty |\eta[\bar{z}](t; 0, y) - \eta[z](t; 0, y)| \|\varphi(\cdot, y)\| \\
& \times \exp \left| \left(\int_0^t M_W(s) ds \right) dy \right. \\
& \left. + \kappa \int_0^\infty \|\varphi(\cdot, y)\| \left| \exp \left(\int_0^t W \left(P_{0,y}^{\bar{u}, \bar{z}}(s) \right) ds \right) \right. \right. \\
& \left. \left. - \exp \left(\int_0^t W \left(P_{0,y}^{u, z}(s) \right) ds \right) \right| dy.
\end{aligned}$$

Applying the Gronwall inequality, we get

$$|\tilde{u}(t, 0) - \tilde{u}(t, 0)| \leq C_0 \|(\bar{u} - u, \bar{z} - z)\|_B \int_0^t \hat{c}(s) B(s) ds$$

with a positive constant C_0 depending on the data. Applying the same technique to (12), we get the following estimate

$$|\tilde{z}(t) - \tilde{z}(t)| \leq C_1 \|(\bar{u} - u, \bar{z} - z)\|_B \int_0^t \hat{c}(s) B(s) ds.$$

Step 5. Estimate of $\tilde{u}(t, x) - \tilde{u}(t, x)$. Denote $\alpha = \alpha[z](t, x)$ and $\bar{\alpha} = \alpha[\bar{z}](t, x)$. If $\alpha = \bar{\alpha} = 0$, then Assumptions $[\varphi]$, $[\lambda, 3]$ and Step 3 imply

$$|\tilde{u}(t, x) - \tilde{u}(t, x)| \leq C_2 \|(\bar{u} - u, \bar{z} - z)\|_B \int_0^t \hat{c}(s) B(s) ds.$$

If $0 < \alpha < \bar{\alpha}$, then Assumption $[\lambda, 3]$, previous Steps 2, 3, 4 and Lemma 3.5 imply

$$|\tilde{u}(t, x) - \tilde{u}(t, x)| \leq C_3 \|(\bar{u} - u, \bar{z} - z)\|_B \int_0^t \hat{c}(s) B(s) ds.$$

In the third case $0 = \alpha < \bar{\alpha}$ (or $0 = \bar{\alpha} < \alpha$) we consider the family of functions

$$(u_\theta, z_\theta) := \theta(u, z) + (1 - \theta)(\bar{u}, \bar{z}) \quad \text{for } \theta \in [0, 1].$$

We analyze the mapping

$$[0, 1] \ni \theta \longmapsto (\alpha[z_\theta](t, x), \eta[z_\theta](\alpha[z_\theta](t, x); t, x)),$$

whose values belong to the axes $0t$ and $0x$. By the continuous dependence there exists $\theta \in [0, 1]$ for which the point $(0, 0)$ is attained.



Then we have $\alpha[z_\theta](t, x) = 0$; thus, $\eta[z_\theta](0; t, x) = 0$. Hence, we get an intermediate point $(\tilde{u}_\theta, \tilde{z}_\theta) = \mathcal{T}(u_\theta, z_\theta)$, for which we have

$$|\tilde{u}_\theta(t, x) - \tilde{u}(t, x)| \leq C_2 \|(u_\theta - u, z_\theta - z)\|_B \int_0^t \hat{c}(s) B(s) ds$$

and

$$|\tilde{\bar{u}}(t, x) - \tilde{u}_\theta(t, x)| \leq C_3 \|(\bar{u} - u_\theta, \bar{z} - z_\theta)\|_B \int_0^t \hat{c}(s) B(s) ds.$$

Due to this observation, we can reduce the estimate of $|\tilde{\bar{u}}(t, x) - \tilde{u}(t, x)|$ to the previous two cases

$$\begin{aligned} |\tilde{\bar{u}}(t, x) - \tilde{u}(t, x)| &\leq |\tilde{\bar{u}}(t, x) - \tilde{u}_\theta(t, x)| + |\tilde{u}_\theta(t, x) - \tilde{u}(t, x)| \\ &\leq (C_2 + C_3) \|(\bar{u} - u, \bar{z} - z)\|_B \int_0^t \hat{c}(s) B(s) ds. \end{aligned}$$

Step 6. The Bielecki norm. Recall that $(\bar{u}, \bar{z}), (u, z) \in \mathcal{X}$. In force of Steps 4 and 5 we derive

$$\frac{|\tilde{\bar{z}}(t) - \tilde{z}(t)|}{B(t)} \leq \|(\bar{u} - u, \bar{z} - z)\|_B \frac{C_1}{B(t)} \int_0^t \hat{c}(s) B(s) ds$$

and

$$\frac{|\tilde{\bar{u}}(t, x) - \tilde{u}(t, x)|}{B(t)} \leq \|(\bar{u} - u, \bar{z} - z)\|_B \frac{C_2 + C_3}{B(t)} \int_0^t \hat{c}(s) B(s) ds.$$

From these relations, we get

$$\begin{aligned} \max \left\{ \frac{|\tilde{\bar{u}}(t, x) - \tilde{u}(t, x)|}{B(t)}, \frac{|\tilde{\bar{z}}(t) - \tilde{z}(t)|}{B(t)} \right\} \\ \leq \|(\bar{u} - u, \bar{z} - z)\|_B \frac{C_1 + C_2 + C_3}{B(t)} \int_0^t \hat{c}(s) B(s) ds. \end{aligned}$$

Since we intend to estimate the right-hand side by $\Theta \|(\bar{u} - u, \bar{z} - z)\|_B$, it suffices to solve the following elementary comparison equation

$$\Theta + (C_1 + C_2 + C_3) \int_0^t \hat{c}(s) B(s) ds = \Theta B(t).$$



Its solution is given by

$$B(t) = \exp\left(\frac{C_1 + C_2 + C_3}{\Theta} \int_0^t \hat{c}(s) ds\right).$$

Now it is seen that

$$\|(\tilde{u} - \tilde{u}, \tilde{z} - \tilde{z})\|_B \leq \Theta \|(\bar{u} - u, \bar{z} - z)\|_B.$$

This completes the proof. □

Remark 4.2. We explain how formula (9) can be regarded as a fixed point equation for the renewal condition (3). Based on (3) and (10), we get

$$\begin{aligned} \tilde{u}(t, 0) &= \int_0^{\eta_0(t)} K(t, x) \tilde{u}_\alpha(\cdot, 0) \\ &\times \exp\left(\int_\alpha^t \lambda(P_{t,x}^{u,z}(s)) ds\right) dx \\ &+ \int_{\eta_0(t)}^\infty K(t, x) \varphi(\cdot, \eta[z](0; t, x)) \\ &\times \exp\left(\int_0^t \lambda(P_{t,x}^{u,z}(s)) ds\right) dx \end{aligned} \tag{11}$$

for $(t, x) \in E$, where $\alpha = \alpha[z](t, x)$. Using the appropriate change of variables, i.e., $\xi = \alpha[z](t, x)$ to the first integral in (11) and $y = \eta[z](0; t, x)$ to the second integral, we obtain (9). Similar arguments apply to the function \tilde{z} , for which we get the explicit formula

$$\begin{aligned} \tilde{z}(t) &= \int_0^t \tilde{u}(\xi, 0) c(\xi, 0, z_\xi) \\ &\times \exp\left(\int_\xi^t W(P_{\xi,0}^{u,z}(s)) ds\right) d\xi \\ &+ \int_0^\infty \varphi(0, y) \\ &\times \exp\left(\int_0^t W(P_{0,y}^{u,z}(s)) ds\right) dy \end{aligned} \tag{12}$$

for $(t, x) \in E$. These representations of $\tilde{u}(t, 0)$ and $\tilde{z}(t)$ are useful in a priori estimates.



Acknowledgments. We are greatly indebted to the anonymous referees for their valuable comments and suggestions. They improved the whole text and made it more reader friendly.

REFERENCES

1. A. Bielecki, *Une remarque sur la méthode de Banach-Caciopoli-Tikhonov dans la théorie des équations différentielles ordinaires*, Bull. Polish Acad. Sci. **4** (1956), 261–268.
2. K.W. Blayneh, *Analysis of age-structured host-parasitoid model*, Far East J. Dyn. Syst. **4** (2002), 125–145.
3. F. Brauer and C. Castillo-Chávez, *Mathematical models in population biology and epidemiology*, Springer-Verlag, New York, 2001.
4. A. Calsina and J. Saldaña, *A model of physiologically structured population dynamics with a nonlinear growth rate*, J. Mat. Biol. **33** (1995), 335–364.
5. J.M. Cushing, *Integro-differential equations and delay models in population dynamics*, Lect. Notes Biomath. **20**, Springer, Berlin, 1977.
6. ———, *The dynamics of hierarchical age-structured populations*, J. Math. Biol. **32** (1994), 705–729.
7. A.L. Dawidowicz and K. Loskot, *Existence and uniqueness of solution of some integro-differential equation*, Ann. Polon. Math. **47** (1986), 79–87.
8. A.L. Dawidowicz and A. Poskrobko, *Age-dependent single-species population dynamics with delayed argument*, Math. Meth. Appl. Sci. **33** (2010), 1122–1135.
9. B. Ebenman and L. Persson, *Size-structured populations: ecology and evolution*, Springer-Verlag, New York, 1988.
10. J.Z. Farkas, D.M. Green and P. Hinow, *Semigroup analysis of structured parasite populations*, Math. Model. Nat. Phenom. **5** (2010), 94–114.
11. G. Feichtinger, A. Prskawetz and V.M. Veliov, *Age-structured optimal control in population economics*, Theor. Pop. Biol. **65** (2004), 373–387.
12. H. von Foerster, *Some remarks on changing populations*, in *The kinetics of cellular proliferation*, Grune and Stratton, New York, 1959.
13. S.A. Gourley and Y. Kuang, *A stage structured predator-prey model and its dependence on maturation delay and death rate*, J. Math. Biol. **49** (2004), 188–200.
14. M.E. Gurtin and R.C. McCamy, *On the diffusion of biological populations*, Math. Biosci. **33** (1977), 35–49.
15. ———, *Non-linear age-dependent population dynamics*, Arch. Rat. Mech. Anal. **54** (1974), 281–300.
16. J. Hale and S.M. Verduyn Lunel, *Introduction to functional differential equations*, Springer Verlag, New York, 1999.
17. N. Kato, *A general model of size-dependent population dynamics with nonlinear growth rate*, J. Math. Anal. Appl. **297** (2004), 234–256.
18. V.B. Kolmanovskii and A. Myshkis, *Introduction to the theory and applica-*



tions of functional differential equations, MIA **463**, Kluwer Academic Publishers, Dordrecht, 1999.

19. E.A. Kraev, *Existence and uniqueness for height structured hierarchical population models*, Nat. Res. Model. **14** (2001), 45–70.

20. Y. Kuang, *Delay differential equations with applications in population dynamics*, Academic Press, Boston, 1993.

21. A. Lasota, M.C. Mackey and M. Ważewska-Czyżewska, *Minimizing therapeutically induced anemia*, J. Math. Biol. **13** (1981), 149–158.

22. P.H. Leslie, *The use of matrices in certain population mathematics*, Biometrika **33** (1945), 183–212.

23. H. Leszczyński, *Differential functional von Foerster equations with renewal*, Cond. Mat. Phys. **54** (2008), 361–370.

24. H. Leszczyński and P. Zwierkowski, *Existence of solutions to generalized von Foerster equations with functional dependence*, Ann. Polon. Math. **83** (2004), 201–210.

25. M.C. Mackey and R. Rudnicki, *Global stability in a delayed partial differential equation describing cellular replication*, J. Math. Biol. **33** (1994), 89–109.

26. J.A.J. Metz and O. Diekmann, *The dynamics of physiologically structured populations*, Lect. Notes Biomath. **68**, Springer-Verlag, Berlin, 1986.

27. B. Perthame, *Transport equations in biology*, Frontiers Math., Birkhauser, 2007.

28. J.A. Powell, I. Slapničar and W. van der Werf, *Epidemic spread of a lesion-forming plant pathogen analysis of a mechanistic model with infinite age structure*, J. Linear Alg. Appl. **398** (2005), 117–140.

29. J.M. Tchuente, *An age-structured model with delay mortality*, BioSystems **81** (2005), 255–260.

30. W. Walter, *Ordinary Differential Equations*, Springer-Verlag, New York, Berlin, Heidelberg, 1998.

31. G.F. Webb, *Theory of nonlinear age-dependent population dynamics*, Marcel Dekker, New York, 1985.

32. J. Wu, *Theory and applications of partial functional equations*, Springer-Verlag, New York, 1996.

FACULTY OF APPLIED PHYSICS AND MATHEMATICS, GDAŃSK UNIVERSITY OF TECHNOLOGY, GABRIELA NARUTOWICZA 11/12, 80-233 GDAŃSK, POLAND

Email address: agnes@mif.pg.gda.pl

INSTITUTE OF MATHEMATICS, UNIVERSITY OF GDAŃSK, WITA STWOSZA 57, 80-952 GDAŃSK, POLAND

Email address: hleszcz@mat.ug.edu.pl

INSTITUTE OF MATHEMATICS, UNIVERSITY OF GDAŃSK, WITA STWOSZA 57, 80-952 GDAŃSK, POLAND

Email address: zwierkow@mat.ug.edu.pl

