

Simultaneous Determination of Indolic Compounds in Plant Extracts by Solid-Phase Extraction and High-Performance Liquid Chromatography with UV and Fluorescence Detection

Tadeusz Pilipczuk · Natalia Dawidowska ·
Barbara Kusznierevich · Jacek Namieśnik ·
Agnieszka Bartoszek

Abstract A high-performance liquid chromatographic method with UV and fluorescence detection (HPLC-DAD-FLD) was developed for simultaneous determination of indolic compounds in plant material. Indole-3-carbinol (I3C), indole-3-acetic acid (I3AA), indole-3-acetonitrile (I3ACN), and 3,3'-diindolylmethane (DIM) were used as representative compounds that cover a wide spectrum of indole structures occurring in nature. For concentration and purification of the analytes, a solid-phase extraction (SPE) pretreatment was used. The separation utilized a Zorbax XDB-C8 reversed-phase column, acetonitrile-water containing 0.01 % formic acid gradient and UV (280 nm) in series with fluorescence (ex. 280 nm; em. 360 nm) detection. Good linearities of calibration curves were found within the ranges of 0.1–100 nmol/mL for I3C and DIM, 0.4–100 nmol/mL for I3AA and I3ACN for UV detection and 0.003–10 nmol/mL for I3C, 0.003–5 nmol/mL for DIM, and 0.02–10 nmol/mL for I3AA and I3ACN for fluorescence detection. The lowest detection limits (LOD) for UV detection were 0.03 nmol/mL for I3C, 0.1 nmol/mL for I3AA and I3ACN, and 0.04 nmol/mL for DIM, while for fluorescence detection were 0.001 nmol/mL for I3C and DIM, and 0.006 nmol/mL for I3AA and I3ACN. The usefulness of HPLC-DAD-FLD method with SPE

pretreatment is illustrated by the analysis of indolic compounds in extracts from different parts of *Brassica* vegetables.

Keywords Indolic compounds · HPLC-DAD-FLD · SPE · Savoy cabbage · Brussels sprouts · Auxins

Introduction

Indolic compounds are widespread throughout the plant kingdom. Plants can synthesize indoles by several independent biosynthetic pathways. Most of them start from tryptophan resulting in such secondary metabolites as indole-3-pyruvic acid, tryptamine, indole-3-acetaldoxime, indole-3-acetamide, or indolic glucosinolates (Kobayashi et al. 1993; Mashiguchi et al. 2011; Sugawara et al. 2009). There are also pathways independent of tryptophan in which biosynthesis of indole-3-acetic acid (I3AA) begins with indole-3-glycerol phosphate (Ouyang et al. 2000; Zhang et al. 2008) or indole (Mano and Nemoto 2012; Sitbon et al. 2000).

Diversified structures and physicochemical properties of indoles are behind their varied biological activities. Indolic phytohormones belonging to the auxins, such as indole-3-butyric acid, 4-chloroindole-3-acetic acid, and indole-3-acetic acid (I3AA), contain both hydrophilic acidic moiety (carboxyl group) and hydrophobic moiety (indole ring) in their structures (Barkawi et al. 2008). Another type of indoles is represented by indolic β -thioglucosides—glucosinolates (GLs)—a defense-related secondary plant metabolites that include glucobrassicin and its derivatives: 1-methoxyglucobrassicin (Pedras and Hossain 2011), 4-hydroxyglucobrassicin (West et al. 2004), 4-

methoxyglucobrassicin (Kim and Ishii 2006), and 1,4-dimethoxyglucobrassicin (Agerbirk et al. 2001). These phytochemicals following plant tissue disruption are degraded by endogenous enzyme myrosinase and other protein factors into new aglycon indolic compounds, which also differ in their physicochemical properties from precursor GLs. Glucobrassicin breakdown product—indole-3-carbinol (I3C) which possesses C=NH group—has different polarity than its dimer 3,3'-diindolylmethane (DIM) or its derivative *N*-methoxyindole-3-carbinol with *N*-methoxy group (He 1999). It can be presumed that the mentioned above modifications in the structure of indole ring determine differences in anticarcinogenic (Acharya et al. 2010; Kim et al. 2013; Neave et al. 2005; Wang et al. 2013), antioxidative (Shertzer et al. 1986), antimicrobial (Sung and Lee 2007b), or antifungal (Sung and Lee 2007a) properties. For example, I3C and DIM are well known for their multiple therapeutic and preventive effects on prostate (Zhang et al. 2014), colon, or breast (Maruthanila et al. 2014) cancers. Both compounds per se inhibit the growth of different cancer cell lines in vitro and in vivo, e.g., hepatocellular carcinoma cells (Wang et al. 2015), while I3AA requires combination with horseradish peroxidase or activation with UVB/Vis light to induce apoptosis of melanoma cells (Kim et al. 2009). I3ACN does not inhibit cancer cell growth but exhibits inhibitory effect against Gram-negative bacteria. I3C has no effect on this type of bacteria, even at a high dose (Aires et al. 2009), but has inhibitory activity against Gram-positive bacteria such as *Staphylococcus aureus* or *Escherichia coli* (Monte et al. 2014). These data pointing to various structure dependent health-promoting properties of some plant indoles have been confirmed in numerous studies and emphasize nutritional importance of indolic food components.

The significance of indoles in agriculture, as well as in human nutrition, has encouraged the development of several analytical methods for determination of these substances in plant samples (Durgbanshi et al. 2005; Hou et al. 2008). Unfortunately, plant indoles are difficult to analyze because they naturally occur in very low amounts in plants that, to make matters worse, are very rich in interfering components. To cope with this problem, before indoles can be reliably measured, the plant extracts must undergo several purification steps. For example, liquid-liquid extraction of I3AA metabolites was proposed by Liu et al. (2002). It occurred, however, that when the plant samples were analyzed, the interference from the impurities such as pigments allowed to identify only one analyte, so the additional purification step was necessary. In a different method, plant hormone analysis was preceded by HPLC column purification (Kowalczyk and Sandberg 2001). In this case, samples were injected into the LC/MS system and analytes were adsorbed on a C18 precolumn. After washing procedure, precolumn was connected to analytical column, and the analysis was performed in which as an

internal standard heavy-labeled compounds had to be used. Moreover, purification on the C18 column was preceded by two liquid-liquid extractions, evaporation of the solvents and pH adjustment. When GC-MS technique is applied, yet, another additional step, such as derivatization with diazomethane (Barkawi et al. 2008) or methyl chloroformate (Liu et al. 2012) or trimethylsilyldiazomethane (Schmelz et al. 2003), is required. In the latter protocol, methylated derivatives must be additionally heated. After evaporation, the compounds of interest are collected on a polymeric adsorbent and then eluted into the vials (Schmelz et al. 2003). Purification of I3AA and its conjugates with amino acids is possible on an immunoaffinity column with immobilized polyspecific rabbit polyclonal antibodies against I3AA (Pencik et al. 2009). In this approach, sample extract is injected onto the immunoaffinity column, then washed with buffer, eluted, and analyzed by HPLC coupled with MS/MS detector. However, when applied to crude extracts, cross-reactivity or inhibition of antibodies may impair quantification results. These time-consuming methods were modified, and for example, liquid-liquid extraction of phytohormones was replaced by time-efficient single-drop liquid-liquid-liquid microextraction (Bai et al. 2012).

As can be concluded from the above survey of literature, a number of very specific techniques have been proposed for the determination of individual indolic compounds in plants. Nonetheless, there is no single fast and reliable method, which would enable quantitative and qualitative analysis of a wide spectrum of indolic compounds that can be expected to occur at highly variable amounts in the same plant sample. The main objectives of the present study were to simplify the procedures of indole purification from plant extracts and to develop a rapid and sensitive liquid chromatography method enabling the simultaneous separation of structurally divergent analytes. Spectrophotometric detection with diode array detector (DAD) is proposed to pinpoint a broad spectrum of even not identified indoles taking advantage of their characteristic spectral properties, while the use of fluorometric detection ensures high sensitivity of determinations. The applicability of the developed analytic procedure was verified for model indoles I3C, I3AA, I3ACN, and DIM, which cover a wide range of structural and physicochemical variations observed for this group of phytochemicals.

Materials and Methods

Chemicals and Reagents

The standard indolic compounds were purchased from Sigma (Steinheim, Germany), indole-3-carbinol (I3C; ≤ 100 %) and 3,3'-diindolylmethane (DIM; ≥ 98 %), or Merck (Darmstadt, Germany), indole-3-acetic acid (I3AA; ≥ 99 %) and indole-3-acetonitrile (I3ACN; ≥ 98 %). HPLC-grade solvents



(acetonitrile and methanol) and analytical-grade formic acid were obtained from Merck (Darmstadt, Germany). Bakerbond SPE Octadecyl C₁₈ (500 mg, 3 mL) and Bakerbond C₁₈ Polar Plus® (500 mg, 3 mL) cartridges were obtained from J.T. Baker (Greisheim, Germany).

Plant Material

Seeds of *Brassica* plants produced by PNOS (Ożarówice Mazowieckie, Poland) were placed in germination plates and then transferred to the phytotron with controlled temperature (25 °C) and photoperiod (16 h of light, 8 h in the dark). During germination, the seeds were sprinkled with water twice a day. After 7 days, sprouts were harvested, washed to remove seed husks, and kept freeze-dried until investigation.

Brussels sprouts (*Brassica oleracea* L. var. *gemmifera*) and savoy cabbage (*Brassica oleracea* L. var. *sabauda* L.) were harvested from organic plantation in Czapielsk (Poland); white cabbage (*Brassica oleracea* var. *capitata* f. *alba*) was grown in the vicinity of Gdańsk University of Technology, Gdańsk (Poland). The fruit papaya (*Carica papaya* L.) was obtained from a local store. Freshly collected vegetables were lyophilized, ground, and stored at 4 °C until investigation. The seeds were ground right before use.

Extraction and Purification Procedures

The lyophilized samples were ground to a fine powder. Then, 0.2 g of plant material was mixed with 5 mL of sodium phosphate buffer (0.01 M; pH 7.4) and incubated for 3 h at 37 °C to enable full enzymatic degradation of glucosinolates (GLs). The reaction mixture pH was monitored, and when its level became stable for at least 20 min, the hydrolysis of GLs was regarded as complete. Subsequently, the samples were centrifuged (4 °C; 4000 rpm; 10 min; Eppendorf, Microcentrifuge 5415 R, Germany) and supernatants passed through a Bakerbond SPE C₁₈ 500-mg cartridges preconditioned with 3 mL of methanol and 6 mL of water. Stationary phase was washed with 1 mL of sodium phosphate buffer (0.01 M, pH 6.0). Finally, the retained indoles were eluted with 1 mL of methanol. The optimization of the purification conditions is described in the “Results and Discussion” section.

HPLC Analyses

An Agilent 1200 Series HPLC-DAD-FLD system (Agilent Technologies, USA) was employed in the study. All modules were controlled by HPLC ChemStation data acquisition software. The column employed was Zorbax Eclipse XDB-C8 (150×4.60 mm, 3.5 μm) fitted with a suitable precolumn. The mobile phase consisted of (A) 0.01 % formic acid in water and (B) acetonitrile; the flow rate was set at 1 mL/min and the injection volume of extracts was 20 μL. The mobile phase

gradient was programmed as follows: initially, 20 % B increased linearly to 44 % B over 15.4 min, then to 100 % B until 22 min, and finally was kept at 100 % B until 25 min. Finally, the system was equilibrated for 7 min. Indoles were monitored by UV detection at 280 nm (Agilent 1200 series diode array detector) and/or fluorescence detection at 280/360 (ex./em.) (Agilent 1260 fluorescence detector). The contents of indolic compounds in different samples, for which standards were available, were calculated from the calibration curves generated by the integration of areas of absorption peaks determined during analysis of serial dilutions of I3C, I3AA, I3ACN, or DIM.

Results and Discussion


In order to achieve the optimum chromatographic conditions for analysis of indoles, variable parameters including mobile phase composition, injection volume, mobile phase flow-rate, and detection wavelength were tested. Only about 10 % of organic molecules have fluorophore structures. One of such groups embraces indolic compounds, which are therefore often used in experiments as fluorescence probes (Borsarelli et al. 2001; Callis 1997). This property of indole ring makes it possible to employ fluorescence detectors during chromatographic analyses, which offers high selectivity, combined with superior limits of detection (LOD) compared to UV detectors. For the four indolic standards selected for the current study, the optima for excitation and emission wavelengths were determined using combining a diode array detector with a fluorescence detector in series. This enables monitoring UV-VIS and fluorescence spectra in a single run. Such data sets were used to select the optimum fluorescence excitation and emission wavelengths for indolic compounds. Based on these results, the excitation and emission wavelength suitable for the simultaneous detection of all of tested analytes was set at 280 and 360 nm, respectively. The four indole derivatives were separated using a C8 column with water/acetonitrile gradient elution. With the addition of formic acid, chromatographic separation was improved greatly providing sharper peak shapes and better peak symmetry of the indolic standards. Figure 1 shows the representative chromatogram obtained during HPLC resolution of the standard substances with retention times of 6.16, 9.24, 14.95, and 21.22 min for I3C, I3AA, I3ACN, and DIM, respectively.

Optimization of the Procedure of Extraction and Purification

In the developed method, solid-phase extraction (SPE) has been proposed as an alternative to popular liquid-liquid extraction (LLE) of indoles because of simplicity and easy automation. Reversed-phase sorbents (C18 in our study) are commonly used in SPE when aqueous samples are involved.



Fig. 1 The representative HPLC chromatograms obtained for a mixture of standards of indolic compounds occurring most abundantly in *Brassica* plants followed by DAD ($\lambda=280$ nm) (a) or FLD (ex./em.—280/360 nm) (b). I3C indole-3-carbinol, I3AA indole-3-acetic acid, I3ACN indole-3-acetonitrile, DIM 3,3'-diindolylmethane. The concentration of indoles in a mixture solution was 25 nmol mL^{-1}



To determine the most suitable medium for washing indole-containing sample retained on SPE cartridges, a series of methanol solutions of different concentrations were tested. In our study, 1 mL of mixed standard solution of I3C, I3AA, I3ACN, and DIM was loaded onto preconditioned C_{18} cartridges. The concentration of the mixed standard solution was 10 nmol mL^{-1} for I3C, I3AA, and I3ACN and 1 nmol mL^{-1} for DIM. In order to eliminate the contaminating interferences as much as possible, the cartridges were washed by 1 mL of water, 1 mL of sodium phosphate buffer (0.01 M, pH 6.0), or 1 mL of 10, 20, 30, and 40 % methanol solutions and the acidified ones (0.1 % formic acid) of corresponding concentrations, respectively. The indoles were eluted with 1 mL of methanol. Results presented in Fig. 2 indicate that the subsequent recoveries of the indolic compounds from a stationary phase that was washed with methanol were better than those observed for acidified solutions of methanol. Moreover, acidified washing solutions caused decomposition of I3C, and new peak on chromatograms was observed (results not shown). Therefore, for further analyses, non-acidified washing medium was selected.

Because of the low recoveries of I3C and I3AA during SPE washing step, another stationary phase—Bakerbond C_{18} Polar plus[®]—was tested. The obtained results presented in Fig. 2c indicate that such a type of cartridges did not improve recoveries of analytes studied and even additionally caused significant losses of DIM during washing procedure. Among six different washing solutions, the one containing sodium

phosphate buffer (0.01 M, pH 6.0) ensured the highest retention of all four standards on the bed. Considering the above, for further analyses, Bakerbond C_{18} cartridges and sodium phosphate buffer (0.01 M, pH 6.0) for washing the cartridges were chosen.


In order to completely elute the adsorbed indoles from stationary phase while leaving the strongly bound impurities behind, a series of methanol solutions of high concentrations was tested (Fig. 3). Aliquots (1 mL) of mixed standard solution of the I3C, I3AA, I3ACN (10 nmol mL^{-1}), and DIM (1 nmol mL^{-1}) were loaded onto preconditioned four Bakerbond C_{18} columns and washed with 1 mL of sodium phosphate buffer (0.01 M, pH 6.0). Then, the eluting operation was carried out using 1 mL of 70, 80, 90, or 100 % methanol solutions. The results showed that 90 % methanol is sufficient to elute I3C, I3AA, and I3ACN, but in the case of DIM, 100 % methanol was required. Therefore, 100 % methanol was used as the eluting medium in further experiments.

Method Evaluation

The linearity of detector response for each analyte was evaluated by using a series of standard solutions. For each standard solution, measurement was repeated three times. The calibration curves were constructed based on peak area ratio versus concentration of the analyte. The precisions of HPLC-DAD-FLD were examined by comparing the results of three times repeated analysis of the same mixed standard solution of



Fig. 2 The comparison of retention efficiency of different indolic standard compounds on stationary phases: **a, b** Bakerbond C₁₈ or **c** Bakerbond C₁₈ Polar Plus. The washing solutions contained either MeOH+0.1 % formic acid in water (a) or MeOH in water (b, c) or 0.01 M phosphate buffer, pH 6.0 (b). *I3C* indole-3-carbinol, *I3AA* indole-3-acetic acid, *I3ACN* indole-3-acetonitrile, *DIM* 3,3'-diindolylmethane. The results are means ± SD of two independent determinations



indoles. Table 1 gives the summary of linear range, calibration curve, and detection limit for all analyzed indoles.

For FLD detector, good linearities were found in the ranges of 0.003–10 nmol mL⁻¹ for I3C, 0.003–5 nmol mL⁻¹ for DIM, 0.02–10 nmol mL⁻¹ for I3AA and I3ACN. Spectrophotometric detector gave linear signal response in




Fig. 3 The influence of MeOH in water proportion in elution solvent on the recovery of indolic standard compounds from Bakerbond C₁₈ SPE column. *I3C* indole-3-carbinol, *I3AA* indole-3-acetic acid, *I3ACN* indole-3-acetonitrile, *DIM* 3,3'-diindolylmethane. The results are means ± SD of two independent determinations

the range of 0.1–100 nmol mL⁻¹ for I3C and DIM or 0.4–100 nmol mL⁻¹ for I3AA and I3ACN. The relative standard deviations of retention times and peak areas were all under 0.21 % for FLD detector and under 0.52 % for UV detector. The detection limits based on signal-to-noise ratio of three to one were lower for I3C and DIM than for I3AA and I3ACN. The LOD and LOQ for analytes were about 17–40 times lower in the case of fluorescence detection and they were in the range of 0.001–0.006 and 0.003–0.02 nmol mL⁻¹, respectively. Other laboratories also observed higher sensitivity of fluorescence detection. For example, the values of LOD for I3AA were determined in the case of fluorescence detection at the level of 0.05 nmol mL⁻¹ (Szkop and Bielawski 2013) and for MS/MS detection at the level of 13 nmol mL⁻¹ (Hou et al. 2008). The Anderton et al. (2003) reported LOD for I3C and DIM equal to 0.20 and 0.12 nmol mL⁻¹ when the UV detector was used.

In order to examine the recovery efficiency of indolic analytes by the developed method, papaya extracts were spiked with four standard indoles at a different concentration



Table 1 Calibration data for standard indoles determined by HPLC-DAD-FLD method using Zorbax Eclipse XDB-C₈ (150×4.60 mm, 3.5 μm) column

Detector	Analyte	Concentration range (nmol mL ⁻¹)	Calibration curve ^a	LOD (nmol mL ⁻¹)	LOQ (nmol mL ⁻¹)	RSD (%)	
						Retention time	Peak area
FLD	I3C	0.003–10	y=152.59x+0.0033	0.001	0.003	0.06	0.15
	I3AA	0.02–10	y=56.95x+0.0556	0.006	0.02	0.04	0.21
	I3ACN	0.02–10	y=128.44x+0.0395	0.006	0.02	0.02	0.16
	DIM	0.003–5	y=195.88x+0.0607	0.001	0.003	0.01	0.03
UV	I3C	0.1–100	y=5.2813x+0.0391	0.03	0.1	0.05	0.26
	I3AA	0.4–100	y=6.1444x+0.0094	0.1	0.3	0.04	0.52
	I3ACN	0.4–100	y=5.9937x+0.0015	0.1	0.3	0.02	0.11
	DIM	0.1–100	y=12.7430x+0.0044	0.04	0.1	0.01	0.20

I3C indole-3-carbinol, I3AA indole-3-acetic acid, I3ACN indole-3-acetonitrile, DIM 3,3'-diindolylmethane, LOD limit of detection, LOQ limit of quantification, RSD (%) relative standard deviation

^a y is the peak area ratio of standard, x is the concentration (nmol mL⁻¹), for each calibration curve correlation coefficient R² > 0.999

level (0.003, 50, 100 nmol mL⁻¹ for I3C and DIM or 0.02, 50, 100 nmol mL⁻¹ for I3AA and I3ACN) and then extracted by the optimized extraction method. Every sample was analyzed in triplicate. The recoveries summarized in Table 2 were always higher than 94.5 % for I3C, 92.9 % for I3AA, 95.1 % for I3ACN, and 93.2 % for DIM considering three levels of spiking. Additionally, the stability of indoles was investigated by analyzing the plant extract spiked with standards of indolic compounds. The I3C, I3AA, I3ACN, and DIM turned out to be stable at -20 °C for at least 2 days (data not shown).

Quantitative Analysis of Indoles in Plant Material

The HPLC-DAD-FLD method developed was applied to analyze I3C, I3AA, I3ACN, and DIM in plant material derived from different parts of savoy cabbage and brussels sprouts. In Fig. 4, the exemplary chromatogram obtained for cabbage sample is shown. In Table 3, the obtained levels of four indolic compounds are presented. The content of individual indoles

Table 2 Recoveries obtained by HPLC-DAD-FLD analysis of papaya extract spiked with standard solutions of four indoles

Analyte	Recovery (%) ^a		
	0.003 ^b (0.02) ^c nmol mL ⁻¹	50 nmol mL ⁻¹	100 nmol mL ⁻¹
I3C	94.5±6.3	113.8±1.4	113.6±7.3
I3AA	92.9±7.3	104.8±2.7	103.4±0.4
I3ACN	95.1±8.9	102.4±3.6	99.0±0.8
DIM	93.2±4.0	100.5±6.1	103.2±1.7

I3C indole-3-carbinol, I3AA indole-3-acetic acid, I3ACN indole-3-acetonitrile, DIM 3,3'-diindolylmethane

^a Results are mean±RSD (%) for three independent samples

^b The concentration of I3C and DIM


^c The concentration of I3AA and I3ACN (in parenthesis)

differed significantly among various parts of *Brassica* plants. In the case of I3C and its dimer DIM, the highest content was observed in roots of both plants studied. The amount of I3C was 181 and 171 nmol g⁻¹ dw in the case of savoy cabbage and brussels sprouts, respectively. It is in accord with the observations that the highest concentration of the precursor indolic glucosinolates occurs in this organ of plant (Rosa 1997; Kirkegaard and Sarwar 1998). However, Lee et al. (2010) reported similar levels of I3C (630±210 nmol g⁻¹ dw) after autolysis in edible part of Chinese cabbage and lower amount of this analyte in mustard (130±40 nmol g⁻¹ dw). The high level of indoles in roots makes this usually discarded part of *Brassica* plants an attractive raw material for functional food and supplement production.

The data assembled in Table 3 reveal also other interesting interdependencies. The contents of I3C and DIM are significantly correlated in all samples studied (Pearson coefficient $r=0.999$ and $r=0.891$ in the case of savoy cabbage and brussels sprouts, respectively). This suggests that all DIM forms from I3C and conditions in all parts of the plants under study are promoting the condensation reaction to a similar extent. Different situation was described by Ciska et al. (2009) who observed lower I3C and higher DIM amounts in cooked fermented cabbage where the content of these compounds was respectively 0.85 and 1.70 nmol g⁻¹ fw. These results are in agreement with the observations by Bradlow and Zeligs (2010) who suggested alleviated conversion of I3C into DIM under acidic conditions.

In contrast to DIM, no correlation between I3C and I3AA contents is seen (Pearson coefficient $r=-0.172$ and $r=0.236$ for savoy cabbage and brussels sprouts, respectively). This is not surprising as the physiologically most active auxin in plants, that is I3AA, in the case of brassicas can be formed both as a product of degradation of indolic glucosinolates via

Fig. 4 The representative HPLC chromatograms of white cabbage extract followed by DAD ($\lambda=280$ nm) (a) or FLD (ex./em.—280/360 nm) (b). *I3C* indole-3-carbinol, *I3AA* indole-3-acetic acid, *I3ACN* indole-3-acetonitrile, *DIM* 3,3'-diindolylmethane, * unknown indolic compound, *LU* luminescence units, *mAU* milli-arbitrary units



I3ACN or be synthesized in a different pathway (De Vos et al. 2008). Apparently, both pathways are active in samples analyzed. The highest concentration of *I3AA* and *I3ACN* in sprouts can be explained by their high potential to fast growth. Interestingly, high concentration of *I3AA* is also present in stalk of brussels sprouts. It seems that *I3AA* auxin is needed to induce “sprouting” of heads from the stalk. As can be seen, the possibility of detection of several indole metabolites simultaneously may lead to an array of conclusions with regard to plant physiology.

Conclusions

In this study, we have developed the HPLC-DAD-FLD method that enables simultaneous determination of structurally different indolic compounds in plant extract. In addition, the SPE method has been proposed as an alternative to popular liquid-liquid extraction of indoles, because of its simplicity and easy automation. The obvious advantage of this analytical procedure is the reduction of amounts of solvents used and also the efficiency of removal of interferences from extracts, while

Table 3 The content of four indoles in *Brassicaceae* samples determined by HPLC-FLD method

Sample		Contents nmol g ⁻¹ dw (mean±SD) ^a			
		<i>I3C</i>	<i>I3AA</i>	<i>I3ACN</i>	<i>DIM</i>
Savoy cabbage	Roots	181.2±4.6	12.90±0.50	0.98±0.03	0.2660±0.0058
	Seed	31.5±1.2	7.50±0.25	0.32±0.01	0.0853±0.0021
	Sprouts	2.3±0.1	18.69±0.741	155.93±7.37	0.0501±0.0022
Brussels sprout	Roots	171.5±6.7	11.52±0.34	1.32±0.03	0.4073±0.0180
	Seed	1.2±0.1	3.00±0.03	0.56±0.02	0.0660±0.0030
	Sprouts	2.5±0.1	18.41±0.83	172.62±8.56	0.1531±0.0080
	Stalk	77.9±2.5	18.86±1.05	12.30±0.05	0.1182±0.0028
	Leaves (heads)	24.1±1.2	2.92±0.10	0.64±0.02	0.1020±0.0057

I3C indole-3-carbinol, *I3AA* indole-3-acetic acid, *I3ACN* indole-3-acetonitrile, *DIM* 3,3'-diindolylmethane

^a The results are mean ± SD for three independent samples



concentrating indoles, which is particularly important for the analytes occurring at the low level. The developed and optimized chromatographic conditions can be applied for routine analyses of plant extracts. The incorporation of fluorescence detection improved the sensitivity of detection, so the limit of detection (LOD) of indoles has become 40 times lower than in the case of commonly used UV-VIS-only detection. The satisfactory results of analyses of *Brassica* extracts demonstrated that this method may be suitable for the determination of indolic compounds in agriculture and plant physiology studies, as well as in food and food supplement industry.

Acknowledgments This study was realized in the framework of the project “The exploitation of white cabbage for phytoremediation and biofumigation of soils (AGROBIOKAP)” cofinanced by European Union from European Regional Development Fund in a framework of the Innovative Economy Operational Programme 2007–2013 and by the Polish Ministry of Science and Higher Education, grant no. 312244336.

Conflict of Interest Tadeusz Pilipczuk declares that he has no conflict of interest. Natalia Dawidowska declares that she has no conflict of interest. Barbara Kusznierecz declares that she has no conflict of interest. Jacek Namieśnik declares that he has no conflict of interest. Agnieszka Bartoszek declares that she has no conflict of interest. This article does not contain any studies with human or animal subjects.

References

- Acharya A, Das I, Singh S, Saha T (2010) Chemopreventive properties of indole-3-carbinol, diindolylmethane and other constituents of cardamom against carcinogenesis. *Recent Pat Food Nutr Agric* 2:166–177
- Agerbirk N, Petersen BL, Olsen CE, Halkier BA, Nielsen JK (2001) 1,4-Dimethoxyglucobrassicin in *Barbarea* and 4-hydroxyglucobrassicin in *Arabidopsis* and *Brassica*. *J Agric Food Chem* 49:1502–1507. doi:10.1021/Jf001256r
- Anderton MJ, Jukes R, Lamb JH, Manson MM, Gescher A, Steward WP, Williams ML (2003) Liquid chromatographic assay for the simultaneous determination of indole-3-carbinol and its acid condensation products in plasma. *J Chromatogr B Analyt Technol Biomed Life Sci* 787:281–291
- Aires A, Mota VR, Saavedra MJ, Rosa EA, Bennett RN (2009) The antimicrobial effects of glucosinolates and their respective enzymatic hydrolysis products on bacteria isolated from the human intestinal tract. *J Appl Microbiol* 106:2086–2095. doi:10.1111/j.1365-2672.2009.04180.x
- Bai Y, Zhang J, Bai Y, Liu H (2012) Direct analysis in real time mass spectrometry combined with single-drop liquid-liquid-liquid microextraction for the rapid analysis of multiple phytohormones in fruit juice. *Anal Bioanal Chem* 403:2307–2314. doi:10.1007/s00216-012-5728-x
- Barkawi LS et al (2008) A high-throughput method for the quantitative analysis of indole-3-acetic acid and other auxins from plant tissue. *Anal Biochem* 372:177–188. doi:10.1016/j.ab.2007.08.009
- Borsarelli CD, Bertolotti SG, Previtalli CM (2001) Exciplex-type behavior and partition of 3-substituted indole derivatives in reverse micelles made with benzylhexadecyldimethylammonium chloride, water and benzene. *Photochem Photobiol* 73:97–104
- Bradlow HL, Zeligs MA (2010) Diindolylmethane (DIM) spontaneously forms from indole-3-carbinol (I3C) during cell culture experiments. *In Vivo* 24:387–391
- Callis PR (1997) L-1(a) and L-1(b) transitions of tryptophan: applications of theory and experimental observations to fluorescence of proteins. *Method Enzymol* 278:113–150
- Ciska E, Verkerk R, Honke J (2009) Effect of boiling on the content of ascorbigen, indole-3-carbinol, indole-3-acetonitrile, and 3,3'-diindolylmethane in fermented cabbage. *J Agric Food Chem* 57:2334–2338. doi:10.1021/jf803477w
- De Vos M, Kriksunov KL, Jander G (2008) Indole-3-acetonitrile production from indole glucosinolates deters oviposition by *Pieris rapae*. *Plant Physiol* 146:916–926. doi:10.1104/Pp.107.112185
- Durgbanshi A, Arbona V, Pozo O, Miersch O, Sancho JV, Gomez-Cadenas A (2005) Simultaneous determination of multiple phytohormones in plant extracts by liquid chromatography-electrospray tandem mass spectrometry. *J Agric Food Chem* 53:8437–8442. doi:10.1021/Jf050884b
- He H (1999) Studies for growth adaptation and identification of glucosinolates on Chinese brassicas. *Herbert Utz Verlag Wissenschaft, München*
- Hou S, Zhu J, Ding M, Lv G (2008) Simultaneous determination of gibberellic acid, indole-3-acetic acid and abscisic acid in wheat extracts by solid-phase extraction and liquid chromatography-electrospray tandem mass spectrometry. *Talanta* 76:798–802. doi:10.1016/j.talanta.2008.04.041
- Kim EK, Kim YS, Milner JA, Wang TT (2013) Indole-3-carbinol and 3', 3'-diindolylmethane modulate androgen's effect on C-C chemokine ligand 2 and monocyte attraction to prostate cancer cells. *Cancer Prev Res (Phila)* 6:519–529. doi:10.1158/1940-6207.CAPR-12-0419
- Kim SJ, Ishii G (2006) Glucosinolate profiles in the seeds, leaves and roots of rocket salad (*Eruca sativa* Mill.) and anti-oxidative activities of intact plant powder and purified 4-methoxyglucobrassicin. *Soil Sci Plant Nutr* 52:394–400. doi:10.1111/J.1747-0765.2006.00049.X
- Kim SY, Kim MK, Kwon SB, Na JI, Park KC, Kim DS (2009) Tumor apoptosis by indole-3-acetic acid/light in B16F10 melanoma-implanted nude mice. *Arch Dermatol Res* 301:319–322. doi:10.1007/s00403-009-0938-6
- Kobayashi M, Izui H, Nagasawa T, Yamada H (1993) Nitrilase in biosynthesis of the plant hormone indole-3-acetic-acid from indole-3-acetonitrile—cloning of the *alcaligenes* gene and site-directed mutagenesis of cysteine residues. *Proc Natl Acad Sci U S A* 90:247–251. doi:10.1073/Pnas.90.1.247
- Kirkegaard JA, Sarwar M (1998) Biofumigation potential of Brassicas. *Plant Soil* 201:71–89. doi:10.1023/A:1004364713152
- Kowalczyk M, Sandberg G (2001) Quantitative analysis of indole-3-acetic acid metabolites in *Arabidopsis*. *Plant Physiol* 127:1845–1853. doi:10.1104/Pp.010525
- Lee SY, Chu SM, Lee SM, Kim HJ, Cho HS, Yu CY, Kim JK (2010) Determination of indole-3-carbinol and indole-3-acetonitrile in Brassica vegetables using high-performance liquid chromatography with fluorescence detection. *J Korean Soc Appl Biol Chem* 53:249–252. doi:10.3839/Jksabc.2010.039
- Liu BF, Zhong XH, Lu YT (2002) Analysis of plant hormones in tobacco flowers by micellar electrokinetic capillary chromatography coupled with on-line large volume sample stacking. *J Chromatogr A* 945:257–265. doi:10.1016/S0021-9673(01)01503-5
- Liu X, Hegeman AD, Gardner G, Cohen JD (2012) Protocol: high-throughput and quantitative assays of auxin and auxin precursors from minute tissue samples. *Plant Methods* 8:31. doi:10.1186/1746-4811-8-31
- Mano Y, Nemoto K (2012) The pathway of auxin biosynthesis in plants. *J Exp Bot* 63:2853–2872. doi:10.1093/Jxb/Ers091
- Maruthanila VL, Poomima J, Mirunalini S (2014) Attenuation of carcinogenesis and the mechanism underlying by the influence of indole-



3-carbinol and its metabolite 3,3'-diindolylmethane: a therapeutic marvel. *Adv Pharmacol Sci* 2014:832161. doi:10.1155/2014/832161

- Mashiguchi K et al (2011) The main auxin biosynthesis pathway in *Arabidopsis*. *Proc Natl Acad Sci U S A* 108:18512–18517. doi:10.1073/pnas.1108434108
- Monte J, Abreu AC, Borges A, Simoes LC, Simoes M (2014) Antimicrobial activity of selected phytochemicals against *Escherichia coli* and *Staphylococcus aureus* and their biofilms. *Pathogens* 3:473–498. doi:10.3390/pathogens3020473
- Neave AS, Sarup SM, Seidelin M, Duus F, Vang O (2005) Characterization of the N-methoxyindole-3-carbinol (NI3C)—induced cell cycle arrest in human colon cancer cell lines. *Toxicol Sci* 83:126–135. doi:10.1093/toxsci/kfi008
- Ouyang J, Shao X, Li J (2000) Indole-3-glycerol phosphate, a branchpoint of indole-3-acetic acid biosynthesis from the tryptophan biosynthetic pathway in *Arabidopsis thaliana*. *Plant J* 24:327–333
- Pedras MS, Hossain S (2011) Interaction of cruciferous phytoanticipins with plant fungal pathogens: indole glucosinolates are not metabolized but the corresponding desulfo-derivatives and nitriles are. *Phytochemistry* 72:2308–2316. doi:10.1016/j.phytochem.2011.08.018
- Pencik A et al (2009) Isolation of novel indole-3-acetic acid conjugates by immunoaffinity extraction. *Talanta* 80:651–655. doi:10.1016/j.talanta.2009.07.043
- Rosa EAS (1997) Daily variation in glucosinolate concentrations in the leaves and roots of cabbage seedlings in two constant temperature regimes. *J Sci Food Agric* 73:364–368. doi:10.1002/(SICI)1097-0010(199703)73:3<364::AID-JSFA742>3.0.CO;2-O
- Schmelz EA, Engelberth J, Alborn HT, O'Donnell P, Sammons M, Toshima H, Tumlinson JH 3rd (2003) Simultaneous analysis of phytohormones, phytotoxins, and volatile organic compounds in plants. *Proc Natl Acad Sci U S A* 100:10552–10557. doi:10.1073/pnas.1633615100
- Shertzer HG, Niemi MP, Tabor MW (1986) Indole-3-carbinol inhibits lipid peroxidation in cell-free systems. *Adv Exp Med Biol* 197:347–356
- Sitbon F, Astot C, Edlund A, Crozier A, Sandberg G (2000) The relative importance of tryptophan-dependent and tryptophan-independent biosynthesis of indole-3-acetic acid in tobacco during vegetative growth. *Planta* 211:715–721. doi:10.1007/S004250000338
- Sugawara S et al (2009) Biochemical analyses of indole-3-acetaldoxime-dependent auxin biosynthesis in *Arabidopsis*. *Proc Natl Acad Sci U S A* 106:5430–5435. doi:10.1073/pnas.0811226106
- Sung WS, Lee DG (2007a) The candidacidal activity of indole-3-carbinol that binds with DNA. *IUBMB Life* 59:408–412. doi:10.1080/15216540701422373
- Sung WS, Lee DG (2007b) In vitro antimicrobial activity and the mode of action of indole-3-carbinol against human pathogenic microorganisms. *Biol Pharm Bull* 30:1865–1869
- Szkop M, Bielawski W (2013) A simple method for simultaneous RP-HPLC determination of indolic compounds related to bacterial biosynthesis of indole-3-acetic acid. *Antonie Van Leeuwenhoek* 103:683–691. doi:10.1007/s10482-012-9838-4
- Wang X et al (2015) Indole-3-carbinol inhibits tumorigenicity of hepatocellular carcinoma cells via suppression of microRNA-21 and upregulation of phosphatase and tensin homolog. *Biochim Biophys Acta* 1853:244–253. doi:10.1016/j.bbamcr.2014.10.017
- Wang YQ et al (2013) Indole-3-carbinol inhibits cell proliferation and induces apoptosis in Hep-2 laryngeal cancer cells. *Oncol Rep* 30:227–233. doi:10.3892/or.2013.2411
- West LG, Meyer KA, Balch BA, Rossi FJ, Schultz MR, Haas GW (2004) Glucoraphanin and 4-hydroxyglucobrassicin contents in seeds of 59 cultivars of broccoli, raab, kohlrabi, radish, cauliflower, brussels sprouts, kale, and cabbage. *J Agric Food Chem* 52:916–926. doi:10.1021/Jf0307189
- Zhang R, Wang B, Jian OY, Li JY, Wang YH (2008) *Arabidopsis* indole synthase, a homolog of tryptophan synthase alpha, is an enzyme involved in the Trp-independent indole-containing metabolite biosynthesis. *J Integr Plant Biol* 50:1070–1077. doi:10.1111/J.1744-7909.2008.00729.X
- Zhang WW, Feng Z, Narod SA (2014) Multiple therapeutic and preventive effects of 3,3'-diindolylmethane on cancers including prostate cancer and high grade prostatic intraepithelial neoplasia. *J Biomed Res* 28:339–348. doi:10.7555/JBR.28.20140008