



# Turán numbers for odd wheels

Tomasz Dzido<sup>a</sup>, Andrzej Jastrzębski<sup>b,\*</sup>

<sup>a</sup> Institute of Informatics, Faculty of Mathematics, Physics and Informatics, University of Gdańsk, 80-308 Gdańsk, Poland

<sup>b</sup> Department of Algorithms and System Modeling, Faculty of Electronics, Telecommunications and Informatics, Gdańsk University of Technology, 80-233 Gdańsk, Poland

## ARTICLE INFO

### Article history:

Received 28 May 2015

Received in revised form 3 October 2017

Accepted 5 October 2017

Available online 6 November 2017

### Keywords:

Turán numbers

Odd wheels

## ABSTRACT

The Turán number  $\text{ex}(n, G)$  is the maximum number of edges in any  $n$ -vertex graph that does not contain a subgraph isomorphic to  $G$ . A *wheel*  $W_n$  is a graph on  $n$  vertices obtained from a  $C_{n-1}$  by adding one vertex  $w$  and making  $w$  adjacent to all vertices of the  $C_{n-1}$ . We obtain two exact values for small wheels:

$$\text{ex}(n, W_5) = \left\lfloor \frac{n^2}{4} + \frac{n}{2} \right\rfloor,$$

$$\text{ex}(n, W_7) = \left\lfloor \frac{n^2}{4} + \frac{n}{2} + 1 \right\rfloor.$$

Given that  $\text{ex}(n, W_6)$  is already known, this paper completes the spectrum for all wheels up to 7 vertices. In addition, we present the construction which gives us the lower bound  $\text{ex}(n, W_{2k+1}) > \lfloor \frac{n^2}{4} \rfloor + \lfloor \frac{n}{2} \rfloor$  in general case.

© 2017 Elsevier B.V. All rights reserved.

## 1. Introduction

In this paper, all graphs considered are undirected, finite and contain neither loops nor multiple edges. Let  $G$  be such a graph. The vertex set of  $G$  is denoted by  $V(G)$ , the edge set of  $G$  by  $E(G)$ , and the number of edges in  $G$  by  $e(G)$ . Let  $d_G(v)$  be the degree of vertex  $v$  in  $G$ ,  $\delta(G)$  and  $\Delta(G)$  be the minimum and maximum degree of vertices of  $G$ ,  $\omega(G)$  be the clique number of a graph  $G$  and  $\chi(G)$  be the chromatic number of graph  $G$ . Define  $G[S]$  to be a subgraph of  $G$  induced by a set of vertices  $S \subseteq V(G)$  and  $G[S, R]$  to be a bipartite subgraph of  $G$  with the bipartition  $\{S, R\}$ .  $G_1 \cup G_2$  denotes the graph which consists of two disconnected subgraphs  $G_1$  and  $G_2$ . We will use  $G_1 + G_2$  to denote the join of  $G_1$  and  $G_2$  defined as  $G_1 \cup G_2$  together with all edges between  $G_1$  and  $G_2$ .  $C_m$  denotes the cycle of length  $m$ . A *wheel*  $W_n$  is a graph on  $n$  vertices obtained from a  $C_{n-1}$  by adding one vertex  $w$  and making  $w$  adjacent to all vertices of the  $C_{n-1}$ .

The *Turán number*  $\text{ex}(n, G)$  is the maximum number of edges in any  $n$ -vertex graph that does not contain a subgraph isomorphic to  $G$ . A graph on  $n$  vertices is said to be *extremal with respect to*  $G$  if it does not contain a subgraph isomorphic to  $G$  and has exactly  $\text{ex}(n, G)$  edges.  $\text{EX}(n, G)$  is the set of all extremal graphs of order  $n$  with respect to  $G$ .

A main motivation for proving results for Turán numbers is that they are often useful in Ramsey Theory where the original extremal statements would not suffice (see [3] for example). Our goal is to determine the Turán numbers of wheels  $W_k$  for odd  $k$ . We describe families of extremal graphs for  $k = 5, 7$  and present a very simple lower bound for all odd  $k$ .

\* Corresponding author.

E-mail addresses: [tdz@inf.ug.edu.pl](mailto:tdz@inf.ug.edu.pl) (T. Dzido), [jendrek@eti.pg.edu.pl](mailto:jendrek@eti.pg.edu.pl) (A. Jastrzębski).

## 2. Known results

First, we recall the result which was proved by Mantel in 1907.

**Theorem 1** (Mantel, [5]). *The maximum number of edges in an  $n$ -vertex triangle-free graph is  $\lfloor \frac{n^2}{4} \rfloor$ .*

By **Theorem 1** and since  $W_3 = C_3$ , it is easy to have the property that for all integers  $n, n \geq 3$ ,  $\text{ex}(n, W_3) = \lfloor \frac{n^2}{4} \rfloor$ . The famous Turán's theorem may be stated as follows.

**Theorem 2** (Turán, [8]). *Let  $G$  be any subgraph of  $K_n$  such that  $G$  is  $K_{r+1}$ -free. Then the number of edges in  $G$  is  $e(G) = \lfloor \frac{(r-1)n^2}{2r} \rfloor$ . In particular,  $\text{ex}(n, K_4) = \lfloor \frac{n^2}{3} \rfloor$ .*

As a special case, for  $r = 2$ , one obtains Mantel's theorem. Since  $W_4 = K_4$ , we obtain that for all integers  $n, n \geq 3$ ,  $\text{ex}(n, W_4) = \lfloor \frac{n^2}{3} \rfloor$ . In 1964 Erdős proved the following theorem.

**Theorem 3** (Erdős, [4]). *Let  $G$  be any graph such that  $|E(G)| \geq \lfloor \frac{n^2}{4} \rfloor + \lfloor \frac{n}{4} \rfloor + \lfloor \frac{n+1}{4} \rfloor + 1$ . Then  $G$  contains a  $W_5$ .*

By **Theorem 3** we immediately obtain the upper bound for  $\text{ex}(n, W_5)$ , namely  $\text{ex}(n, W_5) \leq \lfloor \frac{n^2}{4} \rfloor + \lfloor \frac{n}{4} \rfloor + \lfloor \frac{n+1}{4} \rfloor + 1$ . The first author [2] proved that for all  $k \geq 3$  and  $n \geq 6k - 10$ , if  $G$  is a graph that contains no subgraph isomorphic to  $W_{2k}$ , then  $\text{ex}(n, W_{2k}) = \lfloor \frac{n^2}{3} \rfloor$ . In addition, he showed that  $\text{ex}(n, W_6) = \lfloor \frac{n^2}{3} \rfloor$ .

If  $G$  is an arbitrary graph whose chromatic number is  $r > 2$ , then by Erdős–Stone–Simonovits theorem [7] we have that  $\text{ex}(n, G) = (\frac{r-2}{r-1} + o(1)) \binom{n}{2}$ . This result determines the asymptotic behavior of  $\text{ex}(n, W_k)$ .

It is interesting that exact values for  $\text{ex}(n, C_4)$  and  $\text{ex}(n, C_6)$ , i.e. for rims of wheels  $W_5$  and  $W_7$  remain unknown in general. Even in the case of the  $C_4$  cycle values are known only for  $n \leq 32$  (the last result being  $\text{ex}(32, C_4) = 92$ , obtained in 2009 by Shao, Xu and Xu), whereas for larger  $n$  only the upper or lower bounds are known.

## 3. Progress on $\text{ex}(n, W_{2k+1})$

### 3.1. $\text{ex}(n, W_5)$

If  $G$  and  $H$  have maximum degree 1, then the join  $G+H$  does not contain  $W_5$ . So define  $M_n$  by taking  $K_{\lfloor n/2 \rfloor, \lfloor n/2 \rfloor}$  and adding a maximum matching within each partite set.

**Lemma 4.** *The graph  $M_n$  does not contain a  $W_5$  as a subgraph.*

**Proof.** Every subgraph induced on 3 vertices of  $W_5$  is connected. If  $i, j, k$  have the same parity then, by definition of  $M_n$ , graph  $M_n[v_i, v_j, v_k]$  has at most one edge, so it is a disconnected graph. If we assume that  $M_n$  has a subgraph  $W_5$ , then at least 3 vertices of this subgraph  $W_5$  are indexed by numbers which have the same parity (we denote the vertices of  $M_n$  as in the definition). A graph induced in  $W_5$  by these three vertices is connected, but a graph induced in  $M_n$  by these vertices is not connected. This means that  $M_n$  does not contain a subgraph  $W_5$ .  $\square$

**Theorem 5.** *The graph  $M_n$  is an extremal graph with respect to  $W_5$ .*

**Proof.** We know that  $M_1 = K_1, M_2 = K_2, M_3 = K_3$  and  $M_4 = K_4$  are extremal. Assume that each  $M_n$  is extremal for  $n < N$ . We will show that  $M_N$  is also extremal. Let  $G$  be an extremal graph of order  $N$ . Let  $H$  be a 4-vertex subgraph of  $G$  with maximum possible number of edges.  $\square$

**Lemma 6.** *A graph  $G$  of order 5 contains  $W_5$  as a subgraph if and only if  $\delta(G) \geq 3$ .*

**Proof.** If  $G$  contains  $W_5$ , it must be a spanning subgraph and so  $\delta(G) \geq 3$ . If  $\delta(G) \geq 3$ , then  $G$  contains a vertex of degree 4 and  $G$  contains a  $W_5$ .  $\square$

Consider the graph  $G \setminus V(H)$ . From **Lemma 6** we know that each vertex from  $G \setminus V(H)$  is adjacent to at most 2 vertices from  $H$ . If any  $v \in G \setminus V(H)$  was adjacent to three vertices of  $H$ , then the graph  $G[V(H) \cup \{v\}]$  would contain  $W_5$  as a subgraph or a 4-vertex subgraph with a greater number of edges than  $H$ . From the above it follows that

$$\begin{aligned} e(G) &\leq e(H) + 2 \cdot |V(G \setminus V(H))| + e(G \setminus V(H)) \\ &\leq \binom{4}{2} + 2 \cdot (N - 4) + \text{ex}(N - 4, W_5) = e(M_N). \end{aligned}$$

If  $G$  is extremal, then  $M_N$  does not contain  $W_5$ . In addition,  $e(M_N) \geq e(G)$ , so  $M_N$  is also extremal.  $\square$

**Table 1**

The values of  $ex(n, W_7)$  and  $|EX(n, W_7)|$  for all  $7 \leq n \leq 26$ .

$n$	7	8	9	10	11	12	13	14	15	16
$ex(n, W_7)$	17	21	25	31	37	43	50	57	65	73
$ EX(n, W_7) $	2	1	5	1	1	2	1	2	1	2
$n$	17	18	19	20	21	22	23	24	25	26
$ex(n, W_7)$	82	91	101	111	122	133	145	157	170	183
$ EX(n, W_7) $	1	2	1	2	1	3	2	3	1	2

**Corollary 7.**

$$ex(n, W_5) = \left\lfloor \frac{n^2}{4} + \frac{n}{2} \right\rfloor.$$

Bataineh, Jaradat and Jaradat [1] presented a very extensive characterization of all extremal  $W_5$ -free graphs.

3.2.  $ex(n, W_7)$

It is not hard to verify that if  $G$  has maximum degree 1 and  $H$  has maximum degree 2 and does not contain  $P_5$ , then the join  $G + H$  does not contain  $W_7$ . So let  $G_m$  be the graph formed from  $m$  isolated vertices by adding a maximum matching. Further, let  $H_m$  be any 2-regular  $m$ -vertex graph formed by the disjoint union of copies of 3- or 4-cycles. (It can be checked that  $H_m$  exists for  $m \geq 6$ .) Then define the graph  $N_n$  as  $G_{k-1} + H_{k+1}$  if  $n = 2k$ , and  $G_k + H_{k+1}$  if  $n = 2k + 1$ . It can be checked that  $N_n$  has  $k^2 + k + 1$  edges if  $n = 2k$ , and  $k^2 + 2k + 2$  edges if  $n = 2k + 1$ .

From this construction we see that  $ex(2k, W_7) \geq k^2 + k + 1$  and  $ex(2k + 1, W_7) \geq k^2 + 2k + 2$ .

**Theorem 8.** For all  $k \geq 5$ , if  $ex(2k, W_7) = k^2 + k + 1$ , then  $ex(2k + 1, W_7) \leq k^2 + 2k + 2$ .

**Proof.** Let  $G$  be a graph of order  $2k + 1$  which does not contain  $W_7$  and assume that  $e(G) = k^2 + 2k + 3$ .

Observe that  $\delta(G) \geq e(G) - ex(2k, W_7) = k + 2$ . Since  $e(G) \geq \frac{(2k+1)(k+2)}{2} > k^2 + 2k + 3 = e(G)$  for all  $k \geq 5$ , we deduce the result.  $\square$

**Theorem 9.** For all  $k \geq 5$ ,  $ex(2k, W_7) = k^2 + k + 1$ .

**Proof.** The cases  $5 \leq k \leq 8$  were checked by computational calculations (see Table 1).

Suppose that  $k > 8$  is the smallest number such that  $ex(2k, W_7) > k^2 + k + 1$ , then for all  $5 \leq l < k$  we have  $ex(2l, W_7) = l^2 + l + 1$  and by Theorem 8  $ex(2l + 1, W_7) = l^2 + 2l + 2$ .

Let  $G$  be a graph of order  $2k$  with  $e(G) = k^2 + k + 2$  edges and  $G$  does not contain  $W_7$  as a subgraph. We see that  $\delta(G) \geq e(G) - ex(2k - 1, W_7) = k + 1$ . If  $\delta(G) \geq k + 2$ , then  $e(G) \geq \frac{2k(k+2)}{2} > e(G)$  for all  $k > 2$ . So we have  $\delta(G) = k + 1$ .

The remaining part of the proof is divided into four cases according to the value of  $\omega(G)$ . Clearly  $\omega(G) < 7$ .

**Case 1.**  $\omega(G) = 6$

Let  $K$  be a clique of order 6 in  $G$  and  $W = V(G) \setminus V(K)$ . To avoid  $W_7$ , every vertex in  $W$  is joined to  $K$  by at most two edges. We have

$$\binom{6}{2} + 2(2k - 6) + ex(2k - 6, W_7) = k^2 - k + 10 < e(G),$$

a contradiction.

**Case 2.**  $\omega(G) = 5$

Let  $K = \{v_1, v_2, v_3, v_4, v_5\}$  be a maximum clique and  $W = V(G) \setminus K$ . Consider the edges of the bipartite graph  $H = G[K, W]$ . Let  $W^4 = \{v \in W : d_H(v) = 4\}$ ,  $W^3 = \{v \in W : d_H(v) = 3\}$  and  $W^r = W - W^4 - W^3$ , obviously if  $v \in W^r$  then  $d_H(v) < 3$ .

One can easily verify that if  $|W^4| \geq 2$ , then we immediately have  $W_7$ . If  $|W^4| = 1$ , then to avoid  $W_7$  in  $G$  we have that  $|W^3| = 0$ . Since  $e(H) \leq 4 + 2(2k - 6) < 5(k - 3) = 5(\delta(G) - 4) \leq e(H)$  for  $k > 7$ , we obtain that in fact  $W^4 = \emptyset$ . Note that  $W^3$  in  $G$  is an independent set and each edge in  $G[K, W^3]$  is adjacent to the same three vertices of  $K$ , say  $\{v_1, v_2, v_3\}$ . From  $\delta(G) = k + 1$ , it follows that  $|W^r| + 3 \geq \delta(G)$ , so  $|W^3| \leq k - 3$ . In fact  $|W^3| = k - 3$  because of the inequality  $e(G) \leq 10 + 3|W^3| + 2|W^r| + ex(2k - 5, W_7) = k^2 + 5 + |W^3|$ .

Note that for every vertex  $v$  in  $W^3$  we have that  $d_G(v) = k + 1$ . The bipartite graph  $G[W^r, W^3]$  is complete, therefore  $\Delta(G[W^r]) \leq 2$ . If not, then we have  $W_7$  in  $G[W]$ . Hence,  $e(G[W]) \leq |W^3||W^r| + \frac{2|W^r|}{2} = k^2 - 4k + 4$  and  $e(G) \leq 10 + 3|W^3| + 2|W^r| + e(G[W]) \leq k^2 + k + 1$ , a contradiction.

We have  $W^4 = W^3 = \emptyset$ ,  $|W^r| = 2k - 5$  but  $e(G[K, W]) \leq 2(2k - 5) < 5(k - 3) = 5(\delta(G) - 4) < e(G[K, W])$  for  $k > 5$ , a contradiction.

**Case 3.**  $\omega(G) = 4$ 

Let  $K = \{v_1, v_2, v_3, v_4\}$  be a maximum clique and  $W = V(G) \setminus K$ .

Let  $U_i$  be the set of vertices from  $W$  such that they are adjacent to all vertices from  $V(K) \setminus \{v_i\}$ . This means that if  $v \in U_i$  then  $d_{G[K, W]}(v) = 3$ . To avoid  $K_5$  all  $U_i$  are independent. Let the remaining vertices of  $W$  be  $W^r$ .

First observe that if  $U_i, U_j, U_l$  are not empty for  $i \neq j \neq l \in \{1, 2, 3, 4\}$ , then we immediately have  $W_7$ . Without loss of generality, let us assume that  $U_3, U_4$  are empty. Observe that if  $|U_1 \cup U_2| > 2$ , then the set  $U_1 \cup U_2$  is independent.

**Subcase 3.1**  $U_1 = U_2 = \emptyset$ 

We have  $e(G) \leq \text{ex}(2k - 4, W_7) + 6 + 2(2k - 4) = k^2 + k + 1 < e(G)$ , a contradiction.

**Subcase 3.2**  $|U_1 \cup U_2| = 1$ 

Without loss of generality, let  $w \in U_1$ . To avoid a contradiction similar to the previous subcase, for all vertices  $v \in W^r$  we have  $d_{G[K, W]}(v) = 2$ . This means that one vertex from  $K$  has degree  $k + 2$  and the remaining three vertices have degree  $k + 1$  in  $G$ , so at least one vertex from  $W$  has degree greater than or equal to  $k + 2$  in  $G$ .

Let  $X$  be all vertices from  $W^r$  adjacent to  $w$  and  $Y = W^r \setminus X$ . Obviously  $|X| \geq k - 2$ . It is not hard to see that if  $G[X]$  contains  $P_4$  or  $K_3$  as a subgraph, then  $G[K \cup U_1 \cup X]$  contains  $W_7$  as a subgraph. If  $|X| \geq 4$ , then there exist at least 3 vertices of degree 1 in  $G[X]$ . These vertices are adjacent to all vertices in  $Y$ , therefore  $\Delta(G[Y]) \leq 2$ ,  $|X| = k - 2$ ,  $|Y| = k - 3$ , subsequently  $\delta(G[X]) = 1$ ,  $\delta(G[Y]) \geq 1$  and  $\Delta(G[Y]) \leq 2$ , so each vertex from  $Y$  is adjacent to all or all except one vertex from  $X$ .

If there exists a vertex  $p \in Y$  such that  $d_G(p) > k + 1$ , then  $d_{G[Y]}(p) = 2$  and  $p$  is adjacent to every vertex in  $X$ . Let  $p_1, p_2$  be the vertices adjacent to  $p$  in  $Y$ . If there exists  $P_3$  in  $G[X]$ , then one end-vertex of the path is adjacent to  $p_1$  and the other to  $p_2$ , then the graph induced by the path,  $p_1, p_2, p$  and an additional vertex from  $X$  adjacent to  $p_1$  and  $p_2$  contains  $W_7$  as a subgraph. Contrary, there exist two independent edges in  $G[X]$  such that their vertices are adjacent to  $p_1$  or  $p_2$ . These edges with  $p_1, p_2$  and  $p$  induce a graph with  $W_7$  as a subgraph.

If there exists a vertex  $p \in X$  such that  $d_G(p) > k + 1$ , then  $d_{G[X]}(p) \geq 2$ . If  $d_{G[X]}(p) = 2$  then  $p$  is adjacent to every vertex in  $Y$ . Let  $p_1$  and  $p_2$  be the adjacent vertices to  $p$  in  $X$ . Note that  $p_1, p_2$  have degree 1 in  $G[X]$ . There exist two independent edges in  $G[Y]$ . Since  $p, p_1$  and  $p_2$  are adjacent to vertices incident to these independent edges, then they both with  $w$  induce a graph with a subgraph  $W_7$ . If  $d_{G[X]}(p) > 2$ , then vertex  $w$ , three vertices adjacent to  $p$  in  $X$  and two vertices adjacent to  $p$  in  $Y$  induce a graph with a subgraph  $W_7$ .

From the above arguments, every vertex of  $W$  has degree  $k + 1$  in  $G$ , so  $e(G[W]) < \text{ex}(2k - 5, W_7)$ , a contradiction.

**Subcase 3.3**  $|U_1 \cup U_2| = 2$ 

Let  $w_1, w_2 \in U_1$ . There exists a vertex  $p \in W^r$  adjacent to  $w_1$  and two vertices of  $K$ ,  $v_1$  and another vertex. A graph induced by  $K \cup U_1$  and  $p$  contains  $W_7$  as a subgraph.

Let  $w_1 \in U_1, w_2 \in U_2$  and  $Q_1, Q_2$  be the set of neighbors of  $w_1, w_2$  in  $W^r$ , respectively. Every vertex of  $W^r$  is adjacent to at least one vertex of  $K$ .

Let  $s_1 \in Q_1 \cap Q_2$  such that  $s_1$  is adjacent to a vertex in  $K$  and  $s_2 \in Q_1$  is adjacent to two vertices in  $K$ . The graph induced by  $K \cup U_1 \cup U_2 \cup \{s_1, s_2\}$  contains a subgraph  $W_7$ .

If there are no vertices in  $Q_1 \cap Q_2$  adjacent to one vertex in  $K$  then every vertex in  $Q_1$  or  $Q_2$  is adjacent to two vertices in  $K$ . Without loss of generality, let  $Q_1$  be such a set. It is easy to see that the set  $Q_1$  is independent. The maximal degree of  $w_1$  in  $G[K \cup U]$  is 4. From the assumption  $\delta(G) \geq k + 1$ , we conclude  $|Q_1| \geq k - 3$ . Let  $X = W^r \setminus Q_1$ . Since each vertex from  $Q_1$  has degree at least  $k + 1$  in  $G$  and  $Q_1$  is independent, we conclude  $|X| \geq k - 3$ ,  $|Q_1| = |X| = k - 3$  and  $w_1$  is adjacent to  $w_2$ . If  $Q_1 \neq Q_2$ , then a vertex from  $Q_2 \setminus Q_1$ , any two vertices from  $Q_1$ , vertices  $w_1, w_2$  and  $K$  induce a graph which contains  $W_7$  as a subgraph. Since  $k \geq 7$ , we have that  $\Delta(G[X]) \leq 2$ . From all previous considerations we have  $e(G) \leq 6 + 7 + 2(2k - 6) + 2(k - 3) + (k - 3) + (k - 3)(k - 3) = k^2 + k + 1$ , a contradiction.

**Subcase 3.4**  $|U_1 \cup U_2| > 2$ 

Let  $W^2 = \{v \in W : d_{G[K, W]}(v) = 2\}$ ,  $W^1 = \{v \in W : d_{G[K, W]}(v) \leq 1\}$  and  $U = U_1 \cup U_2$ . At least one of the sets  $U_1, U_2$  has order greater than or equal to 2, say  $U_1$  is such a set. Let  $u_1, u_2 \in U_1$ . If there exist vertices  $w_1, w_2 \in W^2$  (not necessarily different) such that  $u_1$  is adjacent to  $w_1$  and  $u_2$  is adjacent to  $w_2$ , then the graph  $G[K \cup \{u_1, u_2, w_1, w_2\}]$  contains  $W_7$ . In the opposite case, one of the vertices  $u_1, u_2$  is not adjacent to any vertex from  $W^2$  and since  $U$  is an independent set, we have  $|W^1| \geq k - 2$ . By the inequalities  $e(K) + 3|U| + 2|W^2| + |W^1| + \text{ex}(2k - 4, W_7) \geq e(G)$  and  $|U| + |W^2| + |W^1| = 2k - 4$ , we have  $|U| \geq |W^1| + 1$ , so  $|U| + |W^1| \geq 2k - 3$ , a contradiction.

**Case 4.**  $\omega(G) = 3$ 

Let  $K = \{v_1, v_2, v_3\}$  be the clique in  $G$  and the remaining vertices are  $W$ . Let  $U_i$  be a set of all vertices from  $W$  such that they are adjacent to vertices  $K - v_i$ . This means that if  $v \in U_i$  then  $d_{G[K, W]}(v) = 2$ . To avoid  $K_4$  all  $U_i$  are independent. Let the remaining vertices of  $W$  be  $W^r$  and  $U_1 \cup U_2 \cup U_3 = U$ .

First observe that if there is a  $K_2 \cup K_2$  between  $U_i$  and  $U_j$  where  $i \neq j \in \{1, 2, 3\}$ , then we immediately have  $W_7$ . Since  $3(\delta(G) - 2) \leq e(G[K, W]) \leq (2k - 3 - |U|) + 2|U|$ , we have  $|U| \geq k$ . There exists a vertex in  $U$  adjacent to at most two vertices in  $U$ . This vertex is adjacent to at least  $k - 3$  vertices in  $W^r$ . The equalities  $|U| = k$  and  $|W^r| = k - 3$  are obtained by the above inequalities and the property  $|W^r| + |U| = 2k - 3$ .

If there is a vertex of degree at most 1 in  $U$ , then we have a contradiction with  $\delta(G) = k + 1$ . Since graphs  $G[U_i \cup U_j]$  do not contain  $K_2 \cup K_2$ , the only graph with the property is  $K_{k-2, 1, 1}$ .

Note that all vertices of degree 2 in  $U$  are joined to every vertex of  $W^T$  but none of the vertices of degree  $k - 1$  in  $U$  are joined to any of vertices  $W^T$ . Moreover, to avoid  $W_7$  we have  $\Delta(G[W^T]) \leq 2$ , so none of the vertices in  $G$  has degree greater than  $k + 1$ , a contradiction.  $\square$

### Corollary 10.

$$ex(n, W_7) = \left\lfloor \frac{n^2}{4} + \frac{n}{2} + 1 \right\rfloor.$$

At the end of this subsection we enumerate all of the extremal graphs for  $7 \leq n \leq 26$ . An important property to generate these graphs is that if they exist, then they can be selected from the sets of all  $W_7$ -free graphs with the number of edges greater than or equal to  $\lceil \frac{n^2}{4} + \frac{n}{2} - 1 \rceil$ . The sets were generated using the modified McKay's graph generation program `geng` [6].

For the cases when  $n \in \{7, 8, 9\}$ , the example of the extremal graph is  $C_4 + (K_2 \cup (n - 6)K_1)$ . More precisely, the sets  $EX(n, W_7)$  for these values of  $n$  are as follows:

- $EX(7, W_7) = \{C_4 + (K_2 \cup K_1), K_2 + (K_4 \cup K_1)\}$
- $EX(8, W_7) = \{C_4 + (K_2 \cup 2K_1)\}$
- $EX(9, W_7) = \{C_4 + (K_2 \cup 3K_1), (K_3 \cup K_2) + (K_2 \cup 2K_1), (C_4 \cup K_1) + (K_2 \cup 2K_1), C_5 + 4K_1, 2C_3 + (K_2 \cup K_1)\}$ .

### 3.3. $ex(n, W_{2k+1})$ , where $n \geq 2k + 1$ and $k \geq 4$

Let us recall that we denote by  $aG$  the graph consisting of  $a$  disconnected subgraphs  $G$ . It is not hard to see that the graph  $(K_2 \cup aK_1) + bK_k$  does not contain  $W_{2k+1}$  as a subgraph for all  $a, b \in \mathbb{N}$ . We will try to maximize the number of its edges. We need to determine the number of disconnected copies of  $K_k$ . Consider the situation when  $b = \lfloor \frac{n+k+1}{2k} \rfloor$ . In this case,  $a = n - 2 - k \lfloor \frac{n+k+1}{2k} \rfloor$  and  $ex(n, W_{2k+1}) \geq e(K_k)b + kb(n - kb) + 1$ .

**Theorem 11.** Assume that  $k \geq 4$  and  $n \geq 2k + 1$ . Then

$$ex(n, W_{2k+1}) \geq \left\lfloor \frac{n+k+1}{2k} \right\rfloor \left( \binom{k}{2} + kn - \left\lfloor \frac{n+k+1}{2k} \right\rfloor \right) + 1 > \left\lfloor \frac{n^2}{4} \right\rfloor + \left\lfloor \frac{n}{2} \right\rfloor.$$

### Acknowledgments

We would like to thank the anonymous referees for their careful reading of this manuscript and many helpful comments. This research was partially supported by the Polish National Science Centre grant 2011/02/A/ST6/00201.

### References

- [1] M. Bataineh, M. Jaradat, A. Jaradat, Edge maximal graphs containing no specific wheels, *Jordan J. Math. Statist.* 8 (2) (2015) 107–120.
- [2] T. Dzido, A note on Turán numbers for even wheels, *Graphs Combin.* 29 (5) (2013) 1305–1309.
- [3] T. Dzido, M. Kubale, K. Piwakowski, On some Ramsey and Turán-type numbers for paths and cycles, *Electron. J. Combin.* 13 (2006) #R55.
- [4] P. Erdős, Extremal problems in graph theory, *Theory of graphs and its applications*, in: Proc. Sympos. Smolenice, 1964, pp. 29–36.
- [5] W. Mantel, Problem 28, soln. by H. Gouventak, W. Mantel, J. Teixeira de Mattes, F. Schuh and W.A. Wythoff, *Wiskundige Opgaven* 10 (1907) 60–61.
- [6] B.D. McKay, A. Piperno, Practical graph isomorphism, {II}, *J. Symbolic Comput.* 60 (2014).
- [7] M. Simonovits, A method for solving extremal problems in graph theory, stability problems, in: *Theory of Graphs*, Academic Press, New York, 1968, pp. 279–319.
- [8] P. Turán, On an extremal problem in graph theory, *Mat. Fiz. Lapok* 48 (1941) 436–452.