

## Storm petrels as indicators of pelagic seabird exposure to chemical elements

### in the Antarctic marine ecosystem

Aneta Dorota Pacyna<sup>1\*</sup>, Dariusz Jakubas<sup>2</sup>, Anne N.M.A Ausems<sup>2</sup>, Marcin Frankowski<sup>3</sup>,  
Żaneta Polkowska<sup>1\*</sup>, Katarzyna Wojczulanis-Jakubas<sup>2</sup>

<sup>1</sup>Gdańsk University of Technology, Faculty of Chemistry, Department of Analytical  
Chemistry, Gdańsk, Poland

<sup>2</sup>University of Gdańsk, Faculty of Biology, Department of Vertebrate Ecology and Zoology,  
Gdańsk, Poland

<sup>3</sup>Adam Mickiewicz University Poznań, Faculty of Chemistry, Poznań, Poland

\*Corresponding authors:

Aneta Dorota Pacyna, email: [an.pacyna90@gmail.com](mailto:an.pacyna90@gmail.com),

Żaneta Polkowska, email: [zanpolko@pg.edu.pl](mailto:zanpolko@pg.edu.pl)

### Abstract

Data on trace element bioavailability in the south-polar marine ecosystem is still scarce, compared to that relating to temperate zones. Seabirds can be used as indicators of ecosystem health and sentinels of environmental pollution, constituting a link between marine and terrestrial environments. Here, we analysed the concentration of 17 elements (with special emphasis on mercury, Hg) in feathers of adults and chicks of two pelagic seabirds – the Wilson’s storm petrel *Oceanites oceanicus* and the black-bellied storm petrel *Fregetta tropica* – breeding sympatrically in the maritime Antarctic. Since adult feathers are formed during the non-breeding period away from the breeding grounds, but down and body feathers of chicks

25 grow at the breeding sites, we were able to evaluate the birds' exposure to contaminants at  
26 various stages of their annual life cycle and in various marine zones. We found that of the two  
27 studied species, adult black-bellied storm petrels had significantly higher mercury, selenium  
28 and copper levels ( $5.47 \pm 1.61$ ;  $5.19 \pm 1.18$ ;  $8.20 \pm 0.56 \mu\text{g g}^{-1}$  dw, respectively) than Wilson's  
29 storm petrels ( $2.38 \pm 1.47$ ;  $1.81 \pm 0.98$ ;  $2.52 \pm 2.35 \mu\text{g g}^{-1}$  dw, respectively). We found that  
30 Wilson's storm petrel chicks had a significantly different contaminant profile than adults.  
31 Arsenic, bismuth and antimony were detected exclusively in the chick feathers, and the Se:Hg  
32 molar ratio was higher in chicks than in adults. Our study also suggests considerable maternal  
33 transfer of Hg (to down feathers) in both species. As global contaminant emissions are  
34 expected to increase, birds inhabiting remote areas with sparse anthropogenic pollution can  
35 indicate the temporal trends in global contamination.

36 **Keywords:** contamination, feather, toxic metals, ICP-MS, mercury, Procellariiformes

37

## 38 1. Introduction

39 Organisms living in Antarctica are exposed to a number of environmental factors that may  
40 affect their health and survival. Of those, the most influential are harsh climate conditions,  
41 competition for food, and predation, but pollutants may also play an important role (Santos et  
42 al. 2006, Metcheva et al. 2006). Contaminants in the polar zone may originate from natural  
43 processes [i.e. volcanic activity, the input of sea-spray, mechanical and chemical rock  
44 weathering (Malandrino et al. 2009)] and biota [e.g. mammals or bird colonies can be a  
45 source of nutrients/organic matter (N, F) and several elements such as Cd, Hg, As, Se and Zn  
46 to the terrestrial and coastal ecosystem (Cipro et al. 2018)]. Anthropogenic sources of  
47 Antarctic ecosystem contamination are often located outside the region, e.g. lead from  
48 industrial emissions is transported from South America (Sañudo-Wilhelmy et al. 2002, Gaiero

49 et al. 2003, Bargagli 2008). However, local sources, too, may contribute due to increasing  
50 research and tourism activities resulting in fuel combustion, accidental oil spills, waste  
51 disposal sites, sewage and paint residues (Bargagli 2008, Jerez et al. 2011, Mão de Ferro et al.  
52 2013).

53 Pelagic seabirds living in the southern polar zone can be used in ecotoxicological studies to  
54 assess trace element pollution and marine ecosystem health (Carravieri et al. 2014). They  
55 constitute a valuable link between terrestrial and marine zones of the Antarctic (Santos et al.  
56 2006). As they often cover vast distances in search of suitable foraging areas, they are  
57 exposed to pollutants in various geographical locations. They may also carry these  
58 contaminants between wintering and/or stop-over staging and breeding, due to migratory  
59 connectivity (Webster et al. 2002).

60 Seabirds' feathers are often used to evaluate their exposure to contaminants (e.g. Jerez  
61 et al. 2011, Bustamante et al. 2016, Philpot et al. 2019), providing a record of contaminant  
62 uptake at the time of feather growth and development (Bearhop et al. 2002, Jaspers et al.  
63 2004). High metals affinity to sulfhydryl groups of the feather structural proteins are making  
64 them a suitable biomonitoring tool (Thompson et al. 1998). Elemental deposition in feather  
65 tissue is species-specific and depends on multiple factors, including diet, age, detoxification  
66 abilities and moulting pattern (Burger and Gochfeld 1997, Evers et al. 2008, Cipro et al. 2014,  
67 Pacyna et al. 2017). Knowledge of avian moulting sequences is essential to the reconstruction  
68 of the contamination period (Bustamante et al. 2016, Cherel et al. 2018). Adult feathers may  
69 provide a wider perspective on metal exposure over the annual cycle, but as seabirds may  
70 cover a vast area during the moulting period it is challenging to properly interpret their  
71 exposure over time. Also seasonal shifts in element concentrations can occur (Øverjordet et  
72 al. 2015). By contrast, chick feathers may provide information over a shorter period of  
73 exposure for a more defined area (Evers et al. 2005). Chick down is formed in the egg from

74 maternal nutrients and as such represents female contamination during the pre-laying period  
75 (Ackermann et al. 2016). Thus, analysis of feathers collected at various life stages allows the  
76 elemental concentrations of various areas to be reconstructed, indicating temporal and spatial  
77 trends in pollution in the ecosystems being occupied at the time (Becker, 2003).

78 Despite the growing number of studies on Antarctic and sub-Antarctic food web  
79 contamination, still little is known about elemental concentrations in seabirds feeding at low  
80 trophic levels (i.e. preying on zooplankton and krill), likely due to their relatively lower  
81 exposure to contaminants compared to top predators. For instance, petrels (i.e. species from  
82 three families of Procellariiformes: Procellariidae, Oceanitidae, and Hydrobatidae) are still  
83 one of the most poorly examined seabird groups, mostly due to their small body size, their  
84 nesting predominantly on isolated and inaccessible islands, and their high mobility at sea  
85 (Rodríguez et al. 2019). However, even this group is exposed to a multitude of contaminants  
86 (e.g., Anderson et al. 2000, Bocher et al. 2003, Cipro et al. 2014, Fromant et al. 2016, Philpot  
87 et al. 2019).

88 In this study we determined levels of elements in feathers of two storm-petrel species  
89 breeding in the maritime Antarctica, the Wilson's storm petrel (*Oceanites oceanicus*, hereafter  
90 WSP) and the black-bellied storm petrel (*Fregetta tropica*, hereafter BBSP). We focused both  
91 on elements of wider ecotoxicological interest (i.e. arsenic [As], cadmium [Cd], chromium  
92 [Cr], copper [Cu], lead [Pb], mercury [Hg], selenium [Se], and zinc [Zn]) and on those rarely  
93 studied in avian tissues (i.e. antimony [Sb], bismuth [Bi], calcium [Ca], cobalt [Co], iron [Fe],  
94 nickel [Ni], magnesium [Mg], molybdenum [Mo], and strontium [Sr]). Gathering data about  
95 the concentration of various elements in tissues of living animals is crucial in order to  
96 properly assess ecosystem health and to comprehend pollutants' abilities for potential  
97 bioaccumulation and biomagnification. By studying rarely analysed elements, the results



98 should provide background data for research detecting future inputs of elements in remote  
99 polar regions (Santos et al. 2006).

100 We aimed to:

101 1) present reference values for the concentrations of 17 elements that can be used in the future  
102 for monitoring contamination level in Antarctic marine predators;

103 2) compare elemental concentrations between feathers collected from different age groups  
104 representing various life-history stages (i.e. chick feathers representing the chick-growth  
105 period, chick down representing maternal input, and adult feathers representing part of the  
106 non-breeding period, when the feathers grew); by considering the spatial and temporal  
107 differences in feather growth between these groups, we expected to detect differences in  
108 elemental concentrations between the various types of feathers;

109 3) compare elemental concentrations in feathers grown during the non-breeding season  
110 between adults of the two species, with special emphasis on Hg and Se:Hg molar ratio (linked  
111 to protective action against Hg bioaccumulation and toxicity by creation of Hg-Se compounds  
112 [Nigro and Leonzio 1996, Khan and Wang 2009]). Considering inter-specific differences in  
113 trophic level (see Materials and Methods) and in the location of non-breeding areas (Fig. 1),  
114 we expected to detect differences in elemental concentrations between the species;

115 4) identify patterns in concentrations of elements, and thus identify possible common sources  
116 of contamination.

117

## 118 **2. Materials and Methods**

### 119 *2.1. Studied species*

120 The two study species – the Wilson’s storm petrel and the black-bellied storm petrel – are  
121 small pelagic seabirds, with circumpolar breeding distributions including sub-Antarctic  
122 islands and the maritime Antarctic. Both species breed sympatrically in the study area (see  
123 below) during the austral summer (from December to March), with similar breeding biology:  
124 single-egg clutch, incubation lasting 38–44 days, and chick rearing up to 71 days. Although  
125 both species are among the smallest endotherms living in the Antarctic, they play an  
126 important role as predators preying on Antarctic krill, myctophid fish and amphipods (Hahn  
127 1998, Quillfeldt 2002, Quillfeldt et al. 2005, Wasilewski 1986). Preying on fish and  
128 crustaceans in equal proportions (Hahn 1998), BBSP feeds at a higher trophic level than  
129 WSP, which eats mainly crustaceans (80–90% of meals) (Quillfeldt 2002, Quillfeldt et al.  
130 2017). After breeding, both species migrate northwards, where they spend the non-breeding  
131 period at open sea and complete their moult (Beck and Brown 1972). They moult in the  
132 Atlantic Ocean in a wide range of habitats: WSP from sub-Antarctic to subtropical waters and  
133 BBSP primarily either in sub-Antarctic–subtropical waters or at the continental shelf (Phillips  
134 et al. 2009) (Fig. 1).

135

## 136 *2.2. Sample collection*

137 We studied the two storm-petrel species in the breeding colonies located in the vicinity of  
138 Henryk Arctowski Station in Admiralty Bay, King Gorge Island, South Shetlands, Antarctica  
139 (62°02'S 58°21'W, Fig. 1) in 2017. King George Island is the largest island in the South  
140 Shetlands Archipelago, 90% ice-covered, with rocks mainly formed by andesitic and basaltic  
141 magma (Santos et al. 2006). We captured adult birds in the breeding colony (in their nests,  
142 using mist-nets spread in the colony area) during the incubation period and collected 4–5  
143 body feathers from the back. Back body feathers represent mostly trace element input from  
144 food and water intake during part of the non-breeding period when the feathers grew, which

145 takes place outside the colony in the Atlantic Ocean (Fig. 1). To sample chicks we caught  
146 them by hand in the nest and collected down (at the time they were starting to lose it, i.e.  
147 when their body feathers were well grown, thus minimising the risk of affecting  
148 thermoregulation), and 4–5 body feathers from the back (when the nestlings were about to  
149 fledge). Down feathers represent trace elements passed on by the female to the embryo,  
150 reflecting their input during the pre-laying period, probably from areas around the breeding  
151 colony (and likely predominantly reflecting the food intake). Chick body feathers represent  
152 the nesting period and input from marine environments (as food and water intake). We stored  
153 all the samples in individual plastic zip-lock bags until chemical analysis.

154

### 155 *2.3. Analytical Procedure*

156 Prior to chemical analysis, we cleaned all feather samples to remove external contamination,  
157 firstly with acetone (Sigma-Aldrich, USA) and then two times with deionised water (Mili-Q  
158 Gradient A10, Milipore, France) (procedure of Jaspers et al. 2004, modified). We air-dried the  
159 washed feathers for 24 h. If the total mass of the sample permitted, we used an aliquot of the  
160 collected material for the analysis of all trace elements, using Inductively Coupled Plasma  
161 Mass Spectrometry (ICP-MS).

162 We determined concentrations of 17 trace elements by ICP-MS analytical technique in  
163 the following feather samples types: adult WSP (n=12), chick body WSP (n=4), chick down  
164 WSP (n=4) and adult BBSP (n=4). Mean feather mass was: for adults 10 mg (4–18 mg), for  
165 chick body feathers 6 mg (5–8 mg) and for chick down feathers 16 mg (6–39 mg). Due to  
166 insufficient amount of chick body BBSP and down BBSP feathers, we measured only Hg  
167 content in these samples using the cold vapour technique. In total, we determined Hg  
168 concentration in the following types of feather samples: adult WSP (n=35), chick body WSP

169 (n=10), down WSP (n=16), adult BBSP (n=11), chick body BBSP (n=6) and down BBSP  
170 (n=6).

171

### 172 2.3.1. Trace element concentration

173 We homogenised dry feathers by cutting them up, then weighed them to the nearest 0.01 mg,  
174 and placed them in a clean Teflon vessel with 7 ml 65% HNO<sub>3</sub> (Merck, Suprapur). We carried  
175 out digestion using a high-pressure microwave emitter (Microwave Digestion System, Anton  
176 Paar). We increased the temperature from the ambient value to 90°C (approximately 6–  
177 8°C/min). We maintained these conditions for 25 minutes, after which we gradually lowered  
178 the temperature. Subsequently, we diluted the fully mineralised samples with deionised water  
179 to 25 ml in clean plastic flasks. To ensure quality control and check background  
180 contamination, we ran blank samples with every batch. To ensure accuracy of obtained results  
181 we ran certified reference material (CRM, Human hair ERM-DB001) in triplicate. We  
182 analysed samples using an ICP-MS 2030 (Shimadzu, Japan) (for measurement conditions and  
183 parameters see Table 1, Supplementary material)

### 184 2.3.2. Mercury concentration (cold vapour technique)

185 We weighed the dry samples to the nearest 0.01 mg in a ceramic boat, then we covered them  
186 with activated Al<sub>2</sub>O<sub>3</sub> and analysed them using the thermal vaporisation atomic absorption  
187 method (MA-3000 Nippon Instruments Corporation). We analysed at least two feather  
188 aliquots (1–10 mg dry weight) for each individual. The details of the program used and the  
189 equipment specification are described in Pacyna et al. (2018). We determined total Hg  
190 concentration in duplicates or triplicates when possible, taking sub-samples of the  
191 homogenised feathers. We calculated the coefficient of variation (CV) based on these. If the  
192 CV was above 15%, we excluded samples from the analyses, deeming the estimation of Hg



193 concentration unreliable. Thus, for statistical analysis we used: adult WSP (n=25), chick body  
194 WSP (n=5), down WSP (n=16), adult BBSP (n=8), chick body BBSP (n=5) and down BBSP  
195 (n=6). Mean CV was  $8.64\pm 4.65\%$  for adults,  $6.32\pm 5.40\%$  and  $2.92\pm 2.64\%$  for chick body  
196 feathers and down, respectively. To check background contamination, we performed a quality  
197 control including blank samples every 5–6 subsamples. We analysed CRM every 10th  
198 subsample run.

199

#### 200 2.4. *Quality control*

201 We found that results for CRM analysis were in agreement with the certified values (mass  
202 used for analysis: for trace elements 200 mg, for mercury 14-28 mg). Recoveries were high:  
203 As 98%, Cd 106%, Cu 104%, Hg 92%, Pb 96%, Se 92%, Zn 97%. To check accuracy and  
204 recoveries of other elements absent in this CRM we applied a treatment used before by  
205 Pacyna et al. (2019). We blank-corrected samples analysed on the ICP-MS (by a mean value  
206 of all blank samples). For Hg analysis, we found that background contamination was  
207 negligible and we did not perform blank correction. The limit of detection (LOD) and  
208 quantification (LOQ) values were calculated as the concentrations corresponding to signals  
209 equal to three and ten times the standard deviation of blank solution signal, respectively. For  
210 Hg LOD/LOQ were calculated based on the standard deviation of the response (s), and the  
211 slope of the calibration curve (b) according to the following formulas:  $LOD\ 3.3(s/b)$ ,  $LOQ$   
212  $10(s/b)$ . Method LOD/LOQ were in range of 0.004-0.92 and 0.013-3.07 ng/g respectively. We  
213 reported our results as  $\mu\text{g}\cdot\text{g}^{-1}$  dry weight (dw). For statistical analysis results below the LOD  
214 we assigned half of the LOD value.

215 For calibration of the ICP-MS we used the ICP IV multi-element standard (Merck,  
216 USA) and As, Sb, Se, Mo and V (Sigma-Aldrich, USA), Hg (Merck, USA) as single  
217 standards. As internal standards we used: Sc, Rh, Tb and Ge in supra pure 1% HNO<sub>3</sub> (Merck,

218 USA). For sample pre-treatment and sample dilution we used deionised water obtained from  
219 the Milli-Q Direct 8 Water Purification System.

## 220 2.5. Statistical analyses

221 To investigate variation in the qualitative and quantitative composition of trace elements in  
222 feathers, we firstly performed multivariate analyses for all elements to find general patterns  
223 and then we did univariate analyses for particular elements.

224 To compare the qualitative and quantitative compositions of all trace elements in  
225 feathers among the life-history stages and species, we applied the following multivariate  
226 methods:

227 1) a multivariate (for all trace elements together) PERMANOVA (non-parametric MANOVA  
228 based on the Bray–Curtis measure; Anderson 2001) ) with concentrations of all elements as a  
229 response variable and birds' age (adult WSP, adult BBSP, chick down WSP, chick body  
230 WSP) as the explanatory variable;

231 2) A similarity percentage breakdown procedure (SIMPER) to assess the average percentage  
232 contribution of individual elements to the dissimilarity in all elements concentrations between  
233 age groups in a Bray–Curtis dissimilarity matrix (Clarke 1993).

234 To compare the qualitative and quantitative compositions of particular trace elements  
235 in feathers between the life-history stages and species, we used an unimodal Kruskal–Wallis  
236 test with a *U* Mann–Whitney test as a *post-hoc* test for all group pairs, excluding adult BBSP  
237 vs chick down WSP and adult BBSP vs chick body WSP. In a separate analysis, we compared  
238 Hg concentration among all categories for a larger sample size.

239 Then, to find the groups of elements with high degrees of association in feather  
240 elemental concentrations, we performed a Hierarchical Cluster Analysis (HCA). A high  
241 degree of association between element concentrations, expressed by clustering in one group,  
242 can be used to identify common sources of elements (e.g. Hashmi et al. 2013), but it does not

243 require the formulation of any *a priori* hypothesis considering the nature of the relationships  
244 (Bianchi et al. 2008). We performed HCA with Euclidean distance as a distance measure, and  
245 the paired group method as the linkage method. For each cluster obtained, we calculated  
246 Bootstrap Probability (BP) using multiscale bootstrap resampling. BP of a cluster may take a  
247 value between 0 and 100, indicating how well the data supported the cluster, with a higher  
248 value indicating a better fit (Hammer et al. 2001). We only considered clusters with  $BP \geq 95$ .  
249 To determine how well the generated clusters represented dissimilarities between objects, we  
250 calculated the cophenetic correlation coefficient. Values close to 0 indicate poor clustering,  
251 and values close to 1 show strong clustering.

252 We performed PERMANOVA, SIMPER and HCA analyses on  $\log(x+1)$  transformed  
253 data. We classified the strength of the correlation according to Hinkle et al. (2003): strong  
254 correlation with  $r = |0.90-1.00|$ , high correlation with  $r = |0.70-0.90|$ , moderate correlation with  
255  $r = |0.50-0.70|$ , and low correlation with  $r = |0.30-0.50|$ .

256 We performed separate SIMPER and PERMANOVA analyses for three groups of  
257 elements:

- 258 1. all elements;
- 259 2. essential elements, i.e. As, Ca, Cr, Cu, Fe, Mg, Mo, Se and Zn; and
- 260 3. non-essential elements, i.e. Bi, Cd, Hg, Pb, Sb, and Sr.

261 We calculated Se:Hg molar ratios based on the mean Hg values and the mean Se  
262 values from our study. The Se:Hg molar ratio was obtained using the molecular weight  
263 (200.59 and 78.9 for Hg and Se, respectively) (Burger et al. 2013). We compared Se:Hg  
264 molar ratios between the studied age categories in both species using a chi-squared test.

265 We performed PERMANOVA, SIMPER, and HCA analyses in PAST software  
266 (Hammer et al. 2001) and the Kruskal–Wallis and Mann U Whitney test in R software (R  
267 Core Team 2018), using the ggpubr package (Kassambara 2018).

268

### 269 3. Results

#### 270 3.1. Element concentrations

271 Of all the metals examined, Ni and Co were below the limit of detection in all samples, and  
272 were thus excluded from further analysis. As, Bi and Sb were detected exclusively in chick  
273 feathers, both body and down. Concentrations of all elements (mean±SD) are presented in  
274 Table 1 and Table 4 (for Hg). Concentration chain for particular groups are:

275 a) For adult WSP Mg>Zn>Ca>Fe>Sr>Cu>Mo>Hg>Se>Cr>Pb

276 b) For adult BBSP Mg>Zn>Ca>Fe>Cu>Hg>Se>Sr>Cr>Mo>Pb

277 c) For chicks body feathers Mg>Ca>Fe>Zn>Bi>Sr>Mo>Cu>As>Se>Cr>Pb>Sb>Hg>Cd

278 d) For chicks down feathers Mg>Ca>Fe>Zn>Sr>Se>Mo>Pb=Hg=As>Bi=Cr=Cu>Cd>Sb

279

#### 280 3.2. Inter-group differences in all elements concentration

281 **All elements.** The concentrations of all combined studied elements differed significantly  
282 between adult WSP and all other categories (PERMANOVA, Bonferroni-corrected  $p<0.04$ )  
283 (Table 2). SIMPER analysis showed that the overall average dissimilarity was 18.5%. Fe and  
284 Bi contributed most (14% and 11%, respectively) to the pattern of overall dissimilarity (Table  
285 3). Bi, As and Fe contributed most (19%, 14% and 14%, respectively) to the pattern of  
286 dissimilarity in elemental concentrations observed between adult WSP and chick WSP body  
287 feathers. Fe and Se contributed most (14% and 12%, respectively) to the pattern of  
288 dissimilarity in elemental concentrations observed between adult WSP and chick WSP down.  
289 Cu, Fe, Hg and Se contributed most (16%, 15%, 14% and 12%, respectively) to the pattern of  
290 dissimilarity in elemental concentrations observed between adult WSP and BBSP.



291 **Essential elements.** The concentrations of all combined studied elements differed  
292 significantly between adult WSP and all WSP chick categories (PERMANOVA, Bonferroni-  
293 corrected  $p < 0.006$ ). We found no differences between adult WSP and BBSP ( $p = 0.136$ ) (Table  
294 2). The SIMPER analysis showed that the overall average dissimilarity was 14.2%. More than  
295 half of the pattern of overall dissimilarity observed in elemental concentrations was explained  
296 by Fe, As, Cu and Mo (22%, 15%, 13% and 13%, respectively) (Table 3). As, Fe and Mo  
297 together contributed over 50% (23%, 22% and 15%, respectively) to the pattern of  
298 dissimilarity observed in elemental concentrations between adult WSP and chick WSP body  
299 feathers. Fe, Ca, As, Zn and Mo together contributed over 50% (21%, 13%, 12%, 11% and  
300 11%, respectively) to the pattern of dissimilarity in elemental concentrations observed  
301 between adult WSP and chick WSP down (Table 3). Cu, Fe, Se and Mo together contributed  
302 over 50% (23%, 21%, 18% and 13%, respectively) to the pattern of dissimilarity in elemental  
303 concentrations observed between adult WSP and BBSP (Table 3).

304 **Non-essential elements.** The concentrations of all combined studied elements differed  
305 significantly between adult WSP and all other categories (PERMANOVA, Bonferroni-  
306 corrected  $p < 0.05$ ; Table 2). The SIMPER analysis showed that the overall average  
307 dissimilarity was 39.3%. Bi, Sr, Hg and Pb together contributed over 50% (29%, 25%, 18%  
308 and 14%, respectively) to the pattern of overall dissimilarity observed in elemental  
309 concentrations (Table 3). Bi, Sb, Sr and Pb were responsible for 50%, 14%, 13% and 11%,  
310 respectively, of the dissimilarity pattern in elemental concentrations observed between adult  
311 WSP and chick WSP body feathers. Sr, Pb, Bi and Hg together produced the majority of  
312 dissimilarity in elemental concentrations observed between adult WSP and chick WSP down  
313 (36%, 19%, 19% and 14%, respectively). Hg, Sr and Pb together contributed over 50% (49%,  
314 30% and 16%, respectively) to the pattern of dissimilarity in elemental concentrations  
315 observed between adult WSP and BBSP (Table 3).

### 316 3.3. Intergroup differences for particular elements

317 Kruskal–Wallis inter-group tests comparing the concentration of particular elements revealed  
318 significant differences for all elements ( $p < 0.05$ ) (Supplementary Materials2, Fig. ES1–ES7)  
319 except Mg ( $p = 0.44$ ) and Mo ( $p = 0.12$ ). *Post-hoc* tests revealed the following pattern of  
320 significant inter-group differences (Supplementary Materials2, Fig. ES1– ES7):

- 321 1. lower concentration of Cu, Hg and Se in adult WSP compared to adult BBSP;
  - 322 2. lower concentration of As, Bi, Ca, Cr, Fe, Sb and Se in adult WSP compared to chick WSP  
323 body feathers
  - 324 3. higher concentration of Zn in adult WSP compared to chick WSP down
  - 325 4. higher concentration of As, Bi, Ca, Cd, Fe, Hg, Pb, Sb, Se and Sr in chick WSP down  
326 compared to adult WSP
  - 327 5. higher concentration of As, Bi, Cr, Cu, Sb and Zn in chick WSP body feathers compared to  
328 chick WSP down
  - 329 6. lower concentration of Ca and Sr in chick WSP body feathers compared to WSP down
- 330 Other studied inter-group differences were not significant ( $p > 0.05$ ).

331

### 332 3.4. Inter-group differences for Hg concentration determined by cold vapour technique

333 The Kruskal–Wallis test revealed significant inter-group differences ( $p < 0.05$ ) in the  
334 concentration of Hg determined by cold vapour technique. *Post-hoc* tests results and pattern  
335 of significant inter-group differences are presented in Fig. 2.

336

### 337 3.5. Grouping of elements

338 The Hierarchical Cluster Analysis for all studied groups combined (cophenetic correlation  
339 0.902) recognised four main significant clusters grouping the trace elements (Fig. 3). The first  
340 cluster included Ca and Zn (BP=100), while the second cluster contained Cd-Sb (BP=99).

341 Then, the third was Bi-As (BP=96) and the fourth was Cu-Hg-Se (BP=96), with a subcluster  
342 of Hg-Se (BP=98).

343

#### 344 **4. Discussion**

345 Our study provides reference values for concentration of 17 elements in feathers of two  
346 pelagic seabird species from the maritime Antarctic. We revealed several differences in  
347 elemental concentrations between the two species, as well as differences in exposure between  
348 life-cycle stages. We also identified some patterns in concentrations of particular elements.

349

#### 350 **4.1. Contaminant patterns of selected elements and comparisons with other seabirds** 351 **from south polar areas**

352 Although reference values of 17 elements are provided in our study, below, we discuss only  
353 the those considered most relevant in terms of possible effects on birds' health and survival.

##### 354 *4.1.1. Mercury*

355 Hg is an endocrine disruptor associated with several adverse effects, including decreased body  
356 condition, immune responses and hormonal secretion (Wolfe et al. 1998, Scheuhammer et al.  
357 2007, Tartu et al. 2014, 2015). As such, it affects birds reproduction and survival, and so may  
358 impact birds' population dynamics (Tartu et al. 2013, Goutte et al. 2014). Bird feathers are  
359 perceived as the main route for Hg excretion (Monteiro and Furness 1995, Santos et al. 2006),  
360 but its level would depend on multiple factors, including diet, excretion capacities in the  
361 feathers and moulting pattern (Becker et al. 2016, Bustamante et al. 2016). The Hg  
362 concentration reported in our results for BBSP adults ( $5.47 \pm 1.61 \mu\text{g g}^{-1} \text{ dw}$ ) are in a range of  
363 values reported previously by Carravieri et al. (2014;  $4.22 \pm 2.53 \mu\text{g g}^{-1} \text{ dw}$ ). However, for

364 adult WSP, our values ( $2.38 \pm 1.47 \mu\text{g g}^{-1} \text{ dw}$ ) were much higher compared to other studies  
365 ( $0.42 \pm 0.13 \mu\text{g g}^{-1} \text{ dw}$ ; Carravieri et al. 2014). Nevertheless, Hg levels in adult WSP from our  
366 study were comparable to mean levels reported for another low-trophic-level seabird, the  
367 Antarctic prion *Pachyptila desolata* ( $1.73\text{--}2.80 \mu\text{g g}^{-1} \text{ dw}$ ). In general, there is a high  
368 variability between petrel species ( $0.42\text{--}12.43 \mu\text{g g}^{-1} \text{ dw}$ ; Table 4). Here, the inter-species  
369 difference in Hg concentration is most likely associated with diet (Thompson and Furness  
370 1989a, Bustamante et al. 2016, Blévin et al. 2013) as BBSP feeds at a higher trophic level  
371 than WSP (Quillfeldt et al. 2017). Such a dietary explanation was suggested in the study of  
372 Blévin et al. (2013), where chicks of 21 various species breeding in the Southern Ocean were  
373 been found to vary greatly in terms of Hg concentration (from  $0.05 \pm 0.01 \mu\text{g g}^{-1}$  in the South  
374 Georgian diving petrel *Pelecanoides georgicus* to  $5.31 \pm 1.12 \mu\text{g g}^{-1}$  in the northern giant petrel  
375 *Macronectes halli*).

376 Examining Hg concentrations in age groups, in both species we found that it was  
377 significantly higher in adults than in chicks of the same species (excluding WSP down; Fig. 2)  
378 probably due to the longer exposure time of adults. This is similar to white-chinned petrels  
379 *Procellaria aequinoctialis* (Carvalho et al. 2013), for which the same explanation has been  
380 suggested. In contrast, in the wandering albatross *Diomedea exulans*, Hg contamination was  
381 higher in immatures than adults, which may be associated with moulting intensity and  
382 detoxification capacities varying between adults and immatures (Bustamante et al. 2016).

383

#### 384 4.1.2 Selenium and its interaction with mercury

385 Se is an essential trace element for proper organism functioning, including thyroid function  
386 (Burger et al. 2013), and it is known for its protective action against Hg bioaccumulation and



387 toxicity through the creation of Hg-Se compounds (Nigro and Leonzio 1996, Khan and Wang  
388 2009). However, excess Se may as well have toxic effects on vertebrates (Burger et al. 2013).

389 We found that Se levels significantly differ between the two studied species, with  
390 almost three times higher values found in adult BBSP compared to adult WSP ( $5.19 \pm 1.18 \mu\text{g}$   
391  $\text{g}^{-1}$  dw vs  $1.81 \pm 0.98 \mu\text{g g}^{-1}$  dw). These values add to a wide range reported so far from other  
392 seabirds of the Southern Ocean ( $3.40\text{--}19.40 \mu\text{g g}^{-1}$  dw; Anderson et al. 2010, Fromant et al.  
393 2016, Philpot et al. 2019). Interestingly, Se levels in *Pygoscelis* sp. penguins living in King  
394 George Island were similar to values found in our study ( $2.46\text{--}6.37 \mu\text{g g}^{-1}$  dw; Jerez et al.  
395 2011), while for penguin *Pygoscelis* from other area Hg levels were lower, from  $<0.80$  to  $2.0$   
396  $\mu\text{g g}^{-1}$  (Metcheva et al. 2006).

397 The studied WSP chicks had higher Se levels compared to adults (Table 1). This trend  
398 was not observed in gadfly petrels *Pterodroma spp*, where Se levels were significantly lower  
399 in chicks than in adults (Philpot et al. 2019). These differences are difficult to explain given  
400 the currently limited knowledge about Se distribution and metabolism.

401 Worldwide studies quantifying Hg-Se co-exposure and interaction in seabirds are still  
402 rare, but have increased in recent years, and show that seabirds' ability to deal with high  
403 mercury and selenium levels is still not fully understood, and may depend on age and species  
404 (e.g., Carvalho et al. 2013, Cipro et al. 2014, González-Solís et al. 2002, Carravieri et al.  
405 2017, Philpot et al. 2019). Se-Hg molar ratios in our study differed between chicks and adults,  
406 being highest in WSP chick body feathers (Table 1). However, all ratios reported here were  
407  $>1$ , suggesting activation of a defence mechanism against high Hg concentrations and a health  
408 impact associated with potential Se toxicity (Lucia et al. 2016).

409

410 *4.1.3. Cadmium*



411 Cd is another toxic element that readily bioaccumulates in food webs (Cipro et al. 2014), and  
412 Cd has been reported at even higher levels in Antarctic species including plankton, marine  
413 benthic invertebrates, fishes, seabirds and marine mammals (see references in Jerez et al.  
414 2011), than in their counterparts sampled in polluted coastal areas (Petri and Zauke 1993). In  
415 our study, Cd levels in adults were mostly below the quantification limit (Table 1), but it is  
416 not exceptional (e.g. penguin feathers were also generally low ( $<LOD-0.10 \mu\text{g g}^{-1} \text{ dw}$ , Jerez  
417 et al. [2011];  $<0.15-0.21 \mu\text{g g}^{-1} \text{ dw}$  [Metcheva et al. 2006]) and may be related to relatively  
418 low deposition of Cd in feathers (Lucia et al. 2010, Cipro et al. 2014). Indeed, in Antarctic  
419 prions, Cd levels in internal tissues (kidney  $105\pm37 \mu\text{g g}^{-1} \text{ dw}$ ) were considerably higher than  
420 in feathers (mean  $0.06\pm0.03 \mu\text{g g}^{-1} \text{ dw}$ ) (Fromant et al. 2016). Thus, feathers would give only  
421 partial information of bird exposure; i.e. only when it reaches high levels.

422

#### 423 4.1.4. Lead

424 After Hg, Pb is another major contaminant of toxicological concern (Burger and Gochfeld  
425 2009), affecting breeding success, migratory behaviour and survival of animals at various  
426 trophic levels (Burger, 1995). It may affect food web dynamics e.g. by decreasing the  
427 abundance and availability of food prey, or by interfering with its natural hiding or escape  
428 behaviour (Burger, 1995). Pb is accumulated in feathers at higher rate compared to Cd (Jerez  
429 et al. 2011), but can be elevated due to exogenous contamination (Jaspers et al. 2004).  
430 Adverse effects from lead toxicity might occur at levels of  $4 \mu\text{g g}^{-1}$  in feathers (Burger and  
431 Gochfeld 2000) but levels in adult seabirds are usually lower ( $0.51-1.68 \mu\text{g g}^{-1} \text{ dw}$ , Mendes et  
432 al. 2008, Burger and Gochfeld 2009). In our study the Pb concentration was generally low in  
433 adults ( $<1.17 \mu\text{g g}^{-1} \text{ dw}$ , with one outlier reaching  $5.06 \mu\text{g g}^{-1} \text{ dw}$ ), and higher in chicks ( $0.36-$   
434  $3.67 \mu\text{g g}^{-1} \text{ dw}$ ). Similarly low values of Pb concentration have been reported for other  
435 seabirds from the Southern Ocean (Metcheva et al. 2006, Anderson et al. 2010, Jerez et al.

436 2011, Fromant et al. 2016). However, for penguins breeding on King George Island, high Pb  
437 values have been also reported, which has been explained by local human activities (many  
438 scientific bases and a small airport in the study area; Jerez et al. 2011).

439

#### 440 4.1.5. Zinc

441 Zn can be bioaccumulated in polar organisms, but most likely does not biomagnify (Santos et  
442 al. 2006). Variation of this element concentration in adult storm petrels was relatively low  
443 (WSP  $109.20 \pm 18.50 \mu\text{g g}^{-1} \text{ dw}$ , BBSP  $99.95 \pm 13.01 \mu\text{g g}^{-1} \text{ dw}$ ), and in chicks was even lower,  
444 and with small inter-individual variability (WSP down and body feathers  $48.30 \pm 7.08$  and  
445  $93.00 \pm 11.3 \mu\text{g g}^{-1} \text{ dw}$ , respectively). The reported Zn concentration falls well within the range  
446 reported from other seabirds ( $6.95\text{--}301 \mu\text{g g}^{-1} \text{ dw}$ ) (Anderson et al. 2010, Cipro et al. 2014,  
447 Fromant et al. 2016, Philpot et al. 2019, Metcheva et al. 2006, Jerez et al. 2011, Santos et al.  
448 2006).

#### 449 4.1.6. Copper

450 Cu, like Zn, is also an essential element, with concentrations in seabird tissues controlled  
451 mostly in homeostasis processes (Bocher et al. 2003). Variation in the concentration variation  
452 of Cu in the two species was much larger than for Zn (adults, WSP:  $2.52 \pm 2.35 \mu\text{g g}^{-1} \text{ dw}$ ,  
453 BBSP:  $8.12 \pm 0.56$ ; chick body, WSP:  $6.68 \pm 3.15$ , chick down WSP:  $1.52 \pm 0.60 \mu\text{g g}^{-1} \text{ dw}$ ).  
454 These values also seem to fall well within the range reported so far for other seabirds ( $6.0\text{--}$   
455  $12.7 \mu\text{g g}^{-1} \text{ dw}$ ; Metcheva et al. 2006, Jerez et al. 2011).

456

## 457 4.2. Potential sources of elements

458 The contamination of Antarctic biota may have both natural and anthropogenic sources (Jerez  
459 et al. 2011, Lu et al. 2012, Deheyn et al. 2005). Our cluster analysis revealed some interesting  
460 groupings of elements that suggested common source of contamination.

461 The Bi-As cluster suggests a volcanic origin of the two elements. Worldwide  
462 emissions from volcanoes are deemed a considerable source of atmospheric Bi and As  
463 (Candelone et al. 1995, Kabata-Pendias and Szteke 2015), and the soils on King George  
464 Island are mostly composed of mineral and rock fragments with some volcanic ashes (Lee et  
465 al. 2004). The ashes were blown from Deception Island, a volcanic island located ~130 km  
466 south-west of King George Island (Jeong and Yoon 2001), where the most recent eruption  
467 occurred in the late 1960s (Orheim 1972). Storm petrels may additionally gain As from food  
468 sources, as low-trophic organisms (as petrels diet items) easily assimilate this element  
469 (Rahman et al. 2012). Mão de Ferro et al. (2013) found As enrichment in several Antarctic  
470 abiotic and biotic samples to probably be a result of past volcanic activity and sediment  
471 petrologic characteristic, as well as As leaching processes. They also indicated that during a  
472 high tide, leaching processes of As can occur to shore and semi-submerged areas, thus being  
473 available to aquatic organisms (Mão de Ferro et al. 2013). All feathers were cleaned by the  
474 exact same procedure, but we cannot exclude the possibility of external contamination by soil  
475 particles, as both As and Bi were only detected in chicks feathers.

476 Both elements of the Ca-Zn cluster are necessary components in the synthesis of the  
477 feather pigment melanin (McGraw et al. 2003). They also play an essential role in multiple  
478 physiological body functions (Bogden & Klevay 2000). Thus, this cluster may reflect both co-  
479 exposure from diet and the similar co-regulation mechanisms responsible for element  
480 deposition. Both Ca and Zn accumulation in feathers may depend on melanin type and  
481 content, as shown by element enrichment in pigmented feather parts (Niecke et al. 1999  
482 2003).

483           The Cd-Sb cluster may represent common food and/or water input. Cd may originate  
484 from anthropogenic pollution but also from rock weathering and/or natural sources, as it is  
485 more mobile in seawater than in other water bodies and is easily absorbed by aquatic biota  
486 (Kabata-Pendias and Szeke 2015). Natural sources (diffusive fluxes, upwelling and  
487 continental weathering) can be responsible for higher abundance of Cd in Antarctic water  
488 samples (Sañudo-Wilhelmy et al. 2002). High Cd concentrations were found in Antarctic  
489 krill *Euphasia superba*, which is the main dietary component for adults and chicks of both  
490 storm-petrel species (Wasilewski 1986, Petri and Zauke 1993, Hahn 1998, Nygård et al. 2001,  
491 Quillfeldt 2002). The natural sources of Sb and its compounds are volcanic eruptions, sea  
492 spray, forest fires and wind-blown dust, suggesting a non-anthropogenic source (Kabata-  
493 Pendias and Szeke 2015). Considering its clustering with Cd, we would suggest a natural  
494 source of both elements in the feathers of the studied birds.

495           The common clustering of Cu-Hg-Se may be explained by the properties of Se and the  
496 high concentration of all these elements in aquatic organisms, including fish. Marine aerosols  
497 are enriched in Se resulting from the formation of volatile Se-organic compounds. Volcanic  
498 emissions were suggested as a prevalent source of Hg in Deception Island (Mão de Ferro et al.  
499 2014). Also, summer input from the Southern Ocean may be a net source for the gaseous  
500 element Hg in the marine boundary layer (Wang et al. 2017).

501

### 502 **4.3. Species and age differences in elemental concentrations**

503 Significant differences in concentration of various elements (i.e. Cu, Hg, and Se) between the  
504 WSP and BBSP found in our study are most likely to be associated with inter-species  
505 differences in foraging (different trophic levels with a different contribution of fish in their  
506 diet [Quillfeldt et al. 2017]).

507 Significant differences in concentrations of various elements (Supplementary  
508 Materials2, Fig. ES1–ES7) between WSP age groups are also likely to be associated with diet,  
509 although here not with the difference in diet composition but more with the location of food  
510 resources exploited during the period of growth of relevant feathers. Chick feathers are more  
511 suitable for local exposure assessment, as levels are not affected by moulting patterns,  
512 because chicks receive food collected by parents in the vicinity of the breeding colony, and  
513 have a shorter exposure time. Thus, in cases when adult and offspring diet do not differ  
514 significantly, chick feathers may also be used to reconstruct adults' foraging ecology and  
515 adults' exposure to several pollutants during the chick-rearing period (Blévin et al. 2013).

516 Down feathers have been successfully used to estimate Hg concentrations in eggs  
517 (Santos et al. 2017), suggesting its potential as a suitable proxy for contaminant  
518 determination. A strong correlation between the levels of both Hg and Se in eggs and the liver  
519 of incubating females has been found in *Charadriiformes* (Ackermann et al. 2016). In our  
520 study, all examined elements except Ni and Co were detected in down feathers, enabling  
521 exposure assessment at the earliest phase of life. The highest Ca level was found in WSP  
522 down, probably because the developing embryo absorbs Ca and other elements, initially from  
523 the yolk and subsequently from the eggshell (Castilla et al. 2010). We found the lowest Zn  
524 level in down ( $48.30 \pm 7.08 \mu\text{g/g dw}$ ), at almost two times lower than the level in chick body  
525 feathers ( $93.00 \pm 11.30 \mu\text{g/g dw}$ ). Other elements, such as Pb, Hg, Se, Mg and Sr, were higher  
526 in down than in chick body feathers, suggesting that exposure changes over time. Maternal  
527 transfer of contaminants may be a reason for the increased levels of several metals in chick  
528 down feathers, as the maternal transfer is species- and element-specific (Ackermann et al.  
529 2016).

530

## 531 **5. Conclusions**

532 Our study provides a reference values for concentration of 17 elements in feathers of two  
533 pelagic seabird species from the maritime Antarctic. Such data may serve to monitor  
534 contaminant levels in marine systems and to evaluate variability in contaminant levels in  
535 tissues throughout birds' annual cycle (Rodríguez et al. 2019). We also revealed several  
536 differences in elemental concentrations between the two species, as well as differences in  
537 exposure between life-cycle stages. These inter-species and inter-age differences are  
538 attributed to the various diet compositions and geographic areas of feather growth. Finally, we  
539 identified some patterns in concentration of particular elements that suggest a primarily  
540 natural origin of most elements. We believe that our study contributes to understanding spatial  
541 and temporal patterns of contaminant accumulation in the Maritime Antarctic ecosystem. As  
542 emissions and global transport of elements such as Hg, Pb and Cd are expected to increase in  
543 the future, monitoring studies on seabirds breeding in the Southern Hemisphere may be a  
544 warning system for global changes and the consequences of elevated emissions into the  
545 marine food web. Despite the limitations of our study, such as: the relatively small sample  
546 size; sample collection being restricted to one site and one season; and the lack of data on  
547 elemental concentrations in prey items, our study delivered important reference values for  
548 elemental concentrations in various age groups of the two study species. The Antarctic Treaty  
549 members urge long-term monitoring and sustained observations of the Antarctic environment  
550 and the associated data management, to detect, understand and forecast the impacts of  
551 climate-change-driven environmental variability (ATCM 2007).

552  
553 **Acknowledgements.** The birds were handled and the feather samples collected with  
554 permission from the Department of Antarctic Biology of the Polish Academy of Sciences.  
555 The study was supported by the National Science Centre, Poland, through grant No.  
556 2015/19/B/NZ8/01981. We are grateful to all members of the 41<sup>st</sup> Polish Antarctic Expedition

557 to Arctowski Station for their help and logistical support. Special thanks are given to the  
558 Polish Ministry of Science and Higher Education for financial support of research with project  
559 No. DWM.WKE.183.77.2017

560

## 561 **6. References**

562 Ackerman JT, Eagles-Smith CA, Herzog MP, Hartman CA (2016) Maternal transfer of  
563 contaminants in birds: Mercury and selenium concentrations in parents and their eggs,  
564 Environ Pollut 210, 145-154

565 Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance,  
566 Austral Ecol 26, 32–46

567 Anderson ORJ, Phillips RA, Shore RF, McGill RAR, McDonald RA, Bearhop S (2010)  
568 Element patterns in albatrosses and petrels: influence of trophic position, foraging range,  
569 and prey type, Environ Pollut 158, 98–107

570 ATCM (2007) Resolution 3: Long-term Scientific Monitoring and Sustained Environmental  
571 Observation in Antarctica. In: Final Report ATCM XXX New Delhi, Annex C.  
572 Resolutions

573 Bargagli R (2008) Environmental contamination in Antarctic ecosystems, Sci Total Environ  
574 400, 212-226

575 Bearhop S, Waldron S, Votier SC, Furness RW (2002) Factors that influence assimilation  
576 rates and fractionation of nitrogen and carbon stable isotopes in avian blood and feathers,  
577 Physiol Biochem Zool 75, 451-458

578 Beck JR, Brown DW (1972) The biology of Wilson's Storm petrel, *Oceanites oceanicus*  
579 (Kuhl), at Signy Island, South Orkney Islands, Br Antarct Surv Sci Rep 69,1–54

580 Becker, P. H. (2003). Biomonitoring with birds. In Trace Metals and other Contaminants in  
581 the Environment (Vol. 6, pp. 677-736). Elsevier





582 Becker PH, Goutner V, Ryan PG, González-Solís J (2016) Feather mercury concentrations in  
583 Southern Ocean seabirds: Variation by species, site and time, *Environ Pollut* 216, 253-263  
584 Beyer W, Heinz G, Redmon-Norwood AW (1996) Environmental contaminants in wildlife.  
585 Interpreting tissue concentrations. CRC press, Boca Raton, FL  
586 Bianchi N, Ancora S, di Fazio N, Leonzio C (2008) Cadmium, lead and mercury levels in  
587 feathers of small passerine birds: noninvasive sampling strategy, *Environ Toxicol Chem*  
588 27, 2064–2070  
589 BirdLife International and Handbook of the Birds of the World 2018. Bird species distribution  
590 maps of the world. Version 7.0.  
591 Blévin P, Carravieri A, Jaeger A, Chastel O, Bustamante P, Cheryl Y (2013) Wide Range of  
592 Mercury Contamination in Chicks of Southern Ocean Seabirds. *PLoS ONE* 8(1): e54508.  
593 doi:10.1371/journal.pone.0054508  
594 Bocher P, Caurant F, Miramand P, Cherel Y, Bustamante P (2003) Influence of the diet on the  
595 bioaccumulation of heavy metals in zooplankton-eating petrels at Kerguelen archipelago,  
596 Southern Indian Ocean, *Polar Biol* 26, 759–67  
597 Bogden JD, Klevay LM (2000) Clinical nutrition of the essential trace elements and minerals.  
598 Totowa, NJ: Humana Press. <https://doi.org/10.1007/978-1-59259-040-7>  
599 Burger J (1995) A risk assessment for lead in birds, *J Toxicol Environ Health* 45, 369-396  
600 Burger J, Gochfeld M (1997) Risk, mercury levels, and birds: relating adverse laboratory  
601 effects to field biomonitoring. *Environ Res* 75, 160–172  
602 Burger J, Gochfeld M. (2000) Effects of lead on birds (*Laridae*): a review of laboratory and  
603 field studies. *J Toxicol Environ Health*, 3, 59–78  
604 Burger J, Gochfeld M. (2009) Mercury and other metals in feathers of Common Eider  
605 *Somateria mollissima* and Tufted Puffin *Fratercula cirrhata* from the Aleutian chain of  
606 Alaska, *Arch Environ Contam Toxicol* 56, 596-606



607 Burger J, Jeitner Ch, Donio M, Pittfield T, Gochfeld M (2013) Mercury and selenium levels,  
608 and selenium:mercury molar ratios of brain, muscle and other tissues in bluefish  
609 (*Pomatomus saltatrix*) from New Jersey, USA, *Sci Total Environ* 443, 278–286

610 Bustamante P, Carravieri A, Goutte A, Barbraud Ch, Delord K, Chastel O, Weimerskirch H,  
611 Cherel Y (2016) High feather mercury concentrations in the wandering albatross are  
612 related to sex, breeding status and trophic ecology with no demographic consequences,  
613 *Environ Res* 144, 1–10

614 Candelone JP, Bolshov MA, Rudniev SN, Hong S, Boutron CF (1995). Bismuth in recent  
615 snow from central Greenland: preliminary results. *Atmospheric Environ* 29, 1843-1848

616 Carravieri A, Cherel Y, Blévin P, Brault-Favrou M, Chastel O, Bustamante P (2014) Mercury  
617 exposure in a large subantarctic avian community. *Environ Pollut* 190, 51-57

618 Carravieri A, Cherel Y, Brault-Favrou M, Churlaud C, Peluhet L, Labadie P, Budzinski H,  
619 Chastel O, Bustamante P (2017) From Antarctica to the subtropics: Contrasted  
620 geographical concentrations of selenium, mercury, and persistent organic pollutants in skua  
621 chicks (*Catharacta* spp.), *Environ Pollut* 228, 464-473

622 Carvalho PC, Bugoni L, McGill RAR, Bianchini A (2013) Metal and selenium concentrations  
623 in blood and feathers of petrels of the genus *Procellaria*, *Environ Toxicol Chem* 32, 1641-  
624 1648

625 Castilla AM, Van Dongen S, Herrel A, Francesch A, Martinez de Aragon J, Malone J, Negro  
626 J (2010) Increase in membrane thickness during development compensates for eggshell  
627 thinning due to calcium uptake by the embryo in falcons. *Naturwissenschaften* 97, 143–  
628 151

629 Cherel Y, Barbraud C, Lahournat M, Jaeger A, Jaquemet S, Wanless RM, Phillips RA,  
630 Thompson DR, Bustamante P. (2018) Accumulate or eliminate? Seasonal mercury



631 dynamics in albatrosses, the most contaminated family of birds, Environ Pollut 241, 124-  
632 135

633 Cipro CVZ, Cherel Y, Caurant F, Miramand P, Méndez-Fernandez P, Bustamante P (2014)  
634 Trace elements in tissues of white-chinned petrels (*Procellaria aequinoctialis*) from  
635 Kerguelen waters, Southern Indian Ocean. Polar Biol 37, 763–771

636 Cipro CVZ, Bustamante P, Petry MV, Montone RC (2018) Seabird colonies as relevant  
637 sources of pollutants in Antarctic ecosystems: Part 1 - Trace elements, Chemosphere 204,  
638 535-547

639 Clarke KR (1993) Non-parametric multivariate analysis of changes in community structure.  
640 Aust J Ecol 18, 117–143

641 Deheyn DD, Gendreau P, Baldwin RJ, Latz MI (2005) Evidence for enhanced bioavailability  
642 of trace elements in the marine ecosystem of Deception Island, a volcano in Antarctica.  
643 Mar Environ Res 60, 1-33

644 Eisler R (1998) Copper hazards to fish, wildlife, and invertebrates: a synoptic review. U.S.  
645 Department of the Interior US, Geological Survey, Washington, DC

646 Evers DC, Burgess NM, Champoux L, Hoskins B, Major A, Goodale WM, Taylor R,  
647 Poppenga R, Daigle T (2005) Patterns and interpretation of mercury exposure in  
648 freshwater avian communities in northeastern North America. Ecotoxicology 14, 193-221

649 Evers DC, Savoy LJ, DeSorbo CR, Yates DE, Hanson W, Taylor KM, Siegel LS, Cooley JH  
650 Jr, Bank MS, Major A, Munney K, Mower BF, Vogel HS, Schoch N, Pokras M, Goodale  
651 MW, Fair J (2008) Adverse effects from environmental mercury loads on breeding  
652 common loons. Ecotoxicology 17, 69–81

653 Fromant A, Carravieri A, Bustamante P, Labadie P, Budzinski H, Peluhet L, Churlaud C,  
654 Chastel O, Cherel Y (2016) Wide range of metallic and organic contaminants in various



655 tissues of the Antarctic prion, a planktonophagous seabird from the Southern Ocean. *Sci*  
656 *Total Environ* 544, 754–764

657 Furness RW, Camphuysen CJ (1997) Seabirds as monitors of the marine environment. *ICES J*  
658 *Marine Sci* 54, 726-737

659 González-Solís J, Sanpera C, Ruiz X (2002) Metals and selenium as bioindicators of  
660 geographic and trophic segregation in giant petrels *Macronectes* spp. *Mar Ecol Prog Ser*  
661 244, 257-264

662 Grotti M, Soggia F, Lagomarsino C, Dalla Riva S, Goessler W, Francesconi KA (2008)  
663 Natural variability and distribution of trace elements in marine organisms from Antarctic  
664 coastal environments, *Antarctic Sci* 20, 39-51

665 Hahn S (1998) The food and chick feeding of black bellied storm-petrel (*Fregetta tropica*) at  
666 King George Island, South Shetlands. *Polar Biol.* 19, 354–357

667 Hammer Ø, Harper DAT, Ryan, PD (2001) PAST: paleontological statistics software package  
668 for education and data analysis. *Palaeontologia Electronica*, 4,9.

669 Hashmi MZ, Malik RN, Shahbaz M (2013) Heavy metals in eggshells of cattle egret  
670 (*Bubulcus ibis*) and little egret (*Egretta garzetta*) from the Punjab province. *Pakistan.*  
671 *Ecotox Environ Saf* 89, 158–165

672 Hinkle DE, Wiersma W, Jurs SG (2003) *Applied Statistics for the Behavioral Sciences.*  
673 Houghton Mifflin press, Boston, USA

674 Jaspers V, Dauwe T, Pinxten R, Bervoets L, Blust R, Eens M (2004) The importance of  
675 exogenous contamination on heavy metal levels in bird feathers. A field experiment with  
676 free-living great tits, *Parus major*, *J Environ. Monit* 6, 356 – 360

677 Jeong GY, Yoon HI (2001) The origin of clay minerals in soils of King George Island, South  
678 Shetland Islands, West Antarctica, and its implications for the clay-mineral compositions  
679 of marine sediments. *J Sedimentary Res* 71, 833-842



- 680 Jerez S, Motas M, Palacios MJ, Valera F, Cuervo JJ, Barbosa A (2011) Concentration of trace  
681 elements in feathers of three Antarctic penguins: Geographical and interspecific  
682 differences, *Environ Pollut* 159, 2412–2419
- 683 Kabata-Pendias A, Szeke B (2015). Trace elements in abiotic and biotic environments. CRC  
684 Press
- 685 Kassambara Alboukadel (2018) ggpubr: ‘ggplot2’ Based Publication Ready Plots. R package  
686 version 0.2. in press.
- 687 Khan MAK, Wang F (2009) Mercury-selenium compounds and their toxicological  
688 significance: toward a molecular understanding of the mercury-selenium antagonism,  
689 *Environ Toxicol Chem* 28, 1567-1577
- 690 Lee YI, Lim HS, Yoon HI. (2004) Geochemistry of soils of King George Island, South  
691 Shetland Islands, West Antarctica: implications for pedogenesis in cold polar regions.  
692 *Geochimica et Cosmochimica Acta*, 68, 4319-4333
- 693 Lu Z, Cai M, Wang J, Yang H, He J (2012) Baseline values for metals in soils on Fildes  
694 Peninsula, King George Island, Antarctica: the extent of anthropogenic pollution, *Environ*  
695 *Mon Asses* 184, 7013-7021
- 696 Lucia M, André JM, Gontier K, Diot N, Veiga J, Davail S (2010) Trace Element  
697 Concentrations (Mercury, Cadmium, Copper, Zinc, Lead, Aluminium, Nickel, Arsenic,  
698 and Selenium) in Some Aquatic Birds of the Southwest Atlantic Coast of France, *Arch*  
699 *Environ Contam Toxicol* 58, 844-853
- 700 Lucia M, Strøm H, Bustamante P, Gabrielsen GW (2016) Trace Element Concentrations in  
701 Relation to the Trophic Behaviour of Endangered Ivory Gulls (*Pagophila eburnea*) During  
702 Their Stay at a Breeding Site in Svalbard, *Arch Environ Contam Toxicol* 71, 518–529



- 703 Malandrino M, Abollino O, Buoso S, Casalino CE, Gasparon M, Giacomino A, La Gioia C,  
704 Mentasti E (2009) Geochemical characterisation of antarctic soils and lacustrine sediments  
705 from Terra Nova Bay, *Microchem J* 92, 21-31
- 706 Mão de Ferro A, Mota AM, Canário J (2013) Sources and transport of As, Cu, Cd and Pb in  
707 the environmental compartments of Deception Island, Antarctica, *Mar Pollut Bull* 77, 341–  
708 348
- 709 Mão de Ferro A, Mota AM, Canário J (2014) Pathways and speciation of mercury in the  
710 environmental compartments of Deception Island, Antarctica, *Chemosphere* 95, 227-233
- 711 McGraw KJ, Beebee MD, Hill GE, Parker RS. (2003) Lutein-based plumage coloration in  
712 songbirds in a consequence of selective pigment incorporation into feathers, *Comp Bioch*  
713 *Physiol* 135, 689–696
- 714 Mendes P, Eira C, Torres J, Soares AMVM, Melo P, Vingada J. (2008) Toxic element  
715 concentration in the Atlantic Gannet *Morus bassanus* (Pelecaniformes, Sulidae) in  
716 Portugal, *Arch Environ Contam Toxicol* 55, 209-503
- 717 Metcheva R, Yurukova L, Teodorova S, Nikolova E. (2006) The penguin feathers as  
718 bioindicator of Antarctica environmental state, *Sci Total Environ* 362, 259-265
- 719 Monteiro LR, Furness RW (1995) Seabirds as monitors of mercury in the marine  
720 environment. *Water Air Soil Pollut* 80, 851–870
- 721 Monteiro LR, Furness RW (2001) Kinetics, dose-response, and excretion of methylmercury in  
722 free-living adult Cory's Shearwaters. *Environ Sci Tech* 35, 739–746
- 723 Niecke M, Heid M, Krüger A (1999) Correlations between melanin pigmentation and element  
724 concentration in feathers of White-tailed Eagles (*Haliaeetus albicilla*). *J Ornithol* 140,  
725 355–362



- 726 Niecke M, Rothlaender S, Roulin A (2003) Why do melanin ornaments signal individual  
727 quality? Insights from metal element analysis of barn owl feathers. *Oecologia* 137, 153–  
728 158
- 729 Norheim G, Kjos-Hanssen B. (1984) Persistent chlorinated hydrocarbons and mercury in  
730 birds caught off the west coast of Spitsbergen. *Environ Pollut A* 33, 143-152
- 731 Nygård T, Lie E, Røv N, Steinnes E (2001) Metal dynamics in an Antarctic food chain, *Mar*  
732 *Pollut Bull*, 42, 598– 602
- 733 Orheim O (1972) Volcanic activity on Deception Island, South Shetland Islands In: R.J. Adie  
734 (Ed.), *Antarctic Geology and Geophysics*, Universitetsforlaget, Oslo, pp. 117-120
- 735 Øverjordet IB, Kongsrud MB, Gabrielsen GW, Berg T, Ruus A, Evenset A, Borgå K,  
736 Christensen G, Jenssen BM. (2015) Toxic and essential elements changed in black-legged  
737 kittiwakes (*Rissa tridactyla*) during their stay in an Arctic breeding area, *Sci Total Environ*  
738 502, 548–556
- 739 Pacyna AD, Martínez CZ, Miguélez D, Jiguet F, Polkowska Ż, Wojczulanis-Jakubas K.  
740 (2017) Mercury contamination, a potential threat to the globally endangered aquatic  
741 warbler *Acrocephalus paludicola*, *Environ Sci Pollut Res* 24, 26478- 26484
- 742 Pacyna AD, Kozirowska K, Chmiel S, Mazerski J, Polkowska Ż (2018) Svalbard reindeer as  
743 an indicator of ecosystem changes in the Arctic terrestrial ecosystem, *Chemosphere* 203,  
744 209-218
- 745 Pacyna AD, Frankowski M, Koziół K, Węgrzyn MH, Wietrzyk-Pełka P, Lehmann-Konera S,  
746 Polkowska Ż. (2019) Evaluation of the use of reindeer droppings for monitoring essential  
747 and non-essential elements in the polar terrestrial environment, *Sci Total Environ* 658,  
748 1209-1218
- 749 Petri G, Zauke GP (1993) Trace-metals in crustaceans in the Antarctic ocean. *Ambio* 22, 529-  
750 536

- 751 Phillips RA, Bearhop S, McGill RAR, Dawson DA (2009) Stable isotopes reveal individual  
752 variation in migration strategies and habitat preferences in a suite of seabirds during the  
753 nonbreeding period. *Oecologia* 160, 795–806
- 754 Philpot SM, Lavers JL, Nugegoda D, Gilmour ME, Hutton I, Bond AL. (2019) Trace element  
755 concentrations in feathers of seven petrels (*Pterodroma* spp.). *Environ Sci Pollut Res*, 1-9,  
756 doi.org/10.1007/s11356-019-04406-9
- 757 Quillfeldt P (2002) Seasonal and annual variation in the diet of breeding and non-breeding  
758 Wilson’s storm-petrels on King George Island, South Shetland Islands. *Polar Biol* 25, 216–  
759 221
- 760 Quillfeldt P, McGill RAR, Furness RW (2005) Diet and foraging areas of Southern Ocean  
761 seabirds and their prey inferred from stable isotopes: Review and case study of Wilson’s  
762 storm-petrel. *Mar Ecol Prog Ser* 295, 295–304
- 763 Quillfeldt P, Thorn S, Richter B, Nabte M, Coria N, Masello JF, Massaro M, Neves VC,  
764 Libertelli M (2017) Testing the usefulness of hydrogen and compound-specific stable  
765 isotope analyses in seabird feathers: a case study in two sympatric Antarctic storm-petrels.  
766 *Mar Biol* 164, 192
- 767 R Core Team 2018. A language and environment for statistical computing. Vienna, Austria: R  
768 Foundation for Statistical Computing. <http://www.R-project.org/>
- 769 Rahman MA, Hasegawa H, Lim RP. (2012) Bioaccumulation, biotransformation and trophic  
770 transfer of arsenic in the aquatic food chain, *Environ Res* 116, 118-135
- 771 Rodríguez A, Arcos JM, Bretagnolle V, Dias MP, Holmes ND, Louzao M, Provencher J,  
772 Raine AF, Ramírez F, Rodríguez B, Ronconi RA, Taylor RS, Bonnaud E, Borrelle SB,  
773 Cortés V, Descamps S, Friesen VL, Genovart M, Hedd A, Hodum P, Humphries GRW, Le  
774 Corre M, Lebarbenchon C, Martin R, Melvin EF, Montevecchi WA, Pinet P, Pollet IL,  
775 Ramos R, Russell JC, Ryan PG, Sanz-Aguilar A, Spatz DR, Travers M, Votier SC,



776 Wanless RM, Woehler E and Chiaradia A (2019) Future Directions in Conservation  
777 Research on Petrels and Shearwaters. *Front Mar Sci* 6- 94, doi: 10.3389/fmars.2019.00094  
778 Santos IR dos, Silva-Filho EV, Schaefer C, Sella SM, Silva CA, Gomes V, Passos MJ de  
779 ACR, Ngan PV (2006) Baseline mercury and zinc concentrations in terrestrial and coastal  
780 organisms of Admiralty Bay, Antarctica. *Environ Pollut* 140, 304-311  
781 Santos CSA, Blondel L, Sotillo A, Müller W, Stienen EWM, Boeckx P, Soares AMWM,  
782 Monteiro MS, Loureiro S, de Neve L, Lens L (2017) Offspring Hg exposure relates to  
783 parental feeding strategies in a generalist bird with strong individual foraging  
784 specialization. *Sci Total Environ* 601–602, 1315–1323  
785 Sañudo-Wilhelmy SA, Olsen KA, Scelfo JM, Foster TD, Flegal AR (2002) Trace metal  
786 distributions off the Antarctic Peninsula in the Weddell Sea. *Mar Chem* 77, 157-170  
787 Scheuhammer AM, Meyer MW, Sandheinrich MB, Murray MW (2007). Effects of  
788 environmental methylmercury on the health of wild birds, mammals, and fish. *AMBIO J.*  
789 *Hum. Environ.* 36, 12-19  
790 Seco Pon JP, Beltrame O, Marcovecchio J, Favero M, Gandini P. (2011) Trace metals (Cd,  
791 Cr, Cu, Fe, Ni, Pb, and Zn) in feathers of black-browed albatross *Thalassarche*  
792 *melanophrys* attending the Patagonian Shelf, *Mar. Environ. Res.* 72, 40–45  
793 Selin NE, Jacob DJ, Yantosca, RM, Strode S, Jaeglé L, Sunderland, EM (2008). Global 3D  
794 land- ocean- atmosphere model for mercury: Present- day versus preindustrial cycles and  
795 anthropogenic enrichment factors for deposition. *Global biogeochemical cycles*, 22(2).]  
796 Tartu S, Goutte A, Bustamante P, Angelier F, Moe B, Clément-Chastel C, Bech C, Gabrielsen  
797 GW, Bustnes JO, Chastel O (2013) To breed or not to breed: endocrine response to  
798 mercury contamination by an Arctic seabird. *Biol Lett* 9, 20130317

799 Tartu S, Bustamante P, Goutte A, Cherel Y, Weimerskirch H, Bustnes JO, Castel O (2014)  
800 Age-Related Mercury Contamination and Relationship with Luteinizing Hormone in a  
801 Long-Lived Antarctic Bird. PLoS ONE 9, e103642

802 Tartu S, Angelier F, Wingfield JC, Bustamante P, Labadie P, Budzinski H, Weimerskirch H,  
803 Bustnes JO, Chastel O. (2015) Corticosterone, prolactin and egg neglect behavior in  
804 relation to mercury and legacy POPs in a long-lived Antarctic bird, Sci Total Environ 505,  
805 180-188

806 Thompson DR, Furness RW (1989) Comparison of the levels of total and organic mercury in  
807 seabird feathers. Mar Pollut Bull 20, 557–579

808 Thompson DR, Bearhop S, Speakman JR, Furness RW (1998) Feathers as a means of  
809 monitoring mercury in seabirds: Insights from stable isotope analysis. Environ Pollut 101,  
810 193-200

811 Wang J, Xie Z, Wang F, Kang H (2017) Gaseous elemental mercury in the marine boundary  
812 layer and air-sea flux in the Southern Ocean in austral summer. Sci Total Environ 603-604,  
813 510-518

814 Wasilewski A (1986) Ecological aspects of the breeding cycle in the Wilson's storm petrel,  
815 *Oceanites oceanicus* (Kuhl), at King George Island (South Shetland Islands, Antarctica).  
816 Polish Polar Res 7, 173–216

817 Webster MS, Marra PP, Haig SM, Bensch S, Holmes RT (2002) Links between worlds:  
818 unravelling migratory connectivity. Trends Ecol Evol 17, 76–83

819 Wolfe MF, Schwarzbach S, Sulaiman RA. (1998) Effects of mercury on wildlife: a  
820 comprehensive review. Environ. Toxicol Chem 17, 146-160

821

822

823

824

825

**Table 1.** Elemental concentrations of the studied elements in feathers of storm-petrels, mean  $\pm$  SD (min–max)  $\mu\text{g g}^{-1}$  dw, N = the number of individuals sampled, LOD = detection limit, LOQ= quantification limit, Se:Hg – Se:Hg molar ratio

Element	Adult WSP (N=12)	Adult BBSP (N=4)	Chick down WSP (N=4)	Chick body WSP (N=4)
As	75% <LOD	100%<LOQ	1.76 $\pm$ 1.10 (0.45–3.16)	6.11 $\pm$ 1.43 (4.77–8.38)
Bi	92%<LOD	100%<LOD	1.57 $\pm$ 1.78 (<LOD–3.92)	14.16 $\pm$ 5.92 (6.50–20.50)
Ca	96.0 $\pm$ 21.0 (77.7–156.0)	80.0 $\pm$ 11.2 (72.8– 99.3)	242.8 $\pm$ 60.4 (165.0–326.0)	136.50 $\pm$ 11.79 (124.9– 156.0)
Cd	<LOD (<LOD– 0.45)	<LOQ	0.45 $\pm$ 0.22 (<LOQ–0.68)	0.51 $\pm$ 0.24 (<LOD– 0.70)
Cr	0.67 $\pm$ 0.45 (0.11–4.36)*	0.71 $\pm$ 0.45 (0.09– 1.22)	1.54 $\pm$ 0.48 (0.88–2.08)	3.46 $\pm$ 0.78 (2.73–4.78)
Cu	2.52 $\pm$ 2.35* (<LOD–13.9)	8.12 $\pm$ 0.56 (7.52– 9.06)	1.52 $\pm$ 0.60 (0.67–2.26)	6.68 $\pm$ 3.15 (2.49–10.96)
Fe	20.40 $\pm$ 18.00 (<LOD–263.0)*	10.73 $\pm$ 5.04 (<LOD–16.16)	74.30 $\pm$ 19.30 (50.80–102.4)	131.7 $\pm$ 81.8 (63.1–270.0)
Mg	478 $\pm$ 130 (315– 773)	429 $\pm$ 101 (306– 529)	538 $\pm$ 354 (316– 1152)	378 $\pm$ 91 (285–513)
Mo	2.41 $\pm$ 4.09 (<LOD–14.6)	0.56 $\pm$ 0.27 (0.16– 0.84)	1.92 $\pm$ 1.77 (0.36–4.90)	7.13 $\pm$ 7.42 (1.52–19.79)
Pb	0.33 $\pm$ 0.37* (<LOD–5.06)	0.36 $\pm$ 0.24 (0.11– 0.74)	1.77 $\pm$ 0.91 (0.75–2.76)	1.43 $\pm$ 1.32 (0.36–3.67)
Sb	100%<LOD	75% <LOD	0.22 $\pm$ 0.15 (<LOD–0.43)	1.17 $\pm$ 0.53 (0.58–1.92)

Element	Adult WSP (N=12)	Adult BBSP (N=4)	Chick down WSP (N=4)	Chick body WSP (N=4)
Se	1.81 ± 0.98 (<LOD–4.65)	5.19 ± 1.18 (3.74– 6.62)	4.06 ± 0.50 (3.29–4.69)	3.63 ± 1.01 (2.39–5.13)
Sr	5.77 ± 3.62 (2.47–13.53)	2.79 ± 1.21 (1.35– 4.58)	23.19 ± 8.75 (13.4–37.1)	9.57 ± 1.020 (8.63–11.27)
Zn	109.20 ± 18.50 (70.3–141)	99.95 ± 13.01 (85.5–120)	48.30 ± 7.08 (40.30–59.60)	93.00 ± 11.30 (79.4–110.8)
Se:Hg	1.92	2.75	6.00	13.90

\* for Cr, Cu, Fe and Pb one outlier was excluded from the mean calculation, but it is shown as the maximal value

**Table 2** Intergroup differences (one-way PERMANOVA, Bonferroni-corrected p values) of elemental concentration,  $\log(x + 1)$  transformed, in the feathers of the four studied groups of storm-petrels: body feathers of adult Wilson's (WSP\_Ad) and back-bellied (BBSP\_Ad) storm-petrels, down from Wilson's storm-petrel chicks (WSP\_down) and body feathers from Wilson's storm petrel fledglings (WSP\_CHF)

PERMANOVA, F = 11.48, p = 0.0001				
All elements	WSP_CHF	WSP_down	BBSP_Ad	WSP_Ad
WSP_CHF	-	0.182	0.176	0.007
WSP_down		-	0.157	0.004
BBSP_Ad			-	0.037
WSP_Ad				-
PERMANOVA, F = 9.26, p = 0.0001				
Essential	WSP_CHF	WSP_down	BBSP_Ad	WSP_Ad
WSP_CHF	-	0.160	0.181	0.003
WSP_down		-	0.166	0.005
BBSP_Ad			-	0.136
WSP_Ad				-
PERMANOVA, F = 12.93, p = 0.0001				
Non-essential	WSP_CHF	WSP_down	BBSP_Ad	WSP_Ad
WSP_CHF	-	0.160	0.181	0.003
WSP_down		-	0.166	0.012
BBSP_Ad			-	0.043
WSP_Ad				-

**Table 3** Sources of variability (average percentage dissimilarity) in the elemental concentrations ( $\log(x+1)$  transformed) in: body feathers of adult Wilson's (WSP\_Ad) and back-bellied (BBSP\_Ad) storm-petrels, down from Wilson's storm-petrel chicks (WSP\_down) and body feathers from Wilson's storm petrel fledglings (WSP\_CHF), according to the SIMPER analysis. Only elements with a contribution > 10% are shown. ADis - Average Dissimilarity, Contr. (%) – percentage contribution, Overall - overall average similarity

Overall		WSP_Ad vs				WSP_Ad vs				WSP_Ad vs					
dissimilarity		WSP_CHF				WSP_down				BBSP_Ad					
ADis		Contr.		ADis		Contr.		ADis		Contr.		ADis		Contr.	
All elements															
Fe	2.57	13.9	Bi	4.18	19.1	Fe	2.67	14.0	Cu	2.12	16.1				
Bi	2.07	11.2	As	3.06	14.0	Sr	2.36	12.4	Fe	1.94	14.7				
			Fe	3.02	13.8				Hg	1.89	14.3				
									Se	1.63	12.3				
Overall	18.46			21.87			19.30			13.23					
Essential															
Fe	3.10	21.9	As	3.72	22.6	Fe	3.24	21.5	Cu	2.43	22.8				
As	2.06	14.5	Fe	3.68	22.4	Ca	1.94	12.9	Fe	2.22	20.8				
Cu	1.83	12.9	Mo	2.40	14.6	As	1.85	12.3	Se	1.87	17.6				
Mo	1.83	12.9	Cr	1.87	11.3	Zn	1.71	11.3	Mo	1.41	13.3				
			Cu	1.82	11.0	Mo	1.66	11.0							
Overall	14.16			16.46			15.07			10.65					
Non-Essential															
Bi	11.47	29.2	Bi	23.67	50.3	Sr	13.96	36.4	Hg	14.42	49.1				

	Overall		WSP_Ad vs				WSP_Ad vs				
	dissimilarity		WSP_CHF				WSP_down				
	ADis	Contr.	ADis	Contr.	ADis	Contr.	ADis	Contr.			
Sr	10.01	25.5	Sb	6.70	14.2	Pb	7.43	19.4	Sr	8.68	29.5
Hg	7.02	17.9	Sr	6.26	13.3	Bi	7.27	19.0	Pb	4.74	16.1
Pb	5.59	14.2	Pb	5.38	11.4	Hg	5.25	13.7			
Overall	39.31			47.09			38.33			29.41	



**Table 4** Variability of mercury (Hg) levels in feathers of *Procellariiformes*. N – number of individuals


Species	Study area	Tissue	N	Age	Concentrat. mean± SD µg g <sup>-1</sup> dw	Reference
<b>Antarctic prion</b> ( <i>Pachyptila desolata</i> )	Kerguelen	body feathers	10	unknown	2.8±1.2	Fromant et al. 2016
	archipelago		10	adult	1.73±0.50	
<b>White-headed petrel</b> ( <i>Pterodroma lessonii</i> )	Kerguelen	body feathers	10	adult	12.43 ± 2.01	Carravieri et al. 2014
	archipelago		10	chicks	1.54 ± 0.34	
<b>spectacled petrel</b> ( <i>Procellaria conspicillata</i> )	southwestern Atlantic	contour	38	unknown	11.17 ± 3.78	Carvalho et al. 2013
	Ocean off the Brazilian coast	feathers				
<b>white-chinned petrel</b> ( <i>Procellaria aequinoctialis</i> )	Kerguelen	body feathers	22	unknown	7.63 ± 3.87	Cipro et al. 2014
	archipelago					
	southwestern Atlantic	contour	9	adults	3.45 ± 2.84	Carvalho et al. 2013
	Ocean off the Brazilian coast	feathers	21	juveniles	1.14 ± 2	
	Kerguelen	body feathers	14	chicks	1.82 ± 0.51	Blévin et al. 2013
	archipelago					
<b>Leach's storm-petrel</b>	Machias Seal Island,	Breast	15	adult	7.01*	Bond and Diamond,
	New Brunswick,	feathers	20	chicks	1.42*	

<i>(Oceanodroma leucorhoa)</i>	Canada						2009
<b>Wilson's storm-petrel</b>	Kerguelen Islands	body feathers	12	adult		$0.42 \pm 0.13$	Carravieri et al. 2014
<i>(Oceanites oceanicus)</i>	King Gorge Island,	body feathers	25	adult		$2.38 \pm 1.47$	present
	South Shetlands,	body feathers	5	chick		$0.67 \pm 0.27$	study
	Antarctica	down	16	chick		$1.72 \pm 0.65$	
<b>Black-bellied storm-petrel</b>	Kerguelen Islands	body feathers	10	adult		$4.22 \pm 2.53$	Carravieri et al. 2014
<i>(Fregetta tropica)</i>	King Gorge Island,	body feathers	8	adult		$5.47 \pm 1.61$	present
	South Shetlands,	body feathers	5	chick		$1.87 \pm 0.29$	study
	Antarctica	down	6	chick		$3.99 \pm 1.07$	


\* *Estimated marginal mean*

# Figure


[Click here to download Figure: Fig.1.doc](#)



**Fig. 1** Range of the studied species and possible areas of elemental input: triangle – study area, King George Island (KGI), grey rectangles – moulting latitudes for adult storm-petrels (dotted for black-bellied storm-petrel (BBSP) and solid for Wilson's storm-petrel (WSP); according to isotopic data from Quillfeldt et al. 2005 and Phillips et al. 2009 calculated based on equation proposed by Quillfeldt et al. 2005). Storm-petrels non-breeding range map source: BirdLife International and Handbook of the Birds of the World 2018. Photos by DJ



**Fig. 2** Concentration of Hg ( $\mu\text{g}\cdot\text{g}^{-1}$  dw) in feathers of the six studied groups of storm-petrels: body feathers of adult Wilson's (WSP\_Ad) and back-bellied (BBSP\_Ad) storm-petrels, down from Wilson's (WSP\_down) and black-bellied (BBSP\_down) storm-petrel chicks and body feathers from Wilson's (WSP\_CHF) and black-bellied (BBSP\_CHF) storm-petrel chicks. Boxplots show the median (band inside the box), the first (25%) and third (75%) quartile (box), and the lowest and the highest values within 1.5 interquartile range (whiskers)



**Fig. 3** Hierarchical dendrogram of the studied elements in the feathers of the studied storm-petrels (all age and feather type groups combined), obtained using a paired group method and Euclidean distance matrix (the distance reflects degree of association between different elements). Numbers below branches indicate bootstrap probability values (bootstrap  $n = 1000$ ). Clusters with bootstrap support  $\geq 95$  denoted with a dotted rectangle.

**Supplementary material for on-line publication only**

**[Click here to download Supplementary material for on-line publication only: Supplementary materials1.doc](#)**

**Supplementary material for on-line publication only**

**[Click here to download Supplementary material for on-line publication only: Supplementary materials2.doc](#)**

**Supplementary material for on-line publication only**

[Click here to download Supplementary material for on-line publication only: Supplementary materials3.doc](#)