

Offshore surface waters of Antarctica are free of microplastics, as revealed by a circum-Antarctic study.

Piotr Kuklinski ^{a,b,c,*}, Leszek Wicikowski ^d, Mariusz Koper ^c, Tomasz Grala ^c, Hanna Leniec-Koper ^c, Michał Barasiński ^c, Magdalena Talar ^c, Ireneusz Kamiński ^{c,e}, Robert Kibart ^c, Wojciech Małecki ^c

^a *Institute of Oceanology, Polish Academy of Sciences, Powstancow Warszawy 55, Sopot 81-712, Poland.*

^b *Natural History Museum, Department of Life Sciences, London SW7 5BD, United Kingdom*

^c *Katharsis II, World Ocean*

^d *Technical University of Gdansk, Faculty of Applied Physics and Mathematics, Narutowicza 11/12, 80-233 Gdansk, Poland*

^e *Polish Naval Academy, ul. inż. Śmidowicza 69, 81-127 Gdynia, Poland*

* Corresponding author at: Institute of Oceanology, Polish Academy of Sciences, Powstancow Warszawy 55, Sopot 81-712, Poland

E-mail address: kuki@iopan.pl (P.Kuklinski)

ABSTRACT

In 2018, during a circumnavigation of Antarctica below 62° S by the sailing boat Katharsis II, the presence of plastics was investigated with surface sampling nets at ten evenly spaced locations (every 36° of longitude). Although fibres that appeared to be plastic (particles up to 2 cm) were found in numbers ranging from 1 particle (0.002 particles per m³) to 171 particles (1.366 particles per m³) per station, a Fourier-transform infrared spectroscopy (FT-IR) analysis indicated that these particles were not composed of plastic. The fibres which superficially reminded plastic were composed of silica and are of biological origin most likely generated by phytoplankton (diatoms). Therefore, the offshore Antarctic locations were proven to be free of floating microplastics.

Key words: Antarctic, Southern Ocean, pollution, microplastic, surface waters

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Currently, pollution caused by plastics is the focus of media attention. This coverage has very efficiently increased the public's awareness of the problem. There is no doubt that it is time to have a global discussion about the influence of plastics on our ecosystems, as plastics have been discovered in every investigated location. In this century, plastics have been discovered in the most remote marine habitats on the planet, including deep-sea sediments (Tekman et al. 2017) and Arctic ice (Bergmann et al. 2017). With current knowledge, there are still debates about whether plastics have an impact on marine life (see Guzzetti et al. 2018). Plastics were found to be ingested by various organisms, including both invertebrates and vertebrates (e.g., Wright et al 2013, Besseling et al. 2015, Law 2017, Fang et al. 2018, Hernandez-Gonzalez et al. 2018, McGoran et al. 2018). Marine biota were found to be entangled in floating macroplastic products (Barnes et al. 2010, Waluda and Staniland 2013). Tiny fragments of degraded plastics can carry toxins (Guzzetti et al. 2018). As plastics are ingested by organisms from various trophic levels, they might be transferred through food chains (Wright et al 2013, Guzzetti et al. 2018). Therefore, plastics is a quickly emerging problem for humans as a broad pollution issue (Wright et al 2013, Law 2017, Guzzetti et al. 2018).

Antarctic is still considered the most pristine environment on Earth. Humans' impact here is still probably the lowest on the planet, as our density in that area is low and very localized (mostly limited to research bases). In addition, the Antarctic continent, which is surrounded by the Southern Ocean, is isolated from the rest of the world by a convergence zone (a transition between cold Antarctic surface waters and warmer sub-Antarctic water masses), which effectively bars marine life and pollutants exchange from more industrialized areas in lower latitudes (Foster 1984, Chown et al. 2015). One would believe that these factors make the area immune to human impact. Although recent studies indicate that Southern Ocean is not biologically isolated as previously it was believed. It was found that storm-forced surface waves and ocean eddies can enhance oceanographic connectivity between Antarctica and mid-latitudes (Fraser et al. 2018). Since the 1960s, DDT (dichlorodiphenyltrichloroethane) and, later, PCBs (polychlorinated biphenyls) have been found in Antarctica (Risebrough et al. 1976, Bonner 1984). Some pollutants are global in nature, and plastics seem to be one of them. Indeed, plastics are being discovered around the globe, including in the Arctic and Antarctic marine environments (see Lusher et al. 2015, Isobe et al. 2017, Waller et al. 2017 or Kanhai et al. 2018 and the references therein).

In recent years, plastics were not only discovered at locations in Antarctic, such as the Antarctic Peninsula, where the human population is highest and the area is closest to other

1 continents (South America) (see Waller et al. 2017), but also in such remote locations such as
2 in the inner basin of the Ross Sea (Cincinelli et al. 2017, Munari et al. 2017). The highest
3 density of plastics in Antarctic was recorded around the Antarctic Peninsula (Waller et al.
4 2017), which is most likely due to the effect of the higher sampling effort in the area
5 compared to the rest of the Southern Ocean. However, we cannot rule out that this
6 observation is the effect of higher human activity in that sector.
7
8
9

10
11 The goal of this investigation was to study the potential presence of microplastic
12 particles (particles less than 5 mm in size, as defined by the National Oceanic and
13 Atmospheric Administration [NOAA]) in the surface waters of the Southern Ocean. The
14 study aimed to locate sampling sites around the whole continent, ranging from areas with
15 relatively higher human populations, such as the Antarctic Peninsula (due to research, fishing
16 and tourism), to areas where humans were likely never present or were only sporadically
17 present: offshore open ocean.
18
19
20
21
22
23
24

25 This investigation collected samples during the first ever successful human attempt to
26 circumnavigate the Antarctic continent south of the 62nd parallel by sailing boat (Katharsis
27 II). The cruise started on 23.12.2017 in Cape Town, South Africa and finished on 05.04.2018
28 in Hobart, Australia. Ten sampling locations were selected within the convergent zone
29 between 63° and 69° S (Fig. 1, Table 1). The first station was selected when the boat crossed
30 63° S, and weather-permitting, evenly distributed samples were taken around the Antarctic
31 continent at approximately every 36° of longitude (Fig. 1, Table 1).
32
33
34
35
36
37
38

39 Samples were obtained by using a specifically designed Hydro-Bios net for
40 microplastic sampling with lifting buoys (product no. 438 214). The opening of the net was
41 70 x 40 cm, and the mesh size was 300 µm. We aimed for 30 minutes of net towing behind
42 the boat at 1 to 2 knots, depending on wind conditions, but everything had to be modified
43 according to the ice, wind and algal bloom conditions. At stations 3, 4 and 6, algal blooms
44 clogged the net; therefore, the net was towed for 10 minutes. The date, time and location of
45 start and end point of the net towing as well as speed of the boat during the sampling were
46 recorded with GPS, enabling the estimation of the water volume sampled by the net. All these
47 data are summarized in Table 1.
48
49
50
51

All the samples were concentrated on board the boat immediately after sampling by
filtering the collected samples through a metal sieve (mesh size of 200 µm) and then by using
a metal funnel placed in a glass borax bottle. Although these bottles had plastic leads, they

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

were bright blue, which made any contamination easy to recognize. Therefore we took all possible precautions to avoid self-contamination with plastic. The water samples were preserved in 2% formalin.

All the samples taken in Antarctic were analysed in the laboratory by a single investigator to ensure consistent treatment and no introduction of bias into the analysis. In the laboratory, all the samples were treated with hydrogen peroxide (30% H₂O₂), which is an oxidizing agent that enables the digestion of organic matter with no degradation of plastic polymers (Nuelle et al. 2014). The samples treated with hydrogen peroxide were kept in the oven for 24 hours at 60° C and then left in the bottles for one week at room temperature to ensure the digestion of the largest amount of organic matter. The samples were examined visually by using a stereomicroscope equipped with a fluorescent light, enabling the efficient exclusion of organic particles (Qiu et al. 2016). The microplastic contamination was quantified by counting. Fibres which were found in the samples were measured using an eye piece with graticules (± 0.1 cm) and were classified according to size: (i) 0-0.5 cm, (ii) 0.5-1.0 cm, and (iii) longer than 1 cm.

Laboratory equipment made only of glass and metal was used to ensure the lack of contamination by local sources of plastics. The analysis was performed only when the person analysing the samples was present alone in the laboratory. This procedure was also performed to reduce the probability of the potential contamination of the samples. Additionally, before each Antarctic sample was analysed, a control sample was prepared by using local tap water and all laboratory equipment used in the Antarctic sample analyses. All procedures were performed in the laboratory where the Antarctic samples were analysed. Such an approach was applied to analyse whether the laboratory conditions and the analytic procedures contaminated the samples with plastic particles. During that procedure, no particles of any type were detected.

A Fourier-transform infrared spectroscopy (FT-IR) analysis of the samples was performed with a Frontier FT-IR (PerkinElmer) instrument. The measurements were carried out in an attenuated total reflectance (ATR) configuration with ZnSe cells. The tests were performed at 25 °C in dry air. Particles were identified by comparing the FT-IR absorbance spectra of the microplastics to those in a self-collected, polymer reference library.

Generally, the whole methodology followed that described in Nuelle et al. (2014), Qiu et al. (2016), and Prata et al. (2019).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

In total, 3147.7 m³ of surface water was filtered by the deployed net at the ten sampling sites (Table 1). Within this water volume, 488 transparent fibres that appeared to be plastic were recorded (Table 2). Fibres were present at all ten investigated locations (Table 2). The fibres glowed during the examination with fluorescent light, which suggested that they were made of plastic. Although this methodology enables the exclusion of organic particles, Fourier-transform infrared spectroscopy (FT-IR) analysis confirmed that these particles were not plastic fibres. All together 107 particles were randomly selected from the samples and analysed with FT-IR. The fibres that often exceeded the defined microplastic size (above 5 µm) ranged from 1 particles (0.002 particles per m³) to 171 particles (1.366 particles per m³) per station with overall mean concentration for all the stations 48.8 ± 62.81 (standard deviation) (0.262 ± 0.435 particles per m³) (Table 2). Detailed analysis of obtained FT-IR spectrum of the fibres indicated organic origin (silica) of these (Fig. 2). And as comparison of the FT-IR spectrum with literature source shows (Camargo et al. 2016) these fibres originate from planktonic algae – diatoms which could generate those types of structures. Therefore fibre density and size from station to station probably exhibits local abundance of these organisms during period of our sampling.

Based on the obtained results, as sampling sites were rather evenly spread around the Antarctic continent (Fig. 1, Table 1), we can assume that microplastics might still be absent in the offshore marine system of Antarctica, at least at the locations covered by this study. Even the area around the Antarctic Peninsula, which has the most tourist boat traffic, the highest density of research bases (Hansom and Gordon 1998) and is potentially the area most polluted with all types of plastic, had no traces of microplastic in the samples. Yet we had just one sampling station next to Antarctic Peninsula (station no 7) therefore more intense sampling effort is needed to confirm whether surface offshore waters are free of microplastic,

Our results are surprising, as each previous investigation of the Southern Ocean recorded the presence of microplastics. The locations where microplastics were recorded in the Southern Ocean range from along the Antarctic Peninsula to remote locations in the Ross Sea as well as the open waters of the Southern Ocean (see Cincinelli et al. 2017, Isobe et al. 2017, Munaria et al. 2017, Waller et al. 2017, Reed et al. 2018). Even a study by Isobe et al. (2017), which included sampling the open waters of the Southern Ocean with a similar methodology to ours (i.e., a surface net and a comparable volume of water sampled), recorded the presence of plastics. At locations below 60° S, Isobe et al. (2017) recorded concentrations of particles ranging from 0.046 to 0.099 particles per m³ at selected locations. Generally, those values are

1 rather low compared with the amount of microplastics found elsewhere in the world ocean,
2 especially in marine zones close to more industrialized areas (see Wright et al 2013, Isobe et
3 al. 2017 and the references therein). Currently, plastics in Antarctica, especially
4 microplastics, are mostly concentrated close to locations with direct, intense human activities
5 (e.g., the Antarctic Peninsula) (Waller et al. 2017). Therefore more investigations are needed
6 to recognize scale of this pollution in the Southern Ocean. Additionally, this study indicates
7 that materials that visually appear to be plastic must be analysed with specialized equipment
8 (e.g., FT-IR) before drawing final conclusions concerning the composition of the material.
9
10
11
12
13
14

15 The sampling took place during the successful, first-ever circumnavigation of the
16 Antarctic continent below 62° S by sailing boat. This study is an example of a positive
17 citizen-science activity that can generate valuable data.
18
19
20
21
22
23

24 **Acknowledgements**

25
26 We would like to thank all the people who helped organize this exceptional expedition,
27 enabling us to not only achieve the Guinness World Record as the first circumnavigation of
28 Antarctica south of the 62nd parallel in a sailboat but also, most importantly, gather valuable
29 scientific data. We would also like to thank the anonymous reviewers who greatly improved
30 the quality of the manuscript.
31
32
33
34
35
36
37
38

39 **Funding**

40
41 This study was commenced during the PANIC project (grant no. 2016/23/B/ST10/01936)
42 funded by the National Science Centre, Poland.
43
44
45
46
47

48 **Author contributions**

49
50 PK conceived of and designed the study and analysed the data. PK, MK, TG, HL, MB, MT,
51 IK, RK and WM collected samples. LW performed the Fourier-transform infrared
spectroscopy (FT-IR) analysis. The paper was written by PK and LW and was approved by
all authors: PK, MK, TG, HL, MB, MT, IK, RK and WM.

Conflicts of interest

The authors express no conflicts of interest, neither direct nor indirect, in the publication of this manuscript. All funding sources have been referenced in the manuscript.

Author declaration

All authors have approved the submission of this manuscript, which is an original work conducted by all the authors and has not been published prior to its submission. It is not under consideration by any other publishers.

References

- Barnes, D.K.A., Walters, A., Gonçalves, L., 2010. Macroplastics at sea around Antarctica. *Mar. Environ. Res.* 70, 250–252.
- Bergmann, M., Peeken, I., Beyer, B., Krumpen, T., Primpke, S., Tekman, M.B., Gerdts, G., 2017. Vast quantities of microplastics in Arctic sea ice—a prime temporary sink for plastic litter and a medium of transport. *MICRO 2016*. Elsevier, pp. 75–76.
- Besseling, E., Foekema, E.M., Van Franeker, J.A., Leopold, M.F., Kühn, S., Bravo Rebolledo, E.L., Heße, E., Mielke, L., Ijzer, J., Kamminga, P., Koelmans, A.A., 2015. Microplastic in a macro filter feeder: Humpback whale *Megaptera novaeangliae*. *Mar. Pollut. Bull.* 95, 248-252.
- Bonner, W.N., 1984. Conservation and the Antarctic. In *Antarctic Ecology Vol. 2*. Edited R., M., Laws, Academic Press, London. Pp. 821 – 850.
- Camargo, E., Jaime, P.C.J., Lin, C-F., Lin, M-S., Yu, T-Y, Wu, M-C., Lai, S-Y., Wang, M-Y, 2016. Chemical and optical characterization of *Psammodictyon panduriforme* (Gregory) Mann comb. nov. (Bacillariophyta) frustules. *Opt. Mater. Express* 6: 1437.
- Cincinelli, A., Scopetani, C., Chelazzi, D., Lombardini, E., Martellini T., Katsoyiannis A., Fossi, M.C., Corsolini, S., 2017. Microplastic in the surface waters of the Ross Sea (Antarctica): Occurrence, distribution and characterization by FTIR. *Chemosphere* 175, 391-400.

1 Chown, S.L., Clarke, A., Ceridwen I. Fraser, C.I., Cary, S.C, Moon K.L., McGeoch M.A.,
2 2015. The changing form of Antarctic Biodiversity. *Nature* 522, 431-438.

3
4 Fang, C., Zheng, R., Zhang, Y., Hong, F., Mu, J., Chen, M., Song, P., Lin, L., Lin, H., Le, F.,
5 Bo, J., 2018. Microplastic contamination in benthic organisms from the Arctic and sub-Arctic
6 regions. *Chemosphere* 209, 298-306.

7
8
9
10
11 Foster, T.D., 1984. The marine environment. In *Antarctic Ecology Vol. 2*. Edited by R., M.,
12 Laws, Academic Press, London. Pp. 345–371.

13
14
15
16 Fraser, C.I., Morrison, A.K., McC Hogg, A., Macaya, E.C., van Sebille, E., Ryan, P.G.,
17 Padovan, A., Jack, C., Valdivia, N., Waters, J.M., 2018. Antarctica's ecological isolation will
18 be broken by storm-driven dispersal and warming. *Nat. Clim. Chang.* 8, 704-708.

19
20
21
22
23 Hansom, J.D., Gordon, J.E., 1998. Antarctic environments and resources a geographical
24 perspective. Longman, London. Pp 402.

25
26
27
28 Hernandez-Gonzalez, A., Saavedra, C., Gago, J., Covelo, P., Santos, B.M., Pierce, G.J., 2018.
29 Microplastics in the stomach contents of common dolphin (*Delphinus delphis*) stranded on
30 the Galician coasts (NW Spain, 2005–2010). *Mar. Pollut. Bull.* 137, 526-532.

31
32
33
34
35 Guzzetti, E., Sureda, A., Tejada, S., Faggio, C., 2018. Microplastic in marine organism:
36 Environmental and toxicological effects. *Environ. Toxicol. Pharmacol.* 64, 164-171.

37
38
39
40
41 Isobe, A., Uchiyama-Matsumoto, K., Uchida, K., Tokai, T., 2017. Microplastics in the
42 Southern Ocean. *Mar. Pollut. Bull.* 114, 623-626.

43
44
45
46
47
48
49
50
51
Kanhai L.K., Gårdfeldt K., Lyashevskaya, O., Hassellöv M., Thompson, R.C., O'Connor I.,
2018. Microplastics in sub-surface waters of the Arctic Central Basin. *Mar. Pollut. Bull.* 130,
8-18.

Law, K.L., 2017. Plastics in the marine environment. *Annu. Rev. Mar. Sci.* 9, 205–229.

Lusher, A.L., Tirell, V., O'Connor, I., Officer, R., 2015. Microplastics in Arctic polar waters:
the first reported values of particles in surface and sub-surface samples. *Sci. Rep.* 5, 14947.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
McGoran, A.R., Cowie, P.R., Clark, P.F., McEvoy, J.P., Morritt, D., 2018. Ingestion of plastic by fish: A comparison of Thames Estuary and Firth of Clyde populations. *Mar. Pollut. Bull.* 137, 12-23.

Munari, C., Infantini, V., Scoponi, M., Rastelli, E., Corinaldesi, C., Mistri, M., 2017. Microplastics in the sediments of Terra Nova Bay (Ross Sea, Antarctica). *Mar. Pollut. Bull.* 122, 161-165.

Nuelle, M.T., Dekidd, J.H., Remy, D., Fries, E., 2014. A new analytical approach for monitoring microplastics in sediments, *Environ. Pollut.* 184, 161-169.

Qiu, Q., Tan, Z., Wang, J., Peng, J., Li, M., Zhan, Z., 2016. Extraction, enumeration and identification methods for monitoring microplastics in the environment, *Estuar. Coast Shelf Sci.* 176, 102-109.

Prata, J.C., da Costa, J.P., Duarte, A.C., Rocha-Santos, T., 2019. Methods for sampling and detection of microplastics in water and sediment: A critical review. *Trend. Anal. Chem.* 110, 150-159.

Reed, S., Clark, M., Thompson, R., Kevin Hughes, K.A., 2018. Microplastics in marine sediments near Rothera Research Station, Antarctica. *Mar. Pollut. Bull.* 133, 460-463.

Risebrough, R.W., Walker, W., Schmidt, T.T., De Lappe, B.W., Connors, C.W., 1976. Transfer of chlorinated biphenyls to Antarctica. *Nature* 264, 738-739.

Waller, C.L., Griffiths, H.J., Waluda, C.M., Thorpe, S.E., Loaiza, I., Moreno, B., Pacherres, C.O., Hughes, K.A., 2017. Microplastics in the Antarctic marine system: an emerging area of research. *Sci. Total Environ.* 598, 220-227.

Waluda, C.M., Staniland, I.J., 2013. Entanglement of Antarctic fur seals at Bird Island, South Georgia. *Mar. Pollut. Bull.* 74, 244-252.

Wright, S.L., Thompson, R.C., Galloway, T.S., 2013. The physical impacts of microplastics on marine organisms: A review. *Environ. Pollut.* 178, 483-492.

Tekman, M.B., Krumpen, T. , Bergmann, M., 2017. Marine litter on deep Arctic sea floor continues to increase and spreads to the North at the HAUSGARTEN observatory. Deep-Sea Res Part I 120, 88–99.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

1
2
3
4
5
6
Fig. 1. Schematic view of the expedition route of the sailing boat Katharsis II with the location of sampling sites (for detailed coordinates, see Table 1), map sourced and modified from <https://data.aad.gov.au>.

7
8
9
10
11
12
13
14
Fig 2. Chemical and optical characterization of fibres sampled during this study (black line) and phytoplankton group Bacillariophyta - diatoms (grey line) obtained from literature (Camargo et al. 2016). Spectrum obtained with use of a Fourier-transform infrared spectroscopy in an attenuated total reflectance.

15
16
17
18
19
20
21
Table 1. Sampling locations with detailed timing information, the coordinates of the starting and ending points of the net deployment and the volume of samples (n/a – data not available due to a GPS malfunction).

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
Table 2. Quantitative fibre spectrum sizes found at each sampling location. Fibres within the size class “longer than 1 cm” never exceeded 2 cm (n/a – data not available due to a GPS malfunction, for details on the sampling sites see Fig. 1 and Table 1).

Table 1

Station no	Date of sampling	Latitude	Longitude	Time (local)	Volume of water filtered by sampling net [m³]
1	10.01.2018	start: 63° 27' 31.0" S end: 63° 27' 35.7" S	start: 82° 47' 08.2" E end: 82° 49' 05.1" E	16:25 16:57	448.0
2	21.01.2018	start: 63° 49' 22.8" S end: 63° 49' 28.6" S	start: 118° 43' 05.5" E end: 118° 44' 48.1" E	08:28 08:58	392.0
3	28.01.2018	start: 65° 32' 30.9" S end: 65° 31' 33.7" S	start: 155° 28' 50.8" E end: 155° 28' 48.6" E	05:27 05:42	504.0
4	03.02.2018	start: 68° 42' 45.9" S end: 68° 42' 49.3" S	start: 171° 48' 40.0" W end: 171° 48' 05.7" W	16:42 16:53	111.4
5	08.02.2018	start: n.a. end: 69° 45' 25.7" S	start: n.a. 133° 09' 18.4" W	15:30 16:00	n.a.
6	14.02.2018	start: 67° 17' 44.9" S end: 67° 17' 57.5" S	start: 98° 21' 17.7" W end: 98° 20' 59.3" W	05:29 05:41	125.2
7	23.02.2018	start: 63° 23' 42.2" S end: 63° 23' 35.5" S	start: 60° 53' 56.4" W end: 60° 52' 20.6" W	04:23 04:54	364.0
8	02.03.2018	start: 65° 39' 07.2" S end: 65° 38' 47.7" S	start: 24° 05' 59.8" W end: 24° 05' 07.7" W	03:20 03:50	251.2
9	08.03.2018	start: 67° 47' 48.2" S end: 67° 48' 33.8" S	start: 11° 15' 18.4" E end: 11° 16' 18.1" E	16:34 17:05	448.0
10	17.03.2018	start: 63° 42' 21.7" S end: 63° 41' 57.1" S	start: 46° 49' 42.3" E end: 46° 51' 41.4" E	04:29 05:03	504.0

Table 2

Station no	Fibres < 0.5 cm (per m ³)	Fibres 0.5 - 1.0 cm (per m ³)	Fibres > 1.0 cm (per m ³)
1	1 (0.002)		
2		74 (0.188)	5 (0.012)
3	1 (0.001)		2 (0.003)
4	6 (0.053)		
5		19 (n.a.)	1 (n.a.)
6		171 (1.366)	
7			1 (0.002)
8		162 (0.645)	
9		14 (0.031)	2 (0.004)
10	19 (0.037)	10 (0.019)	
Total number of fibres (per m³)	27 (0.091)	450 (2.249)	11 (0.021)

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)

