



Generic invariant measures for iterated systems of interval homeomorphisms

WOJCIECH CZERNOUS AND TOMASZ SZAREK

To the memory of Professor Józef Myjak.

Abstract. It is well known that iterated function systems generated by orientation preserving homeomorphisms of the unit interval with positive Lyapunov exponents at its ends admit a unique invariant measure on $(0, 1)$ provided their action is minimal. With the additional requirement of continuous differentiability of maps on a fixed neighbourhood of $\{0, 1\}$, we present a metric in the space of such systems which renders it complete. Using then a classical argument (and an alternative uniqueness proof), we show that almost singular invariant measures are admitted by systems lying densely in the space. This allows us to construct a residual set of systems with unique singular stationary distribution. Dichotomy between singular and absolutely continuous unique measures is assured by taking a subspace of systems with absolutely continuous maps; the closure of this subspace is where the residual set is found.

Mathematics Subject Classification. 37E05, 60G30, 37C20.

Keywords. Markov operators, Semigroups of interval homeomorphisms, Absolute continuity, Singularity, Minimal actions.

1. Introduction. Ergodic properties of random dynamical systems have been extensively studied for many years (see [3] and the references therein). The most important concepts related to these properties are attractors and invariant measures (see [4, 6, 10, 11]). Among random systems, iterated function systems and related skew products are studied (see [1, 2, 5]). This note is concerned with iterated function systems generated by orientation preserving homeomorphisms on the interval $[0, 1]$. It was proved (see [7]) that under quite general conditions the system has a unique invariant measure on the open interval

Tomasz Szarek was supported by the Polish NCN Grant 2016/21/B/ST1/00033.

(0, 1). (Clearly, such systems also have trivial invariant measures supported on the endpoints.) It is well known that this measure is either absolutely continuous or singular with respect to the Lebesgue measure. The methods that would allow us to distinguish each of these cases are still unknown. The main purpose of our paper is to support the conjecture that typically such invariant measures should be singular (see [1]). In fact, we prove that the set of iterated function systems that have a unique invariant measure on the interval (0, 1) which is singular is generic in the natural class of systems.

2. Notation. By \mathcal{M}_1 and \mathcal{M}_{fin} we denote the set of all probability measures and all finite measures on the σ -algebra of Borel sets $\mathcal{B}([0, 1])$, respectively. By $C([0, 1])$ we denote the family of all continuous functions equipped with the supremum norm $\| \cdot \|$.

An operator $P : \mathcal{M}_{fin} \rightarrow \mathcal{M}_{fin}$ is called a *Markov operator* if it satisfies the following two conditions:

- (1) positive linearity: $P(\lambda_1\mu_1 + \lambda_2\mu_2) = \lambda_1P\mu_1 + \lambda_2P\mu_2$ for $\lambda_1, \lambda_2 \geq 0$; $\mu_1, \mu_2 \in \mathcal{M}_{fin}$;
- (2) preservation of measure: $P\mu([0, 1]) = \mu([0, 1])$ for $\mu \in \mathcal{M}_{fin}$.

A Markov operator P is called a *Feller operator* if there is a linear operator $P^* : C([0, 1]) \rightarrow C([0, 1])$ (dual to P) such that

$$\int_{[0,1]} P^*f(x)\mu(dx) = \int_{[0,1]} f(x)P\mu(dx) \text{ for } f \in C([0, 1]), \mu \in \mathcal{M}_{fin}.$$

Note that, if such an operator exists, then $P^*(\mathbf{1}) = \mathbf{1}$. Moreover, $P^*(f) \geq 0$ if $f \geq 0$ and

$$\|P^*(f)\| \leq \|P^*(|f|)\| \leq \|f\|,$$

so P^* is a continuous operator. A measure μ_* is called *invariant* if $P\mu_* = \mu_*$.

Let $\Gamma = \{g_1, \dots, g_k\}$ be a finite collection of nondecreasing absolutely continuous functions, mapping $[0, 1]$ onto $[0, 1]$, and let $p = (p_1, \dots, p_k)$ be a probability vector. Clearly, it defines a probability distribution p on Γ by putting $p(g_i) = p_i$. Put $\Sigma_n = \{1, \dots, k\}^n$, and let $\Sigma_* = \bigcup_{n=1}^\infty \Sigma_n$ be the collection of all finite words with entries from $\{1, \dots, k\}$. For a sequence $\omega \in \Sigma_*$, $\omega = (i_1, \dots, i_n)$, we denote by $|\omega|$ its length (equal to n). Finally, let $\Omega = \{1, \dots, k\}^\mathbb{N}$. For $\omega = (i_1, i_2, \dots) \in \Omega$ and $n \in \mathbb{N}$, we set $\omega|_n = (i_1, i_2, \dots, i_n)$. Let \mathbb{P} be the product measure on Ω generated by the initial distribution on $\{1, \dots, k\}$. To any $\omega = (i_1, \dots, i_n) \in \Sigma_*$, there corresponds a composition $g_\omega = g_{i_n, i_{n-1}, \dots, i_1} = g_{i_n} \circ g_{i_{n-1}} \circ \dots \circ g_{i_1}$. The pair (Γ, p) , called an *iterated function system*, generates a Markov operator $P : \mathcal{M}_{fin} \rightarrow \mathcal{M}_{fin}$ of the form

$$P\mu = \sum_{i=1}^k p_i g_i\mu,$$

where $g_i\mu(A) = \mu(g_i^{-1}(A))$ for $A \in \mathcal{B}([0, 1])$; which describes the evolution of distribution due to the action of randomly chosen maps from the collection Γ . It is a Feller operator, that is, its dual operator P^* , given by the formula



$$P^* f(x) = \sum_{i=1}^k p_i f(g_i(x)) \quad \text{for } f \in C([0, 1]), x \in [0, 1],$$

has the property $P^*(C[0, 1]) \subset C([0, 1])$.

Definition 1. We denote by C_β^+ , $\beta > 0$, the space of continuous functions $g : [0, 1] \rightarrow [0, 1]$ satisfying the following properties:

- (1) g is nondecreasing, $g(0) = 0$, and $g(1) = 1$,
- (2) g is continuously differentiable on $[0, \beta]$ and $[1 - \beta, 1]$.

We introduce the space \mathcal{F}_0 of pairs (Γ, p) such that $\Gamma = \{g_1, \dots, g_k\} \subset C_\beta^+$ is a finite collection of homeomorphisms, and $p = (p_1, \dots, p_k)$ is a probabilistic vector. We endow this space with the metric

$$d((\Gamma, p), (\Delta, q)) = \sum_{i=1}^k \left(|p_i - q_i| + \|g_i - h_i\| + \|g_i^{-1} - h_i^{-1}\| + \sup_{[0, \beta] \cup [1 - \beta, 1]} |g'_i - h'_i| \right),$$

where $\Delta = \{h_1, \dots, h_k\}$ and $q = (q_1, \dots, q_k)$. It is easy to check that (\mathcal{F}_0, d) is a complete metric space.

Definition 2. Let $\Gamma = \{g_1, \dots, g_k\} \subset C_\beta^+$ be a finite collection of homeomorphisms, and let $p = (p_1, \dots, p_k)$ be a probability vector such that $p_i > 0$ for all $i = 1, \dots, k$. The pair (Γ, p) is called an *admissible iterated function system* if

- (1) for any $x \in (0, 1)$, there exist $i, j \in \{1, \dots, k\}$ such that $g_i(x) < x < g_j(x)$;
- (2) the Lyapunov exponents at both common fixed points 0, 1 are positive, i.e.,

$$\sum_{i=1}^k p_i \log g'_i(0) > 0 \quad \text{and} \quad \sum_{i=1}^k p_i \log g'_i(1) > 0.$$

We consider the space $\mathcal{F} \subset \mathcal{F}_0$ of admissible iterated function systems (Γ, p) with all maps $g \in \Gamma$ absolutely continuous. The closure $\overline{\mathcal{F}}$ of \mathcal{F} in (\mathcal{F}_0, d) is clearly complete. Recall that a subset of a complete metric space is called *residual* if its complement is a set of the first Baire category.

Let $\mathcal{M}_1(0, 1)$ denote the space of $\mu \in \mathcal{M}_1$ supported in the open interval $(0, 1)$, i.e., satisfying $\mu((0, 1)) = 1$. Clearly, each $(\Gamma, p) \in \mathcal{F}$ has two invariant measures: $\nu_1 = \delta_0$ and $\nu_2 = \delta_1$. As we will show, it also admits a unique invariant measure in $\mathcal{M}_1(0, 1)$.

Since $P\mu$ is singular for singular $\mu \in \mathcal{M}_{fin}$, a unique invariant measure is either singular or absolutely continuous (see [9]). The aim of this paper is to show that the set of all $(\Gamma, p) \in \overline{\mathcal{F}}$, which have singular unique invariant measure $\mu \in \mathcal{M}_1(0, 1)$, is residual in $\overline{\mathcal{F}}$.



3. Invariant measure. The paper [7] deals with a special case of an IFS, namely for $k = 2$, with $g_1(x) < x$, $g_2(x) > x$, on $(0, 1)$, being twice continuously differentiable. The proof of [7, Lemma 3.2] gives the following result.

Lemma 1. *Let (Γ, p) be an admissible iterated function system. Then there exists an ergodic invariant measure $\mu \in \mathcal{M}_1(0, 1)$.*

The main ingredient of the proof of Lemma 1 is the following subset of $\mathcal{M}_1(0, 1)$: for small $0 < \alpha < 1$ and positive M , one defines

$$\mathcal{N}_{M, \alpha} = \{\mu \in \mathcal{M}_1(0, 1) : \mu([0, x]) \leq Mx^\alpha \text{ and } \mu([1 - x, 1]) \leq Mx^\alpha\}. \quad (1)$$

For each admissible IFS, thanks to the positivity of the Lyapunov exponents, the corresponding Markov operator P keeps an $\mathcal{N}_{M, \alpha}$ invariant for some M and α . Then, among the invariant measures for P , given by the Krylov-Bogolyubov procedure, one can find an ergodic one, due to the convexity of $\mathcal{N}_{M, \alpha}$.

For later use, we give below an easy extension of the invariance part of [7, Lemma 3.2] and its proof.

Lemma 2. *Let (Γ, p) be an admissible iterated function system and let P be the Markov operator corresponding to (Γ, p) . There exist $M > 0$ and $\alpha > 0$ such that $\mathcal{N}_{M, \alpha} \neq \emptyset$ and*

$$P(\mathcal{N}_{M, \alpha}) \subset \mathcal{N}_{M, \alpha}.$$

Proof. Note that, by Definitions 1 and 2, we can find $x_0 > 0$ and $\lambda_1, \dots, \lambda_k > 0$ such that

$$\Lambda := \sum_{i=1}^k p_i \log \lambda_i > 0$$

and

$$g_i^{-1}(x) \leq x/\lambda_i \quad \text{for all } x \in [0, x_0].$$

Let $F(\alpha) = \sum_i p_i e^{-\alpha \log \lambda_i}$. Writing the Taylor expansion at 0, we obtain

$$\begin{aligned} F(\alpha) &= \sum_i p_i (1 - \alpha \log \lambda_i + \mathcal{O}(\alpha^2)) = 1 - \alpha \sum_i p_i \log \lambda_i + \mathcal{O}(\alpha^2) \\ &= 1 - \alpha \Lambda + \mathcal{O}(\alpha^2) < 1 \end{aligned}$$

for α small enough; choose such $\alpha > 0$. Choose also $M > 0$ large enough so that

$$Mx_0^\alpha \geq 1.$$

Assume that $x \in [0, x_0]$ and let $\mu \in \mathcal{N}_{M, \alpha}$. Then

$$\begin{aligned} P\mu([0, x]) &= \sum_i p_i \mu([0, g_i^{-1}(x)]) \leq \sum_i p_i \mu([0, x/\lambda_i]) \leq M \sum_i p_i x^\alpha / \lambda_i^\alpha \\ &= Mx^\alpha \sum_i p_i e^{-\alpha \log \lambda_i} = Mx^\alpha \cdot F(\alpha) \leq Mx^\alpha. \end{aligned}$$

On the other hand, if $x > x_0$, then $P\mu([0, x]) \leq 1 \leq Mx_0^\alpha \leq Mx^\alpha$. In the same way, we prove that $P\mu([1 - x, 1]) \leq Mx^\alpha$ (with possibly larger M and smaller α). This completes the proof. \square



Using ideas from [8, Lemma 3], one can prove the following:

Lemma 3. *Let (Γ, p) be an admissible iterated function system. Assume that there exist $g_1, g_2 \in \Gamma$ such that the ratio $\ln g'_1(0)/\ln g'_2(0)$ is irrational and $g'_1(0) > 1 > g'_2(0)$; let g'_1 be Hölder continuous in a neighborhood of zero. Then the IFS (Γ, p) is minimal, that is, $\{g_{i_n, i_{n-1}, \dots, i_1}(x)\}_n$ is dense in $[0, 1]$ for each x in $(0, 1)$ and \mathbb{P} -almost all ω .*

Uniqueness may be shown under the conditions of the preceding lemma, for instance using an argument given in [7, Lemma 3.4]; we give another proof, which does not rely on the injectivity of each $g \in \Gamma$.

Lemma 4. *Under the conditions of the preceding lemma, there is a unique stationary measure in $\mathcal{M}_1(0, 1)$.*

Proof. Let $M > 0$ be such that $P(\mathcal{N}_{M, \alpha}) \subset \mathcal{N}_{M, \alpha}$ and let $\nu \in \mathcal{N}_{M, \alpha}$. Put

$$\mu_n = \frac{\nu + P\nu + \dots + P^{n-1}\nu}{n}$$

and note that, by (1), $\mu_n \in \mathcal{N}_{M, \alpha}$. Let μ be an accumulation point of the sequence (μ_n) in the weak- $*$ topology in $C([0, 1])^*$. Then it is easy to check that also $\mu \in \mathcal{N}_{M, \alpha}$. Moreover, since P is a Feller operator, every accumulation point of the sequence μ_n is an invariant measure for the process P .

We now prove the uniqueness. Let $\mu \in \mathcal{M}_1(0, 1)$ be an arbitrary invariant measure. Fix $f \in C([0, 1])$. We define a sequence of random variables $(\xi_n^{f, \mu})_{n \in \mathbb{N}}$ by the formula

$$\xi_n^{f, \mu}(\omega) = \int_{[0, 1]} f(g_{i_1, \dots, i_n}(x)) \mu(dx) \quad \text{for } \omega = (i_1, i_2, \dots).$$

Since μ is an invariant measure for P , we easily check that $(\xi_n^{f, \mu})_{n \in \mathbb{N}}$ is a bounded martingale with respect to the natural filtration. Note that this martingale depends on the measure μ . From the martingale convergence theorem, it follows that $(\xi_n^{f, \mu})_{n \in \mathbb{N}}$ is convergent \mathbb{P} -a.s. and since the space $C([0, 1])$ is separable, there exists a subset Ω_0 of Ω with $\mathbb{P}(\Omega_0) = 1$ such that $(\xi_n^{f, \mu})_{n \in \mathbb{N}}$ is convergent for any $f \in C([0, 1])$ and $\omega \in \Omega_0$. Therefore, for any $\omega \in \Omega_0$, there exists a measure $\omega(\mu) \in \mathcal{M}_1$ such that

$$\lim_{n \rightarrow \infty} \xi_n^{f, \mu}(\omega) = \int_{[0, 1]} f(x) \omega(\mu)(dx) \quad \text{for every } f \in C([0, 1]).$$

We are now ready to show that for any $\varepsilon > 0$, there exists $\Omega_\varepsilon \subset \Omega$ with $\mathbb{P}(\Omega_\varepsilon) = 1$ satisfying the following property: for every $\omega \in \Omega_\varepsilon$, there exists an interval I of length $|I| \leq \varepsilon$ such that $\omega(\mu)(I) \geq 1 - \varepsilon$. Hence we obtain that $\omega(\mu) = \delta_{v(\omega)}$ for all ω from some set $\tilde{\Omega}$ with $\mathbb{P}(\tilde{\Omega}) = 1$. Here $v(\omega)$ is a point from $[0, 1]$.

Fix $\varepsilon > 0$ and let $a, b \in (0, 1)$ be such that $\mu([a, b]) > 1 - \varepsilon$. Let $\ell \in \mathbb{N}$ be such that $1/\ell < \varepsilon/2$. Since for any $x \in (0, 1)$, there exists $i \in \{1, \dots, k\}$ such



that $g_i(x) < x$, we may find a sequence $(\mathbf{j}_n)_{n \in \mathbb{N}}$, $\mathbf{j}_n \in \Sigma_*$, such that $g_{\mathbf{j}_n}(b) \rightarrow 0$ as $n \rightarrow \infty$. Therefore, there exist $\mathbf{i}_1, \dots, \mathbf{i}_\ell$ such that

$$g_{\mathbf{i}_m}([a, b]) \cap g_{\mathbf{i}_n}([a, b]) = \emptyset \quad \text{for } m, n \in \{1, \dots, \ell\}, m \neq n.$$

Put $n^* = \max_{m \leq \ell} |\mathbf{i}_m|$ and set $J_m = g_{\mathbf{i}_m}([a, b])$ for $m \in \{1, \dots, \ell\}$. Each J_m is a closed interval. (If we skip the requirement of injectivity of g_i 's, J_m would possibly be a singleton; which does not spoil the proof; see Remark 5 afterwards). Now observe that for any sequence $\mathbf{u} = (u_1, \dots, u_n) \in \Sigma_*$, there exists $m \in \{1, \dots, \ell\}$ such that $\text{diam}(g_{\mathbf{u}}(J_m)) < 1/\ell < \varepsilon/2$. This shows that for any cylinder in Ω , defined by fixing the first initial n entries (u_1, \dots, u_n) , the conditional probability, that $(u_1, \dots, u_n, \dots, u_{n+k})$ are such that $\text{diam}(g_{u_1, \dots, u_n, \dots, u_{n+k}}([a, b])) \geq \varepsilon/2$ for all $k = 1, \dots, n^*$, is less than $1 - q$ for some $q > 0$ independent of n . Hence there exists $\Omega_\varepsilon \subset \Omega$ with $\mathbb{P}(\Omega_\varepsilon) = 1$ such that for all $(u_1, u_2, \dots) \in \Omega_\varepsilon$, we have $\text{diam}(g_{u_1, \dots, u_n}([a, b])) < \varepsilon/2$ for infinitely many n . Since $[0, 1]$ is compact, we may additionally assume that for infinitely many n 's, the set $g_{u_1, \dots, u_n}([a, b])$ is contained in a set I (depending on $\omega = (u_1, u_2, \dots)$) with $\text{diam}(I) \leq \varepsilon$. Observe that this I does not depend on μ ; if only μ_1, μ_2 are two stationary distributions, and if a, b are chosen in such a way that $\mu_1([a, b]) > 1 - \varepsilon$ and $\mu_2([a, b]) > 1 - \varepsilon$, then we obtain that both $\omega(\mu_1)(I) \geq 1 - \varepsilon$ and $\omega(\mu_2)(I) \geq 1 - \varepsilon$, \mathbb{P} -a.s. Hence $\omega(\mu_1) = \omega(\mu_2) = \delta_{v(\omega)}$ for \mathbb{P} -almost every $\omega \in \Omega$. Consequently, for any $f \in C([0, 1])$ and for $i = 1, 2$, we have

$$\int_{[0,1]} f(x) \mu_i(dx) = \lim_{n \rightarrow \infty} \int_{[0,1]} f(x) P^n \mu_i(dx) = \int_{\Omega} \lim_{n \rightarrow \infty} \xi_n^{f, \mu_i}(\omega) \mathbb{P}(d\omega).$$

Since the last integral equals $\int_{\Omega} f(v(\omega)) \mathbb{P}(d\omega)$ in both cases, and since $f \in C([0, 1])$ was arbitrary, we obtain $\mu_1 = \mu_2$ and the proof is complete. \square

Remark 5. The proofs of Lemmata 2, 3, and 4 do not use the fact that g_i , restricted to $(\beta, 1 - \beta)$, is injective. Precisely, these results hold even if we replace the word “homeomorphisms” by “functions” in Definition 2.

4. Auxiliary results and main theorem. Recall that the weak- $*$ topology in $C([0, 1])^*$ is induced by the Fortet-Mourier norm

$$\|\nu\|_{\text{FM}} = \sup \{ \langle f, \nu \rangle : f \in BL_1 \},$$

where BL_1 is the space of all functions $f : [0, 1] \rightarrow \mathbb{R}$ such that $|f(x)| \leq 1$ and $|f(x) - f(y)| \leq |x - y|$ for $x, y \in [0, 1]$.

Denote by d_0 the distance between arbitrary (not necessarily admissible) iterated function systems given by

$$d_0((\{S_1, \dots, S_k\}, p), (\{T_1, \dots, T_k\}, q)) = \sum_{i=1}^k |p_i - q_i| + \|S_i - T_i\|.$$

Lemma 6. Let (S, p) and (T, q) be given iterated function systems and let P_S and P_T be the corresponding Markov operators. Then for every $\mu_1, \mu_2 \in \mathcal{M}_{\text{fin}}$,



we have

$$\|(P_S - P_T)(\mu_1 - \mu_2)\|_{FM} \leq (\mu_1([0, 1]) + \mu_2([0, 1])) d_0((S, p), (T, q)). \quad (2)$$

A proof of this lemma may be found in [12].

Lemma 7. *Suppose that the iterated function systems (S, p) , and (S_1, p) , $(S_2, p), \dots$, are such that, for some $M > 0$ and $\alpha \in (0, 1)$,*

$$\mu_1, \mu_2, \dots \in \mathcal{N}_{M, \alpha} \text{ and } \lim_{n \rightarrow \infty} d_0((S_n, p), (S, p)) = 0,$$

where μ_n is a stationary distribution for (S_n, p) . Then $(\mu_n)_n$ admits a subsequence weakly convergent to a stationary distribution μ for (S, p) ; moreover, $\mu \in \mathcal{N}_{M, \alpha}$.

Proof. By the Prohorov theorem, there is a subsequence μ_{n_k} converging in $(\mathcal{M}_1, \|\cdot\|_{FM})$ to some $\mu \in \mathcal{M}_1$. We denote it by μ_n again, for convenience, so $\|\mu_n - \mu\|_{FM} \rightarrow 0$. Applying Lemma 6, we get

$$\|P\mu_n - \mu_n\|_{FM} = \|(P - P_n)\mu_n\|_{FM} \leq \mu_n([0, 1]) d_0((S_n, p), (S, p)) \rightarrow 0,$$

where P, P_n are the Markov operators corresponding to $(S, p), (S_n, p)$, respectively. The weak continuity of P implies $P\mu_n \rightarrow P\mu$ in $\|\cdot\|_{FM}$, so that $\|P\mu - \mu\|_{FM} = 0$. Now, μ is an invariant measure for P , but we still have to check $\mu \in \mathcal{N}_{M, \alpha}$. Indeed, let $x \in (0, 1)$ be arbitrary. Then

$$\mu_n([x, 1 - x]) \geq 1 - 2Mx^\alpha \quad \text{for all } n \in \mathbb{N}.$$

By the Alexandrov theorem, $\mu([x, 1 - x]) \geq \limsup_n \mu_n([x, 1 - x]) \geq 1 - 2Mx^\alpha$, which means that $\mu \in \mathcal{N}_{M, \alpha}$. □

Let $\mathcal{M}_1^\varepsilon$ be the set of those $\mu \in \mathcal{M}_1$, for which there is a Borel set A with the Lebesgue measure $m(A) < \varepsilon$ and such that $\mu(A) > 1 - \varepsilon/2$.

Remark 8. Note that, if $\mu \in \mathcal{M}_1$ is singular, then for every $\varepsilon > 0$, there is $\delta > 0$ such that for $\nu \in \mathcal{M}_1, \|\mu - \nu\|_{FM} < \delta$, we have $\nu \in \mathcal{M}_1^\varepsilon$. For a proof, see [13, Lemma 2.4.1].

For every $n \in \mathbb{N}$, let $\mathcal{F}_n \subset \mathcal{F}$ be the set of all $(\Gamma, p) \in \mathcal{F}$ with invariant measure $\mu \in \mathcal{M}_1(0, 1) \cap \mathcal{M}_1^{1/n}$.

Lemma 9. *For every $n \in \mathbb{N}$, the set \mathcal{F}_n is dense in the space \mathcal{F} endowed with the metric d .*

Proof. Fix $(\Gamma, p) \in \mathcal{F}, n \in \mathbb{N}$, and $\varepsilon > 0$. We may assume that (Γ, p) satisfies the requirements of Lemma 3. It is easy to construct absolutely continuous modifications of g_1 (on $(\beta, 1 - \beta)$) in such a way that we obtain a sequence $(\Gamma_m, p)_{m \in \mathbb{N}} \subset \mathcal{F}$ satisfying $d((\Gamma_m, p), (\Gamma, p)) < \varepsilon$ and

$$\lim_{m \rightarrow \infty} d_0((\Gamma_m, p), (S_0, p)) = 0$$

where $S_0 = \{g_0, g_2, \dots, g_k\}$ and $g_0(x) = x_0$ for $x \in [u, v] \subset (\beta, 1 - \beta)$, provided $v - u < 2\varepsilon$. As we have noted in Remark 5, due to the inclusion $[u, v] \subset (\beta, 1 - \beta)$, the system (S_0, p) has a unique stationary distribution $\mu_0 \in \mathcal{M}_1(0, 1)$, and μ_0 has full support in $[0, 1]$. Since μ_0 is invariant, we obtain $\mu_0(\{x_0\}) > 0$,



so μ_0 is not absolutely continuous. As remarked earlier, the uniqueness of μ_0 implies that it is either absolutely continuous or singular.

Now, μ_0 is singular and we would like to apply Remark 8, to find $(\Gamma_m, p) \in \mathcal{F}_n$ for some $m \in \mathbb{N}$. In the light of Lemma 7, it suffices to show that all (Γ_m, p) have stationary distributions belonging to the same set $\mathcal{N}_{M,\alpha}$ for some $M, \alpha > 0$. But this follows easily from the proof of Lemma 2 by taking account of the fact that the corresponding elements of Γ_m are identical on $(0, \beta) \cup (1 - \beta, 1)$. \square

Theorem 10. *The set of all $(\Gamma, p) \in \overline{\mathcal{F}}$ which have a unique singular stationary distribution in $\mathcal{M}_1(0, 1)$ is residual in $\overline{\mathcal{F}}$.*

Proof. For $(\Gamma, p) \in \mathcal{F}_n$, we find $\delta_0 > 0$ such that

$$|x - \min_{1 \leq i \leq k} g_i(x)| > \delta_0 \text{ and } |x - \max_{1 \leq i \leq k} g_i(x)| > \delta_0 \quad \text{on } (1/n, 1 - 1/n).$$

Moreover, to (Γ, p) we adjoin a compact set $Z_{(\Gamma,p)} \subset [0, 1]$ such that

$$\mu_{(\Gamma,p)}(Z_{(\Gamma,p)}) > 1 - \frac{1}{2n} \text{ and } m(Z_{(\Gamma,p)}) < \frac{1}{n},$$

where $\mu_{(\Gamma,p)} \in \mathcal{M}_1(0, 1)$ is the invariant measure for (Γ, p) . Further, due to the regularity of the Lebesgue measure, we can find a positive number $r_{(\Gamma,p)}$ such that

$$m(O(Z_{(\Gamma,p)}, r_{(\Gamma,p)})) < \frac{1}{n},$$

where $O(Z_{(\Gamma,p)}, r_{(\Gamma,p)})$ is the open neighborhood of $Z_{(\Gamma,p)}$ in $[0, 1]$ with the radius $r_{(\Gamma,p)}$. Denote by $A_{(\Gamma,p)}$ the set $[0, 1] \setminus O(Z_{(\Gamma,p)}, r_{(\Gamma,p)})$ and consider the classical Tietze function $f_{(\Gamma,p)} : [0, 1] \rightarrow \mathbb{R}$ for the sets $Z_{(\Gamma,p)}$ and $A_{(\Gamma,p)}$ given by the formula

$$f_{(\Gamma,p)}(x) = \frac{\|x, A_{(\Gamma,p)}\|}{\|x, A_{(\Gamma,p)}\| + \|x, Z_{(\Gamma,p)}\|},$$

where $\|x, A\|$ stands for the distance of the point x from the set A . We have $f_{(\Gamma,p)} = 0$ for $x \in A_{(\Gamma,p)}$ and $f_{(\Gamma,p)}(x) = 1$ for $x \in Z_{(\Gamma,p)}$. Obviously, $|f_{(\Gamma,p)}| \leq 1$ and $f_{(\Gamma,p)}$ is Lipschitz, with the Lipschitz constant $l_{(\Gamma,p)} > 1$.

By Lemma 7, and by the proof of Lemma 2, the map taking $(\Gamma, p) \in \mathcal{F}$ into its invariant measure $\mu_{(\Gamma,p)} \in \mathcal{M}_1(0, 1)$ is continuous with respect to (\mathcal{F}, d) and $(\mathcal{M}_1(0, 1), \|\cdot\|_{\text{FM}})$. Hence, there is $\delta_1 > 0$ such that for $(\Delta, q) \in \mathcal{F}$ satisfying $d((\Delta, q), (\Gamma, p)) < \delta_1$, we have

$$\|\mu_{(\Gamma,p)} - \mu_{(\Delta,q)}\|_{\text{FM}} < \frac{1}{2n \cdot l_{(\Gamma,p)}}, \tag{3}$$

where $\mu_{(\Delta,q)} \in \mathcal{M}_1(0, 1)$ is the invariant measure for (Δ, q) .

Additionally, since (Γ, p) is admissible, it follows from Definition 2 that there is $\delta_2 > 0$ with the property

$$\sum_{i=1}^k p_i \log(g'_i(0) - \delta_2) > 0 \text{ and } \sum_{i=1}^k p_i \log(g'_i(1) - \delta_2) > 0.$$



Now, for $(\Gamma, p) \in \mathcal{F}_n$, with the aid of $\delta_{(\Gamma,p)} = \min\{\delta_0/2, \delta_1, \delta_2\}$, we define

$$\widehat{\mathcal{F}} = \bigcap_{n=1}^{\infty} \bigcup_{(\Gamma,p) \in \mathcal{F}_n} B_{\mathcal{F}}((\Gamma, p), \delta_{(\Gamma,p)}),$$

where $B_{\mathcal{F}}((\Gamma, p), \delta_{(\Gamma,p)})$ is an open ball in $(\overline{\mathcal{F}}, d)$ with center at (Γ, p) and radius $\delta_{(\Gamma,p)}$. Clearly, $\widehat{\mathcal{F}}$ as an intersection of open dense subsets of $\overline{\mathcal{F}}$ is residual.

We will show that if $(T, q) \in \widehat{\mathcal{F}}$, then (T, q) has the unique singular stationary distribution supported on $(0, 1)$. Let $(T, q) \in \widehat{\mathcal{F}}$, be fixed. From the definition of $\widehat{\mathcal{F}}$, it follows that there exists a sequence $((\Gamma, p)_n)_{n \in \mathbb{N}}$ such that $(\Gamma, p)_n \in \mathcal{F}_n$ and $(T, q) \in B_{\mathcal{F}}((\Gamma, p)_n, \delta_{(\Gamma,p)_n})$ for $n \in \mathbb{N}$. As we have noted, $T = \{h_1, \dots, h_k\}$ is a collection of homeomorphisms due to the convergence in the metric d . By the same token, the functions h_i are even of class C^1 on $[0, \beta]$ and on $[1 - \beta, 1]$. It is easy to check that (T, q) satisfies condition (1) from Definition 2 due to the proper choice of $\delta_{(\Gamma,p)} < \delta_0$. Moreover, by the inequality $\delta_{(\Gamma,p)} \leq \delta_2$, the Lyapunov exponents for (T, q) are kept positive. This means that (T, q) is an admissible iterated function system. Hence it admits a unique invariant measure $\mu_{(T,q)} \in \mathcal{M}_1(0, 1)$. Again by Lemma 7, there is a subsequence $(\Gamma, p)_{n_k}$ with invariant measures converging to $\mu_{(T,q)}$ in $\|\cdot\|_{\text{FM}}$. For brevity, let us denote the elements of this subsequence again by $(\Gamma, p)_n$, the corresponding invariant measures by μ_n , and let $Z_n = Z_{(\Gamma,p)_n}$, $O(Z_n, r_n) = O(Z_{(\Gamma,p)_n}, r_{(\Gamma,p)_n})$ for $n \in \mathbb{N}$.

The obvious inequality $\text{Lip}(l_n^{-1} f_n) \leq 1$, where $f_n = f_{(\Gamma,p)_n}$ and $l_n = l_{(\Gamma,p)_n}$, implies that

$$\left| \int_{[0,1]} f_n(x) \mu_n(dx) - \int_{[0,1]} f_n(x) \mu_{(T,q)}(dx) \right| \leq l_n \|\mu_n - \mu_{(T,q)}\|_{\text{FM}} < \frac{1}{2n}.$$

However, since $f_n(x) = 1$ for $x \in Z_n$ and $f_n(x) = 0$ for $x \notin O(Z_n, r_n)$,

$$\mu_n(Z_n) - \mu_{(T,q)}(O(Z_n, r_n)) < \frac{1}{2n}.$$

Thus

$$\mu_{(T,q)}(O(Z_n, r_n)) > \mu_n(Z_n) - \frac{1}{2n} > 1 - \frac{1}{2n} - \frac{1}{2n} = 1 - \frac{1}{n}.$$

This and the inequality $m(O(Z_n, r_n)) < \frac{1}{n}$, $n \in \mathbb{N}$, prove that $\mu_{(T,q)}$ is singular. The proof is completed. \square

Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



References

- [1] Alsedá, L., Misiurewicz, M.: Random interval homeomorphisms. *Publ. Math.* **58**, 15–36 (2014)
- [2] Alsedá, L., Misiurewicz, M.: Skew product attractors and concavity. *Proc. Am. Math. Soc.* **143**(2), 703–716 (2015)
- [3] Arnold, L.: *Random Dynamical Systems*. Springer, Berlin (2010)
- [4] Barnsley, M.F., Demko, S.G., Elton, J.H., Geronimo, J.S.: Invariant measures arising from iterated function systems with place dependent probabilities. *Ann. Inst. Henri Poincaré* **24**, 367–394 (1988)
- [5] Bonifant, A., Milnor, J.: Schwarzian derivatives and cylinder maps. In: *Holomorphic Dynamics and Renormalization*, Fields Institute communications, vol. 53, pp. 1–21. American Mathematical Soc., Providence, RI (2008)
- [6] Crauel, H., Flandoli, F.: Attractors for random dynamical systems. *Probab. Theory Relat. Fields* **100**(3), 365–393 (1994)
- [7] Gharaei, M., Homburg, A.J.: Random interval diffeomorphisms. *Discrete Contin. Dyn. Syst. Ser. S* **10**(2), 241–272 (2017)
- [8] Ilyashenko, Yu.: Thick attractors of step skew products. *Regul. Chaotic Dyn.* **15**(2–3), 328–334 (2010)
- [9] Lasota, A., Myjak, J.: Generic properties of fractal measures. *Bull. Polish Acad. Sci. Math.* **42**, 283–296 (1994)
- [10] Díaz, L.J., Matias, E.: Stability of the Markov operator and synchronization of Markovian random products. *Nonlinearity* **31**(5), 1782–1806 (2018)
- [11] Scheutzwow, M.: Comparison of various concepts of a random attractor: a case study. *Arch. Math. (Basel)* **78**(3), 233–240 (2002)
- [12] Szarek, T.: Generic properties of continuous iterated function systems. *Bull. Polish Acad. Sci. Math.* **47**, 77–89 (1999)
- [13] Szarek, T.: Invariant measures for nonexpansive Markov operators on Polish spaces. *Diss. Math.* **415**, 1–62 (2003)

WOJCIECH CZERNOUS
University of Gdańsk
Al. Marszałka Piłsudskiego 46
81-378 Gdynia
Poland
e-mail: wojciech.czernous@ug.edu.pl



TOMASZ SZAREK
Faculty of Physics and Applied Mathematics
Gdańsk University of Technology
ul. Gabriela Narutowicza 11/12
80-233 Gdańsk
Poland
e-mail: szarek@intertele.pl

Received: 31 May 2019