

PAPER • OPEN ACCESS

Low energy elastic electron scattering from benzonitrile (C_6H_5CN)

To cite this article: G Tatreau *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 182007

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Low energy elastic electron scattering from benzonitrile (C_6H_5CN)

G Tatreau¹, B Diaz¹, F Alharthi¹, M Zawadzki^{1,2*} and M A Khakoo^{1†}

¹Department of Physics, California State University, Fullerton, California 92831, USA

²Atomic Physics Division, Department of Atomic, Molecular and Optical Physics, Faculty of Applied Physics and Mathematics, Gdańsk University of Technology, ul. Gabriela Narutowicza 11/12, 80-233 Gdańsk, Poland

Synopsis We present experimental differential elastic scattering cross sections (DCSs) for low energy electron scattering from benzonitrile along with integral and momentum-transfer cross sections that are determined from these DCSs. The measurements of DCSs are obtained using the relative flow method with helium as the standard gas, in a crossed electron-molecular gas beam arrangement. Our measurements are made at incident electron energies in the range of 1 – 30 eV and scattering angles in the range of (10°-130°).

Recently, in many previous theoretical and experimental electron collision studies the molecular targets containing a ring structure were of great interest due to its relevance in many fields, i.e. medicine, biology and bio-physics [1]. In present work we focus on providing the experimental insight into integral cross sections (ICSs) and momentum transfer cross sections (MTCSs) for benzonitrile.

This aromatic organic compound is a phenyl group bonded to a cyanide group (see figure). Such configuration, where a triple CN bond is present is very interesting from electron-molecule interaction point of view in terms of electron attachment to a system different from a C=C bond commonly found in organic compounds.

Our apparatus consists of a high energy-resolution electron gun/analyzer system, both are equipped with double hemispherical energy selectors to provide well-defined electron beam energy profiles and finesse. Typical electron currents ranged around 19-26 nA, with a corresponding energy resolution of between 40 and 55 meV, full width at half maximum. The experimental setup is located in a magnetically shielded vacuum chamber with a base pressure of 10^{-8} torr.

In order to evaluate ICSs and MTCSs for benzonitrile we performed series of measurements of elastic differential cross section (DCS) for a range of scattering angles (10°-130°) and energies (1 eV to 30 eV). ICSs and MTCSs were calculated from the measured DCS by extrapolating the DCS to 0° and 180° and applying standard numerical integration methods to them. Each DCS set was extrapolated to 0° and 180° by a

*E-mail: mateusz.zawadzki@pg.edu.pl

†E-mail: mkhakoo@fullerton.edu

polynomial curve [2]. The present results are presented in Fig. 1; both the behavior of the elastic ICSs and MTCSs, indicate a possible resonance located around 3.5 eV.




Figure 1. ICSs and MTCSs for elastic electron scattering by benzonitrile.

Acknowledgments: The work carried out at California State University, Fullerton was funded by the National Science Foundation under grant NSF-RUI-PHY 1606905 and by a Fulbright Fellowship to M Zawadzki.

References

- [1] Gorfinkiel J D and Ptasinska S 2017 *J. Phys. B: At. Mol. Opt. Phys.* **50** 182001
- [2] Fedus K, Navarro C, Hargreaves L R, Khakoo M A, Silva F M, Bettega M H F, Winstead C and McKoy V 2014 *Phys. Rev. A* **90** 032708