

PAPER • OPEN ACCESS

Low energy differential elastic electron scattering from trichloromethane

To cite this article: B Diaz *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 182016

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Low energy differential elastic electron scattering from trichloromethane

B Diaz¹, G Tatreau¹, B Hlousek¹, M Zawadzki^{1,2*} and M A Khakoo^{1†}

¹Department of Physics, California State University, Fullerton, California 92831, USA

²Atomic Physics Division, Department of Atomic, Molecular and Optical Physics, Faculty of Applied Physics and Mathematics, Gdańsk University of Technology, ul. Gabriela Narutowicza 11/12, 80-233 Gdańsk, Poland

Synopsis Experimental differential cross sections for low energy electron scattering from trichloromethane is measured utilizing a crossed electron-molecular beam experiment via the relative flow method, for the incident electron energies in the range of $E_0 = 0.5\text{ eV}-30\text{ eV}$ and the scattering angles in the range of $\theta = 10^\circ - 130^\circ$.

Low-energy electron collision processes play an important role in several areas including astrophysics, radiation biology, atmospheric chemistry and technology. When considering the inelastic and elastic collisions, the former constitutes a dominant process of low-energy electron transport through gaseous media, plasmas and condensed matter. In our present work we provide the elastic differential cross sections (DCSs) for low energy electron scattering from trichloromethane - the organic compound with formula CHCl_3 .

The experimental setup employs a crossed electron/target beam arrangement. The energy separation of electrons is made at high resolution (40-50 meV FWHM) using electrostatic lenses combined with hemispherical analyzers [1]. In order to accurately determine background scattered electrons we employ a movable collimated gas beam source [2] developed in our laboratory.

The DCSs for trichloromethane, $Q_A(E_0, \theta)$ are obtained from the relative-flow formula:

$$Q_A(E_0, \theta) = Q_{\text{He}}(E_0, \theta) \frac{\mathcal{R}_{\text{He}} I_{S,A}}{\mathcal{R}_A I_{S,\text{He}}} \sqrt{\frac{M_{\text{He}}}{M_A}}, \quad (1)$$

where \mathcal{R} stands for the relative flow rates, I_S represents the scattering rates and M represents the molar masses where the subscripts indicate the gas species, with A being the unknown gas (trichloromethane) whose DCS is to be determined and He the standard gas (helium) whose DCSs are known; thus our measured trichloromethane DCSs were normalized to elastic DCSs for helium.

*E-mail: mateusz.zawadzki@pg.edu.pl

†E-mail: mkhakoo@fullerton.edu

We will present the experimental data on differential cross sections for elastic electron scattering on CHCl_3 for range of energies (0.5, 1, 2, 3, 4, 5, 7, 10, 15, 20, 30 eV) and the scattering angles ($10^\circ - 130^\circ$). An example of the data for four incident electron energies is shown in Fig. 1.




Figure 1. Differential cross sections for elastic electron scattering by trichloromethane.

Acknowledgments: The work carried out at California State University, Fullerton was funded by the National Science Foundation under grant NSF-RUI-PHY 1606905. M.Z. acknowledges the Fulbright Program to conduct this work at California State University Fullerton.

References

- [1] Zawadzki M and Khakoo M A 2018 *J. Chem. Phys.* **149** 124304
- [2] Hughes M, James Jr K E, Childers J G and Khakoo M A 2003 *Meas. Sci. Technol.* **14** 1

