

PAPER • OPEN ACCESS

Methylation effect in e^- — scattering on methyl-substituted ethylenes

To cite this article: S Stefanowska-Tur *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 182001

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Methylation effect in e^- – scattering on methyl-substituted ethylenes

S Stefanowska-Tur^{1*}, Cz Szmytkowski¹, E Ptasńska-Denga¹ and P Możejko¹

¹Department of Atomic, Molecular and Optical Physics, Faculty of Applied Physics and Mathematics, Gdańsk University of Technology, Gabriela Narutowicza 11/12, Gdańsk, 80-233, Poland

Synopsis Methylation effect has been observed and studied in electron-scattering from selected hydrocarbon molecules. In measured total cross section (TCS) functions we have noticed energy shifts and changes in the intensity of observed structures.

Electron-scattering cross sections data for hydrocarbons are of great importance in many fields of science and technology. Systematic studies of TCS for electron scattering from molecules allow also to notice relationships between shape and magnitude of TCS energy dependence and some physical properties of targets.

In this work, we present how methylation can affect TCS for ethylene and its methylated derivatives. All data have been obtained in our laboratory using the electrostatic 127° electron spectrometer working in the linear transmission mode [1].

energy structure is also observed, near 25 eV, which becomes more pronounced as the number of CH_3 groups in a target molecule increases.




Figure 1. TCS for methyl-substituted ethylenes: ethylene [2], propene [3], 2-methylpropene [4], 2-methyl-2-butene [5] and 2,3-dimethyl-2-butene [5].

Table 1. Location of the first maximum in TCS for studied targets.

Molecule	E_{1max} [eV]
Ethylene	1.9
Propene	2.2
2-Methylpropene	2.4
2-Methyl-2-butene	2.6
2,3-Dimethyl-2-butene	2.7

To investigate how the replacement of H atoms in target molecule with CH_3 affects TCS energy dependence, we compared results for ethylene [2], propene [3], 2-methylpropene [4], 2-methyl-2-butene [5] and 2,3-dimethyl-2-butene [5] molecules (fig. 1). Table 1 gives the location of low-energy TCS maxima: increase in the number of methyl groups in the molecule causes a shift of this structures toward higher energies. This effect can be associated with different redistribution of electric charge in the methyl-substituted molecules. A weak high-

This work has been supported in part by the Polish Ministry of Science and Higher Education (MNiSzW Project 2018-2019). S. Stefanowska-Tur kindly acknowledge the support of the Polish Ministry of Science and Higher Education within the Diamond Grant program (Project No. DI2015 018945).

References

- [1] Szmytkowski Cz *et al* 2001 *Vacuum* **63** 549
- [2] Szmytkowski Cz *et al* 2003 *Phys. Rev. A* **68** 032715
- [3] Szmytkowski Cz *et al* 2002 *J. Phys. B: At. Mol. Opt. Phys.* **35** 3781
- [4] Możejko P *et al* 2012 *J. Phys. B: At. Mol. Opt. Phys.* **45** 145203
- [5] Szmytkowski Cz *et al* 2015 *J. Chem. Phys.* **143** 064306

*E-mail: s141549@student.pg.edu.pl

