

1 This is a postprint version of the article (please cite as):

2 Postprint of: Ruman M., Kosek K., Koziol K., Ciepły M., Kozak-Dylewska K., Polkowska Ż., A High-Arctic
3 flow-through lake system hydrochemical changes: Revvatnet, southwestern Svalbard (years 2010–
4 2018), CHEMOSPHERE, Vol. 275 (2021), 130046, DOI: [10.1016/j.chemosphere.2021.130046](https://doi.org/10.1016/j.chemosphere.2021.130046)

5 Published version available at:

6 <https://www.sciencedirect.com/science/article/pii/S0045653521005154?via%3Dihub>

7 © 2021. This manuscript version is made available under the CC-BY-NC-ND 4.0 license

8 <https://creativecommons.org/licenses/by-nc-nd/4.0/>

9

10 **A High-Arctic flow-through lake system hydrochemical changes: Revvatnet,**
11 **southwestern Svalbard (years 2010-2018)**

12 Marek Ruman¹, Klaudia Kosek^{2*}, Krystyna Koziół³, Michał Ciepły¹, Katarzyna Kozak-
13 Dylewska³, Żaneta Polkowska³

14 ¹Faculty of Natural Sciences, University of Silesia, 60 Będzińska St., Sosnowiec 41-200, Poland

15 ²Faculty of Civil and Environmental Engineering, Gdansk University of Technology, 11/12 Narutowicza St.,
16 Gdansk 80-233, Poland

17 ³Faculty of Chemistry, Gdansk University of Technology, 11/12 Narutowicza St., Gdansk 80-233, Poland

18 *Corresponding author: K.Kosek (e-mail address: klaudia.kosek@pg.edu.pl)

19

20 **Abstract:** Lake ecosystems are strongly coupled to features of their surrounding landscapes
21 such as geomorphology, lithology, vegetation and hydrological characteristics. In the 2010-
22 2018 summer seasons, we investigated an Arctic flow-through lake system Revvatnet, located
23 in the vicinity of the coastal zone of Hornsund fjord in Svalbard, characterising its
24 hydrological properties and the chemical composition of its waters. The lake system
25 comprises of a small upper lake and a large lower one, the latter cone-shaped, with -29.1 m
26 maximum depth. With near-neutral pH (full range 6.5 – 8.4) and low EC (7 to 147 $\mu\text{S cm}^{-1}$),
27 the lake has rather similar characteristics to many Arctic lakes. Metal and metalloid
28 concentrations were either similar across the lake system or increased downstream (except Zn,
29 which has important ore-bearing veins in the upper part of the catchment), which is consistent
30 with the likely slow dissolution of suspended particles within the lakes. The ΣPAHs
31 concentrations ranged from <MDL to 2151 ng L^{-1} , and according to the indicator PAHs
32 concentration ratios, they originated from a mixture of combustion processes (they were not
33 petrogenic). Principal component analysis showed that seasonal variability was the most
34 characteristic feature of the chemical composition of these waters, although there appear to be

35 consistent changes with time (sampling year) as well. Future research should explore the
36 occurrence of high maxima in the concentrations of priority pollutants, such as PAHs, metals
37 and metalloids (e.g. As).

38

39 **Keywords:** Arctic; flow-through lake; bathymetry; freshwater; chemical pollution

40

41 **1. Introduction**

42 A characteristic and often dominant feature of polar landscapes is the great diversity and
43 abundance of standing surface waters (Kling, 2009). Many lakes owe their origin to glacial
44 action that shaped their lake basin and surrounding landscape, and that distributed glaciogenic
45 deposits of different sources and mineral composition across their catchments, a fact
46 referenced by other studies, e.g. Marszałek and Górnjak (2017). In the postglacial landscape,
47 lake density is more than four times higher than in terrain which was not glaciated (Smith et
48 al., 2007). Furthermore, lake densities and area fractions are on average approximately 100 –
49 170 % greater in permafrost-influenced terrain (compared with permafrost-free) (Vincent and
50 Laybourn-Parry, 2008).

51 High-latitude lakes are likely to show ongoing responses to present climate instability that
52 leads to deglaciation, variations in chemical composition of freshwater, and, for lakes located
53 close to coastal waters, changes in their linkages with the sea. All of these processes are major
54 controls on the structure and functioning of aquatic ecosystems (Van Hove et al., 2006;
55 Vincent and Laybourn-Parry, 2008). Geological variety of the polar regions, including the
56 differences in substrate lithology, may also affect the extent of rock weathering and the
57 chemical composition of soil water that finally discharges into polar lakes (Hamilton et al.,
58 2001; Szumińska et al., 2018; Vincent and Laybourn-Parry, 2008). In brief, Arctic lake



59 ecosystems are strongly coupled to the features of their surrounding landscapes, such as
60 hydrological characteristics, geomorphology, lithology and vegetation, and due to this, there
61 are several significant differences in lake characteristics in various parts of the Arctic (Wetzel,
62 2001).

63 High solar radiation levels, reaching high latitudes in spring, result in rapid snowmelt. Runoff
64 in late spring typically comprises 80 – 90 % of the annual total in the Arctic, while it lasts
65 only a few weeks (Prowse and Ommanney, 1990). Summer sources of freshwater include
66 perennial and durable snow patches, rain, glaciers, melting of the active permafrost layer and
67 some cases of groundwater discharge (Vincent and Laybourn-Parry, 2008). Groundwater
68 levels and distribution within polar regions are greatly influenced by bedrock geology,
69 permafrost layers and soil thickness (Prowse and Ommanney, 1990).

70 Freshwater is crucial to the unique and fragile ecosystems of the Arctic. Climate change
71 enhances the moisture transport from lower latitudes towards the pole (Mueller et al., 2009).
72 This is contributing to increases in precipitation in the Arctic, falling either as rain or snowfall
73 and transporting at the same time a variety of persistent organic pollutants (POPs),
74 characterised by durability and resistance to degradation. The residence time of those
75 pollutants is long enough for them to be transported thousands of kilometres by air and ocean
76 water (Halsall et al., 2001). Consequently, the compounds that have not been produced for the
77 past few decades still appear in the environment (including polar regions). The quantities of
78 them may impact negatively on the functioning of ecosystems, animal and human health
79 (Kozak et al., 2013). In many parts of the Arctic, the proportion of precipitation that falls as
80 rain, as opposed to snow, has increased, and the period of snow-cover has become shorter
81 (Kling, 2009; Olichwer et al., 2013). These processes have a significant impact on the
82 hydrological changes of the entire Arctic lake catchments and they are expected to continue,
83 although to variable degrees between different parts of the Arctic and over different seasons.

84 Among polar regions, the Svalbard archipelago is distinguished by its location as the gateway
85 to the Arctic. Relatively close location of Svalbard archipelago to Europe makes this sensitive
86 region particularly exposed to the influence of pollutants (Kozak et al., 2016). Additionally,
87 the landscape of the Revvatnet lake area with its prevalence of mountains, favours the
88 accumulation of pollutants transported by air masses from Europe and Asia. Therefore, it has
89 been chosen for comprehensive studies on the pollutants deposition (e.g., Laing et al., 2014;
90 Wojtuń et al., 2013). Worldwide, countries are striving to reduce emissions of potentially
91 toxic pollutants to reduce their levels in all environmental media. Therefore, studying the
92 pollution in polar waters is a timely and important research task.

93 Since the Revvatnet is a flow-through lake, near the river estuary in the Hornsund fjord, it can
94 be expected that some of its pollution will be transported to the coastal zone, causing adverse
95 effects. Many contaminants are particle-bound, thus sedimentation and resuspension
96 processes will determine whether they reach the coastal zone and the sea or if they are
97 deposited along the way. When a river flows slowly, particles usually settle, and bottom
98 sediments can become enriched in persistent organic pollutants (e.g. the PAHs described
99 here). However, the river bottom is usually only a temporary trap, since turbulent flow would
100 lead to the resuspension of particles (CliC/AMAP/IASC, 2016). The objective of the present
101 study was to characterise the hydrological properties of the High-Arctic flow-through
102 Revvatnet lake in Svalbard and to evaluate its chemical composition, especially of
103 contaminants, in an eight-year period (2010-2018), taking into account geomorphology . The
104 latter, as well as pollutant accumulation, both have the potential to affect lake evolution
105 through a variety of processes, especially in the polar regions which are sensitive to small
106 alterations in air temperatures and the chemical composition of water (Kozak et al., 2016;
107 Szumińska et al., 2018).

108



109 **2. Materials and Methods**

110 2.1. Study area

111 Revvatnet is located in Wedel-Jarlsberg Land, in southwestern Spitsbergen (15°36'E,
112 77°02'N), in the vicinity of the Polish Polar Station in Hornsund. The studied lake is situated
113 at 40 m a.s.l. within the glacial Revdalen catchment. The Revvatnet lake is fed by atmospheric
114 precipitation, snow melt water streams and the Revelva river with its tributaries. Revvatnet is
115 covered with ice for about 8-9 months a year. It is usually free from ice cover from mid-June
116 to the end of September. However, the lake does not freeze to the bottom.

117 The upper part of the Revdalen catchment is characterised by the streams originating from the
118 slopes of Eimfjellet and Skålfjellet. The catchment is characterised by many tributaries
119 located on its left side, of which the proglacial Ariebekken is the largest (Kosek et al., 2019a).
120 The Revvatnet, as an example of a High-Arctic flow-through lake system, has been chosen for
121 a study on pollution transfer through such a system. For that purpose, both lake bathymetry
122 and water chemistry have been investigated. Sampling locations for chemical measurements
123 have been selected at the left shore of the lake, mainly to determine inputs from tributaries on
124 that side.

125 2.2. Bathymetric measurements

126 Bathymetric measurements were collected in August 2010 using a Lowrance LCX-17MT
127 echo sounder system with a teleCover-Lok telescopic handle mounted on the transom of the
128 rubber boat, recording the path of the boat movement with an integrated GPS receiver (LGC-
129 2000). Bathymetric measurements were made according to the methodology proposed by
130 Lange (1993) and presented in the guide for hydrographic fieldwork (Gutry-Korycka and
131 Werner-Więckowska, 1996). The development of the bathymetric plan of the lake required a
132 series of cross-sections and longitudinal sections, and then the transfer of information about

133 the distribution of depth in these profiles to the digital map of the lake. The total length of all
134 bathymetric profiles was over 20 km (2700 measuring points), which allowed to draw up a
135 bathymetric plan as well as a three-dimensional model of the lake basin. ArcGIS 10.6.1
136 (ESRI) and Surfer 18 softwares (Golden Software) were used to make the bathymetric plan
137 and the lake model.

138 2.3. Freshwater sampling

139 Water samples were collected from the Revvatnet lake in the summer periods between 2010
140 and 2018 (all samples were collected from the same sampling points, Figure 1). The timing of
141 the sampling campaigns was adjusted to the seasonality of the atmospheric conditions in
142 Hornsund, yet logistical constraints prevented us from repeating the sampling every year, and
143 sometimes the month of sampling shifted. The summer is considered to be a period of first
144 snow, then glacier ice and permafrost melting, and through these processes pollutants enter
145 the studied lake. Moreover, there have been significant trends noted in atmospheric
146 precipitation sums across the summer period, which may modify the delivery of pollutants
147 through wet deposition: a decreasing trend in June, and a rising one in July and August
148 (Wawrzyniak and Osuch, 2020). Finally, as summer biomass burning events tend to have a
149 stronger impact on the atmospheric particulate matter in Svalbard (Zielinski et al., 2020),
150 there is a risk of increasing polycyclic aromatic hydrocarbons (PAHs) delivery in that period.

151 **Figure 1.** Location of the sampling points Rev 1-5. For a colour figure, please refer to the
152 online version of the article

153 Samples were collected manually from the Revvatnet lake at a distance of 1.5 m from the
154 shore with no headspace into air-tight, chemically clean bottles (the purity of the procedure
155 has been verified by daily blank sample collection). Pre-cleaning procedure for the bottles
156 included week-long soaking with Milli-Q deionised water and removing the water from the

157 sampling containers several times. The water was sampled from a depth between 20 and 50
158 cm below water level.

159 2.4. Chemical analyses

160 Milli-Q deionised water (18 M Ω) was used during the determination of the various target
161 analyte groups to determine instrumental background (by inserting it in the analysis queue
162 once every six samples) and to prepare sample container blanks. The concentrations of
163 chemical compounds in the collected samples were determined by: 1) for metals and
164 metalloids: Inductively Coupled Plasma – Mass Spectrometry (the element concentration CVs
165 of the obtained triplicate results ranged from 0.5 to 1.5%); 2) for total organic carbon (TOC):
166 TOC-VCSH/CSN Analyser (Shimadzu, Japan) with NDIR detector; 3) for formaldehyde and
167 the sum of phenols: a spectrophotometer, and 4) for PAHs: a gas chromatograph coupled to a
168 single quadrupole mass spectrometer. All the applied analytical procedures have been
169 validated against certified reference materials (CRMs) concordant with ISO Guide 34:2009
170 and ISO/IEC 17025:2005, and the data obtained here were subject to strict QC procedures.
171 Prior to pH measurements, a three-point electrode calibration was performed with temperature
172 compensation, using MERCK Millipore Certipur® buffer solutions of pH 4.00, 7.00 and 9.00
173 (at 25°C). In the analysis of metal and metalloid concentrations, we applied the Standard
174 Reference Material (RM) NIST 1643e Trace Elements in Water, and RM Enviro MAT ES-L-
175 2CRM, ES-H-2 CRM SCP SCIENCE. The calibration of the apparatus was based on RMs by
176 Inorganic ventures ANALITYK: CCS-4, CCS-6, CCS-1, IV-ICPMS-71A. Potassium
177 hydrogen phthalate by NacalaiTesque (Japan) was used for the calibration of the TOC
178 Analyser. Each sample was analysed in triplicate. Sensitivity tests were performed by
179 injecting standard analyte mixtures within the measured concentration range, and linear
180 calibration curves of the peak area against concentration showed correlation coefficients (R^2)
181 in the range of 0.898–0.999 for all standards. Detailed technical specifications of the applied

182 determination techniques, as well as the validation parameters of the analytical procedures,
183 are summarised in Table S1 (Supplementary information).

184 2.5. Statistical analyses

185 The significance of temporal trends and inter-group differences was checked in STATISTICA
186 13 (TIBCO Software Inc.), and due to the lack of normal distribution for most of the data, we
187 applied non-parametric statistics, such as Kruskal-Wallis ANOVA. The same software was
188 used for exploring data distributions and the drawing of box-and-whisker plots.

189 Principal Component Analysis (PCA) is a multivariate statistical analysis that allows
190 revealing internal relations in the data set. PCA finds linear combinations of the original
191 variables, referred to as principal components, which provide better descriptors of the data
192 pattern than the original measurements and account for most of the dataset variation. Principal
193 component analysis (PCA) has been conducted in R software (version 3.6.2.), using the
194 *FactoMineR* package. Missing data were omitted by case. Prior to analysis, values <MDL
195 have been replaced with $\frac{1}{2}$ of the MDL value, subsequently all variables except pH (which
196 already is a logarithm) were log-transformed to bring their distribution closer to normal.
197 Departures from normality were the most pronounced where there were several <MDL
198 values. Due to the exploratory character of the performed PCA, we accepted minor departures
199 from the normal distribution without further data transformation. All variables were scaled
200 (i.e. z-transformed) as part of the PCA calculation procedure (by using the parameter
201 `scale.unit=TRUE`). R was also used to calculate the Wilcoxon signed-rank test results (with
202 the *wilcox.test* function) for the matched samples (i.e. from two sites in the same lake system,
203 which are connected by water flow).

204 205 **3. Results**



206 3.1. Hydrological characteristics of the lake (morphometry)

207 The Revvatnet lake, with a surface area of 0.9 km², has an elongated shape – its axis runs
208 NW-SE. Its maximum length (l) is 2.40 km, and the lake elongation index (λ) equals 6.4. The
209 mean width (B_{av}) of the lake is 0.38 km, while the maximum width (B_{max}) reaches 0.60 km.
210 The lake shoreline is moderately developed, with a shoreline development index (K) of 1.6
211 and the shoreline length (l) of 5.37 km. The mean lake depth was approximately 9.0 m, while
212 the maximum depth, recorded during bathymetric measurements, reached 29.1 m (Figure 2).
213 The volume of the lake (V_0) was calculated at 0.0086 km³. The shape of the lake resembles a
214 cone, which can be estimated from its relative depth (h_w) of 30.6% and the lake depth
215 indicator (W_g) of 0.33. This morphometric feature is a characteristic of glacial lakes (Figure
216 S1, Supplementary Information).

217 For the context of geomorphological situation of the lake, we refer the reader to the
218 geomorphological map (Jania et al., 1984). In brief, the river valley form, within which the
219 lake is located, is surrounded by plains of raised marine terraces (in the upper part of the
220 catchment only by flat ground moraine plains to the east of the lake). Frost-wedge polygons
221 occur in the part of the catchment surrounding and below the lower part of the lake.

222 **Figure 2.** Bathymetry of the Revvatnet lake. For a colour figure, please refer to the online
223 version of the article

224 3.2. Chemical composition of freshwater samples collected in summer periods between 2010 225 and 2018

226 The chemistry of polar freshwater reservoirs is shaped by the geological substratum, as well
227 as by atmospheric deposition, marine aerosols, chemical weathering, biological processes,
228 glacial and periglacial activity. Depending on the geological structures and soil cover on the
229 raised marine terraces and in the surrounding areas, the waters that reach tundra lakes

230 transport varying proportions of specific chemicals (Mazurek et al., 2012). The results
231 obtained for the Revvatnet samples from the summer periods of 2010 to 2018 are shown in
232 Table S3 (Supplementary information).

233 3.2.1. pH and electrical conductivity (EC) measurements

234 Basic physico-chemical parameters, such as pH can vary greatly in aquatic ecosystems (Bååth
235 and Kritzberg, 2015). Lake, river and stream waters vary in pH values from below 4 to above
236 9, even within small geographical areas. In highly productive lakes, pH at the surface may be
237 2 units higher than in bottom waters. The variation of the values is driven by vertical
238 differences in photosynthesis, respiration, and redox conditions (Wetzel, 2001). pH can also
239 fluctuate rapidly. For example, during snow melt and rain storms, pH values in streams can
240 decrease several units, sometimes within a few hours (Lawrence, 2002). On the other hand,
241 sunny days can result in high photosynthetic activity, leading to an increase in water pH.
242 Accordingly, changes of 2-3 pH units may be found in highly productive aquatic
243 environments (Tank et al., 2009). In our study, the pH values in the samples collected over the
244 years 2010-2018 were differing significantly, their annual medians ranging from 6.7 to 7.8
245 (Figure 3).

246 Another basic parameter is electrical conductivity (EC), which in freshwater is affected
247 primarily by the geology of the area through which the water flows. The Arctic lake types
248 range from very dilute waters with EC approaching rainwater, to waters concentrated by
249 evaporation to beyond the salinity of seawater. Within this range, however, the majority of
250 lakes are relatively diluted, with $EC < 300 \mu S cm^{-1}$ (Kling, 2009). The EC values detected in
251 freshwater samples collected in the summer periods of 2010–2018 differed from each other
252 significantly in some years (Figure 3), ranging from 7 to $147 \mu S cm^{-1}$. This places the studied
253 lake system within the spectrum of typical Arctic lakes. Between the two lakes which form



254 the studied system, there was a significant increase in EC noted at the transition from upper to
255 lower (and larger) lake (Wilcoxon signed-rank test $p < 0.05$). The low EC values may be due
256 to both the predominant substratum geology with siliceous rocks (of limited solubility in
257 water) and the underlying permafrost, which isolates surface waters and soils from weathering
258 interactions with deeper mineral soils and rocks. The unfrozen zone beneath the lake may
259 extend for many meters, but the impact of weathering in this zone on lake chemistry is almost
260 entirely unknown. Usually, in the rest of the catchment, weathering reactions are confined to
261 the very shallow unfrozen layers at the surface (Kosek et al., 2019b; Olichwer et al., 2013).

262 **Figure 3.** Range (whiskers) and quartile distribution (Q1 and Q3 frame the box, Q2 = median
263 shown by the line inside it) of pH and EC [$\mu\text{S cm}^{-1}$] measurements over the 2010, 2012, and
264 2014-2018 summer seasons. (R) and (L) denote value plotted on right and left axis,
265 respectively. Kruskal-Wallis ANOVA results are given in the box at the bottom of the graph.
266 For a colour figure, please refer to the online version of the article

267 3.2.2. Trace elements

268 Metals and metalloids were also determined in the samples collected between 2010 and 2018
269 in the summer periods. Changes in the median concentrations of the elements Al, As, Ba, Be,
270 Co, Cr, Cu, Ga, Li, Mn, Ni, Rb, Sr, Tl, U, V, and Zn, in the period 2010 – 2018 are shown in
271 Figure 4, together with their Kruskal-Wallis ANOVA results for inter-annual differences. Of
272 these elements, the following showed significant differences between the sampling years
273 ($p < 0.05$): As, Co, Cr, Li, Mn, Ni, Rb, Sr, and Zn (9 out of 17, which can be interpreted that
274 the trace metal composition of these waters is not stable in time).

275 **Figure 4.** Concentration levels of trace elements in the collected freshwater samples: box
276 shows the inter-quartile range (50% of all results, 25% on each side of the median, which is
277 marked as line inside the box); whiskers encompass full range of results. All Y scale units are



278 [$\mu\text{g L}^{-1}$]. (R) next to an element symbol means it is plotted on the right Y axis, (L) – on the
279 left. Elements are divided into panels due to their varying range of concentration values.
280 Kruskal-Wallis ANOVA results are printed within each graph. For a colour version of the
281 figure, please refer to the online version of the article

282 The Revvatnet lake waters contain abundant metals and metalloids due to the presence of ore-
283 bearing veins and metamorphic rocks in the surrounding area (Smulikowski, 1965;
284 Wojciechowski, 1964). This geological substratum is more exposed in the upper parts of the
285 catchment feeding the lake. Furthermore, the spatial variability of the underlying rocks in the
286 studied catchment allows for the more abundant occurrence of titanium, possibly also barium,
287 caesium, lithium, rubidium, and zinc in the upper part of the lake waters, of zirconium in the
288 left tributaries of the middle part, and of chromium and vanadium in both these areas. The
289 local rocks are relatively abundant in aluminium and manganese throughout the catchment
290 (Smulikowski, 1965). As for the ore-bearing veins, in the area occur those with chalcopyrite,
291 cuprite, malachite and azurite, which are copper minerals, as well as smaller concentrations of
292 sphalerite (with zinc) and galena (with lead). The specific locations of these veins favour the
293 occurrence of copper near Skoddefjellet mountain and in the left tributaries of the lake, and of
294 both lead and zinc in the left tributaries of the smallest upper part of the Revvatnet lake
295 (Wojciechowski, 1964).

296 These elements (except Cs, Pb and Zr, for which we had no or not sufficient data) were tested
297 with the Wilcoxon signed-rank test between points Rev 1 and Rev 2 (the closest in the two
298 lakes) and Rev 1 and Rev 5 (top and bottom of the lake system) to find if their concentration
299 significantly changed between the small lake and the large one. Significant differences (at p
300 <0.05) were found between the concentrations of Ba, Cu, Li and Zn in the small lake (Rev 1)
301 and the output from the large lake (Rev 5); for Al, Ba and Li, the difference was significant
302 between Rev 1 and Rev 2. Among other elements, with less clear geological divisions in the

303 catchment, statistically significant differences (at $p>0.05$) were found between Rev 1 and Rev
304 2 for As and Sr, and between top and bottom of the lake system (Rev 1 and Rev 5) for Ga and
305 Ni. All the noted significant differences were downstream increases, except Zn – it could be
306 interpreted than only in the case of Zn the local input from ore-bearing veins was more
307 prominent than the slow dissolution processes of suspended particles which would normally
308 increase downstream concentrations. This is an opposite pattern to the one noted for lake
309 bottom sediment upper layer in another flow-through lake system of the temperate zone
310 (Kuriata-Potasznik et al., 2016), which highlights that different processes govern lake water
311 and sediment chemistry, therefore direct comparisons may be misleading.

312 Slight changes in the individual concentration levels of metals and metalloids could have been
313 caused by various intensification of geological processes related to the temperature changes
314 and to the mineral surface reactions, increased or reduced amount of precipitation in a
315 selected summer sampling period, complexation, chemical weathering and sorption to solid-
316 phase soil organic matter (Colombo et al., 2018), as well as due to the variability of
317 hydrological processes occurring in the research area. Moreover, the elemental concentration
318 variations may be caused by the occurrence of groundwater associated with the active layer of
319 permafrost, which gains more importance in the hydrological regime of the Revvatnet lake as
320 snow patches disappear in the catchment. Apart from the local natural occurrence of metals
321 and metalloids, they are assumed to be derived to the Arctic mostly from long-range
322 atmospheric transport (AMAP, 2009), and we consider this issue further in the Discussion
323 section 4.3.

324 3.2.3. Organic compounds as water pollution of the Revvatnet lake

325 Among the chemical characteristics of the Revvatnet lake, the presence of organic compounds
326 contaminating the study area should also be emphasized. The Arctic is no longer considered a
327 highly pristine environment, although the air, water, soil and sediment concentrations reveal

328 considerably lower levels of contaminants compared to those found in temperate regions
329 (Kallenborn et al., 2012). Despite this, some characteristic features of the Arctic, e.g. low
330 temperatures, precipitation, ice coverage, and extended periods of darkness during winter,
331 mean that it has the potential to accumulate certain contaminants including persistent organic
332 pollutants (POPs) in all components of the environment (Hung et al., 2010). There are only a
333 few local sources of contaminants in the Arctic, such as military installations, industrial
334 outlets and waste from the old mines, settlements, airports and ships. However, the majority
335 of Arctic pollution problem has arisen through a combination of long-range transport of
336 pollutants and the Arctic haze phenomenon, locking the contaminated air in the area for
337 months (Kallenborn et al., 2012). The cold condensation of POPs enhances the problem of
338 their deposition in the Arctic (Mackay and Wania, 1995).

339 The Arctic contamination research has mainly focused on POPs (including the PAHs
340 described in this paper) because they are bioaccumulative, resistant to degradation and toxic.
341 In addition, other organic chemicals (sum of phenols and formaldehyde) were also determined
342 in collected freshwater samples, which could testify to their global impact (if they are
343 confirmed as anthropogenic). Figure 5 shows the average concentration levels of the sum of
344 phenols, formaldehyde, and total organic carbon (TOC) in the freshwater samples collected
345 between 2010 and 2018.

346 **Figure 5.** Average concentration levels of the sum of phenols, formaldehyde and total organic
347 carbon in the collected freshwater samples. For a colour figure, please refer to the online
348 version of the article

349 Σ Phenols and formaldehyde concentrations levels showed no trend over the studied years
350 (Figure 5), ranging from <0.025 to 0.261 mg L^{-1} and from <0.005 to 0.60 mg L^{-1} ,
351 respectively. However, in the case of HCHO, inter-annual variability was significant (as

352 expressed by Kruskal-Wallis ANOVA $p = 0.00005$). For HCHO, out of 55 measurements,
353 there were two exceptionally high values noted above 0.30 mg L^{-1} (0.60 mg L^{-1} at Rev 1 in
354 July 2012 and 0.53 mg L^{-1} at Rev 5 in June 2016), which could impact the interannual
355 variability picture. Σ Phenols maximum was reached at Rev 2 in September 2015, and
356 Kruskal-Wallis ANOVA for interannual changes was not significant. The maxima were
357 formed by rare departures from the much lower concentration levels occurring most
358 frequently, leading to more uncertainty in estimating annual means than in the case of TOC,
359 whose concentrations were more uniform, ranging from 0.45 to 1.95 mg L^{-1} . The interannual
360 variability of TOC concentrations was, nevertheless, statistically significant (Kruskal-Wallis
361 ANOVA $p = 0.0065$). None of the three mentioned parameters has differed significantly
362 between the upper and the lower lake (as measured by the Wilcoxon signed-rank test, run in
363 two sets of paired samples from: Rev 1 - Rev 2 and Rev 1 - Rev 5).

364 Polycyclic aromatic hydrocarbons are another group of compounds found in the waters of the
365 Revvatnet lake in the years 2010 – 2018. The distribution of concentrations of the detected
366 PAHs in the collected lake water samples are shown in Figure 6.

367 **Figure 6.** Concentration levels of determined PAHs in the collected lake water samples: a)
368 percentage contributions of particular PAHs: b) box and whisker plot, showing median
369 (middle line), inter-quartile range (box) and full range of results (whiskers). All PAHs
370 concentrations in $[\text{ng L}^{-1}]$. For a colour figure, please refer to the online version of the article

371 Historically, only 16 PAHs have been prioritized as environmentally significant and thus
372 received the focus of research, however, it is now recognised that aquatic ecosystems may be
373 exposed to, and potentially affected by hundreds of PAHs and the risks related to that are
374 poorly understood. From a mixture of 16 PAHs that were analysed, only those that were
375 detected in the collected samples are shown in Figure 6. Three compounds from the PAH
376 group occurred in the analysed samples at the highest concentration levels in each year:

377 naphthalene (range of results 69 – 451 ng L⁻¹; median 223 ng L⁻¹), anthracene (range 22 – 657
378 ng L⁻¹; median 147 ng L⁻¹) and fluoranthene (range 13 – 977 ng L⁻¹; median 80 ng L⁻¹). In the
379 period 2015 – 2018, these compounds were at similar concentration levels, with no significant
380 differences in single PAH concentrations between the years (measured with Kruskal-Wallis
381 ANOVA).

382 The two parts of the studied lake system differed significantly only with respect to some
383 PAHs, and the changes were not uniform. Wilcoxon signed-rank test yielded significant
384 results at $p < 0.05$ for naphthalene and fluorene for the transition from upper to lower lake
385 (Rev 1 to Rev 2), while acenaphthene and phenanthrene differed significantly in
386 concentrations at the top and bottom of the lake system (Rev 1 to Rev 5). No other PAHs
387 showed a consistent change between those points in the sampling scheme.

388 The Σ PAHs concentrations ranged from <MDL to 2151 ng L⁻¹ in the collected samples, with
389 annual median concentrations between 303 and 956 ng L⁻¹ (median concentration in the whole
390 study period was 565, and average concentration 748 ng L⁻¹). A typical distribution of a single
391 PAH compound concentrations was right-skewed, with the skewness being stronger for PAHs
392 with a higher number of aromatic rings. With such data distribution, there may be locally or
393 temporally enhanced pollution effects due to the occurrence of strong maxima in PAHs
394 concentrations. The pattern and frequency of their occurrence should be subject of a separate
395 study to establish the environmental risk connected to them.

396 Of interest is the origin of PAHs compounds in the studied remote catchment, and this has
397 been investigated here using the diagnostic PAHs concentration ratios (Table S2,
398 Supplementary information). They show a uniformly pyrogenic origin of the detected PAHs.
399 In no sample, the petrogenic ratios of ANT/(ANT+PHE) < 0.1 or FLA/(FLA+PYR) < 0.4
400 have been found (De La Torre-Roche et al., 2009; Pies et al., 2008). Based on the
401 FLA/(FLA+PYR) and FL/(FL+PYR) ratios, further distinctions can be made. The majority of



402 the collected samples exhibited both ratios exceeding 0.5, which for the latter is interpreted as
403 indicating petrol combustion origin (Ravindra et al., 2006), while the former suggests an
404 origin from grass, wood, and coal combustion (De La Torre-Roche et al., 2009) or diesel
405 (Ravindra et al., 2008). Thus, a combination of various combustion processes is likely to have
406 produced the local PAHs composition, especially as there are stable and low local sources of
407 diesel combustion to fuel the station (there is also a waste incinerator and multiple vehicles
408 necessary for the operation of the Polish Polar Station), less stable regional sources, e.g. ships
409 en route near Svalbard, and the long-range transport of both fossil-fuel- and biomass-burning-
410 related PAHs (Granberg et al., 2017; Winther et al., 2014; Zielinski et al., 2020). The lower
411 ratios of FLA/(FLA+PYR) can be interpreted as fossil fuel combustion (De La Torre-Roche
412 et al., 2009), and FL/(FL+PYR) < 0.5 may come from diesel use (Ravindra et al., 2006).
413 These have occurred first in 2010, then in September 2015 and 2016 only in the smaller upper
414 lake (Rev 1 location), and only in 2018 they occurred in more samples again. In the context of
415 the location of the small lake (more distant from the local human activity in the polar station)
416 and these ratios occurring more often in September, they can be treated as the background
417 pollution from long-range transport and remobilized from the melting permafrost. On the
418 other hand, the higher ratios may show a combination of more frequent biomass burning
419 influence and the local fuel exhaust being carried in certain meteorological conditions into the
420 Revelva valley. The strongest conclusion from the calculation of these indicator ratios is the
421 lack of petrogenic source influence in the area, which is consistent with its geological
422 substratum lacking bituminous material.

423

424 **4. Discussion**

425 4.1. Interactions between various chemical factors – principal component analysis



426 The overview of the chemical parameters in the sampling points of the Revvatnet (Figure 7a)
427 highlights the main chemical division between parts of the summer season when samples have
428 been collected. The particular sampling locations did not show any striking differences within
429 the collected dataset – it should be especially noted that the overall composition of the upper,
430 smaller lake at Rev 1 is similar to the chemical parameters of the large Revvatnet proper. The
431 year 2015 differed from the later period, yet this may represent also the different month of
432 sample collection within that year (July instead of June representing the early summer). The
433 highest contribution to overall variability in the dataset was contributed by several metals and
434 metalloids: Al, B, Ba, Ga, Li, Mn, Ni and Sr (Figure 7b), as well as TOC.

435 **Figure 7.** Principal component analysis, exploring the variability in the entire collected
436 chemical dataset in the space defined by two most prominent components (representing >37%
437 of total variability in the dataset): a.) & b.) graph for individual datapoints, with the month
438 and sampling site highlighted by colour in a.) and b.), respectively; c.) graph for variables; For
439 a colour figure, please refer to the online version of the article

440 PCAs were also conducted for subsets of variables of similar character, to explore the possible
441 reduction of the dataset (or to exhibit the variables influencing the differences in the dataset
442 the most). For metals and metalloids, if treated separately, the main variability drivers were:
443 Al, B, Ba, Li, Sr, but also V and Cr. They effectively separated the data points by both
444 sampling month and year (except the 2016-2018 period, which remained relatively similar)
445 (Figure S2, Supplementary information). The variability was spread rather evenly in the
446 PAHs domain, and those could be divided into two closely correlated groups: 1) phenanthrene
447 and pyrene, which shared some variability in concentrations with TOC, 2) the other six
448 frequently detected compounds, changing independently from TOC concentrations. A
449 significant proportion of the variability in PAHs concentrations (approximated by PC2 in
450 Figure S3b&c) appears to have resulted from an intra-seasonal difference in hydrochemical

451 regime (e.g. snowmelt as opposed to permafrost thaw as a source of PAHs in the collected
452 samples). PC1 in Figure S3c, representing 49% of the variability, has been compromised by a
453 clear outlier collected in 2010, which makes it more difficult to interpret. A renewed analysis
454 without the outlier yielded a clearer seasonal division (Figure S3a&b), where PHE and PYR
455 high concentrations aligned with late summer sampling time, while NAP and ACE did so in
456 the early summer samples. The site Rev 5 was also more clearly distinguished by its higher
457 contribution to total variability of the concentrations of PAHs (Figure S3b). Notably, in point
458 Rev 1 occurred the maxima in ANT concentrations, while in Rev 5 the absolute maxima for
459 any single PAH concentration were found (among all sampling points), and these extreme
460 values concerned FLA. Those resulted also in the highest Σ PAHs found at that location. Thus,
461 it appears that the flow of water through the lake does not necessarily facilitate cleansing it of
462 PAHs pollution (Hamilton et al., 2001).

463 4.2. A concise overview of other Arctic lake systems research

464 In the Arctic, the problem of the appearance of pollutants in various elements of the
465 environment is widespread, and therefore it is widely studied by scientists from around the
466 world. Researching lakes and river catchments in terms of their chemical composition, as well
467 as hydrochemical parameters of the catchments, is increasingly the subject of interest of
468 researchers throughout Spitsbergen and other parts of the Arctic (e.g., Franczak et al., 2016;
469 Lehmann-Konera et al., 2019; Mazurek et al., 2012; Szumińska et al., 2018; Wawrzyniak et
470 al., 2020). Lakes in the Arctic are typically ultra-oligotrophic and fed by allochthonous
471 nutrients (Hamilton et al., 2001; Lim and Douglas, 2003).

472 The most frequently studied areas in Spitsbergen in terms of the hydrochemistry of lakes are
473 the areas adjacent to the fjord Hornsund, Bellsund, Isjord and Kongsfjord, which are strongly
474 influenced by marine aerosols, which may introduce several metals and metalloids into those
475 catchments (e.g., Nawrot et al., 2016). The content of the metals such as Sr or Ba was at a



476 similar level in Revvatnet as in the other areas of Svalbard (Szumińska et al., 2018), and Sr
477 was mentioned there to come from geogenic sources in the lakes located on a raised marine
478 terrace, which is a similar feature to the environment surrounding the Revvatnet. The general
479 levels of metal and metalloid concentrations in Revvatner were also comparable to the state of
480 Lake Imandra (in the Russian Arctic) prior to the intensive human influence in the
481 Monchegorsk area (in the 1930s; Moiseenko et al., 2009), highlighting the representativeness
482 of our study site in terms of general hydrochemical features of Arctic lakes. Similarly to the
483 study results from Canadian Arctic (Lim and Douglas, 2003), in Revvatnet, several metals
484 and metalloids concentrations were also influenced by the local geology. From early to late
485 summer, most metal concentrations in Revvatnet increased or remained similar, consistently
486 with a study on seasonal changes in thermokarst lakes by Manasypov et al. (2015).

487 The organic compounds content, including pollutants, has also been tested in many
488 catchments in Spitsbergen and elsewhere in the Arctic. Smaller lakes tend to have higher TOC
489 concentrations (Lim and Douglas, 2003). Similar levels of PAHs concentrations were
490 determined, for example, in the nearby Fuglebekken catchment (average sum of determined
491 PAHs ranged from 4-600 ng L⁻¹ (Kozak et al., 2017). The content of PAHs as well as other
492 pollutants, e.g. phenols and aldehydes, as well as microplastic, was determined in lakes in
493 Bellsund area (Lehmann-Konera et al., 2019, 2018) and those located in Ny-Ålesund area
494 (González-Pleiter et al., 2020; Jiao et al., 2009). Persistent organic pollutants have also been
495 detected extensively in two high Arctic lakes in Bjørnøya (Bear Island) (Evenset et al., 2004) .
496 As the lakes on Svalbard are remote, more than hundreds km from any known point source,
497 the presence of POPs suggests that they have been transported to the area mainly by long-
498 range transport (Evenset et al., 2004), including through the pathway of seabird guano
499 (Evenset et al., 2007). As POPs pose a serious threat to the entire polar ecosystem, their
500 redistribution in the dynamic environment of the Arctic and the combined effects of pollution

501 and climate change warrant further investigation in the times of increased strain on the Arctic
502 environment (AMAP, 2011).

503 PAHs are typically formed during the incomplete combustion of fossil fuels, biomass, and
504 through other industrial activities; they also occur naturally in bituminous rocks. They have
505 been found in the Arctic environment, originating both from the long-range atmospheric
506 transport (Wang et al., 2013) and the local sources. PAHs have been found widely in polar
507 environmental media: the atmosphere, water, sediments and biota (Kosek et al., 2019a, 2019b,
508 2018, 2017, 2016; Kozak et al., 2017; Koziol et al., 2020; Polkowska et al., 2011; Potapowicz
509 et al., 2019).

510 PAHs are generally hydrophobic and many interact strongly with sedimentary organic carbon
511 (Burgess et al., 2003) and as such, PAHs are commonly associated with sediments and
512 particulate matter, which plays a role in their removal from lake water. They also
513 bioaccumulate in aquatic biota, particularly with the lower trophic levels (Besten et al., 2003;
514 Kosek et al., 2019b; Neff et al., 2005), and the ecotoxicological concerns have focused on
515 those. The toxicology of PAHs in the aquatic environments has been well documented and
516 numerous articles/books are available in relation to bioavailability (Besten et al., 2003;
517 Burgess et al., 2003), bioaccumulation (Besten et al., 2003; Meador, 2003) and toxicity
518 (Albers, 2003; Logan, 2007; Malcolm and Shore, 2003).

519

520 4.3. Revvatnet lake system in comparison to the environmental concentrations of pollutants in
521 the Hornsund area

522 As the described flow-through lake system is part of larger landscape, where pollution has
523 been studied by other researchers, we discuss its contamination level also in the context of
524 linkages to other environmental media in the surroundings. While it is impossible to compare
525 directly the metal concentrations in lake water to the concentrations in sea sediment, it is



526 nevertheless valuable information to see whether the metals with elevated concentration in the
527 catchment of the Revvatnet lake are connected to elevated concentrations of the same metals
528 in sediments located downstream from such a catchment. In Zaborska et al. (2017) the Pb
529 concentration levels found in the outside part of the Hornsund fjord (near Revelva catchment)
530 were elevated against the levels further inside the fjord (unlike the levels of Cu, Cd or Zn,
531 which were in a similar range for those sites). Of these four metals, we have complete data
532 only in 2010, later on only Zn and Cu concentrations have been measured. However, the
533 measured concentration at a ppb level or lower in lake water is not likely to influence the
534 variability in marine sediment concentrations, which range from 0.01 to >100 ppm. This is
535 consistent with the conclusion by Zaborska et al. (2017), that large-scale atmospheric and
536 oceanic processes influence the concentrations in ocean sediment, and not the local runoff. A
537 similar comparative material, but for PAHs, can be found in Pouch et al. (2017) article on
538 fjord sediments from Svalbard. The concentrations measured by them ranged 37.3 to 1973
539 ppb for $\Sigma 12$ PAHs. Such levels are more likely to be influenced by the local runoff, however
540 such an influence is far from certain (the concentrations in lake water oscillated around 1 ppb
541 for $\Sigma 8$ PAHs). This is consistent with the interpretation of spatial variability in POPs
542 concentrations in the fjord sediments may be influenced by local secondary sources of
543 contamination, such as melting glaciers, especially as sediment is a preferential deposition
544 media for POPs (in comparison to water) due to their hydrophobicity. The sources of PAHs in
545 both lake water and sea sediment, according to PAHs concentration indicator ratios, were
546 consistently pyrolytic (Table S2 and Pouch et al., 2017). We conclude that the catchments,
547 which are in majority not glaciated, like that of the Revvatnet lake, also contain non-
548 negligible POPs contamination and should be monitored for their concentration levels in the
549 future.



550 A high concentration of the ^{90}Sr radionuclide, which is of anthropogenic descent, was found
551 on the nearby Hans glacier (Łokas et al., 2016), indicating that perhaps the high
552 concentrations of Sr in the waters of Revvatnet (up to 80 ppb) were also influenced by human
553 activity (possibly via long-range transport of contamination). Furthermore, the cryoconite
554 upon Hans glacier was found to have heavy metal concentrations beyond those noted in local
555 rocks, especially for Pb, Cd, Cu, Zn, which testifies to their enhanced deposition from long-
556 range transport or more efficient accumulation in cryoconite holes than elsewhere. While the
557 comparison between water and sediment concentrations is not directly possible, we observe
558 that Zn and Cu were important components of the metal impurity concentrations present in
559 the Revvatnet waters (while Pb and Cd were only measured in 2010, and therefore they
560 cannot be included in this comparison). This observation cautions that at least part of the
561 metal load found in the Revvatnet may be anthropogenic, including long-range transported
562 contamination. However, the concentrations of these metals did not correlate with the
563 anthropogenic radionuclide concentrations in Łokas et al. (2016) study.

564 Since the origin of metals in cryoconite is most likely from atmospheric deposition, this
565 finding is consistent with Kozak et al. (2015), who found Zn and Cu to be provided in excess
566 by atmospheric precipitation, and likely from anthropogenic contamination. However, river
567 and lake waters experience longer contact with the geological substratum, and normally they
568 note higher concentration of soluble components present in parent rock of the area. With
569 respect to precipitation, it is also helpful to explore whether precipitation events bring a
570 significant load of certain metals, which has been shown to be possible especially for Mn, Sr
571 and Zn in 2010 (Kozioł et al., 2020). Furthermore, Sr was found to show a higher
572 concentration in smaller lakes of the area than in Revvatnet (in 2010; Kozioł et al., 2020),
573 which supports it could have originated from precipitation. In Kozak et al. (2015) study on
574 precipitation waters in the area, Mn and Sr were notably changing in concentration, depending

575 on the local wind direction during the rainfall event, which could mean that despite their
576 transfer through rain, these metals have local rather than distant sources in the Hornsund
577 region. Furthermore, in the snow cover of Hans glacier (Koziol et al., 2021; *in press*), both
578 Mn and Sr were found to cluster (by concentration variability) with natural sources in rock
579 and seawater, respectively, and Sr was shown to originate from seawater in 39% (median
580 seawater fraction of Sr in the snow samples collected). On the other hand, Szumińska et al.,
581 (2018) claim that Sr in lakes in Bellsund area is geogenic, while Mn, Co, Ni, Cu, Ga, Ba and
582 Cd in that site are interpreted to be transported there airborne. In the Koziol et al. (2021; *in*
583 *press*) study, Cu also clustered with the natural geogenic elements, as did Mn. In the same
584 study, Zn clustered with the likely antropogenic elements from long-range transport.
585 However, due to its presence in ore-bearing veins in the Revvatnet surroundings, the
586 composition of surface waters may still be impacted importantly by substratum, and not only
587 precipitation, and this is supported by the spatial distribution of Zn concentrations in the lakes
588 of the area (Koziol et al., 2020).

589

590 **5. Summary and conclusions**

591 The particular chemical compositions of Arctic lakes are dependent on several factors,
592 including proximity to the ocean and geology for inorganic materials, and for organic
593 compounds compositions depend on the extent and type of terrestrial vegetation in the
594 surrounding catchment, as well as anthropogenic pollution (ACIA, 2005; AMAP, 2016;
595 Kling, 2009). In this article, we presented the characteristics of one such a lake system, the
596 flow-through Revvatnet in southern Svalbard, composed of a smaller upper lake and a large
597 lower lake, cone-shaped, reaching -29.1 m depth. The system has typical morphometric
598 characteristics of a lake formed by glacial processes. The pH of its waters was near-neutral, its
599 annual medians ranging 6.7 to 7.8, while the EC was rather low (which is typical for many

600 Arctic lakes), ranging 7 to 147 $\mu\text{S cm}^{-1}$. The significant downstream changes in metal and
601 metalloid concentrations consisted of increases downstream (with the exception of Zn), which
602 is likely a result of the slow dissolution of suspended particles in the lake system. However,
603 such changes for PAHs were not consistent in direction and relatively few. The ΣPAHs
604 concentrations ranged from <LOD to 2151 ng L^{-1} , and they were not petrogenic according to
605 indicator PAHs concentration ratios – they appear to come from a mixture of combustion
606 processes, near and far, from fossil fuel and biomass. The chemical composition of these
607 waters was characterised by a strong seasonal variability, and in terms of metals and
608 metalloids especially – also significant inter-annual variability. Of particular concern in terms
609 of pollution in these waters are the high maxima in organic compound and metal or metalloid
610 concentrations, and the patterns of their occurrence requires further study to determine the
611 environmental risk related to them.

612

613 **Acknowledgments:** This work was supported by two National Science Centre of Poland
614 research grants: 2013/09/N/ST10/04191 and 2017/25/N/NZ9/01506. Additionally, the authors
615 would like to thank the Institute of Geophysics, PAS, the staff and other scientists living in
616 the Polish Polar Station, Hornsund during several expeditions (especially Ł. Stachnik, K.
617 Michalczewska, D. Dąbrowska, R. Pogorzelski, M. Czarnecka, I. Kulaszewicz, B. Siepierska,
618 K. Stachniak, A. Piwowarczyk, J. Potapowicz, S. Czapiewski) for supporting the field
619 sampling. Moreover, Stanisław Chmiel from Maria Curie-Skłodowska University, Faculty of
620 Earth Sciences and Spatial Management in Lublin, Poland, is thanked for enabling the
621 determination of trace elements described in this paper.

622

623 **Conflict of Interests:** The authors declare no conflict of interest. The funding institution had
624 no role in the design of the study, in the samples collection, analyses, interpretation of data, in
625 the writing of the manuscript, and in the decision to publish the results.

626

627 **References**

628 ACIA, 2005. Impacts of a warming Arctic: Arctic climate impact assesment.

629 Albers, P.H., 2003. Petroleum and individual polycyclic aromatic hydrocarbons, in: Hoffman,
630 D.J., Rattner, B.A., Jr., G.A.B., Jr., J.C. (Eds.), Handbook of Ecotoxicology. Lewis
631 Publishers, Boca Raton, FL, pp. 342–360.

632 AMAP, 2016. AMAP Assessment 2015: Temporal Trends in Persistent Organic Pollutants in
633 the Arctic. Oslo.

634 AMAP, 2011. Combined Effects of Selected Pollutants and Climate Change in the Arctic
635 Environment. Arctic Monitoring and Assessment Programme, Oslo.

636 AMAP, 2009. Update on Selected Climate Issues of Concern: Observations, Short-lived
637 Climate Forcers, Arctic Carbon Cycle, and Predictive Capability. Oslo.

638 Bååth, E., Kritzberg, E., 2015. pH Tolerance in Freshwater Bacterioplankton: Trait Variation
639 of the Community as Measured by Leucine Incorporation. Appl. Environ. Microbiol. 81,
640 7411–7419. <https://doi.org/10.1128/AEM.02236-15>

641 Besten, P., Ten Hulscher, D., van Hattum, B., 2003. Bioavailability, Uptake and Effects of
642 PAHs in Aquatic Invertebrates in Field Studies. pp. 127–146.
643 <https://doi.org/10.1002/0470867132.ch8>

644 Burgess, R.M., Ahrens, M.J., Hickey, C.W., 2003. Geochemistry of PAHs in aquatic



645 environments: A synthesis of distribution, source, persistence, partitioning and
646 bioavailability, in: Douben, P.E.T. (Ed.), PAHs: An Ecological Perspective. John Wiley
647 & Sons Ltd, Chichester, UK, pp. 36–45.

648 CliC/AMAP/IASC, 2016. The Arctic freshwater system in a changing Climate, The Arctic
649 Freshwater in a changing Climate. WCRP CLimate and Cryosphere (CLiC) Project,
650 Arctic Monitoring and Assessment Programme (AMAP), International Arctic Science
651 Committee (IASC).

652 Colombo, N., Salerno, F., Gruber, S., Freppaz, M., Williams, M., Fratianni, S., Giardino, M.,
653 2018. Review: Impacts of permafrost degradation on inorganic chemistry of surface
654 fresh water. *Glob. Planet. Change* 162, 69–83.
655 <https://doi.org/10.1016/j.gloplacha.2017.11.017>

656 De La Torre-Roche, R.J., Lee, W.Y., Campos-Díaz, S.I., 2009. Soil-borne polycyclic
657 aromatic hydrocarbons in El Paso, Texas: Analysis of a potential problem in the United
658 States/Mexico border region. *J. Hazard. Mater.*
659 <https://doi.org/10.1016/j.jhazmat.2008.07.089>

660 Evenset, A., Carroll, J., Christensen, G.N., Kallenborn, R., Gregor, D., Gabrielsen, G.W.,
661 2007. Seabird guano is an efficient conveyer of persistent organic pollutants (POPs) to
662 Arctic lake ecosystems. *Environ. Sci. Technol.* 41, 1173–1179.
663 <https://doi.org/10.1021/es0621142>

664 Evenset, A., Christensen, G.N., Skotvold, T., Fjeld, E., Schlabach, M., Wartena, E., Gregor,
665 D., 2004. A comparison of organic contaminants in two high Arctic lake ecosystems,
666 Bjørnøya (Bear Island), Norway. *Sci. Total Environ.* 318, 125–141.
667 [https://doi.org/10.1016/S0048-9697\(03\)00365-6](https://doi.org/10.1016/S0048-9697(03)00365-6)

- 668 Franczak, Ł., Kociuba, W., Gajek, G., 2016. Runoff Variability in the Scott River (SW
669 Spitsbergen) in Summer Seasons 2012–2013 in Comparison with the Period 1986–2009.
670 Quaest. Geogr. 35, 39–50. <https://doi.org/10.1515/quageo-2016-0025>
- 671 González-Pleiter, M., Velázquez, D., Edo, C., Carretero, O., Gago, J., Barón-Sola, Á.,
672 Hernández, L.E., Yousef, I., Quesada, A., Leganés, F., Rosal, R., Fernández-Piñas, F.,
673 2020. Fibers spreading worldwide: Microplastics and other anthropogenic litter in an
674 Arctic freshwater lake. *Sci. Total Environ.* 722, 137904.
675 <https://doi.org/10.1016/j.scitotenv.2020.137904>
- 676 Granberg, M.E., Ask, A., Gabrielsen, G.W., 2017. Local contamination in Svalbard.
677 Overview and suggestions for remediation actions, Kortrapport / Brief Report.
678 Norwegian Polar Institute, Tromsø.
- 679 Gutry-Korycka, M., Werner-Więckowska, H., 1996. Przewodnik do hydrologicznych badań
680 terenowych [Hydrological field research guide], 2nd ed. PWN, Warsaw.
- 681 Hamilton, P.B., Gajewski, K., Atkinson, D.E., Lean, D.R.S., 2001. Physical and chemical
682 limnology of 204 lakes from the Canadian Arctic Archipelago. *Hydrobiologia* 457, 133–
683 148. <https://doi.org/10.1023/A:1012275316543>
- 684 Hung, H., Kallenborn, R., Breivik, K., Su, Y., Brorström-Lundén, E., Olafsdottir, K.,
685 Thorlacius, J.M., Leppänen, S., Bossi, R., Skov, H., Manø, S., Patton, G.W., Stern, G.,
686 Sverko, E., Fellin, P., 2010. Atmospheric monitoring of organic pollutants in the Arctic
687 under the Arctic Monitoring and Assessment Programme (AMAP): 1993–2006. *Sci.*
688 *Total Environ.* 408, 2854–2873. <https://doi.org/10.1016/j.scitotenv.2009.10.044>
- 689 Jania, J., Pulina, M., Karczewski, A., [Scientific Editors], 1984. Hornsund. Spitsbergen.
690 Geomorphology. Map 1:75000.

- 691 Jiao, L., Zheng, G.J., Binh, T., Richardson, B., Chen, L., Zhang, Y., Yeung, L.W., Lam,
692 J.C.W., Yang, X., Lam, P.K.S., Wong, M.H., 2009. Persistent toxic substances in remote
693 lake and coastal sediments from Svalbard , Norwegian Arctic : Levels , sources and
694 fluxes. *Environ. Pollut.* 157, 1342–1351. <https://doi.org/10.1016/j.envpol.2008.11.030>
- 695 Kallenborn, R., Reiersen, L.-O., Olseng, C.D., 2012. Long-term atmospheric monitoring of
696 persistent organic pollutants (POPs) in the Arctic: a versatile tool for regulators and
697 environmental science studies. *Atmos. Pollut. Res.* 3, 485–493.
698 <https://doi.org/10.5094/APR.2012.056>
- 699 Kling, G., 2009. Lakes of the Arctic, in: *Encyclopedia of Inland Waters*. pp. 577–588.
700 <https://doi.org/10.1016/B978-012370626-3.00030-2>
- 701 Kosek, K., Jankowska, K., Polkowska, Ż., 2017. Bacterial presence in polar regions
702 associated with environment modification by chemical compounds including
703 contaminants. *Environ. Rev.* 25, 481–491. <https://doi.org/10.1139/er-2017-0007>
- 704 Kosek, K., Kozak, K., Koziół, K., Jankowska, K., Chmiel, S., Polkowska, Ż., 2018. The
705 interaction between bacterial abundance and selected pollutants concentration levels in
706 an arctic catchment (southwest Spitsbergen, Svalbard). *Sci. Total Environ.* 622–623,
707 913–923. <https://doi.org/10.1016/j.scitotenv.2017.11.342>
- 708 Kosek, K., Koziół, K., Luczkiewicz, A., Jankowska, K., Chmiel, S., Polkowska, Ż., 2019a.
709 Environmental characteristics of a tundra river system in Svalbard. Part 2: Chemical
710 stress factors. *Sci. Total Environ.* 653, 1585–1596.
711 <https://doi.org/https://doi.org/10.1016/j.scitotenv.2018.11.012>
- 712 Kosek, K., Luczkiewicz, A., Koziół, K., Jankowska, K., Ruman, M., Polkowska, Ż., 2019b.
713 Environmental characteristics of a tundra river system in Svalbard. Part 1: Bacterial

- 714 abundance, community structure and nutrient levels. *Sci. Total Environ.* 653, 1571–
715 1584. <https://doi.org/10.1016/j.scitotenv.2018.11.378>
- 716 Kosek, K., Polkowska, Ż., Żyszka, B., Lipok, J., 2016. Phytoplankton communities of polar
717 regions–Diversity depending on environmental conditions and chemical
718 anthropopressure. *J. Environ. Manage.* 171, 243–259.
719 <https://doi.org/10.1016/j.jenvman.2016.01.026>
- 720 Kozak, K., Koziół, K., Luks, B., Chmiel, S., Ruman, M., Marć, M., Namieśnik, J.,
721 Polkowska, Ż., 2015. The role of atmospheric precipitation in introducing contaminants
722 to the surface waters of the Fuglebekken catchment, Spitsbergen. *Polar Res.* 34, 24207.
723 <https://doi.org/10.3402/polar.v34.24207>
- 724 Kozak, K., Polkowska, Ż., Stachnik, Ł., Luks, B., Chmiel, S., Ruman, M., Lech, D., Koziół,
725 K., Tsakovski, S., Simeonov, V., 2016. Arctic catchment as a sensitive indicator of the
726 environmental changes: distribution and migration of metals (Svalbard). *Int. J. Environ.*
727 *Sci. Technol.* 13. <https://doi.org/10.1007/s13762-016-1137-6>
- 728 Kozak, K., Ruman, M., Kosek, K., Karasiński, G., Stachnik, Ł., Polkowska, Ż., 2017. Impact
729 of Volcanic Eruptions on the Occurrence of PAHs Compounds in the Aquatic Ecosystem
730 of the Southern Part of West Spitsbergen. *Water (MDPI)* 9, 1–21.
731 <https://doi.org/10.3390/w9010042>
- 732 Koziół, K., Ruman, M., Pawlak, F., Chmiel, S., Polkowska, Ż., 2020. Spatial Differences in
733 the Chemical Composition of Surface Water in the Hornsund Fjord Area: A Statistical
734 Analysis with A Focus on Local Pollution Sources. *Water (MDPI)* 12, 1–22.
735 <https://doi.org/10.3390/w12020496>
- 736 Koziół, K., Uszczyk, A., Pawlak, F., Frankowski, M., Polkowska, Z., 2021. Seasonal and

- 737 Spatial Differences in Metal and Metalloid Concentrations in the Snow Cover of
738 Hansbreen, Svalbard. *Front. Earth Sci.* 8. <https://doi.org/10.3389/feart.2020.538762>
- 739 Kuriata-Potasznik, A., Szymczyk, S., Skwierawski, A., Glińska-Lewczuk, K., Cymes, I.,
740 2016. Heavy metal contamination in the surface layer of bottom sediments in a flow-
741 through lake: A case study of Lake Symsar in Northern Poland. *Water (Switzerland)* 8.
742 <https://doi.org/10.3390/w8080358>
- 743 Laing, J.R., Hopke, P.K., Hopke, E.F., Husain, L., Dutkiewicz, V.A., Paatero, J., Viisanen,
744 Y., 2014. Long-term particle measurements in Finnish Arctic: Part I – Chemical
745 composition and trace metal solubility. *Atmos. Environ.* 88, 275–284.
746 <https://doi.org/10.1016/j.atmosenv.2014.03.002>
- 747 Lange, W., 1993. *Metody badań fizyczno-limnologicznych [en: Methods of physical*
748 *limnological research]*. University of Gdansk, Gdansk.
- 749 Lawrence, G., 2002. Persistent episodic acidification of streams linked to acid rain effects on
750 soil. *Atmos. Environ.* 36, 1589–1598. [https://doi.org/10.1016/S1352-2310\(02\)00081-X](https://doi.org/10.1016/S1352-2310(02)00081-X)
- 751 Lehmann-Konera, S., Franczak, Ł., Kociuba, W., Szumińska, D., Chmiel, S., Polkowska, Ż.,
752 2018. Comparison of hydrochemistry and organic compound transport in two non-
753 glaciated high Arctic catchments with a permafrost regime (Bellsund Fjord,
754 Spitsbergen). *Sci. Total Environ.* 613–614, 1037–1047.
755 <https://doi.org/10.1016/j.scitotenv.2017.09.064>
- 756 Lehmann-Konera, S., Kociuba, W., Chmiel, S., Franczak, Ł., Polkowska, Ż., 2019.
757 Concentrations and loads of DOC, phenols and aldehydes in a proglacial arctic river in
758 relation to hydro-meteorological conditions. A case study from the southern margin of
759 the Bellsund Fjord – SW Spitsbergen. *CATENA* 174, 117–129.



- 760 <https://doi.org/10.1016/j.catena.2018.10.049>
- 761 Lim, D.S.S., Douglas, M.S.V., 2003. Limnological Characteristics of 22 Lakes and Ponds in
762 the Haughton Crater Region of Devon Island, Nunavut, Canadian High Arctic. *Arctic,*
763 *Antarct. Alp. Res.* 35, 509–519. [https://doi.org/10.1657/1523-](https://doi.org/10.1657/1523-0430(2003)035[0509:LCOLAP]2.0.CO;2)
764 [0430\(2003\)035\[0509:LCOLAP\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2003)035[0509:LCOLAP]2.0.CO;2)
- 765 Logan, D.T., 2007. Perspective on Ecotoxicology of PAHs to Fish. *Hum. Ecol. Risk Assess.*
766 *An Int. J.* 13, 302–316. <https://doi.org/10.1080/10807030701226749>
- 767 Łokas, E., Zaborska, A., Kolicka, M., Różycki, M., Zawierucha, K., 2016. Accumulation of
768 atmospheric radionuclides and heavy metals in cryoconite holes on an Arctic glacier.
769 *Chemosphere* 160, 162–172. <https://doi.org/10.1016/j.chemosphere.2016.06.051>
- 770 Mackay, D., Wania, F., 1995. Transport of contaminants to the Arctic: partitioning, processes
771 and models. *Sci. Total Environ.* 160/161, 25–38.
- 772 Malcolm, H.M., Shore, R.F., 2003. Effects of PAHs on Terrestrial and Freshwater Birds,
773 Mammals and Amphibians, in: *PAHs: An Ecotoxicological Perspective*. John Wiley &
774 Sons, Ltd, Chichester, UK, pp. 225–241. <https://doi.org/10.1002/0470867132.ch12>
- 775 Manasyov, R.M., Vorobyev, S.N., Loiko, S. V., Kritzkov, I. V., Shirokova, L.S.,
776 Shevchenko, V.P., Kirpotin, S.N., Kulizhsky, S.P., Kolesnichenko, L.G., Zemtsov, V.A.,
777 Sinkinov, V. V., Pokrovsky, O.S., 2015. Seasonal dynamics of organic carbon and
778 metals in thermokarst lakes from the discontinuous permafrost zone of western Siberia.
779 *Biogeosciences* 12, 3009–3028. <https://doi.org/10.5194/bg-12-3009-2015>
- 780 Marszałek, H., Górnjak, D., 2017. Changes in water chemistry along the newly formed High
781 Arctic fluvial–lacustrine system of the Brattegg Valley (SW Spitsbergen, Svalbard).
782 *Environ. Earth Sci.* 76. <https://doi.org/10.1007/s12665-017-6772-9>

- 783 Mazurek, M., Paluszkiewicz, R., Rachlewicz, G., Zwoliński, Z., 2012. Variability of Water
784 Chemistry in Tundra Lakes, Petuniabukta Coast, Central Spitsbergen, Svalbard. *Sci.*
785 *World J.* 2012, 1–13. <https://doi.org/10.1100/2012/596516>
- 786 Meador, J.P., 2003. Bioaccumulation of PAHs in Marine Invertebrates, in: *PAHs: An*
787 *Ecotoxicological Perspective*. John Wiley & Sons, Ltd, Chichester, UK, pp. 147–171.
788 <https://doi.org/10.1002/0470867132.ch9>
- 789 Moiseenko, T.I., Sharov, A.N., Vandish, O.I., Kudryavtseva, L.P., Gashkina, N.A., Rose, C.,
790 2009. Long-term modification of Arctic lake ecosystems: Reference condition,
791 degradation under toxic impacts and recovery (case study Imandra Lakes, Russia).
792 *Limnologica* 39, 1–13. <https://doi.org/10.1016/j.limno.2008.03.003>
- 793 Mueller, D.R., Van Hove, P., Antoniadis, D., Jeffries, M.O., Vincent, W.F., 2009. High
794 Arctic lakes as sentinel ecosystems: Cascading regime shifts in climate, ice cover, and
795 mixing. *Limnol. Oceanogr.* 54, 2371–2385.
796 https://doi.org/10.4319/lo.2009.54.6_part_2.2371
- 797 Nawrot, A.P., Migala, K., Luks, B., Pakszys, P., Głowacki, P., 2016. Chemistry of snow
798 cover and acidic snowfall during a season with a high level of air pollution on the Hans
799 Glacier, Spitsbergen. *Polar Sci.* 10, 249–261. <https://doi.org/10.1016/j.polar.2016.06.003>
- 800 Neff, J.M., Stout, S.A., Gunster, D.G., 2005. Ecological Risk Assessment of Polycyclic
801 Aromatic Hydrocarbons in Sediments: Identifying Sources and Ecological Hazard.
802 *Integr. Environ. Assess. Manag.* 1, 22. https://doi.org/10.1897/IEAM_2004a-016.1
- 803 Olichwer, T., Tarka, R., Modelska, M., 2013. Chemical composition of groundwaters in the
804 Hornsund region, southern Spitsbergen. *Hydrol. Res.* 44, 117–130.
805 <https://doi.org/10.2166/nh.2012.075>

- 806 Pies, C., Hoffmann, B., Petrowsky, J., Yang, Y., Ternes, T.A., Hofmann, T., 2008.
807 Characterization and source identification of polycyclic aromatic hydrocarbons (PAHs)
808 in river bank soils. *Chemosphere*. <https://doi.org/10.1016/j.chemosphere.2008.04.021>
- 809 Polkowska, Ż., Cichała-Kamrowska, K., Ruman, M., Koziół, K., Krawczyk, W.E.,
810 Namieśnik, J., 2011. Organic pollution in surface waters from the Fuglebekken basin in
811 Svalbard, Norwegian Arctic. *Sensors* 11, 8910–29. <https://doi.org/10.3390/s110908910>
- 812 Potapowicz, J., Szumińska, D., Szopińska, M., Polkowska, Ż., 2019. The influence of global
813 climate change on the environmental fate of anthropogenic pollution released from the
814 permafrost. *Sci. Total Environ.* 651, 1534–1548.
815 <https://doi.org/10.1016/j.scitotenv.2018.09.168>
- 816 Pouch, A., Zaborska, A., Pazdro, K., 2017. Concentrations and origin of polychlorinated
817 biphenyls (PCBs) and polycyclic aromatic hydrocarbons (PAHs) in sediments of western
818 Spitsbergen fjords (Kongsfjorden, Hornsund, and Adventfjorden). *Environ. Monit.*
819 *Assess.* 189. <https://doi.org/10.1007/s10661-017-5858-x>
- 820 Prowse, T.D., Ommanney, C.S.L., 1990. Northern hydrology: Canadian perspectives.
821 National Hydrology Research Institute, Saskatoon, Sask., Canada.
- 822 Ravindra, K., Bencs, L., Wauters, E., De Hoog, J., Deutsch, F., Roekens, E., Bleux, N.,
823 Berghmans, P., Van Grieken, R., 2006. Seasonal and site-specific variation in vapour and
824 aerosol phase PAHs over Flanders (Belgium) and their relation with anthropogenic
825 activities. *Atmos. Environ.* <https://doi.org/10.1016/j.atmosenv.2005.10.011>
- 826 Ravindra, K., Sokhi, R., Van Grieken, R., 2008. Atmospheric polycyclic aromatic
827 hydrocarbons: Source attribution, emission factors and regulation. *Atmos. Environ.*
828 <https://doi.org/10.1016/j.atmosenv.2007.12.010>

- 829 Smith, L.C., Sheng, Y., MacDonald, G.M., 2007. A first pan-Arctic assessment of the
830 influence of glaciation, permafrost, topography and peatlands on northern hemisphere
831 lake distribution. *Permafr. Periglac. Process.* 18, 201–208.
832 <https://doi.org/10.1002/ppp.581>
- 833 Smulikowski, W., 1965. Petrology and some structural data of lower metamorphic formations
834 of the Hecla Hoek Succession in Hornsund, Vestspitsbergen, *Studia Geologica Polonica*.
835 Warsaw.
- 836 Szumińska, D., Szopińska, M., Lehmann-Konera, S., Franczak, Ł., Kociuba, W., Chmiel, S.,
837 Kalinowski, P., Polkowska, Ż., 2018. Water chemistry of tundra lakes in the periglacial
838 zone of the Bellsund Fiord (Svalbard) in the summer of 2013. *Sci. Total Environ.* 624,
839 1669–1679. <https://doi.org/10.1016/j.scitotenv.2017.10.045>
- 840 Tank, S.E., Lesack, L.F.W., McQueen, D.J., 2009. Elevated pH regulates bacterial carbon
841 cycling in lakes with high photosynthetic activity. *Ecology* 90, 1910–1922.
842 <https://doi.org/10.1890/08-1010.1>
- 843 Van Hove, P., Belzile, C., Gibson, J.A., Vincent, W.F., 2006. Coupled landscape-lake
844 evolution in High Arctic Canada. *Can. J. Earth Sci.* 43, 533–546.
845 <https://doi.org/10.1139/e06-003>
- 846 Vincent, W.F., Laybourn-Parry, J., 2008. *Polar Lakes and Rivers: Limnology of Arctic and*
847 *Antarctic Aquatic Ecosystems*. Oxford University Press.
848 <https://doi.org/10.1093/acprof:oso/9780199213887.001.0001>
- 849 Wang, Z., Na, G., Ma, X., Fang, X., Ge, L., Gao, H., Yao, Z., 2013. Occurrence and
850 gas/particle partitioning of PAHs in the atmosphere from the North Pacific to the Arctic
851 Ocean. *Atmos. Environ.* 77, 640–646. <https://doi.org/10.1016/j.atmosenv.2013.05.052>

- 852 Wawrzyniak, T., Majerska, M., Osuch, M., 2020. Hydrometeorological dataset (2014–2019)
853 from the high Arctic unglaciated catchment Fuglebekken (Svalbard). *Hydrol. Process.*
854 *hyp.13974*. <https://doi.org/10.1002/hyp.13974>
- 855 Wawrzyniak, T., Osuch, M., 2020. A 40-year High Arctic climatological dataset of the Polish
856 Polar Station Hornsund (SW Spitsbergen, Svalbard). *Earth Syst. Sci. Data* 12, 805–815.
857 <https://doi.org/10.5194/essd-12-805-2020>
- 858 Wetzel, R.G., 2001. *Limnology: lake and river ecosystems*, 3rd ed. Elsevier, New York.
- 859 Winther, M., Christensen, J.H., Plejdrup, M.S., Ravn, E.S., Eriksson, Ó.F., Kristensen, H.O.,
860 2014. Emission inventories for ships in the arctic based on satellite sampled AIS data.
861 *Atmos. Environ.* 91, 1–14. <https://doi.org/10.1016/j.atmosenv.2014.03.006>
- 862 Wojciechowski, J., 1964. Ore-bearing veins of the Hornsund area, Vestspitsbergen, *Studia*
863 *Geologica Polonica Polonica*. Institute of Geological Sciences, Polish Academy of
864 Sciences, Warsaw.
- 865 Wojtuń, B., Samecka-Cymerman, A., Kolon, K., Kempers, A.J., Skrzypek, G., 2013. Metals
866 in some dominant vascular plants, mosses, lichens, algae, and the biological soil crust in
867 various types of terrestrial tundra, SW Spitsbergen, Norway. *Polar Biol.* 36, 1799–1809.
868 <https://doi.org/10.1007/s00300-013-1399-0>
- 869 Zaborska, A., Beszczyńska-Moeller, A., Włodarska-Kowalczyk, M., 2017. History of heavy
870 metal accumulation in the Svalbard area: Distribution, origin and transport pathways.
871 *Environ. Pollut.* 231, 437–450. <https://doi.org/10.1016/j.envpol.2017.08.042>
- 872 Zielinski, T., Bolzacchini, E., Cataldi, M., Ferrero, L., Graßl, S., Hansen, G., Mateos, D.,
873 Mazzola, M., Neuber, R., Pakszys, P., Posyniak, M., Ritter, C., Severi, M., Sobolewski,
874 P., Traversi, R., Velasco-Merino, C., 2020. Study of chemical and optical properties of

875 biomass burning aerosols during long-range transport events toward the arctic in summer
876 2017. Atmosphere (Basel). 11. <https://doi.org/10.3390/ATMOS11010084>

877