

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <https://link.springer.com/article/10.1007/s00217-022-04012-z>

Postprint of: Ostrowski B., Krawczyk B., Breastfeeding as a regulating factor of the development of the intestinal microbiome in the early stages of life, EUROPEAN FOOD RESEARCH AND TECHNOLOGY (2022), DOI: [10.1007/s00217-022-04012-z](https://doi.org/10.1007/s00217-022-04012-z)

Draft Manuscript for Review

Breastfeeding as a regulating factor of the development of the intestinal microbiome in the early stages of life

Journal:	<i>European Food Research and Technology</i>
Manuscript ID	Draft
Manuscript Type:	Review article
Date Submitted by the Author:	n/a
Complete List of Authors:	Krawczyk, Beata; Gdańsk University of Technology, Department of Molecular Biotechnology and Microbiology Ostrowski , Bartosz; Gdańsk University of Technology, Department of Molecular Biotechnology and Microbiology
Keywords:	microbiota, neonatal gut, breast milk, dysbiosis, enterotypes, probiotics

SCHOLARONE™
Manuscripts

1 **Breastfeeding as a regulating factor of the development of the intestinal**

2 **microbiome in the early stages of life**

3
4
5
6
7 3 Bartosz Ostrowski¹, Beata Krawczyk^{1*}

8
9 4 ¹ Department of Molecular Biotechnology and Microbiology, Faculty of Chemistry, Gdańsk University of Technology,
10
11 5 Gdańsk, Poland

12
13 6 *Correspondence: beata.krawczyk@pg.edu.pl; <https://orcid.org/0000-0001-5528-8898>

14 7
15 8 Bartosz Ostrowski: barostro2@student.pg.edu.pl; <https://orcid.org/0000-0002-6316-9001>
16 9

17 18 10 **Abstract**

19
20 11 Since the first bacterial inhabitants of the human gastrointestinal tract were identified a lot of research into the study of the
21
22 12 human microbiome and its effects on health has been conducted. Currently, it is accepted that humans have a symbiotic
23
24 13 relationship with the gut microbiome, though the specifics of this relationship are not well understood. The microbiome of
25
26 14 neonates constantly changes and appears to influence many facets of the infant's health and predisposition later in life. This
27
28 15 review aims to show how the microbiome develops over time. We discuss its composition, origins and stages of
29
30 16 development of microbiota, the possible health benefits of a proper neonatal microbiome, and the dangers associated with
31
32 17 dysbiosis. We emphasize the shielding, modulating, and stimulating effects breast milk has on the infant microbiota. The
33
34 18 methods commonly used for the study of microbiota are also discussed.
35
36 19

37 20 **Keywords:** microbiota, neonatal gut, breast milk, dysbiosis, enterotypes, probiotics
38
39 21

40 41 42 22 **Statements and Declarations**

43
44 23 **Competing Interests and Funding:** The authors have no relevant financial or non-financial interests to disclose.

45
46 24 **Conflict of interest:** The authors declare that they have no conflict of interest.

47
48 25 **Ethics approval:** Authors declared that this work does not require any ethical clearance.
49
50 26
51
52 27
53
54 28
55
56 29
57
58
59
60

30 **The importance of the gut microbiome**

31 The human microbiota has been investigated thoroughly and its effects on human health are firmly established
32 even though the precise relationship is yet to be understood [1,2]. The gut microbiome is seen as an integral part of the
33 human body and contributes to metabolic functions, protects against pathogens, and educates the immune system [3]. It is
34 often seen as an extension of the human genetic pool, with the gut microbiome encoding over 3 million genes, which
35 eclipses the 23 thousand genes present in the human genome [1]. The microbiome is flexible and can be affected by dietary
36 ingredients and the resulting changes can affect the health of the host. Transplantation of a microbiome from healthy
37 individuals to sick patients can effectively treat *Clostridium difficile* infections [4] and other applications for this procedure
38 are emerging [5]. Research on mice has linked the composition of the microbiome to obesity [6].

39 Microbes present in the gut, metabolize substrates present in consumed food creating nutrients that are usable for
40 the host while also producing bioactive compounds that modulate the immune system, physiology, and gene expression of
41 host cells [7]. Humans only produce a few hydrolases capable of hydrolyzing starches and rely on the enzymes produced by
42 the microbiome to gain energy from complex carbohydrates[8]. Short-chain fatty acids (SCFAs) produced by bacterial
43 metabolism of carbohydrates contributes to approximately 10% of the caloric requirement of humans. Additionally, these
44 fatty acids provide anti-inflammatory effects. Butyrate, a SCFA, improves the integrity of the host's intestinal epithelial
45 cells [9].

46 The gut also hosts microorganisms capable of utilizing the gaseous byproducts of fermentation such as carbon
47 dioxide and hydrogen, and through the removal of these waste products, helps drive metabolism forward [10,11]. The
48 fermentation of amino acids by these bacteria provides additional SCFAs, that can be used as fuel.[12] However, the
49 metabolism of aromatic, sulphur-containing, and basic amino acids produces pro-inflammatory, cytotoxic, and neuroactive
50 compounds [7]. Only a small portion of dietary fat reaches the colon[13] and the relationship between microbial lipid
51 metabolism and the host's health is unknown. However, it is known that free lipids have antimicrobial properties [14].
52 Saturated fatty acids promote inflammation [15], which might be one reason for the chronic inflammation present in obesity
53 [7], while omega-3 unsaturated fatty acids are anti-inflammatory [16]. .

54 Interactions with various antigens play an important role in immune system maturation. It is suggested that
55 exposure to certain microorganisms early in life is a factor in preventing the development of allergies and aids in regulating
56 immune system activity. The gut has the greatest concentration of microorganisms that humans have contact with in their
57 lives. Therefore, it is natural to assume that the gut microbiome plays an important role in immune system regulation [17].

1
2 58 The microbiome is known for modulating the secretion of antibodies and interleukins and the functions of other
3
4 59 immune cells [18]. As suggested by recent studies, the early establishment of symbiosis between the immune system and
5
6 60 the gut microbiome has a large influence on the susceptibility or the resistance to diseases later in life [19]. During the
7
8 61 weaning period, the immune system of infants undergoes rapid development. It has been shown that the microbiome takes
9
10 62 an important part in the development of isolated lymphoid follicles and the regulation of intraepithelial lymphocytes,
11
12 63 macrophages, and invariant killer T cells [18].
13
14 64

15 65 **Changes within the microbiome during pregnancy**

16
17 66 Although the adult microbiome differs between persons, it is fairly stable during life and research has revealed
18
19 67 some generalities. The most common phyla present in healthy individuals are Firmicutes (22.2 +/- 18.66%) and
20
21 68 Bacteroidetes (73.13 +/- 22.16%) followed by Proteobacteria (2.15 +/- 10.39%) and Actinobacteria, which is mostly
22
23 69 represented by the *Bifidobacterium* genus (1.82+/- 3%). A vast majority of Bacteroidetes are members of the *Bacteroides*
24
25 70 genus, with *Bacteroides dorei* being the most dominant (17.44 +/- 8.74%), while *Bacteroides fragilis* is the most
26
27 71 widespread species. The abundance of *Bifidobacteria* varies between 0.004% and 12.21%. In regards to Firmicutes, the
28
29 72 genus *Clostridium* appears to be the most common [2].
30

31 73 During pregnancy, the mother's vaginal, oral, and gut microbiota undergo significant changes, the origin of which
32
33 74 is unknown. Changes in hormonal regulation, immunity, energy homeostasis, and fat storage likely have a role in
34
35 75 influencing the microbiome [20]. The changes in the microbiome happen gradually during pregnancy. An increase in the
36
37 76 abundance of Proteobacteria and Actinobacteria is seen at the cost of *Faecalibacterium* and other SCFA producers [21].
38

39 77 During the third trimester, mothers showed a lower diversity within a single sample, while having the largest
40
41 78 diversity between different mothers. This suggests that pregnancy causes the depletion of microbial diversity, however, it
42
43 79 increases the diversity between individuals. The increased diversity between mothers lasted for up to one month postpartum
44
45 80 [21]. When transferred to germ-free mice, third-term microbiota caused more weight gain, insulin resistance, and
46
47 81 inflammatory responses than first-term microbiota. This shows that the microbiota contributes to the changes occurring
48
49 82 during pregnancy [21]. There is also evidence suggesting that an alternation in maternal microbiota during pregnancy such
50
51 83 as during exposure to antibiotics, influences the neonate's immunity and health [22]. Changes to the vaginal microbiota,
52
53 84 such as the presence of certain fungi like *Candida albicans* [23], a lower *Lactobacillus* abundance, and an increased
54
55 85 *Gardnerella* and *Ureaplasma* abundance[24] are associated with preterm birth .
56
57 86

The “in utero” origin of the microbiome

The exact source of the early microbiome is unknown. The proposed sources of early life gut bacteria are the mother’s vagina during birth, breast milk, and the mother’s gut microflora, however, the mechanism of such transfer is unknown. Recent studies have proposed the idea of in utero colonization [25].

The placenta and the amniotic fluid have always been considered sterile, however recent studies have raised doubts about this assumption. Bacteria have been isolated from the placenta and studies have shown the presence of the microorganisms in amniotic fluid [26, 27]. However, the detected biomass remains low suggesting the detected microbiota are contaminants rather than native inhabitants [28]. It has also been suggested that polymerase chain reaction (PCR) based detection might identify DNA of dead bacteria instead of living ones [29].

Bacteria have also been found in the umbilical cord suggesting the transfer of microbiota between mother and fetus [30]. However, the mechanism for such a transfer is not understood. One theory is that the bacteria are transferred from the mother’s intestine. An experiment in mice showed *Enterococcus faecium* strains fed to the mother orally were later detected in the amniotic fluid supporting this claim [31].

It has also been shown that microbial exposure of the mother during pregnancy might have a significant impact in preventing allergies [32]. Children, whose mothers were exposed to farm animals during pregnancy are less likely to develop allergies, as well as an exposure to other allergens reduced the symptoms of asthma, hay fever, and eczema in the children [22]. However, the evidence supporting the existence of a placental microbiome is still controversial.

The changes in the microbiome associated with type of delivery

One of the first big shifts in the microbial composition of the infant’s gut happens during birth, and the birth mode seems to be a major factor influencing the early microbiome. Children born from cesarean section have lower *Bifidobacteria* abundance and the colonization by *Bifidobacteria* is delayed. This delay is not affected by the form of feeding. They also have an abundance of potentially harmful *Klebsiella* and *Enterococcus*. This increase in *Klebsiella* and *Enterococcus* is also independent of antibiotic exposure, hospitalization time, and feeding.

There is evidence suggesting that children delivered vaginally are seeded by the mother’s fecal microbiota. Furthermore, these children have a more stable early microbiota than children born by cesarean section, who are inhabited by more strains associated with respiratory tract infections during the first year of life. This suggests that the passage through the vaginal canal has an important role in the early colonization of the infant’s gut [33].

1
2 115 Vaginal seeding is a procedure in which a gauze swab is used to transfer vaginal fluid, and the microorganisms
3
4 116 within it, onto an infant born via cesarean section. In theory, this should alter the infant's microbiota towards a more
5
6 117 "natural" composition. However, the evidence regarding the health benefits of this procedure is limited and harbors the
7
8 118 possibility of transferring pathogenic microorganisms. Due to the absence of evidence of the benefits and potential risks,
9
10 119 performing this procedure is currently not recommended [34].
11
12

13 121 **Breast milk composition and bioactive components**

15 122 Breast milk is the most optimal source of nutrients for newborns, but the evidence for its role in preventing health
16
17 123 problems and disease in early childhood is prevalent. Some suggest that the benefits might also apply later in life, though
18
19 124 this is inconclusive [3]. Although the artificial formula has improved since it was first introduced, it is still unable to provide
20
21 125 the same health benefits as natural human breast milk. Breastfed children have lower risks of respiratory tract infections,
22
23 126 neonatal necrotizing enterocolitis (NEC), and gastrointestinal illnesses [35]. As such, breastfeeding remains the
24
25 127 recommended feeding method of newborns, however, in certain cases, such as babies with lactose intolerance or mothers
26
27 128 who cannot breastfeed due to health reasons, it is not possible and must be replaced or supplemented by artificial formula.
28

29 129 The composition of breast milk changes over time and is considered fully mature 4 to 6 weeks after birth. The
30
31 130 colostrum, which is produced in low quantities during the first few days following birth is rich in IgA, lactoferrin,
32
33 131 leukocytes, developmental factors, sodium, magnesium, and chloride[36]. However, it contains relatively lower
34
35 132 concentrations of lactose, calcium, and potassium. This suggests that the main function of colostrum is immunogenic rather
36
37 133 than nutritional [37,38,39]. The composition of macronutrients in breast milk varies between mothers, however, remains
38
39 134 similar across populations despite differences in maternal nutrition [40]. In preterm mothers, breast milk contains higher
40
41 135 concentrations of secretory IgA, likely to compensate for the underdeveloped neonatal immune system [41].
42

43 136 The protein content of breast milk is estimated to be around 0.9 to 1.2 g/dL [36] and can be grouped into 3 major
44
45 137 classes based on where they can be found: caseins (α -casein, β -casein, and κ -casein), whey (α -lactalbumin, lactoferrin,
46
47 138 lysozyme, and secretory IgA), and mucins. Caseins are aggregated in micelles while whey proteins are present in solution
48
49 139 and mucins are incorporated into the milk fat globule membrane (MFGM) [42]. In addition to proteins, breast milk contains
50
51 140 free amino acids with higher concentrations of glutamic acid and glutamine, thought to have an appetite-regulating effect
52
53 141 [43].
54

54 142 Lipids represent 44% of the total energy provided by human milk, being the major contributor. The most common
55
56 143 fatty acids in breast milk are palmitic acid and oleic acid. Palmitic acid is mostly concentrated in the 2nd position of
57
58
59
60

1
2 144 triglycerides, which allows for increased absorption and decreased calcium malabsorption [42]. The fat in human breast
3
4 145 milk is concentrated in globules surrounded by a MFGM, which contains a high amount of bioactive compounds that play a
5
6 146 role in neurocognitive development and immune function [44]. The content of long-chain polyunsaturated fatty acids
7
8 147 (LCPUFA) is largely affected by the mother's diet, and is negatively affected by the high omega-6/omega-3 ratio present in
9
10 148 western diets. [45,46] A higher ratio of omega-6/omega-3 is positively associated with higher body fat percentages between
11
12 149 2 weeks and 4 months of age and may contribute to adiposity [47].

13 150 The main carbohydrate in human breast milk is lactose [36]. It appears at a concentration of 6.7g/100 ml exceeding
14
15 151 the concentration of other species [48]. The concentration of lactose increases in mothers with a higher volume of milk
16
17 152 production [49]. The micronutrient composition of breast milk varies by maternal diet and body stores. Breast milk contains
18
19 153 vitamins A, B1, B2, B6, B12, and D along with iodine and other micronutrients.[50,51] Regardless of diet, vitamin K is low
20
21 154 in human breast milk and should be supplemented [23d]. The effects of the micronutrients in human breast milk on infant
22
23 155 growth are not well known [43].

24
25 156 In addition to macro- and micronutrients, breast milk contains numerous bioactive components including
26
27 157 hormones, growth factors, cytokines, and immune cells. The growth factors present in milk stimulates the development of
28
29 158 the intestines, growth and maturation of neurons, repair of tissues, and protection against damage from hypoxia and
30
31 159 ischemia . T cells, stem cells, lymphocytes, and macrophages are all present in breast milk along with non-cellular immune
32
33 160 components such as immunoglobulins and cytokines. Additionally, it contains compounds such as lactoferrin, lactadherin,
34
35 161 bile salt-stimulating lipase, and mucins which serve a role in protecting the infant against bacteria and viruses [36].

36 162 37 38 163 **The infant microbiome, health risks and benefits associated with microorganisms found in the** 39 40 41 164 **neonatal gut**

42
43 165 The composition of the neonatal microbiome has substantially more plasticity than adults. It changes rapidly with
44
45 166 ageing and depends on various factors such that it is significantly different between formula-fed and breastfed babies
46
47 167 [1,9,14]. Table 1 contains a comparison of the neonatal gut and breast milk microbiota.

48
49 168 During the first week of life, the microbiome is dominated by facultative anaerobes, such as those from the Proteobacteria
50
51 169 family. These bacteria consume oxygen and shape the intestinal environment to be more habitable for obligatory anaerobes
52
53 170 which appear later [54].

54 171 Not only does breast milk contain factors that shield the underdeveloped immune system of newborns but it also appears to
55
56 172 promote the growth of certain microbes, such as *Bifidobacterium* species, due to their ability to metabolize human milk
57
58
59
60

1
2 173 oligosaccharides (HMOs), and *Lactobacilli*. *Enterococci* are also more prevalent in breastfed infants, while formula-fed
3
4 174 children show an increase in the presence of *Clostridium*, *Escherichia*, and *Bacteroides* [1]. The microbiome has a big effect
5
6 175 on infant development. Studies show that mice grown in germ-free environments have poor growth, decreased weight,
7
8 176 intestinal problems, and altered neurodevelopment [55].

9
10 177 Overall, neonates are characterized by lower bacterial diversity than adults with breast-fed infants having less
11
12 178 diversity in their gut than formula-fed infants [19]. It also appears that children on a mixed diet have the bacterial diversity
13
14 179 profile of formula-fed infants. The microbiome shifts quite dramatically when solid foods are introduced to the infants' diet,
15
16 180 with a shift in dominance towards fiber-fermenting *Bacteroides* and *Firmicutes* and moving towards a composition similar
17
18 181 to that of adults [19].

19 182 The most studied members of the gut microbiome are the model organism *Escherichia coli* along with the genera
20
21 183 *Lactobacillus*, *Bifidobacterium*, and potential pathogens such as *Clostridium*. *Escherichia coli* is a microorganism
22
23 184 commonly found in the lower intestine of mammals. Although most strains are harmless and even aid in the health of the
24
25 185 host by producing exogenous vitamins such as K vitamins [56]. Unfortunately, there exist *E.coli* strains that can cause
26
27 186 diarrhoea, respiratory tract infections, pneumonia, and urinary tract infections [57]. The pathogenicity of *E.coli* is dependent
28
29 187 on several virulence factors such as fimbriae, adhesions, toxins, and other elements which can directly interact with
30
31 188 epithelial cells of the intestinal, respiratory, and urinary tract [58]. It has been shown that *E.coli* strains in breast-fed infants
32
33 189 have fewer virulence factors such as the K-capsule and have increased type 1 fimbriae expression. The IgA contained in the
34
35 190 mothers' milk can bind to the same type of fimbriae [59]. It appears that *E.coli* isolated from breast-fed infants show higher
36
37 191 adherence to epithelial cells of the colon compared to those in formula-fed children [60]. Type 1 fimbriae expression has
38
39 192 been shown to enhance the virulence of *E.coli* in the urinary tract [61]. However, breast-fed infants have shown a lower risk
40
41 193 of urinary tract infection [59].

42 194 Bacteria from the *Lactobacillus* genus belong to a broad group called the lactic acid bacteria, defined by their
43
44 195 ability to produce lactic acid as the sole or main byproduct of carbohydrate metabolism. They are known to colonize oral
45
46 196 cavities, gastrointestinal tracts, and vaginas of humans and animals. The presence of *Lactobacilli* in the gut is commonly
47
48 197 regarded as beneficial to the host and are frequently used as probiotics. However, there is little evidence supporting any
49
50 198 major role this genus might have on the human gastrointestinal tract. On the contrary, evidence suggests only a small
51
52 199 number of *Lactobacilli* are true residents of the mammalian gastrointestinal tract, and that most are instead allochthonous
53
54 200 and derived from food or the oral cavity. Recent research, based on the amplification of 16S rRNA genes, shows that
55
56 201 *Lactobacilli* make up only a small fraction of the total microbiota [62]. Attempts to treat infant colic with *Lactobacilli*
57
58 202 supplementation have shown no benefit [63]. However, it has been shown that supplementation with *Lactobacillus*

1
2 203 *rhamnosus* reduces the duration of diarrhea [64]. Studies in animals have shown that treatment with *Lactobacillus* can
3
4 204 improve enteritis recovery [65] and inhibit the colonization of the pathogenic *E.coli* K1 strain [66].

5
6 205 *Clostridium* is a genus of Gram-positive, anaerobic, and spore-forming bacilli. *C. difficile* is a major cause of
7
8 206 diarrhoea and potentially lethal nosocomial infections, especially in the elderly [67]. However, its pathogenicity in infants is
9
10 207 still debated [68]. Up to 70% of healthy newborns can be colonized by *C. difficile* during the first months of life and most
11
12 208 lack any symptoms of infection even when large numbers of toxin-producing bacteria are present. The underdeveloped
13
14 209 intestinal mucosa may lack *C. difficile* toxin receptors or other factors such as the immaturity of the immune system might
15
16 210 also play a role, although the true reason is unknown [68]. It is important to note that *C. difficile* infections still occur
17
18 211 especially in infants with hematological malignancies, inflammatory bowel disease, and cystic fibrosis following lung
19
20 212 transplantation [67]. Colonization by *C. difficile* is more common among formula-fed infants than among breastfed ones
21
22 213 [67] due to the lack of IgA in the formula [69]. The presence of *C. difficile* decreases with ageing and reaches the
23
24 214 prevalence levels similar to adults by 3 years of age [70].

25 215 *Bifidobacterium* is a genus of Gram-positive, anaerobic bacteria that commonly inhabit the gastrointestinal tract,
26
27 216 vagina, and oral cavities of mammals, including humans. Their presence in the gastrointestinal tract is deemed beneficial,
28
29 217 thus they are commonly added to probiotics and functional foods. *Bifidobacteria* rapidly colonize the infant gut during the
30
31 218 first weeks after birth. *Bifidobacteria* have been associated with protection from carcinogens, reduction in inflammation,
32
33 219 and regulation of gut function. They are more prevalent in babies born vaginally suggesting they are acquired from the
34
35 220 vaginal tract of the mother. Furthermore, breastfeeding supports the growth of this genus due to its ability to digest human
36
37 221 breast milk oligosaccharides. As a result, *Bifidobacteria* are a major part of the newborn microbiome. However, their
38
39 222 presence decreases rapidly with ageing and remains low but stable during adulthood [71].

40 223 41 42 224 **Enterotypes in infants and stages of gut microflora development**

43
44
45 225 In recent years, metagenomic studies have suggested that the intestinal microbiome of each human belongs to one
46
47 226 of three types based on the dominating microorganism. These genera are *Bacteroides* (Enterotype 1), *Prevotella* (Enterotype
48
49 227 2), and *Ruminococcus* (Enterotype 3). These enterotypes do not differ in functional abundance and do not correlate with any
50
51 228 factors relating to the host. However, the prevalence of certain genera indicates the use of different routes to generate energy
52
53 229 from fermentation [72]. Although the possible benefits of using the enterotype model are high, there are certain points of
54
55 230 contest when it comes to the theory. The enterotypes are not sharply delineated [72], and apparent clusters may arise from
56
57 231 certain methods of data processing even when they are not factual [73]. Additionally, by focusing on the enterotype model it

1
2 232 is possible to miss smaller changes and individual differences in the microbiota. The long-term stability of a human's
3
4 233 enterotype also comes into question [73]. Some research suggests that there are only two enterotypes, the *Prevotella* and
5
6 234 *Bacteroides* genera [74].

7
8 235 Certain studies seeking to evaluate the presence and importance of enterotypes in infants have been performed.
9
10 236 This research has identified four distinct enterotypes with the dominant microorganisms being either the Firmicutes phylum,
11
12 237 *Bifidobacterium*, *Bacteroides*, or *Prevotella* [75]. Unlike adults, the differences in enterotypes seem to be dependent on the
13
14 238 stage of gut development and can transition from a less mature into a more mature one. In particular, the strains associated
15
16 239 with Firmicutes and *Bifidobacterium* were correlated with the early developmental stages of the gut microbiota, while
17
18 240 *Bacteroides* and *Prevotella* were correlated with later stages [75]. While the enterotypes did not seem correlated with
19
20 241 antepartum or postpartum factors, certain clinical factors seemed to influence them to an extent. Type Firmicutes were more
21
22 242 common in infants delivered by C-section and in infants with lower gestational age, although these factors often appear
23
24 243 together. The duration of breastfeeding was also a factor with Firmicutes being more common in infants breastfed for
25
26 244 shorter durations while breastfeeding longer seemed to promote *Bifidobacterium* [75]. A different study, using two
27
28 245 enterotype models failed to detect a negative correlation between *Prevotella* and *Bacteroides* in infants 9 to 18 months of
29
30 246 age. However, such a correlation appeared at 36 months suggesting stable enterotypes develop between 18 and 36 months
31
32 247 of age [76].
33

34 249 **The dangers of microbial dysbiosis and factors contributing to its occurrence**

35
36 250 Multiple factors affect the composition of the infant microbiome, including but not limited to the mother's diet,
37
38 251 feeding type, and medication [55]. Dysbiosis is a term used to describe a breakdown in the balance between "protective"
39
40 252 and "harmful" intestinal bacteria [77]. Dysbiosis is associated with multiple diseases, such as obesity, type 2 diabetes,
41
42 253 hypertension, NEC, and inflammatory bowel disease, autoimmune diseases [18], asthma, food allergies, autism, and
43
44 254 opportunistic infections [19].

45
46 255 One of the most common causes of dysbiosis is antibiotic treatment. Antibiotics are the most common medication
47
48 256 prescribed for children. Studies have shown that the use of antibiotics in early life is associated with obesity and the
49
50 257 occurrence of diseases later in life. Antibiotic treatment has a long-term effect on the microbial composition and diversity in
51
52 258 the gut. Antibiotic treatment in early life has been associated with allergies, atopic diseases, autoimmune diseases, and
53
54 259 infections such as NEC [77]. Acid blockers are also associated with dysbiosis and NEC [55]. Children of obese mothers
55
56 260 have a different bacterial colonization profile than those born to nonobese mothers. These differences are maintained during
57
58
59
60

1
2 261 the first few years of life. However, the development of obesity may begin in utero due to the obesogenic and inflammatory
3
4 262 maternal environment [78].

5
6 263 Gestational diabetes mellitus (GDM) is associated with changes to the microflora of both mother and child.
7
8 264 Samples taken from GDM positive subjects exhibited less diversity than those from GDM negative patients. In addition, the
9
10 265 meconium of GDM positive mothers exhibits a higher abundance and prevalence of eukaryotic viruses possibly exposing
11
12 266 the child to a greater number of viruses [79].

13 267 Preterm infants are especially susceptible to dysbiosis due to their underdeveloped intestines. The immaturity of
14
15 268 the gastrointestinal tract and immune system coupled with altered gut microbiota can have severe health consequences.
16
17 269 Moreover, pre-terms require hospital treatment which further disturbs the microbiome and exposes the infant to the
18
19 270 influences of the hospital's environmental microbiome [41].
20

21 271 Antibiotics are routinely prescribed for preterm children to prevent infections. Although this treatment decreases
22
23 272 mortality it also alters the microbiota causing reduced bacterial diversity[80], delaying *Bifidobacteria* colonization [81] and
24
25 273 increased presence of multi-drug resistant strains [80]. Furthermore, the time required for the recovery from such
26
27 274 disruptions is positively correlated with the length of antibiotic treatment [80,81]. Additionally, artificial respiration shifts
28
29 275 the microbiome towards aerobic and facultative anaerobic bacteria due to the introduction of oxygen to an otherwise anoxic
30
31 276 gastrointestinal tract [82]. This can result in the weakening of the mucosal barrier [83] and reduced production of energy,
32
33 277 nutrients, and bioactive components [84].
34
35 278

36 279 **Modulation of the gut microbiota by probiotics and breast milk**

37
38
39 280 Probiotics are live microorganisms promoted as having health benefits when taken as food supplements, while
40
41 281 prebiotics are compounds that promote the growth or activity of beneficial microorganisms. There have been several studies
42
43 282 investigating the benefits of pre- and probiotic supplementation for infants.

44 283 Studies on animal models show that *Bifidobacteria* supplementation might counteract the effect of carcinogens,
45
46 284 help reduce diarrhea caused by viral infections or antibiotic treatment, and prevent constipation [71]. There is also evidence
47
48 285 that supplementation with *Bifidobacteria* reduces the occurrence and severity of NEC in low birth or preterm infants [85]. It
49
50 286 also has the potential to reduce the spread of gastroenteritis and diarrhea in infants in residential care units [86].
51

52 287 Attempts to treat infant colic with *Lactobacilli* supplementation has shown no benefit [63]. However,
53
54 288 supplementation with *Lactobacillus rhamnosus* reduces the duration of diarrhea [64]. *Lactobacillus* GG has been shown to
55
56 289 prevent and reduce the duration of diarrhea caused by rotavirus infections in animals [87, 88]. Animal studies have shown
57
58
59
60

1
2 290 that treatment with *Lactobacillus* can improve enteritidis recovery [65] and inhibit the colonization of the pathogenic *E.coli*
3
4 291 K1 strain [66]. *Lactobacilli* have also been shown to modulate Th1/Th2 cytokine balance [89,90] which might help in the
5
6 292 prevention of atopic disease and supplementing breastfeeding mothers or infants has been shown to reduce the incidence of
7
8 293 atopic dermatitis (eczema) [91].

9
10 294 In the absence of a mother's breast milk, donor human milk (DHM) appears to be the best substitute for helping the
11
12 295 development of preterm babies. The microbiota of children fed DHM is similar to breastfed infants, although it shows a
13
14 296 decrease in Bifidobacteriaceae and an increase in Staphylococcaceae, Clostridiaceae, and Pasteurellaceae. The
15
16 297 pasteurization of donated breast milk and the different composition of preterm milk and donated milk might contribute to
17
18 298 this effect [92].

19 299 HMOs are a type of carbohydrate present in breast milk and although they don't have any nutritional value, they
20
21 300 serve as a prebiotic stimulating the growth of proper microbiota and modulating several infant mucosal and systemic
22
23 301 immune functions [36]. These oligosaccharides differ between mothers, but this does not cause any incompatibility issues
24
25 302 [93, 94, 95]. However, it has been shown that one type of HMO, specifically disialyllacto-N-tetraose (DSLNT) is protective
26
27 303 against the risk of NEC in rats, which point to the conclusion that the protective effects of these compounds are dependant on
28
29 304 specific HMO structured [96]. A study in piglets has also shown that HMOs can reduce the symptoms of rotavirus
30
31 305 infections [97].

32
33 306 Research on the benefits of probiotics in infants has been promising and they appear to be safe. However, the
34
35 307 studies have used different strains and administration strategies thus more studies are needed to identify the ideal
36
37 308 combination. As of today, feeding breast milk from either the mother or that has been donated appears to be the best method
38
39 309 of stimulating a beneficial microbial composition.

40 310 41 42 311 **Methods for studying the microbiome**

43
44
45 312 Historically the study of human and animal microbiota has been based around traditional non-molecular methods
46
47 313 involving the isolation of microbes, microscopic observation, and growing them in culture. Although these methods have
48
49 314 been incredibly useful in the early study of the microbiome, they have several limitations. Traditional cultures tend to
50
51 315 underestimate the true variety of microorganisms present as a large number of bacteria cannot be cultivated using currently
52
53 316 known methods or require artificially created environmental conditions for that organism to grow [98,99]. Temperature, pH,
54
55 317 oxygen, and nutrient levels [98] and cultivation time [99] need to be tuned towards the studied microorganism. Furthermore,
56
57 318 the existence of mutual relationships between different bacteria further complicates the issue. In particular, the creation of a

1
2 319 biofilm, which is composed of many different microorganisms, is difficult to replicate in a lab. This limits the variety of
3
4 320 microorganisms that can be studied using traditional methods, which provides a biased view of the microbiome composition
5
6 321 with an overrepresentation of aerobic organisms [100].

7
8 322 Even with these limitations, culture methods have the unique advantage of allowing living microorganisms to be
9
10 323 studied in regards to antibiotic response and susceptibility, antigens, microorganism relationships, biofilm formation, and
11
12 324 the creation of experimental models [99]. New culture methods are still being developed to allow for the growing of
13
14 325 microorganisms previously considered uncultivable. Examples of such methods for the cultivation of hard-to-culture
15
16 326 microorganisms are the use of gnotobiotic animals [101] or the creation of artificial environments simulating the intestinal
17
18 327 environment, such as the SHIME system [102]. These methods come with the additional benefit of being able to study gut
19
20 328 microbe-host and microbe-microbe relationships [103, 104].

21 329 In response to the limitations of traditional methods, molecular methods for studying microorganisms were
22
23 330 developed. These methods involve the study of a microorganism's molecular components such as DNA, RNA, proteins, and
24
25 331 metabolites. These methods are culture-independent, meaning that the studied microorganisms do not need to be isolated
26
27 332 and cultivated in a medium. Rather, they allow for the *in vitro* study of microorganisms considered impossible to be grown.
28
29 333 The basis for most molecular methods is a variant of the DNA PCR [98]. By using PCR, the amount of DNA in a sample
30
31 334 can be increased exponentially allowing for further analysis with techniques such as Southern blotting [105]. With
32
33 335 modifications of the PCR method by using different starters, conditions, or pre-preparation techniques on the samples it is
34
35 336 possible to turn it into a diagnostic method itself. For example, ligation-mediated PCR techniques utilize the selective
36
37 337 amplification of DNA fragments generated by enzymatic restrictions creating a genetic fingerprint for a sample [106]. Other
38
39 338 methods can also provide certain insights, for instance, terminal restriction fragment length polymorphism (T-RFLP) has
40
41 339 suggested that *Clostridium* plays an important role in the pathogenesis of NEC [107]. While variants of gradient gel
42
43 340 electrophoresis have revealed the disruption of the human microbiome by antibiotic administration and identified a
44
45 341 correlation between *Sphingomonas* and NEC in human children [108,109]

46 342 With the rise of DNA sequencing technology, the ability to study complex microbial communities has increased.
47
48 343 Although its use was initially limited due to costs, improvements in the technology have allowed for cheaper, faster, and
49
50 344 more sensitive identification technologies. The increased availability of bioinformatic tools has allowed for the creation of
51
52 345 modern new generation sequencing (NGS) technology and allowed for the development of metagenomics, which is the
53
54 346 study of the total genetic material within an environmental sample. Metagenomics can be used to study microbial diversity
55
56 347 and dysbiosis of the intestine, identify new genes and microbial pathways and identify relationships between the
57
58 348 microbiome and the host's health [110]. Metagenomics aims to catalog all the genes from a microbial community by
59
60

1
2 349 random sequencing of all DNA present in a sample [110]. The gene most commonly used for sequencing is the 16S rRNA
3
4 350 gene as it is present in various microorganisms while having highly variable regions allowing for its differentiation between
5
6 351 species [111]. For fungi the 18S rRNA gene or the Internal Transcribed Spacer sequence is used [112].

7
8 352 Another method of sequencing is whole-genome shotgun sequencing which allows for the identification of viruses
9
10 353 [58g,58h] whose genetic data is missed by 16S sequencing as they lack such sequences [98]. This method can also provide
11
12 354 information regarding gene content and metabolic pathways [113]. However, a major disadvantage of this technique is that
13
14 355 the DNA from the host is also amplified and can often overwhelm the bacterial DNA. Additionally, analysis of the acquired
15
16 356 data is complex and requires a lot of computational power [114]. The sequences obtained by either method can be analyzed
17
18 357 with the assistance of bioinformatic tools and methods, such as databases stemming from data acquired by the Human
19
20 358 microbiome project or the MetaHIT project. This enables a broader understanding of the structure and function of microbial
21
22 359 communities. Metagenomic methods have revealed the relative stability of a healthy individual's microbiome and identified
23
24 360 multiple factors that affect its composition [110].

25 361 Although molecular methods are incredibly useful in the study of microorganisms they do have limitations. DNA
26
27 362 sequencing provides information on the presence of genes but doesn't give any insight into gene expression [98,99].
28
29 363 Additionally, some DNA sequences may amplify more efficiently under given conditions introducing bias to the results.
30
31 364 Furthermore, the PCR reaction does not discriminate between living and dead bacteria or their fragments [114]. The DNA
32
33 365 samples used for metagenomic analysis must be of high quality and in sufficient quantity. However, microbiome samples
34
35 366 are almost universally contaminated by human DNA. Additionally, not all identified sequences can be matched due to the
36
37 367 lack of reference and determining function on sequence homology introduces ambiguity to the results [110, 115].

38 368 A supplementary method to DNA sequencing is RNA sequencing, which allows researchers to look directly at the
39
40 369 transcriptome of microorganisms and gain insight into gene expression [116]. RNA sequencing can be used to study the
41
42 370 effects of environmental perturbations and factors on the function of the gut microbiome and identify a functional change
43
44 371 before a composition change occurs. This could allow one to preemptively detect the signs of dysbiosis [117, 118].
45
46 372 Metatranscriptomics can be used to determine the activity of genes in a defined environment, such as the human gut.
47
48 373 However, this method requires high-quality RNA samples, which are difficult to obtain and often difficult to separate
49
50 374 mRNA from other types of RNA. Additionally, mRNA is unstable and the reference databases are still insufficient [110].

51
52 375 Methods for studying protein (metaproteomics) or metabolite (metabolomics) profiles are also being developed and
53
54 376 can supplement metagenomic analysis. Metaproteomics has greatly benefited from improved methods of protein separation,
55
56 377 high throughput mass spectrometry, increased computing power, and the growth of metagenomic databases. However, such
57
58 378 methods are in their infancy and their development is difficult due to the high complexity of human samples and difficulties

1
2 379 in analyzing the data [114]. Meanwhile, metabolomic profiles of the human gut microbiota combined with other methods
3
4 380 can be used to predict the appearance of dysbiosis [119]. Methods for studying the microbiome are presented in Fig 1.

5 6 381 7 382 **Conclusion**

8
9 383 The human intestinal microbiome is an incredibly complex subject to study. Not only is it one of the richest
10
11 384 microbial ecosystems found on earth but the relationships between the host, the microbiome, and one's health are often not
12
13 385 straightforward, with each influencing the other. Furthermore, the microbiome of babies displays significant plasticity and is
14
15 386 influenced by multiple factors such as mode of birth, type of feeding, medical conditions and treatments, and is shaped by
16
17 387 the development of the infant's gut (Fig 2). However, research has identified several health effects associated with the
18
19 388 microbiome and found ways to influence the developing microbiome, with some of these methods being put into practice.
20
21 389 Although several issues remain unclear.

22 390 The origins and roles of pre- and postpartum factors on the development of an infant's microbiome are still
23
24 391 inconclusive. The specific roles certain classes of microorganisms assume in the gut and the importance of their metabolic
25
26 392 products have yet to be discovered. New methods for studying microorganisms have been crucial in enhancing our current
27
28 393 knowledge base and in conjunction with traditional methods have provided further insight into the ecosystem of the human
29
30 394 gut. With such knowledge, new ways of treating illnesses and improving an infant's health may appear.

31 32 395 33 34 396 **Author contributions**

35
36 397 Bartosz Ostrowski: Conceptualization, Collected data, Writing - Original Draft

37
38 398 Beata Krawczyk: Conceptualization, Visualization, Supervision, Writing – Review & Editing

39
40 399 All authors approved the final version of the manuscript and agreed to be accountable for all aspects of the work.
41
42 400

43 44 401 **References**

- 45
46 402 1. Rinninella E, Raoul P, Cintoni M, et al (2019). What is the Healthy Gut Microbiota Composition? A Changing
47
48 403 Ecosystem across Age, Environment, Diet, and Diseases. *Microorganisms* 7(1):14.
49
50 404 <https://doi.org/10.3390/microorganisms7010014>
51
52 405 2. King CH, Desai H, Sylvetsky AC, et al (2019). Baseline human gut microbiota profile in healthy people and
53
54 406 standard reporting template. *PloS one* 14(9), e0206484. <https://doi.org/10.1371/journal.pone.0206484>

- 1
2 407 3. Stiemsma LT, Michels KB (2018). The Role of the Microbiome in the Developmental Origins of Health and
3
4 408 Disease. *Pediatrics* 141(4), e20172437. <https://doi.org/10.1542/peds.2017-2437>
- 5
6 409 4. Bakken JS. Fecal bacteriotherapy for recurrent *Clostridium difficile* infection. *Anaerobe* 15, 285–289 (2009).
7
8 410 <https://doi.org/10.1016/j.anaerobe.2009.09.007>
- 9
10 411 5. Borody T J, Khoruts A (2011). Fecal microbiota transplantation and emerging applications. *Nat Rev Gastroenterol*
11
12 412 *Hepatol* 9(2): 88–96. <https://doi.org/10.1038/nrgastro.2011.244>
- 13
14 413 6. Turnbaugh P J, Bäckhed F, Fulton L, et al (2008). Diet-Induced Obesity Is Linked to Marked but Reversible
15
16 414 Alterations in the Mouse Distal Gut Microbiome. *Cell Host and Microbe* 3(4):213–223.
17
18 415 <https://doi.org/10.1016/j.chom.2008.02.015>
- 19
20 416 7. Oliphant K, Allen-Vercoe E (2019). Macronutrient metabolism by the human gut microbiome: Major fermentation
21
22 417 by-products and their impact on host health. *Microbiome*, 7(1):1–15. <https://doi.org/10.1186/s40168-019-0704-8>
- 23
24 418 8. Wong JMW, Jenkins DJA (2007). Carbohydrate digestibility and metabolic effects. *J Nutr* 137:2539S–46S.
25
26 419 <https://doi.org/10.1093/jn/137.11.2539S>
- 27
28 420 9. den Besten G, van Eunen K, Groen AK, et al (2013). The role of short-chain fatty acids in the interplay between
29
30 421 diet, gut microbiota, and host energy metabolism. *J Lipid Res* 54:2325–40. <https://doi.org/10.1194/jlr.R036012>
- 31
32 422 10. Krajmalnik-Brown R, Ilhan Z-E, Kang D-W, et al (2012). Effects of gut microbes on nutrient absorption and
33
34 423 energy regulation. *Nutr Clin Pract Off Publ Am Soc Parenter Enter Nutr* 27:201–14.
35
36 424 <https://doi.org/10.1177/0884533611436116>
- 37
38 425 11. Wolf PG, Biswas A, Morales SE, et al (2016). H₂ metabolism is widespread and diverse among human colonic
39
40 426 microbes. *Gut Microbes* 7:235–45. <https://doi.org/10.1080/19490976.2016.1182288>
- 41
42 427 12. Portune KJ, Beaumont M, Davila A-M, et al (2016). Gut microbiota role in dietary protein metabolism and health-
43
44 428 related outcomes: the two sides of the coin. *Trends Food Sci Technol* 57:213–32.
45
46 429 <https://doi.org/10.1016/j.tifs.2016.08.011>
- 47
48 430 13. Cândido FG, Valente FX, Grześkowiak ŁM, et al. (2018). Impact of dietary fat on gut microbiota and low-grade
49
50 431 systemic inflammation: mechanisms and clinical implications on obesity. *Int J Food Sci Nutr* 69:125–43.
51
52 432 <https://doi.org/10.1080/09637486.2017.1343286>
- 53
54 433 14. Desbois AP, Smith VJ (2010). Antibacterial free fatty acids: activities, mechanisms of action and biotechnological
55
56 434 potential. *Appl Microbiol Biotechnol* 85:1629–42. <https://doi.org/10.1007/s00253-009-2355-3>
- 57
58 435 15. Huang S, Rutkowsky JM, Snodgrass RG et al (2012). Saturated fatty acids activate TLR-mediated
59
60 436 proinflammatory signaling pathways. *J Lipid Res* 53:2002–13.

- 1
2 437 16. Calder PC (2013). N-3 fatty acids, inflammation and immunity: new mechanisms to explain old actions. Proc Nutr
3 Soc 72:326–36. <https://doi.org/10.1017/S0029665113001031>
4 438
- 5 439 17. Scudellari M (2017). Cleaning up the hygiene hypothesis. Proc Natl Acad Sci U S A 114(7): 1433–1436.
6 440 <https://doi.org/10.1073/pnas.1700688114>
7 441
- 8 442 18. Shi N, Li N, Duan X, Niu H (2017). Interaction between the gut microbiome and mucosal immune system. Mil
9 443 Med Res 4(1): 1–7. <https://doi.org/10.1186/s40779-017-0122-9>
10 444
- 11 445 19. Sanidad KZ, Zeng MY (2020). Neonatal gut microbiome and immunity. Curr Opin Microbiol. 56:30–37.
12 446 <https://doi.org/10.1016/j.mib.2020.05.011>
13 447
- 14 448 20. Neuman H, Koren O (2017). The Pregnancy Microbiome. Nestle Nutr Inst Workshop Ser , 88: 1-9.
15 449 <https://doi.org/10.1159/000455207>
16 450
- 17 451 21. Koren O, Goodrich J K, Cullender T C et al. Cell, 150(3): 470–480. <https://doi.org/10.1016/j.cell.2012.07.008>
18 452
- 19 453 22. Nyangahu D D, Jaspan H B (2019). Influence of maternal microbiota during pregnancy on infant immunity. Clin
20 454 Exp Immunol 198(1):47–56. <https://doi.org/10.1111/cei.13331>
21 455
- 22 456 23. Farr A, Kiss H, Holzer I, et al (2015). Effect of asymptomatic vaginal colonization with *Candida albicans* on
23 457 pregnancy outcome. Acta Obstet Gynecol Scand 94: 989–996. <https://doi.org/10.1111/aogs.12697>
24 458
- 25 459 24. DiGiulio DB, Callahan BJ, McMurdie PJ, et al (2015). Temporal and spatial variation of the human microbiota
26 460 during pregnancy. Proc Natl Acad Sci U S A 112:11060–11065. <https://doi.org/10.1073>
27 461
- 28 462 25. Singh A, Mittal M (2020). Neonatal microbiome—a brief review. Journal of Maternal-Fetal and Neonatal Medicine,
29 463 33(22): 3841–3848. <https://doi.org/10.1080/14767058.2019.1583738>
30 464
- 31 465 26. Kovalovszki L, Villanyi Z, Pataki I, et al (1982). Isolation of aerobic bacteria from the placenta. Acta Paediatr
32 466 Acad Sci Hung 1982; 23: 357– 360
33 467
- 34 468 27. Aagaard K, Ma J, Antony KM, et al (2014) The placenta harbors a unique microbiome. Sci Transl Med 6: 237ra65.
35 469 <https://doi.org/10.1126/scitranslmed.3008599>
36 470
- 37 471 28. Lauder AP, Roche AM, Sherrill-Mix S, et al (2016). Comparison of placenta samples with contamination controls
38 472 does not provide evidence for a distinct placenta microbiota. Microbiome 4: 29. <https://doi.org/10.1186/s40168-016-0172-3>
39 473
- 40 474 29. Dunn A B, Jordan S, Baker B J, et al (2017). The Maternal Infant Microbiome: Considerations for Labor and Birth.
41 475 MCN Am J Matern Child Nurs 42(6), 318–325. <https://doi.org/10.1097/NMC.0000000000000373>
42 476
- 43 477 30. Matamoros S, Gras-Leguen C, Le Vacon F, et al (2012). Trend Microbiol 21(4): 167–173.
44 478 <https://doi.org/10.1016/j.tim.2012.12.001>
45 479

- 1
2 467 31. Jiménez E, Fernández L, Marín ML, et al (2005). Isolation of commensal bacteria from umbilical cord blood of
3
4 468 healthy neonates born by cesarean section. *Curr Microbiol.* 51(4):270-4. [https://doi.org/10.1007/s00284-005-0020-](https://doi.org/10.1007/s00284-005-0020-3)
5
6 469 3
- 7
8 470 32. Abrahamsson TRY, Jenmalm MC (2015). Gut microbiota and allergy: The importance of the pregnancy period.
9
10 471 *Pediatric Research*, 77(1):214–219. <https://doi.org/10.1038/pr.2014.165>
- 11
12 472 33. Reyman M, van Houten M A, van Baarle D et al (2019). Impact of delivery mode-associated gut microbiota
13 473 dynamics on health in the first year of life. *Nature Communications*, 10(1):1–12. [https://doi.org/10.1038/s41467-](https://doi.org/10.1038/s41467-019-13014-7)
14
15 474 019-13014-7
- 16
17 475 34. Cunnington A J, Sim K, Deierl A, et al (2016). “Vaginal seeding” of infants born by caesarean section: How
18
19 476 should health professionals engage with this increasingly popular but unproved practice? *BMJ (Online)*, 352:1–2.
20
21 477 <https://doi.org/10.1136/bmj.i227>
- 22
23 478 35. Allen J, Hector D (2005). Benefits of breastfeeding. *N S W Public Health Bull*
24
25 479 16(3–4): 42–46. <https://doi.org/10.1071/nb05011>
- 26
27 480 36. Ballard O, Morrow A L (2013). Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am*
28
29 481 60(1):49–74. <https://doi.org/10.1016/j.pcl.2012.10.002>
- 30
31 482 37. Castellote C, Casillas R, Ramirez-Santana C, et al. (2011) Premature delivery influences the immunological
32 483 composition of colostrum and transitional and mature human milk. *J Nutr* 141(6):1181–7.
33
34 484 <https://doi.org/10.3945/jn.110.133652>
- 35
36 485
- 37
38 486 38. Pang WW, Hartmann PE (2007). Initiation of human lactation: secretory differentiation and secretory activation. *J*
39
40 487 *Mammary Gland Biol Neoplasia* 12(4):211–21. <https://doi.org/10.1007/s10911-007-9054-4>
- 41
42 488 39. Kulski JK, Hartmann PE (1981). Changes in human milk composition during the initiation of lactation. *Aust J Exp*
43
44 489 *Biol Med Sci* 59(1):101–14. <https://doi.org/10.1038/icb.1981.6>
- 45
46 490 40. Prentice A. Regional variations in the composition of human milk. In: Jensen RG, editor. *Handbook of milk*
47
48 491 *composition*. San Diego (CA): Academic Press, Inc; 1995. p. 919.
- 49
50 492 41. Henderickx JGE, Zwiittink RD, Van Lingen RA, et al (2019). The preterm gut microbiota: An inconspicuous
51
52 493 challenge in nutritional neonatal care. *Front. Cell. Infect. Microbiol* 9:1–12.
53
54 494 <https://doi.org/10.3389/fcimb.2019.00085>
- 55
56 495 42. Mosca F, Gianni ML (2017). Human milk: composition and health benefits. *Pediatr Med Chir* 39(2): 155.
57
58 496 <https://doi.org/10.4081/pmc.2017.155>
- 59
60

- 1
2 497 43. Eriksen KG, Christensen SH, Lind MV, et al (2018). Human milk composition and infant growth. *Curr Opin Clin*
3
4 498 *Nutr Metab Care* 21(3): 200–206. <https://doi.org/10.1097/MCO.0000000000000466>
- 5
6 499 44. Timby N, Domellöf M, Lönnerdal B, et al (2017). Supplementation of infant formula with bovine milk fat globule
7
8 500 membranes. *Adv Nutr* 8:351–355. <https://doi.org/10.3945/an.116.014142>
- 9
10 501 45. Valentine CJ, Morrow G, Pennell M, et al (2012). Randomized controlled trial of docosahexaenoic acid
11
12 502 supplementation in midwestern U.S. Human milk donors. *Breastfeed Med*
13 503 <http://dx.doi.org/10.1089/bfm.2011.0126>.
14
- 15 504 46. Martin MA, Lassek WD, Gaulin SJ, et al (2012). Fatty acid composition in the mature milk of Bolivian forager-
16
17 505 horticulturalists: controlled comparisons with a US sample. *Matern Child Nutr* 8(3):404–18.
18
19 506 <https://doi.org/10.1111/j.1740-8709.2012.00412.x>
- 20
21 507 47. Much D, Brunner S, Vollhardt C, et al (2013). Breast milk fatty acid profile in relation to infant growth and body
22
23 508 composition: results from the INFAT study. *Pediatr Res* 74:230–237
- 24
25 509 48. Andreas NJ, Kampmann B, Mehring Le-Doare K, et al (2015). Human breast milk: a review on its composition and
26
27 510 bioactivity. *Early Hum Dev* 91:629–35. <https://doi.org/10.1016/j.earlhumdev.2015.08.013>
- 28
29 511 49. Nommsen LA, Lovelady CA, Heinig MJ, et al (1991). Determinants of energy, protein, lipid, and lactose
30
31 512 concentrations in human milk during the first 12 mo of lactation: the DARLING Study. *Am J Clin Nutr*
32 513 53(2):457–65. <https://doi.org/10.1093/ajcn/53.2.457>
- 34 514 50. Greer FR (2001). Do breastfed infants need supplemental vitamins? *Pediatr Clin North Am* 48(2):415–23.
35
36 515
- 38 516 51. Allen LH (2012). B vitamins in breast milk: relative importance of maternal status and intake, and effects on infant
39
40 517 status and function. *Adv Nutr* 3(3):362–369. <https://doi.org/10.3945/an.111.001172>
- 42 518 52. La Tuga M S, Stuebe A, Seed PC (2014). A review of the source and function of microbiota in breast milk. *Semin*
43
44 519 *Reprod Med*. 32(1):68–73. <https://doi.org/10.1055/s-0033-1361824>
- 45
46 520 53. Gritz EC, Bhandari V (2015). The human neonatal gut microbiome: a brief review. *Front Pediatr*
47
48 521 3:17. <https://doi.org/10.3389/fped.2015.00017>
- 49
50 522 54. Shin NR, Whon TW, Bae JW (2015). Proteobacteria: Microbial signature of dysbiosis in gut microbiota. *Trends*
51
52 523 *Biotechnol* 33(9):496–503. <https://doi.org/10.1016/j.tibtech.2015.06.011>
- 53
54 524 55. Underwood MA, Mukhopadhyay S, Lakshminrusimha, S et al (2020). Neonatal intestinal dysbiosis. *J Perinatol*
55
56 525 40(11):1597–1608. <https://doi.org/10.1038/s41372-020-00829-2>
- 57
58
59
60

- 1
2 526 56. Bentley R, Meganathan R (1982). Biosynthesis of vitamin K (menaquinone) in bacteria. *Microbiol Rev* 46(3):241–
3 280. <https://doi.org/10.1128/mr.46.3.241-280.1982>
4 527
5
6 528 57. Percival SL, Williams DW (2013) *Escherichia coli*. In: *Microbiology of Waterborne Diseases: Microbiological*
7 *Aspects and Risks: Second Edition* Elsevier. doi: 10.1016/B978-0-12-415846-7.00006-8
8 529
9
10 530 58. Mainil J (2013). *Escherichia coli* virulence factors. *Vet Immunol Immunopathol* 152(1–2):2–12.
11 <https://doi.org/10.1016/j.vetimm.2012.09.032>
12 531
13 532 59. Hanson LA (2004). Protective effects of breastfeeding against urinary tract infection. *Acta Pædiatr* 93: 154–156.
14 Stockholm. ISSN 0803-5253 <https://doi.org/10.1080/08035250310022586>
15 533
16
17 534 60. Slavikova M, Lodinova-Zadnikova R, Adlerberth I, et al (1995). Increased mannose-specific adherence and
18 colonizing ability of *Escherichia coli* 083 in breastfed infants. *Adv Exp Med Biol* 371(A): 421–423.
19 535 https://doi.org/10.1007/978-1-4615-1941-6_88
20 536
21 537 61. Connell H, Agace W, Klemm P, et al (1996). Type 1 fimbrial expression enhances *Escherichia coli* virulence for
22 the urinary tract. *PNAS*:93(18), 9827–9832. <https://doi.org/10.1073/pnas.93.18.9827>
23 538
24
25 539 62. Walter J (2008). Ecological role of *Lactobacilli* in the gastrointestinal tract: Implications for fundamental and
26 biomedical research. *Appl Environ Microbiol* 74(16):4985–4996. <https://doi.org/10.1128/AEM.00753-08>
27 540
28
29 541 63. Sung V, Hiscock H, Tang MLK, et al (2014). Treating infant colic with the probiotic *Lactobacillus reuteri*: Double
30 blind, placebo controlled randomised trial. *BMJ (Online)*, 348:1–11. <https://doi.org/10.1136/bmj.g2107>
31 542
32
33 543 64. Guarino A, Guandalini S, Lo Vecchio A (2015) Probiotics for prevention and treatment of diarrhea. *J Clin*
34 *Gastroenterol* 49, Suppl 1:S37–S45. <https://doi.org/10.1097/MCG.0000000000000349>
35 544
36
37 545 65. Higgins SE, Higgins JP, Wolfenden AD, et al (2008). Evaluation of a *Lactobacillus*-based probiotic culture for the
38 reduction of *Salmonella enteritidis* in neonatal broiler chicks. *Poultry Science*, 87(1):27–31.
39 <https://doi.org/10.3382/ps.2007-00210>
40 546
41
42 547 66. Lee DJ, Drongowski RA, Coran AG, et al (2000). Evaluation of probiotic treatment in a neonatal animal model.
43 *Pediatr Surg Int* 16(4):237–242. <https://doi.org/10.1007/s003830050736>
44 548
45
46 549 67. Lees E A, Miyajima F, Pirmohamed M, & Carrol E D (2016). The role of *Clostridium difficile* in the paediatric and
47 neonatal gut — a narrative review. *European Journal of Clinical Microbiology and Infectious Diseases*, 35(7),
48 550 1047–1057. <https://doi.org/10.1007/s10096-016-2639-3>
49 551
50 552 68. Elward A, Brady MT, Bryant K, et al (2020). *Clostridioides difficile* in Neonatal Intensive Care Unit Patients: A
51 Systematic Review. *Centers for Disease Control and Prevention National Center for Zoonotic and Emerging*
52 *Infectious Diseases*.
53
54 553
55
56 554
57
58 555
59
60

- 1
2 556 Infectious Diseases Division of Healthcare Quality Promotion. P. 1–10. <https://www.cdc.gov/hicpac/reviews/cdiff->
3
4 557 [nicu/index.html](https://www.cdc.gov/hicpac/reviews/cdiff-nicu/index.html)
- 5
6 558 69. Bridgman SL, Konya T, Azad MB, et al (2016) High fecal IgA is associated with reduced *C. difficile* colonization
7
8 559 in infants, *Microbes Infect* 18(9):543-549. <https://doi.org/10.1016/j.micinf.2016.05.001>.
- 9
10 560 70. Schutze GE, Willoughby RE (2013). *Clostridium difficile* infection in infants and children. *Pediatrics* 131(1): 196–
11
12 561 200. <https://doi.org/10.1542/peds.2012-2992>
- 13 562 71. O’Callaghan A, van Sinderen D (2016). *Bifidobacteria* and their role as members of the human gut microbiota.
14
15 563 *Frontiers Microbiol*, 7(JUN). <https://doi.org/10.3389/fmicb.2016.00925>
- 16
17 564 72. Arumugam M, Raes J, Pelletier E, et al (2011). Enterotypes of the human gut microbiome. *Nature* 473(7346): 174–
18
19 565 180. <https://doi.org/10.1038/nature09944>
- 20
21 566 73. Knights D, Ward TL, Mckinlay CE, et al (2017). Rethinking “Enterotypes”. *Cell Host Microbe* 16(4):433–437.
22
23 567 <https://doi.org/10.1016/j.chom.2014.09.013>
- 24
25 568 74. Wu GD, Chen J, Hoffmann C, et al (2011). Linking long-term dietary patterns with gut microbial enterotypes.
26
27 569 *Science (New York, N.Y.)*: 334(6052), 105–108. <https://doi.org/10.1126/science.1208344>
- 28
29 570 75. Xiao L, Wang J, Zheng J, et al (2021). Deterministic transition of enterotypes shapes the infant gut microbiome at
30
31 571 an early age. *Genome Biol* 22(1):243. <https://doi.org/10.1186/s13059-021-02463-3>
- 32
33 572 76. Bergström A, Skov TH, Bahl MI, et al (2014). Establishment of intestinal microbiota during early life: A
34
35 573 longitudinal, explorative study of a large cohort of Danish infants. *Appl Environ Microbiol* 80(9): 2889–2900.
36
37 574 <https://doi.org/10.1128/AEM.00342-14>
- 38 575
39
40 576 77. Tamboli CP, Neut C, Desreumaux P, et al (2004). Dysbiosis in inflammatory bowel disease. *Gut*, 53(1):1–4.
41
42 577 <https://doi.org/10.1136/gut.53.1.1>
- 43
44 578 78. Li M, Monaco MH, Wang M et al (2014). Human milk oligosaccharides shorten rotavirus-induced diarrhea and
45
46 579 modulate piglet mucosal immunity and colonic microbiota. *ISME J* 8: 1609–1620.
47
48 580 <https://doi.org/10.1038/ismej.2014.10>
- 49
50 581 79. Garcia-Mantrana I, Collado MC (2016). Obesity and overweight: Impact on maternal and milk microbiome and
51
52 582 their role for infant health and nutrition. *Mol Nutr Food Res* 60(8):1865-1875
53
54 583 <https://doi.org/10.1002/mnfr.201501018>
- 55
56 584 80. Dardas M, Gill SR, Grier A, et al (2014). The impact of postnatal antibiotics on the preterm intestinal microbiome.
57
58 585 *Pediatr Res* 76:150–158. doi: 10.1038/pr.2014.69

- 1
2 586 81. Zwiittink RD, Renes IB, van Lingen RA, et al (2018). Association between duration of intravenous antibiotic
3 administration and early-life microbiota development in late-preterm infants. *Eur. J. Clin. Microbiol. Infect. Dis.*
4 587 37:475–483. [https://doi: 10.1007/s10096-018-3193-y](https://doi.org/10.1007/s10096-018-3193-y)
5 588
6 589 82. Shaw AG, Sim K, Randell P, et al. (2015). Late-onset bloodstream infection and perturbed maturation of the
7 gastrointestinal microbiota in premature infants. *PLoS ONE*. 10:e0132923. [https://doi:](https://doi.org/10.1371/journal.pone.0132923)
8 590 10.1371/journal.pone.0132923
9 591
10 592 83. Duffy LC (2000). Interactions mediating bacterial translocation in the immature intestine. *J. Nutr* 130:432S–436S.
11 593 [https://doi: 10.1093/jn/130.2.432S](https://doi.org/10.1093/jn/130.2.432S)
12 594
13 595 84. Brooks B, Mueller RS, Young JC et al (2015). Strain-resolved microbial community proteomics reveals
14 simultaneous aerobic and anaerobic function during gastrointestinal tract colonization of a preterm infant. *Front*
15 596 *Microbiol* 6:654. [https://doi: 10.3389/fmicb.2015.00654](https://doi.org/10.3389/fmicb.2015.00654)
16 597
17 598 85. Deshpande GC, Rao SC, Keil AD, et al (2011). Evidence-based guidelines for use of probiotics in preterm
18 neonates. *BMC Medicine*, 9(92): 1–13. <https://doi.org/10.1186/1741-7015-9-92>
19 599
20 600 86. Bin-Nun A, Bromiker R, Wilschanski M, et al (2005). Oral probiotics prevent necrotizing enterocolitis in very low
21 birth weight neonates. *J Pediatr* 147(2):192–196. <https://doi.org/10.1016/j.jpeds.2005.03.054>
22 601
23 602 87. Zhang Z, Xiang Y, Li N, et al (2013). Protective effects of *Lactobacillus rhamnosus* GG against human rotavirus-
24 induced diarrhoea in a neonatal mouse model. *Pathog Dis*. 67(3):184-191. doi: 10.1111/2049-632X.12030.
25 603
26 604 88. Liu F, Li G, Wen K, et al (2013). *Lactobacillus rhamnosus* GG on rotavirus-induced injury of ileal epithelium in
27 gnotobiotic pigs. *J Pediatr Gastroenterol Nutr* 57(6):750-8. [https://doi: 10.1097/MPG.0b013e3182a356e1](https://doi.org/10.1097/MPG.0b013e3182a356e1).
28 605
29 606 89. Won TJ, Kim B, Song DS, et al (2011), Modulation of Th1/Th2 balance by *Lactobacillus* strains isolated from
30 Kimchi via stimulation of macrophage cell line J774A.1 In Vitro. *J Food Sci* 76:H55-H61.
31 607 <https://doi.org/10.1111/j.1750-3841.2010.02031.x>
32 608
33 609 90. Toori A, Toori S, Fujiwara S, et al (2007). *Lactobacillus acidophilus* strain L-92 regulates the production of Th1
34 cytokine as well as Th2 cytokines. *Allergol Int* 56(3):293-301. <https://doi.org/10.2332/allergolint.O-06-459>
35 610
36 611 91. Isolauri E, Arvola T, Sutas Y, et al. (2000). Probiotics in the management of atopic eczema. *Clin Exp Allergy*
37 612 30:1604–1610. <https://doi.org/10.1046/j.1365-2222.2000.00943.x>
38 613
39 614 92. Parra-Llorca A, Gormaz M, Alcántara C, et al (2018). Preterm gut microbiome depending on feeding type:
40 Significance of donor human milk. *Fron Microbiol* 9: 1376. <https://doi.org/10.3389/fmicb.2018.01376>
41 615
42 616 93. Newburg DS, Ruiz-Palacios GM, Morrow AL. Human milk glycans protect infants against enteric pathogens.
43 *Annu Rev Nutr* 2005;25:37–58. <https://doi.org/10.1146/annurev.nutr.25.050304.092553>
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2 616 94. Morrow AL, Ruiz-Palacios GM, Jiang X, et al (2005). Human-milk glycans that inhibit pathogen binding protect
3 breast-feeding infants against infectious diarrhea. *J Nutr* 135(5):1304–7. <https://doi.org/10.1093/jn/135.5.1304>
4 617
5 618 95. Gabrielli O, Zampini L, Galeazzi T, et al (2011). Preterm milk oligosaccharides during the first month of lactation.
6 619 *Pediatrics* 128(6):e1520–31. <https://doi.org/10.1542/peds.2011-1206>
7
8 620 96. Jantscher-Krenn E, Zharebtsov M, Nissan C, et al (2012). The human milk oligosaccharide disialyllacto-N-tetraose
9 621 prevents necrotising enterocolitis in neonatal rats. *Gut* 61(10):1417–25. <https://doi.org/10.1136/gutjnl-2011-301404>
10
11 622 97. Li M, Monaco M H, Wang M, et al (2014). Human milk oligosaccharides shorten rotavirus-induced diarrhea and
12 623 modulate piglet mucosal immunity and colonic microbiota. *ISME Journal*, 8(8): 1609–1620.
13 624 <https://doi.org/10.1038/ismej.2014.10>
14
15 625 98. Arnold JW, Roach J, Azcarate-Peril MA (2016) Emerging Technologies for Gut Microbiome Research. *Trends*
16 626 *Microbiol* 24(11):887–901. <https://doi.org/10.1016/j.tim.2016.06.008>
17 627
18 628 99. National Academies of Sciences, Engineering, and Medicine, et al (2017) Environmental Chemicals, the Human
19 629 Microbiome, and Health Risk: A Research Strategy. National Academies Press (US) Bookshelf ID: NBK481560
20 630 doi:10.17226/24960
21 631
22 632 100. Sarangi AN, Goel A, Aggarwal R (2019). Methods for studying gut microbiota: a primer for physicians. *J Clin*
23 633 *Exp Hepatol* 9(1):62–73. <https://doi.org/10.1016/j.jceh.2018.04.016>
24 634
25 635 101. Fritz JV, Desai MS, Shah P, et al (2013) From meta-omics to causality: Experimental models for human
26 636 microbiome research. *Microbiome* 1(1):14. <https://doi.org/10.1186/2049-2618-1-14>
27 637
28 638 102. Van de Wiele T, Van den Abbeele P, Ossieur W, et al (2015). The simulator of the human intestinal microbial
29 639 ecosystem (SHIME®). In: Verhoeckx K, Cotter P, López-Expósito I, Kleiveland C, Lea T, Mackie A, Requena T,
30 640 Swiatecka D, Wichers H, editors. *The Impact of Food Bioactives on Health: in vitro and ex vivo models* [Internet].
31 641 Cham (CH): Springer; 2015. Chapter 27. PMID: 29787058.
32 642
33 643 103. Faith JJ, Rey FE, O'Donnell D, et al (2010). Creating and characterizing communities of human gut microbes in
34 644 gnotobiotic mice. *ISME J* 4:1094–1098. <https://doi.org/10.1038/ismej.2010.110>
35 645
36 646 104. Faith JJ, Ahern PP, Ridaura VK, et al (2014) Identifying gut microbe-host phenotype relationships using
37 647 combinatorial communities in gnotobiotic mice. *Sci Transl Med* 6, 220ra11.
38 648 <https://doi.org/10.1126/scitranslmed.3008051>
39 649
40 650 105. Blaut M, Collins MD, Welling GW et al (2002) Molecular biological methods for studying the gut microbiota: the
41 651 EU human gut flora project. *Br. J. Nutr.* 87 (Suppl. 2), S203–S211. <https://doi.org/10.1079/BJNBJN/2002539>
42 652
43 653
44 654
45 655
46 656
47 657
48 658
49 659
50 660

- 1
2 646 106. Krawczyk B, Kur J, Stojowska-Swędryńska K et al (2016). Principles and applications of Ligation Mediated PCR
3
4 647 methods for DNA-based typing of microbial organisms. *Acta Biochim Pol.* 63(1):39-52.
5
6 648 https://doi.org/10.18388/abp.2015_1192
7
8 649 107. Azcarate-Peril MA, Foster DM, Cadenas MB, et al (2011). Acute necrotizing enterocolitis of preterm piglets is
9
10 650 characterized by dysbiosis of ileal mucosa-associated bacteria. *Gut Microbes* 2:234–243.
11
12 651 <https://doi.org/10.4161/gmic.2.4.16332>
13
14 652 108. Donskey CJ, Hujer AM, Das SM, et al (2003) Use of denaturing gradient gel electrophoresis for analysis of the
15
16 653 stool microbiota of hospitalized patients. *J. Microbiol. Methods* 54: 249–256. [https://doi.org/10.1016/s0167-](https://doi.org/10.1016/s0167-7012(03)00059-9)
17
18 654 [7012\(03\)00059-9](https://doi.org/10.1016/s0167-7012(03)00059-9)
19
20 655 109. Stewart CJ, Nelson A, Scribbins D, et al (2013) Bacterial and fungal viability in the preterm gut: NEC and sepsis.
21
22 656 *Arch Dis Child Fetal Neonatal Ed.* 98, F298–F303. <https://doi.org/10.1136/archdischild-2012-302119>
23
24 657 110. Lepage P, Leclerc MC, Joossens M, et al (2013). A metagenomic insight into our gut's microbiome. *Gut*,
25
26 658 62(1):146–158. <https://doi.org/10.1136/gutjnl-2011-301805>
27
28 659 111. Janda JM, Abbott SL (2007). 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory:
29
30 660 pluses, perils, and pitfalls. *J Clin Microbiol.* 45(9):2761-2764. <https://doi.org/10.1128/JCM.01228-07>
31
32 661 112. Houpiqian P, Raoult D (2002). Traditional and molecular techniques for the study of emerging bacterial diseases:
33
34 662 one laboratory's perspective. *Emerg Infect Dis* 8(2): 122–131. <https://doi.org/10.3201/eid0802.010141>
35
36 663 113. Palmero D, Rodríguez JM, de Cara M et al. (2011) Fungal microbiota from rain water and pathogenicity of
37
38 664 *Fusarium* species isolated from atmospheric dust and rainfall dust. *J Ind Microbiol Biotechnol* 38:13–20.
39
40 665 <https://doi.org/10.1007/s10295-010-0831-5>
41
42 666 114. Kolmeder CA, de Vos WM (2014). Metaproteomics of our microbiome - developing insight in function and
43
44 667 activity in man and model systems. *J Proteomics*, 97:3–16. <https://doi.org/10.1016/j.jprot.2013.05.018>
45
46 668 115. Larsen PE, Dai Y (2015). Metabolome of human gut microbiome is predictive of host dysbiosis. *GigaScience*,
47
48 669 4(1):1–16. <https://doi.org/10.1186/s13742-015-0084-3>
49
50 670 116. Morgan XC, Huttenhower C (2014) Meta'omic analytic techniques for studying the intestinal microbiome.
51
52 671 *Gastroenterology* 146:1437-1448.e1. <https://doi.org/10.1053/j.gastro.2014.01.049>
53
54 672 117. Franzosa EA, Morgan XC, Segata N, et al (2014). Relating the metatranscriptome and metagenome of the human
55
56 673 gut. *Proc Natl Acad Sci U S A* 111:E2329–E2338. <https://doi.org/10.1073/pnas.1319284111>
57
58
59
60

- 1
2 674 118. El Aidy S, Kleerebezem M (2013). Molecular signatures for the dynamic process of establishing intestinal host-
3
4 675 microbial homeostasis: potential for disease diagnostics? *Curr Opin Gastroenterol.* 29:621–627.
5
6 676 <https://doi.org/10.1097/MOG.0b013e328365d365>
7
8 677 119. Larsen PE, Dai Y (2015). Metabolome of human gut microbiome is predictive of host dysbiosis. *GigaScience,*
9
10 678 4(1):1–16. <https://doi.org/10.1186/s13742-015-0084-3>
11
12 679
13 680
14
15 681
16
17 682
18
19 683
20
21 684
22
23 685
24
25 686
26
27 687
28
29 688
30
31 689
32
33 690
34
35 691
36
37 692
38
39 693
40
41 694
42
43 695
44
45 696
46
47 697
48
49 698
50
51 699
52
53 700
54
55 701
56
57 702
58
59 703
60

Table 1. Comparison of bacteria commonly found in breast milk [52] and the infant's intestine [53].

Phylum	Genera	Breast milk	Neonatal intestine
Firmicutes	<i>Staphylococcus</i>	+	+
	<i>Streptococcus</i>	+	+
	<i>Veillonella</i>	+	+
	<i>Enterococcus</i>	+	+
	<i>Gemella</i>	+	-
	<i>Clostridium</i>	+	+
	<i>Lactobacillus</i>	+	+
	<i>Eubacterium</i>	-	+
	<i>Ruminococcus</i>	-	+
	<i>Peptostreptococcus</i>	-	+
Actinobacteria	<i>Propionibacterium</i>	+	+
	<i>Actinomyces</i>	+	-
	<i>Corynebacterium</i>	+	+
	<i>Bifidobacterium</i>	+	+
	<i>Streptomyces</i>	-	+
Proteobacteria	<i>Pseudomonas</i>	+	-
	<i>Sphingomonas</i>	+	-
	<i>Serratia</i>	+	-
	<i>Escherichia</i>	+	+
	<i>Enterobacter</i>	+	+
	<i>Ralstonia</i>	+	-
	<i>Bradyrhizobium</i>	+	-
	<i>Klebsiella</i>	-	+
	<i>Acinetobacter</i>	-	+
	<i>Desulfovibrio</i>	-	+
Bacteroidetes	<i>Prevotella</i>	+	+
	<i>Bacteroides</i>	-	+

1

2 706 **Fig 1. Methods used for the study of microbiome.** Legend: T-RFLP - Terminal restriction fragment length

3

4 707 polymorphism; WGS - Whole Genome Sequencing

5

6 708 **Fig 2. Source of the infant microbiome.** The figure shows the influence of the mother's microbiota and the environmental

7

8 709 microbiota on the bacterial colonization of newborns and infants.

9

10

11 710

12

13 711

14

15 712

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

For Peer Review




Fig.1 Methods used for the study of microbiome. Legend: T-RFLP - Terminal restriction fragment length polymorphism; WGS - Whole Genome Sequencing

173x156mm (300 x 300 DPI)




Fig 2. Source of the infant microbiome. The figure shows the influence of the mother's microbiota and the environmental microbiota on the bacterial colonization of newborns and infants.

173x97mm (300 x 300 DPI)