

DEDICATED TO PROFESSOR IWO BIAŁYNICKI-BIRULA ON HIS 90TH BIRTHDAY



Contra Bellum: Bell’s Theorem as a Confusion of Languages

M. CZACHOR*

Institut Fizyki i Informatyki Stosowanej, Politechnika Gdańska, 80-233 Gdańsk, Poland

Published: 14.06.2023

Doi: [10.12693/APhysPolA.143.S158](https://doi.org/10.12693/APhysPolA.143.S158)

*e-mail: marek.czachor@pg.edu.pl

Bell’s theorem is a conflict of mathematical predictions formulated within an infinite hierarchy of mathematical models. Inequalities formulated at level $k \in \mathbb{Z}$ are violated by probabilities at level $k + 1$. We are inclined to think that $k = 0$ corresponds to the classical world, while $k = 1$ — to the quantum one. However, as the $k = 0$ inequalities are violated by $k = 1$ probabilities, the same relation holds between $k = 1$ inequalities violated by $k = 2$ probabilities, $k = -1$ inequalities violated by $k = 0$ probabilities, and so forth. By accepting the logic of the Bell theorem, can we prove by induction that nothing exists?

topics: Bell’s theorem, black holes, non-Newtonian calculus, quantum cryptography

1. Introduction

Is Bell’s theorem a mathematical theorem? If we treat Bell’s theorem [1] as a theorem about the additivity of Lebesgue measures, then yes — this is a mathematical theorem. However, Bell’s theorem is more ambitious. It tells us about reality per se, the security of communication channels, the structure of space and time, and even the freedom of experimental physicists.

Although mathematical theorems cannot have counterexamples, this is not necessarily true for theorems about physical reality. The whole history of science is a series of exceptions to various well-established truths.

One such famous truth about reality was known as Euclid’s fifth axiom, which essentially states that angles in any triangle add up to 180° . It was so self-evident to 19th-century mathematicians that even Gauss himself was not eager to publish his thoughts on the subject.

Bell’s theorem is technically based on another apparently self-evident truth about additivity, namely

$$\int_{\Lambda} d\lambda (f \pm g)(\lambda) = \int_{\Lambda} d\lambda f(\lambda) \pm \int_{\Lambda} d\lambda g(\lambda). \quad (1)$$

In proofs of Bell-type inequalities, one often replaces (1) with a more elementary rule,

$$\frac{n \pm m}{N} = \frac{n}{N} \pm \frac{m}{N}. \quad (2)$$

Thus, (1) occurs in contexts of probability measures, while (2) is typical of frequentist approaches.

However, neither (1) nor (2) are universally true; (1) fails for fuzzy or fractal functions; (2) fails if n, m , represent velocities and N is the velocity of light. In the latter case, what we get is rather

$$n \oplus m = N \tanh \left(\tanh^{-1} \left(\frac{n}{N} \right) + \tanh^{-1} \left(\frac{m}{N} \right) \right) = f^{-1} \left(f(n) + f(m) \right). \quad (3)$$

Of course, nothing can prevent us from adding velocities by means of (2), but this is not what Nature does. The arithmetic of Nature is (3). A relation between \oplus and $+$ is here analogous to the one between a curvature of a general manifold and the flatness of its local chart of coordinates (charts in atlases are flat). Arithmetic in special relativity becomes as physical as geometry in general relativity.

A similar situation occurs with (1). In fuzzy and fractal applications, one often encounters [2–12]

$$\int_{\Lambda} d\lambda f(\lambda) \oplus \int_{\Lambda} d\lambda g(\lambda) = \int_{\Lambda} d\lambda (f \oplus g)(\lambda). \quad (4)$$

The exact meaning of \oplus depends on the way fuzzy sets are constructed [13] or which fractals one is dealing with [14]. A particular example of (4) occurs in Maslov's idempotent analysis [15, 16]. Here, certain optimization problems that are nonlinear in the usual framework become linear with respect to generalized arithmetic operations [17], even though the generalized arithmetic is not isomorphic to the one of \mathbb{R} [18].

The goal of the paper is to show that quantum probabilities typical of a two-particle singlet state (that is, those used by Bell in his argumentation), despite all the wisdom of theoretical physicists, *can* result from a local theory where Einstein–Podolsky–Rosen-type (EPR) elements of reality exist [19–21], with probabilities given by local realistic Clauser–Horne formulas [22], where observers have free will, and their detectors are 100% efficient. The only difference is in the form of the integral, whose linearity is with respect to \oplus , \ominus , \odot , and \oslash appropriately defined.

Violation of Bell-type inequalities is then no more paradoxical than $c+c=2c$, which could be claimed to violate the speed of light limit. Moreover, there is no problem with circumventing the Tsirelson bounds typical of Hilbert-space models of probability [23], still maintaining Bell locality, EPR elements of reality, the free will of observers, and 100% efficient detectors.

I can reassure the readers that the models we are analyzing are *not* an alternative to quantum mechanics. They do not explain why probability amplitudes interfere, but, nevertheless, shift the discussion of linearity to new, unexplored areas. The status of theorems based on algebraic properties of observables, such as the Greenberger–Horne–Zeilinger theorem [24, 25] or its single-particle analogs [26], is still open.

But what the models *do* show is that quantum correlations of an EPR type do not necessarily exclude EPR elements of reality — a conclusion with potentially dramatic implications for quantum cryptography. How serious the consequences are, remains to be investigated. Einstein's views on incompleteness of quantum mechanics receive unexpected support.

We will begin our discussion with the observation that principles of relativity are more general and ubiquitous than Einstein's relativity of uniform motion or Copernican relativity of point of observation. The most fundamental principle occurring in all natural sciences is the relativity of arithmetic [27]. It implies, in particular, principles of the relativity of calculus and the relativity of probability. Both are essential for Bell's theorem.

2. Relativity of probability

Relativity of probability occurs at several levels. The most obvious one is illustrated by the following

example. It can be regarded as a particular case of Einstein's special relativity.

Assume a source emits particles to the right or to the left, with certain probability density $\rho(v)$ of velocities. If N particles have been emitted, let N_+ denote the number of particles propagating to the right. An observer measures N_+/N and compares it with the theoretical prediction, $p_+ = \int_0^\infty dv \rho(v)$.

An observer that moves with velocity V with respect to the previous one will measure a different value of N_+/N , even though both of them analyze the same experiment with the same N .

The example is trivial, but it illustrates an important fact about probability — different observers may associate different probabilities with the same experimental situation and with the same definition of elementary events. In this concrete example, the relativity of probability, $p_+(V) = \int_0^\infty dv \rho(v+V)$, results from the relativity of motion.

As a less trivial relativistic example, consider the gravitational collapse of a star. There are two observers: Alice, who falls with the star, and Bob, who remains at rest at position r . Alice employs a broken clock that randomly fails to work (which happens with probability p_0). The motion of the clock's hand becomes a Poisson process characterized by probability $p_1 = 1 - p_0$ of a forward move.

A single run of experiment lasts a fixed amount τ of the observer's proper time. Alice measures $N = \lfloor \tau/\Delta\tau \rfloor$ bits A_1, \dots, A_N (N_1 events $A_j=1$ when the clock's hand moves; $N_0 = N - N_1$ events $A_j=0$ when it gets stuck). The average amount of proper time measured by the Alice's damaged clock is $p_1\tau$.

The experimental ratio N_1/N observed by Alice gets translated into \tilde{N}_1/\tilde{N} observed by Bob. In general, $\tilde{N}_1 \neq N_1$ and $\tilde{N} \neq N$ because the numbers of observed events differ for Alice and Bob due to the relativity of time and the presence of the horizon. The events observed by Alice after she crosses the Schwarzschild radius at her proper time τ_S will be unavailable to Bob, even though his detectors are 100% efficient.

Bob should cautiously draw conclusions about N_1 and N on the basis of \tilde{N}_1 and \tilde{N} he observes. For example, if he concludes that τ_S is greater than τ because, from his perspective, Alice cannot reach the Schwarzschild radius, this inequality can be "violated" in the world of Alice.

Bob can derive various inequalities about the data of Alice, provided he knows the map g_r that relates her N_1/N with his

$$\frac{\tilde{N}_1}{\tilde{N}} = g_r\left(\frac{N_1}{N}\right). \quad (5)$$

The exact form of g_r is irrelevant to our argument, but it could be derived on the basis of general relativity if needed.

From our perspective, it is important that g_r connects two real probabilistic processes. Both N_1/N and $g_r(N_1/N)$ are true, physically significant

probabilities. The “violation” of Bob’s world $\tau_S > \tau$ by Alice’s world $\tau_S < \tau$ is paradoxical only for those who do not understand Einstein’s theory of gravity.

3. A lemma on relativity of binary probabilities

For binary events, there exists a simple result guaranteeing that both N_1/N and $g(N_1/N)$ are probabilities.

Lemma 1: $g(p) + g(1 - p) = 1$ for any $p \in [0, 1]$, if and only if

$$g(p) = \frac{1}{2} + h\left(p - \frac{1}{2}\right), \quad (6)$$

where $h(-x) = -h(x)$. Any such g has a fixed point at $p = \frac{1}{2}$.

So, any antisymmetric $h(x)$ leads to an acceptable $g(p)$. The proof can be found in [28]. For a discussion of non-binary probabilities, see [29].

As an example, consider the antisymmetric function

$$h(x) = \frac{1}{2} \sin(\pi x). \quad (7)$$

Then $g(p) = \sin^2(\frac{\pi}{2}p)$, and indeed

$$g(p) + g(1-p) = \sin^2\left(\frac{\pi}{2}p\right) + \cos^2\left(\frac{\pi}{2}p\right) = 1 \quad (8)$$

for any p . Now let $p = (\pi - \theta)/\pi$ be the probability of finding a point belonging to the overlap of two half-circles rotated by θ . Then

$$g(p) = \sin^2\left(\frac{\pi}{2} \frac{\pi - \theta}{\pi}\right) = \cos^2\left(\frac{\theta}{2}\right) \quad (9)$$

is the Malus law for spin 1/2 (or Mach–Zehnder interferometers).

Note that g is one-to-one on $[0, 1]$. Moreover, $g(0) = 0$ and $g(1) = 1$ — a property with important implications for the definition of bits: classical, quantum, and intermediate.

The readers should think of p and $\tilde{p} = g(p)$ in categories similar to those that have led us to (5). Both p and \tilde{p} can be physically meaningful. We should be as cautious as Bob in formulating statements about the level of p on the basis of the rules that apply to the level of \tilde{p} .

4. Arithmetic elements of reality

Consider some set \mathbb{X} and a bijection $f_{\mathbb{X}} : \mathbb{X} \rightarrow \mathbb{R}$. Cardinality of \mathbb{X} must be the same as the one of \mathbb{R} . The inverse map is $g_{\mathbb{X}} = f_{\mathbb{X}}^{-1}$, $g_{\mathbb{X}} : \mathbb{R} \rightarrow \mathbb{X}$. The map g from the previous section can be an example of $g_{\mathbb{R}}$ restricted to $[0, 1]$. To put it differently, the bijection $g : [0, 1] \rightarrow [0, 1]$ can be extended to a bijection $g_{\mathbb{R}} : \mathbb{R} \rightarrow \mathbb{R}$, satisfying $g_{\mathbb{R}}(p) = \sin^2(\frac{\pi}{2}p) = g(p)$ when restricted to $p \in [0, 1]$.

We define arithmetic operations in \mathbb{X} ,

$$x \oplus_{\mathbb{X}} y = g_{\mathbb{X}}(f_{\mathbb{X}}(x) + f_{\mathbb{X}}(y)), \quad (10)$$

$$x \ominus_{\mathbb{X}} y = g_{\mathbb{X}}(f_{\mathbb{X}}(x) - f_{\mathbb{X}}(y)), \quad (11)$$

$$x \odot_{\mathbb{X}} y = g_{\mathbb{X}}(f_{\mathbb{X}}(x) \cdot f_{\mathbb{X}}(y)), \quad (12)$$

$$x \oslash_{\mathbb{X}} y = g_{\mathbb{X}}(f_{\mathbb{X}}(x)/f_{\mathbb{X}}(y)). \quad (13)$$

The arithmetic given by (10)–(13) is called projective [12, 30]. Here $f_{\mathbb{X}}$ defines an isomorphism of arithmetics. The neutral elements, $0_{\mathbb{X}} = g_{\mathbb{X}}(0)$ (projective zero in \mathbb{X}), $1_{\mathbb{X}} = g_{\mathbb{X}}(1)$ (projective one in \mathbb{X}) are to some extent analogous to qubits [29, 31].

Indeed, expressions such as $0_{\mathbb{X}} + 0_{\mathbb{Y}}$ are, in general, meaningless if $\mathbb{X} \neq \mathbb{Y}$. Just think of $\mathbb{X} = \mathbb{R}$ and $\mathbb{Y} = \mathbb{R}^2$. Even if $\mathbb{X} = \mathbb{Y}$ and $0_{\mathbb{X}} = 0_{\mathbb{Y}} = 0$, $1_{\mathbb{X}} = 1_{\mathbb{Y}} = 1$, the projective bits can be as incompatible as eigenvalues of non-commuting projectors.

However, in spite of this incompatibility, $0_{\mathbb{X}} = g_{\mathbb{X}}(0)$ and $0_{\mathbb{Y}} = g_{\mathbb{Y}}(0)$ are images of the same $0 \in \mathbb{R}$. This “ordinary zero” can play the role of an EPR-type element of reality for $0_{\mathbb{X}}$ and $0_{\mathbb{Y}}$, i.e., incompatible projective bits can be correlated by means of their elements of reality, in exact analogy to the formulas postulated by Bell in his classic analysis.

Note that (3) is an example of (10). The neutral elements are $0_{\mathbb{X}} = N \tanh(0) = 0$, $1_{\mathbb{X}} = N \tanh(1) = 0.76N$ (hence velocity $0.76c$ is the neutral element of special relativistic multiplication). The velocity of light is literally infinite, of course in the sense of $\infty_{\mathbb{X}} = N \tanh(\infty) = N$. The case $c \oplus c = c$ is an example of $\infty_{\mathbb{X}} \oplus_{\mathbb{X}} \infty_{\mathbb{X}} = \infty_{\mathbb{X}}$. Strictly speaking, a relativistic unit of velocity is not c but $c \tanh(1)$.

5. Clauser–Horne formulas for projective bits

We are interested in singlet-state probabilities,

$$\begin{aligned} P_{0_1} &= P_{1_1} = P_{0_2} = P_{1_2} = \\ \langle \psi | \hat{P}_{0_1} \otimes I | \psi \rangle &= \langle \psi | \hat{P}_{1_1} \otimes I | \psi \rangle = \\ \langle \psi | I \otimes \hat{P}_{0_2} | \psi \rangle &= \langle \psi | I \otimes \hat{P}_{1_2} | \psi \rangle = \frac{1}{2} \end{aligned} \quad (14)$$

with joint probabilities,

$$\begin{aligned} P_{0_1 0_2} &= P_{1_1 1_2} = \langle \psi | \hat{P}_{0_1} \otimes \hat{P}_{0_2} | \psi \rangle = \langle \psi | \hat{P}_{1_1} \otimes \hat{P}_{1_2} | \psi \rangle = \\ &= \frac{1}{2} \sin^2\left(\frac{\alpha - \beta}{2}\right), \end{aligned} \quad (15)$$

$$\begin{aligned} P_{0_1 1_2} &= P_{1_1 0_2} = \langle \psi | \hat{P}_{0_1} \otimes \hat{P}_{1_2} | \psi \rangle = \langle \psi | \hat{P}_{1_1} \otimes \hat{P}_{0_2} | \psi \rangle = \\ &= \frac{1}{2} \cos^2\left(\frac{\alpha - \beta}{2}\right). \end{aligned} \quad (16)$$

We will write them in a Clauser–Horne form [12]

$$P_{A_1 A_2} = \int D x \chi_{A_1}(x) \odot_{\mathbb{X}} \chi_{A_2}(x) \odot_{\mathbb{X}} \rho(x), \quad (17)$$

$$P_A = \int D x \chi_A(x) \odot_{\mathbb{X}} \rho(x) = \frac{1}{2}, \quad (18)$$

where the χ s are characteristic functions and $\rho(x) \geq 0$ is a non-negative probability density normalized to 1,

$$\int D x \rho(x) = 1. \quad (19)$$

Of course, the trick is to work with appropriate forms of the integral and employ the freedom available in possible meanings of $\odot_{\mathbb{X}}$ and $\oplus_{\mathbb{X}}$. We will assume $\mathbb{X} = \mathbb{R}$, and $g_{\mathbb{X}}(0) = 0$, $g_{\mathbb{X}}(1) = 1$. The latter two conditions imply that the values of projective bits will be given by ordinary 0 and 1.

Formulas (17) and (18) implicitly imply that measurements are modeled in the usual way by the products of $\rho(x)$ with characteristic functions,

$$\rho(x) \mapsto \chi_A(x) \odot_{\mathbb{X}} \rho(x), \tag{20}$$

$$\rho(x) \mapsto \chi_{A \cap B}(x) \odot_{\mathbb{X}} \rho(x) =$$

$$\chi_A(x) \odot_{\mathbb{X}} \chi_B(x) \odot_{\mathbb{X}} \rho(x), \tag{21}$$

and so forth. If A' denotes the set-theoretic completion of set A , then

$$\chi_A(x) \oplus_{\mathbb{X}} \chi_{A'}(x) = 1, \tag{22}$$

$$1 \ominus_{\mathbb{X}} \chi_A(x) = \chi_{A'}(x), \tag{23}$$

$$\chi_A(x) \odot_{\mathbb{X}} \chi_{A'}(x) = 0, \tag{24}$$

$$\chi_A(x) \odot_{\mathbb{X}} \chi_A(x) = \chi_A(x), \tag{25}$$

$$\chi_{A'}(x) \odot_{\mathbb{X}} \chi_{A'}(x) = \chi_{A'}(x). \tag{26}$$

The probabilities must add up to 1 in an ordinary way,

$$P_{0_1 0_2} + P_{0_1 1_2} + P_{1_1 0_2} + P_{1_1 1_2} = 1, \tag{27}$$

because this is how experimentalists will use them.

On the other hand, the integral can be additive in a more general sense of (4), similarly to fuzzy, fractal, or idempotent integrals. A dual form of normalization will be a consequence of such a generalized linearity,

$$P_{0_1 0_2} \oplus_{\mathbb{X}} P_{0_1 1_2} \oplus_{\mathbb{X}} P_{1_1 0_2} \oplus_{\mathbb{X}} P_{1_1 1_2} = 1. \tag{28}$$

Note that (27) and (28) must hold simultaneously for any $P_{A_1 A_2}$, a condition, which is not entirely trivial, but whose solution exists.

The choice of arithmetic will naturally define the integral occurring in (17)–(18). Historically the first construction of calculus based on projective arithmetic was given by Grossman and Katz in their 1972 book *Non-Newtonian Calculus* [32]^{†1}.

Bell published his paper in 1964.

^{†1}Grossman and Katz had worked on the problem since the late 1960s, but their little book, as well as its two sequels [33, 34], went practically unnoticed by both mathematicians and physicists. The main idea was rediscovered by Endre Pap and published in 1993 in a local journal of Novi Sad University [35]. Over the next two decades, the formalism developed by Pap matured into a whole new branch of applied mathematics (see [9, 10, 36]). The so-called F^α calculus on fractals [37–39] can be regarded as a special case of non-Newtonian calculus. The same can be said of Maslov's idempotent analysis [15, 16]. In 2014, the ideas of Grossman, Katz, and Pap were once again rediscovered by myself [27] and led to nontrivial applications in physics, just to mention wave equations on Koch curves (a long-standing problem of fractal analysis) [40], elements of Fourier analysis on arbitrary Cantor sets (circumventing a no-go theorem about Fourier transforms on the triadic Cantor set) [11], or the issues of dark energy and matter [41, 42]. The problem of

6. Non-Newtonian Calculus

We need an integral because Clauser–Horne formulas involve integration. In fuzzy or fractal applications, the usual strategy would be to define some measure on a fractal or fuzzy set, and only then start worrying whether the resulting integral is consistent with derivatives, typically defined by means of a completely different procedure than the one that has led to the integral. In effect, the fundamental theorem of calculus often becomes problematic [13].

The approach that starts with arithmetic is much more systematic. First, one defines a derivative by means of a formula which is a straightforward generalization of

$$\frac{dF(x)}{dx} = \lim_{\delta \rightarrow 0} \frac{F(x + \delta) - F(x)}{\delta}. \tag{29}$$

Then one demands that the integral be related to the derivative by means of the fundamental theorem of calculus. The notion of measure appears automatically at the very end, once we know how to integrate. Knowing the measure, we know how to define probability.

The non-Newtonian derivative of a function $F: \mathbb{X} \rightarrow \mathbb{Y}$ depends on the arithmetics of \mathbb{X} and \mathbb{Y} . Denoting $\delta_{\mathbb{X}} = g_{\mathbb{X}}(\delta)$, $\delta_{\mathbb{Y}} = g_{\mathbb{Y}}(\delta)$, one defines

$$\frac{DF(x)}{Dx} = \lim_{\delta \rightarrow 0} (F(x \oplus_{\mathbb{X}} \delta_{\mathbb{X}}) \ominus_{\mathbb{Y}} F(x)) \oslash_{\mathbb{Y}} \delta_{\mathbb{Y}}, \tag{30}$$

whose more practical form reads

$$\frac{DF(x)}{Dx} = g_{\mathbb{Y}} \left(\frac{d\tilde{F}(f_{\mathbb{X}}(x))}{df_{\mathbb{X}}(x)} \right). \tag{31}$$

The argument of $g_{\mathbb{Y}}$ in (31) is the (Newtonian) derivative (29) of \tilde{F} defined by the commutative diagram

$$\begin{array}{ccc} \mathbb{X} & \xrightarrow{F} & \mathbb{Y} \\ f_{\mathbb{X}} \downarrow & & \uparrow g_{\mathbb{Y}} \\ \mathbb{R} & \xrightarrow{\tilde{F}} & \mathbb{R} \end{array} \tag{32}$$

Although (31) makes non-Newtonian differentiation as simple as the Newtonian one, (30) reveals the logical structure behind the derivative. For example, it explains why we find the generalized form of additivity

$$\frac{D(F(x) \oplus_{\mathbb{Y}} G(x))}{Dx} = \frac{DF(x)}{Dx} \oplus_{\mathbb{Y}} \frac{DG(x)}{Dx} \tag{33}$$

and the generalized Leibniz rule

Bell's theorem was reformulated from a non-Newtonian perspective in a series of four papers [28, 29, 31, 43]. The version of non-Newtonian formalism introduced in [27] is based on the weakest assumptions and thus is the most general and flexible so far, at least in my opinion. A review of the formalism can be found in [12].

$$\frac{D(F(x) \odot_{\mathbb{Y}} G(x))}{Dx} = \frac{DF(x)}{Dx} \odot_{\mathbb{Y}} G(x) \oplus_{\mathbb{Y}} F(x) \odot_{\mathbb{Y}} \frac{DG(x)}{Dx}. \quad (34)$$

To define a non-Newtonian integral $\int_a^b Dx F(x)$, we demand its consistency with the derivatives (two fundamental theorems of calculus)

$$\int_a^b Dx \frac{DF(x)}{Dx} = F(b) \ominus_{\mathbb{Y}} F(a), \quad (35)$$

and

$$\frac{D}{Dx} \int_a^x Dy F(y) = F(x). \quad (36)$$

The result is

$$\int_a^b Dx F(x) = g_{\mathbb{Y}} \left(\int_{f_{\mathbb{X}}(a)}^{f_{\mathbb{X}}(b)} dr \tilde{F}(r) \right). \quad (37)$$

The argument of $g_{\mathbb{Y}}$ in (37) is the (Newtonian, hence Lebesgue, Riemann, etc.) integral of \tilde{F} defined by the commutative diagram (32). Such an integral inherits additivity,

$$\int_a^b Dx F(x) \oplus_{\mathbb{Y}} G(x) = \int_a^b Dx F(x) \oplus_{\mathbb{Y}} \int_a^b Dx G(x), \quad (38)$$

and one-homogeneity (for a constant F),

$$\int_a^b Dx F \odot_{\mathbb{Y}} G(x) = F \odot_{\mathbb{Y}} \int_a^b Dx G(x), \quad (39)$$

from the arithmetic that defines the derivative.

It should be now rather clear why non-Newtonian hidden-variable models lead to Bell-type inequalities of basically the usual form, but with the ordinary plus, minus, times, and divided replaced by \oplus , \ominus , \odot , and \oslash .

If one takes this subtlety into account, then quantum mechanical singlet-state probabilities will *not* violate the Bell inequality — not the one that can be derived for the hidden-variable model.

7. Singlet-state probabilities

It remains to construct the projective arithmetic (10)–(13) that implies (14)–(18) by means of the corresponding non-Newtonian integral (37). First, $g_{\mathbb{X}}$ will be constructed via an intermediate g , whose properties are described by the following consequence of Lemma 1.

Lemma 2: Consider four joint probabilities $p_{0_1 0_2}$, $p_{1_1 1_2}$, $p_{0_1 1_2}$, $p_{1_1 0_2}$, satisfying

$$\sum_{AB} p_{AB} = 1, \quad (40)$$

$$\sum_A p_{AA_2} = \sum_A p_{A_1 A} = \frac{1}{2}. \quad (41)$$

A sufficient condition for

$$\sum_{AB} G(p_{AB}) = 1, \quad (42)$$

for any p_{AB} satisfying (40), (41), is given by $G(p) = \frac{1}{2}g(2p)$, where g satisfies Lemma 1. Any such G has a fixed point at $p = 1/4$.

The proof can be found in [43].

Guided by Lemmas 1 and 2, we take $\mathbb{X} = \mathbb{R}$ and define (Fig. 1),

$$g_{\mathbb{X}}(x) = \frac{n}{2} + \frac{1}{2} \sin^2 \left(\pi x - \pi \frac{n}{2} \right), \quad (43)$$

$$f_{\mathbb{X}}(x) = \frac{n}{2} + \frac{1}{\pi} \arcsin \sqrt{2x - n}, \quad (44)$$

for $\frac{1}{2}n \leq x \leq \frac{1}{2}(n+1)$, $n \in \mathbb{Z}$ (for more details, see [26]). Function (43) is, up to the rescaling $g(p) \mapsto \frac{1}{2}g(2p)$ required by Lemma 2, the one we have used as the illustration of Lemma 1 for spins $1/2$, but extended from $[0, 1]$ to the whole of \mathbb{R} . Non-Newtonian integrals (17)–(18) constructed by means of (43)–(44) reconstruct singlet-state probabilities if we appropriately define the characteristic functions. For example,

$$\left(\int_{\alpha_{\mathbb{X}}}^{\beta_{\mathbb{X}}} D\lambda \right) \oslash_{\mathbb{X}} g_{\mathbb{X}}(2\pi) = \frac{1}{2} \sin^2 \left(\frac{\beta - \alpha}{2} \right), \quad (45)$$

$\alpha_{\mathbb{X}} = g_{\mathbb{X}}(\alpha)$, $\beta_{\mathbb{X}} = g_{\mathbb{X}}(\beta)$, which is the standard local hidden-variable expression postulated by Bell. It can be rewritten as a particular case of (17) if we denote

$$\rho(\lambda) = 1 \oslash_{\mathbb{X}} g_{\mathbb{X}}(2\pi) = g_{\mathbb{X}}(1/(2\pi)), \quad (46)$$

and integration is over the circle $0 \leq \lambda < g_{\mathbb{X}}(2\pi)$. The product of characteristic functions is encoded in the integration limits.

The rotational invariance of the probability is a consequence of

$$\int_{\alpha_{\mathbb{X}}}^{\beta_{\mathbb{X}}} Dx = \int_{\alpha_{\mathbb{X}} \oplus_{\mathbb{X}} \gamma_{\mathbb{X}}}^{\beta_{\mathbb{X}} \oplus_{\mathbb{X}} \gamma_{\mathbb{X}}} Dx \quad (47)$$

valid for any $\gamma_{\mathbb{X}} \in \mathbb{X}$ and any non-Newtonian integral defined by means of the arithmetic (10)–(13).

Then, what about the Bell inequality?

Of course, it is *not* violated by (45) despite the exact quantum mechanical form of the probability, and there is nothing paradoxical about this statement. Just try to derive any form of a Bell-type inequality for such a non-Newtonian local hidden-variable model. For example, following the steps of the Clauser–Horne reasoning, one arrives at the projective-arithmetic generalization of the Clauser–Horne inequality,

$$0 \leq 3 \oslash_{\mathbb{X}} P_{1_1 0_2}(\theta) \ominus_{\mathbb{X}} P_{1_1 0_2}(3\theta) \leq 1. \quad (48)$$

Inserting singlet-state probabilities into (48), one finds

$$3 \oslash_{\mathbb{X}} P_{1_1 0_2}(\theta) \ominus_{\mathbb{X}} P_{1_1 0_2}(3\theta) = 1 \quad (49)$$

for any θ , so there is no contradiction.




Fig. 1. One-to-one $f_x : \mathbb{R} \rightarrow \mathbb{R}$ (full line) and its inverse g_x (dotted) defined by (44) and (43), as implied by Lemma 2. Both functions have fixed points at integer multiples of $1/4$. The plots are given in two scales, (a) $-1 \leq x \leq 1$ and (b) $-10 \leq x \leq 10$, explaining the origin of the correspondence principle discussed in Sect. 10.

The inequality that will be indeed violated is

$$0 \leq 3P_{1_1 0_2}(\theta) - P_{1_1 0_2}(3\theta) \leq 1, \quad (50)$$

but it is derived under the *wrong* assumption of additivity (1), which does not hold for this concrete model of non-Newtonian integration. Standard Clauser–Horne inequality (50) cannot be proved for non-Newtonian hidden variables in question, so it is no surprise that it is not satisfied in our model.

The readers should keep in mind that although (27) and (28) are simultaneously valid, this is no longer true for arbitrary linear combinations of probabilities, in particular those occurring in (48) and (50).

We will now show that the relation between p and $\tilde{p} = g_x(p)$, which is at the core of the Bell inequality violation, is, in fact, a very special case of an infinite hierarchy of relations, based on an infinite hierarchy of arithmetics and calculi. What we intuitively regard as the “normal” or “our” arithmetic and calculus can correspond to *any* level of the hierarchy.

This will lead us to the notion of a Copernican hierarchy of models. We call them Copernican because they deprive our human point of view of the aura of uniqueness. Each level of such a hierarchy can be our level.

The standard Bell theorem describes a relation between any two neighboring levels of the hierarchy. A surprising consequence of this relation is that in the same way that Bell proved the non-existence of EPR elements of reality, it is possible to prove the non-existence of ourselves.

Well, at least the author of this paper exists as an element of reality.

8. Copernican hierarchies

Functions g that satisfy Lemma 1 form an interesting structure, closed under the composition of maps [43].

Lemma 3: Consider two functions $g_j : [0, 1] \rightarrow [0, 1]$, $j = 1, 2$, that satisfy assumptions of Lemma 1,

$$g_j(p) = \frac{1}{2} + h_j \left(p - \frac{1}{2} \right), \quad (51)$$

where $h_j(-x) = -h_j(x)$. Then $g_{12} = g_1 \circ g_2$ also satisfies Lemma 1 with $h_{12} = h_1 \circ h_2$,

$$g_{12}(p) = \frac{1}{2} + h_{12} \left(p - \frac{1}{2} \right). \quad (52)$$

Accordingly,

$$g_{12}(p) + g_{12}(1-p) = 1 \quad (53)$$

for any $p \in [0, 1]$.

Lemma 4: Let $g^k = g \circ \dots \circ g$, $g^{-k} = g^{-1} \circ \dots \circ g^{-1}$ (k times), $g^0(x) = x$. If g satisfies Lemma 1,

$$g(p) = \frac{1}{2} + h \left(p - \frac{1}{2} \right), \quad (54)$$

then g^k also satisfies Lemma 1 for any $k \in \mathbb{Z}$,

$$g^k(p) = \frac{1}{2} + h^k \left(p - \frac{1}{2} \right). \quad (55)$$

Accordingly,

$$g^k(p) + g^k(1-p) = 1 \quad (56)$$

for any $p \in [0, 1]$, and any integer k . In particular,

$$g^{-1}(p) + g^{-1}(1-p) = 1. \quad (57)$$

The proofs are straightforward [43].

As an illustration, consider again $g(p) = \sin^2\left(\frac{\pi}{2}p\right)$ and

$$g^2(p) = \sin^2 \left[\frac{\pi}{2} \sin^2 \left(\frac{\pi}{2}p \right) \right], \quad (58)$$

$$g^{-1}(p) = \frac{2}{\pi} \arcsin \sqrt{p}. \quad (59)$$

The cross-check of (56) for (58) is simple but instructive

$$\begin{aligned} g^2(p) + g^2(1-p) &= \\ \sin^2 \left[\frac{\pi}{2} \sin^2 \left(\frac{\pi}{2}p \right) \right] + \sin^2 \left[\frac{\pi}{2} \sin^2 \left(\frac{\pi}{2}(1-p) \right) \right] &= \\ \sin^2 \left[\frac{\pi}{2} \sin^2 \left(\frac{\pi}{2}p \right) \right] + \sin^2 \left[\frac{\pi}{2} \cos^2 \left(\frac{\pi}{2}p \right) \right] &= \\ \sin^2 \left[\frac{\pi}{2} \sin^2 \left(\frac{\pi}{2}p \right) \right] + \sin^2 \left[\frac{\pi}{2} \left(1 - \sin^2 \left(\frac{\pi}{2}p \right) \right) \right] &= \\ \sin^2 \left[\frac{\pi}{2} \sin^2 \left(\frac{\pi}{2}p \right) \right] + \cos^2 \left[\frac{\pi}{2} \sin^2 \left(\frac{\pi}{2}p \right) \right] &= 1. \end{aligned} \quad (60)$$

An analogous proof for (59) is left as an exercise. Figure 2 shows the result.

We are inclined to believe that “our” arithmetic corresponds to $k = 0$. So, consider any binary probabilities from level 0,

$$p_0 + p_1 = 1, \tag{61}$$

and those from level k ,

$$g^k(p_0) + g^k(p_1) = 1. \tag{62}$$

Denoting $P_0 = g^k(p_0)$, $P_1 = g^k(p_1)$, we obtain a symmetric rule,

$$P_0 + P_1 = 1, \tag{63}$$

$$g^{-k}(P_0) + g^{-k}(P_1) = 1. \tag{64}$$

In both cases, k is an arbitrary integer: positive, negative, or zero.

The question is: How do we know that it is p_A and not P_A that defines level-zero probabilities?

We can phrase the same question in arithmetic terms. To this end, assume $g(p)$ is a restriction to $[0, 1]$ of some bijection $g_{\mathbb{R}} : \mathbb{R} \rightarrow \mathbb{R}$ that satisfies $g_{\mathbb{R}}(1) = g(1) = 1$. We can act on both sides of (62) with g^{-k} , while on both sides of (65) with g^k , obtaining

$$g^{-k}(g^k(p_0) + g^k(p_1)) = p_0 \oplus_k p_1 = 1, \tag{65}$$

and

$$g^k(g^{-k}(P_0) + g^{-k}(P_1)) = P_0 \oplus_{-k} P_1 = 1. \tag{66}$$

We have no criterion that could tell us which of the four additions — (61), (63), (65), or (66) — defines the level of description we employ in everyday life. Which of these two probabilities, and which of the several ways of adding them, is our usual way of processing experimental data?

Which of the three additions, $+$, \oplus_k , or \oplus_{-k} , is the one we have learned as kids?

Which of the three derivatives, (29), or

$$\frac{DF(x)}{Dx} = \lim_{\delta \rightarrow 0} (F(x \oplus_k \delta_k) \ominus_k F(x)) \oslash_k \delta_k, \tag{67}$$

or

$$\frac{DF(x)}{Dx} = \lim_{\delta \rightarrow 0} (F(x \oplus_{-k} \delta_{-k}) \ominus_{-k} F(x)) \oslash_{-k} \delta_{-k}, \tag{68}$$

is the one we have mastered during our undergraduate education?

Last but not least, which of the three integrals, $\int Dx F(x)$, is the one that should define a hidden-variable theory?

Numerous fundamental answers are possible.

One possibility is that Nature prefers only one $k \in \mathbb{Z}$ as the true physical arithmetic with some fixed form of g , determined by some unknown physical law. This is the situation we encounter in special relativity when we add velocities by means of $g = \tanh$. In principle, we can detect such a physical g in an experiment. In [42], it is shown that problems with dark energy may indicate that time at cosmological scales involves a nontrivial $g \sim \sinh$. If this conclusion were true, dark energy would be as unreal as the luminiferous aether.




Fig. 2. The results of $g^{-1}(p) = \frac{2}{\pi} \arcsin \sqrt{p}$ (full line), $g^{-1}(1-p)$ (dotted), and their sum (dashed).

The second fundamental possibility is that all these possibilities are simultaneously true. Perhaps there is no preferred k , like there is no preferred rest frame or preferred point of observation of the universe. Only relative k might be observable. Such an option is intriguing and tempting from a theoretical perspective. It could mean, for example, that the same physical law might have its mathematical representations at any level of the hierarchy, and each of these representations might be meaningful. Violation of Bell-type inequalities would then be a conflict of predictions derived at level k but tested at level $k + 1$.

A similar conflict occurs if Bob concludes that Alice will never reach the Schwarzschild radius, and yet she crosses it in a finite time.

It remains to say something about the conflicts that occur between non-neighboring levels of the hierarchy. We will see that other well-known bounds, such as the Tsirelson inequality characterizing Hilbert-space models of probability, can be easily circumvented as well.

9. Beyond Tsirelson’s bounds

The standard Clauser–Horne inequality (50)

$$0 \leq 3P_{1_1 0_2}(\theta) - P_{1_1 0_2}(3\theta) \leq 1, \tag{69}$$

is derived for joint probabilities limited by

$$0 \leq P_{1_1 0_2}(\alpha) \leq 0.5. \tag{70}$$

The absolute bounds for such a linear combination of probabilities are, therefore,

$$-0.5 \leq 3P_{1_1 0_2}(\theta) - P_{1_1 0_2}(3\theta) \leq 1.5. \tag{71}$$

Tsirelson bounds are narrower,

$$-\frac{\sqrt{2}-1}{2} \leq 3P_{1_1 0_2}(\theta) - P_{1_1 0_2}(3\theta) \leq \frac{\sqrt{2}+1}{2}. \tag{72}$$

In order to understand the influence of k on the violation of Clauser–Horne $k = 0$ inequalities, we have to estimate the expression [43]

$$X(g^k, \theta) = 3g^k\left(\frac{\theta}{2\pi}\right) - g^k\left(\frac{3\theta}{2\pi}\right), \tag{73}$$

where $0 \leq \theta < \pi/3$. The singlet-state example corresponds in this range of parameters to $g^1(p) = \frac{1}{2} \sin^2(\pi p)$. For $\theta = \pi/4$, we find

$$X(g^1, \frac{\pi}{4}) = \frac{3}{2} \sin^2\left(\pi \frac{\pi/4}{2\pi}\right) - \frac{1}{2} \sin^2\left(\pi \frac{3\pi/4}{2\pi}\right) - \frac{\sqrt{2}-1}{2} = -0.20711, \tag{74}$$

that is, the maximal left Tsirelson bound. This is what is usually called the maximal (left) violation of the Clauser–Horne inequality by singlet-state quantum probabilities. For other values of k , we find

$$\begin{aligned} X(g^0, \frac{\pi}{4}) &= 0, \\ X(g^1, \frac{\pi}{4}) &= -0.20711, \\ X(g^2, \frac{\pi}{4}) &= -0.39602, \\ X(g^3, \frac{\pi}{4}) &= -0.48669, \\ X(g^4, \frac{\pi}{4}) &= -0.49978, \\ &\vdots \\ X(g^\infty, \frac{\pi}{4}) &= -0.5. \end{aligned} \tag{75}$$

Of course, as stressed before, the choice of $k=0$ as the reference level is arbitrary. Level $k=2022$ probabilities violate level $k=2021$ inequalities in exactly the same way as quantum mechanics violates the standard Clauser–Horne inequality.

More importantly, $k=2$ probabilities violate $k=1$ inequalities in the same way as $k=1$ probabilities violate $k=0$ inequalities. If $k=0$ elements of reality do not exist, then $k=1$ elements of reality do not exist either. Accepting the logic of Bell's theorem, can we prove by induction that nothing exists?

Slightly modifying the experimental configuration, one obtains the maximal right violations. In our formalism, the function to estimate is

$$Y(g^k, \theta) = 3g^k \left(\frac{3\theta}{2\pi}\right) - g^k \left(\frac{\theta}{2\pi}\right). \tag{76}$$

We find

$$\begin{aligned} Y(g^0, \frac{\pi}{4}) &= 1, \\ Y(g^1, \frac{\pi}{4}) &= \frac{1}{2}(\sqrt{2} + 1) = 1.20711, \\ Y(g^2, \frac{\pi}{4}) &= 1.39602, \\ Y(g^3, \frac{\pi}{4}) &= 1.48669, \\ Y(g^4, \frac{\pi}{4}) &= 1.49978, \\ &\vdots \\ Y(g^\infty, \frac{\pi}{4}) &= 1.5. \end{aligned} \tag{77}$$

All these models are local-realistic, observers have free will, and detectors are ideal. The only modification is in the presence of the bijection g that links arithmetics, calculi, and probabilities at various levels of the hierarchy.

Our g^k plays a role analogous to g_r that linked experimental data collected at different neighborhoods of a collapsing star.

Both examples are based on principles of relativity. We have learned to live with special relativity, general relativity, and the Copernican principle.

It is time to learn to live with the arithmetic principle of relativity.

10. Correspondence principles

Trigonometric functions $\cos_{\mathbb{X}} : \mathbb{X} \rightarrow \mathbb{X}$, $\sin_{\mathbb{X}} : \mathbb{X} \rightarrow \mathbb{X}$ are defined by

$$\cos_{\mathbb{X}}(x) = g_{\mathbb{X}}(\cos(f_{\mathbb{X}}(x))), \tag{78}$$

$$\sin_{\mathbb{X}}(x) = g_{\mathbb{X}}(\sin(f_{\mathbb{X}}(x))). \tag{79}$$

They satisfy all the usual trigonometric relations, of course with respect to appropriate arithmetic operations. They also satisfy all the usual differential relations, of course with respect to appropriate non-Newtonian derivatives. In particular, they define circles by

$$\theta \mapsto (r \odot_{\mathbb{X}} \cos_{\mathbb{X}} \theta, r \odot_{\mathbb{X}} \sin_{\mathbb{X}} \theta). \tag{80}$$

Let us now take the bijections (43) and (44), which we have used to reconstruct singlet state probabilities. Figure 3 shows seven circles defined by (80) for decreasing radii. A picture to the right is a close-up of its left neighbor. All the circles are given by the same formula (80), with the same bijection $g_{\mathbb{X}}$ — the greater the radius, the more circular the shape. Simply put, the larger the x argument, the more difficult it is to tell $g_{\mathbb{X}}(x)$ from x . However, the readers must bear in mind that all these circles are *truly* rotationally invariant! They have been generated as homogeneous spaces of the rotation group in 2D — the only nonstandard element being the choice of arithmetic.

The notion of a *hidden* or *internal* symmetry, often used in particle physics, seems especially adequate here. Each of these circles would have looked “normal” if we had reprogrammed Wolfram Mathematica to make the plots in the arithmetic $\{\mathbb{R}, \oplus_1, \ominus_1, \odot_1, \oslash_1\}$.

The limit $r \rightarrow \infty$ plays a role of a correspondence principle with the ordinary, rotational external symmetry. The obvious similarity to the classical limit of quantum mechanics is striking. Other examples of arithmetic correspondence principles can be found in [27] and [44]. An analogous correspondence principle occurs in the idempotent analysis [18].

11. Implications for cryptography

In 1862, more than a century before Bell's paper, George Boole submitted to Philosophical Transactions of the Royal Society the article *On the theory of probabilities*, where he introduced inequalities imposing constraints on our “possible experience” [45].




Fig. 3. Seven circles of different radii described by (80) with the arithmetic $\{\mathbb{R}, \oplus_1, \ominus_1, \odot_1, \oslash_1\}$ defined by (43) and (44). Despite appearances, all these circles are rotationally invariant. Panels to the right are the close-ups of those to the left.

Three decades after Bell’s theorem, in 1994, Itamar Pitowsky noticed that Boole’s inequalities are inequalities of a Bell-type [46].

Boole inequalities defined “possible experience” in common-sense categories appropriate for 1862. Boole’s scientific paradigm has been falsified by quantum mechanics.

If someone had asked Boole if he could give an example of a system that violates his inequalities, he probably would have answered in the negative. Treating his negative answer as the ultimate proof that Nature has to comply with Boole inequalities, we would prove that quantum mechanics is logically impossible.

Bell inequalities defined “possible experience” in common-sense categories appropriate for 1964. Grossman and Katz’s book appeared in 1972, but its implications for Bell’s theorem went unnoticed until very recently.

In light of these results, what is the actual status of all the claims about the fundamental security of quantum cryptography [47–51]? We typically base them on the belief that EPR elements of reality cannot exist. Protocols that are not based on a hidden-variable argumentation (such as the Bennett–Brassard–Mermin one [49], essentially based on rotational invariance of singlet-state probabilities) *can* be successfully attacked in non-Newtonian local hidden variable theories — non-Newtonian hidden variables are rotationally invariant because the rotation group works there by means of the *hidden* representation depicted in Fig. 3.

Furthermore, what if the Newtonian paradigm of contemporary quantum mechanics will one day be falsified by some new theory?

What if it has already been falsified? What if our enemies, whoever they may be, are well ahead of us and know systems that can mimic quantum probabilities by means of non-Newtonian hidden variables? Can they hack entangled-state quantum communication channels?

Can a no-go theorem, based on algebraic rather than probabilistic properties of quantum mechanics, cure the arithmetic loophole in quantum proofs of security?

How to guarantee that we are not in the position of German cryptographers in the 1930s, so happy with their Enigma and its security certified by appropriate theorems, while at the same time, it was systematically hacked by the Polish Cypher Bureau?

The list of open questions is longer.

12. Non-Newtonian quantum mechanics

Non-Newtonian hidden variables are not meant as an alternative to quantum mechanics.

However, non-Newtonian calculus paves the way to natural generalizations of quantum mechanics (quantum mechanics on a Cantor set is an example [42]). The resulting theory is non-Newtonian linear but Newtonian nonlinear. Such a form of nonlinear quantum mechanics [52–55] is isomorphic to the standard textbook theory, so it is free of all the difficulties that have plagued the formalisms based on nonlinear Schrödinger equations [56–59].

Yet, “mathematically isomorphic” is not synonymous with “physically equivalent”.

The following three examples illustrate the idea.

Assume $\mathbb{X} = \mathbb{Y} = \mathbb{R}$ with projective arithmetics defined by some $f_{\mathbb{X}} : \mathbb{R} \rightarrow \mathbb{R}$ and $f_{\mathbb{Y}} : \mathbb{R} \rightarrow \mathbb{R}$. Let $\psi : \mathbb{X} \rightarrow \mathbb{Y}$ be a solution of

$$H\psi(x) = \ominus_{\mathbb{Y}}\psi''(x) \oplus_{\mathbb{Y}} U(x) \odot_{\mathbb{Y}} \psi(x) = E \odot_{\mathbb{Y}} \psi(x), \tag{81}$$

where $\psi''(x)$ is the non-Newtonian second derivative. Normalization of states is assumed in the form

$$\langle \psi | \psi \rangle = \int_{(-\infty)_{\mathbb{X}}}^{\infty_{\mathbb{X}}} Dx |\psi(x)|^{2_{\mathbb{Y}}} = 1_{\mathbb{Y}} = g_{\mathbb{Y}}(1). \tag{82}$$




Fig. 4. Probability (88) of finding a particle in $[-a, a]$ for $0 \leq a \leq 3$, with $\tilde{\psi}(r) \sim \exp(-r^2/2)$ representing the ground state of a quantum harmonic oscillator $\tilde{U}(r) = r^2$ (in dimensionless units) for (i) the ordinary arithmetic (full line), and (ii) projective arithmetics in $\mathbb{X} = \mathbb{R} = \mathbb{Y}$ defined by $f_{\mathbb{X}}(x) = x^3$, $f_{\mathbb{Y}}(x) = x$ (dashed). The ordinary arithmetic is experimentally distinguishable from the projective one because limits of integration are $[-a^3, a^3]$ instead of the usual $[-a, a]$.




Fig. 5. The same situation as in the previous figure, but now with $f_{\mathbb{X}}(x) = x^3 = f_{\mathbb{Y}}(x)$.

(For real-valued $\psi(x)$, the modulus in

$$|\psi(x)|^{2_{\mathbb{Y}}} = \psi(x) \odot_{\mathbb{Y}} \psi(x) \tag{83}$$

is redundant, but we keep it to make the notation less awkward.) The probability of finding a particle in $[a, b] \subset \mathbb{X}$ equals

$$P(a, b) = \int_a^b D_{\mathbb{X}} |\psi(x)|^{2_{\mathbb{Y}}}. \tag{84}$$

As usual, $\psi = g_{\mathbb{Y}} \circ \tilde{\psi} \circ f_{\mathbb{X}}$, $U = g_{\mathbb{Y}} \circ \tilde{U} \circ f_{\mathbb{X}}$ (compare (32)). Let $\tilde{\psi}''(f_{\mathbb{X}}(x))$ be the Newtonian second derivative of $\tilde{\psi}$ with respect to $f_{\mathbb{X}}(x)$ so that the non-Newtonian Schrödinger equation is equivalent to the usual Newtonian equation, but with redefined parameters, i.e.,

$$f_{\mathbb{Y}}(E)\tilde{\psi}(r) = -\tilde{\psi}''(r) + \tilde{U}(r)\tilde{\psi}(r), \tag{85}$$

$$1 = \langle \tilde{\psi} | \tilde{\psi} \rangle = \int_{-\infty}^{\infty} dr |\tilde{\psi}(r)|^2. \tag{86}$$

Now let us consider $f_{\mathbb{X}}(x) = x^3$, $f_{\mathbb{Y}}(x) = x$. Then $\psi = \tilde{\psi} \circ f_{\mathbb{X}}$, $U = \tilde{U} \circ f_{\mathbb{X}}$, and the Schrödinger equation is just

$$E\tilde{\psi}(r) = -\tilde{\psi}''(r) + \tilde{U}(r)\tilde{\psi}(r), \tag{87}$$

so apparently the problem is completely equivalent to the standard one. However, due to the triviality of $f_{\mathbb{Y}}$ and the non-triviality of $f_{\mathbb{X}}$, probability (84) is now explicitly given by

$$P(a, b) = \int_{f_{\mathbb{X}}(a)}^{f_{\mathbb{X}}(b)} dr |\tilde{\psi}(r)|^2 = \int_{a^3}^{b^3} dr |\tilde{\psi}(r)|^2. \tag{88}$$

As we can see, despite the mathematical banality of the problem, the non-Diophantine arithmetic of \mathbb{X} does influence the probability of finding the particle in the interval $[a, b]$ because the integral is over $[a^3, b^3]$. Figure 4 shows the probability of finding the particle in $[-a, a]$ as a function of a .

Taking $f_{\mathbb{X}}(x) = x^3 = f_{\mathbb{Y}}(x)$, we obtain the probability depicted in Fig. 5.

As the third example consider $f_{\mathbb{X}}(x) = x$, $f_{\mathbb{Y}}(x) = x/\sqrt{|x|}$, and $g_{\mathbb{Y}}(x) = f_{\mathbb{Y}}^{-1}(x) = x^3/|x|$. Now,

$$P(a, b) = \left(\int_a^b dr |\tilde{\psi}(r)|^2 \right)^2. \tag{89}$$

The projective addition of probabilities looks here like a superposition principle from quantum mechanics,

$$P(a, c) = P(a, b) \oplus_{\mathbb{Y}} P(b, c) = \left(\sqrt{P(a, b)} + \sqrt{P(b, c)} \right)^2. \tag{90}$$

Theories based on non-Newtonian calculi involve the same physical principles, but their mathematical forms may differ from one another.

Is there any natural law that determines the form of arithmetic and calculus?

13. Towards a new paradigm

Paul Benioff, a pioneer of quantum computation, was among those physicists who believed that physics and mathematics should be logically formulated at a unified level [60, 61]. According to Benioff, physical or geometric quantities do not possess numerical values per se, but these values are introduced through “value maps”. Natural numbers are elements of any well-ordered set, and in themselves do not possess numerical values. A value map takes a number and turns it into an object with concrete numerical properties. This is somewhat similar

to the idea that “zero”, the neutral element of addition, can be, in fact, an arbitrary point $0_{\mathbb{X}} \in \mathbb{X}$, provided that \mathbb{X} can be bijectively mapped onto \mathbb{R} by means of some $f_{\mathbb{X}}$ that fulfills $f_{\mathbb{X}}(0_{\mathbb{X}}) = 0 \in \mathbb{R}$. Benioff considered only linear value maps but allowed for the possibility of value-map fields. One of his conclusions was that scalar value maps are in many respects analogous to the Higgs field [62–64].

A fundamental role of the set of bijections occurs also in Etesi’s recent reformulation of black-hole entropy [65]. The “arithmetic continuum” \mathbb{R} plays there a role of a gas subject to thermodynamic laws, while the black hole entropy is a purely set-theoretic notion, related to Gödel’s first incompleteness theorem.

The approach advocated in our work involves arithmetics and calculi constructed by means of bijections that can be regarded either as (global) non-linear Benioff’s maps or as compositions of value maps with some bijections. More precisely, the “nonlinear” bijection $f_{\mathbb{X}} : \mathbb{X} \rightarrow \mathbb{R}$ is always linear here, but with respect to $\oplus_{\mathbb{X}}$, $\ominus_{\mathbb{X}}$, $\odot_{\mathbb{X}}$, and $\oslash_{\mathbb{X}}$. The non-Newtonian formalism in its most general form demands only the bijectivity of $f_{\mathbb{X}}$. One does not impose continuity or topological conditions on either $f_{\mathbb{X}}$ or \mathbb{X} . Note that $f_{\mathbb{X}}$ is always smooth in the topology and calculus it induces from \mathbb{R} , even if \mathbb{X} is as weird as Cantor or Sierpiński fractals. Non-Newtonian derivatives of $f_{\mathbb{X}}$ and $g_{\mathbb{X}}$ are “trivial” (equal to 1 and $1_{\mathbb{X}}$, respectively [12]) because from the point of view of the projective arithmetic in \mathbb{X} , the map $f_{\mathbb{X}}$ behaves as the identity map.

The duality between non-Newtonian linearity and Newtonian nonlinearity is one of the trademarks of the new paradigm. This is not the usual linearization of a nonlinear problem by a nonlinear change of variables. The idea can be traced back to Maslov’s superposition principle and its application to nonlinear optimization problems [15].

The resulting structure is incredibly flexible. It automatically leads to well-behaved calculi on all sets whose cardinalities are the same as the cardinality of the continuum. The resulting relativity principle (relativity of arithmetic and calculus) is much more general than the principle of general covariance.

Non-Newtonian calculus has a huge potential for the unification and systematization of various ideas scattered over mathematical and physical literature [29]. It is quite typical, however, that even if some elements of non-Newtonian thinking can be identified in those works, their arithmetic aspects are not exploited in their full generality.

For example, velocities in special relativity are added and subtracted in a projective way, but it is difficult to find a paper where repeated addition would be replaced by multiplication and its inverse — division. I have found only one place in relativistic physics where velocity $v = c \tanh(1)$, the “one” in special-relativistic projective multiplication, plays a distinguished role [66].

Kolmogorov–Nagumo averages [67, 68], the departure point of Rényi’s studies on generalized entropies [69], are exactly the averages in the sense of projective arithmetic. However, when Rényi discussed the additivity of his α -entropies, he did not think of additivity in the same sense as the one he implicitly used in Kolmogorov–Nagumo averaging. Various forms of projective arithmetic operations and derivatives have been studied in the context of generalized statistical physics and thermodynamics by Kaniadakis [70–73], but only some of the derivatives he invented were non-Newtonian, whereas the others were neither Newtonian nor non-Newtonian, a fact explaining why only the non-Newtonian ones have found applications [29]. The whole field of psychophysics is implicitly based on projective addition (see Chapter 7 in [12]). Typically, we are unaware that decibels and star magnitudes correspond to logarithmic scales because our sensory systems induce projective arithmetic in our brains, based on approximately logarithmic bijections (the Weber–Fechner law). However, although projective subtraction is here essential, the remaining three arithmetic operations are not employed. Certain elements of non-Newtonian integral calculus are present in cepstral analysis and homomorphic filtering of images [74]. Fractional derivatives can be regarded as non-Newtonian first derivatives, but only when formulated in the so-called F^{α} formalism [39]. Fuzzy calculus is non-additive but not necessarily fully non-Newtonian, and this is why the fundamental theorem of calculus does not necessarily work. Non-additive probability, somewhat similar to our non-Newtonian hidden variables (as based on non-additive Vitali and Choquet integrals), is a standard element of modern decision theory [5, 8].

Perhaps the most radical view on generalized arithmetics is due to Mark Burgin, who studied arithmetics that are *not* isomorphic to the arithmetic of natural numbers [75]. One of his goals was to replace inconsistent arithmetics (e.g., the computer arithmetic based on the notion of “machine infinity”: $\infty_M < \infty$, $\infty_M + 1 = \infty_M$) with arithmetics that are consistent but non-Diophantine.

Similarly radical is the approach of Sergeev [76], where infinities and infinitesimals are reformulated in a more intuitive and, essentially, non-Diophantine way. Here the infinity of integers is twice bigger than the infinity of natural numbers, while events of zero probability cannot happen (as opposed to the Kolmogorovian formalism based on measures) [77]. Such a new arithmetic and probabilistic paradigm often turns out to be more practical than the usual Kolmogorovian framework, just to mention Sergeev’s Infinity Calculator software.

The interference of probabilities is one of the greatest puzzles of quantum mechanics. Quantum probabilities sometimes behave as if they were negative, a situation known from projective arithmetics based on, say, $g_{\mathbb{X}}(p) = \ln(p)$. Risk aversion paradoxes in economics can be modeled by

non-additive integrals [5], but a new tendency appears where the same effects are modeled by quantum probabilities [78]. Ironically, while here we have shown that quantum probabilities can be classical but non-Newtonian, some authors are starting to replace non-additive integrals from classical economics with quantum probabilities [79].

Another aspect of interference is the superposition principle and the problem of linearity of quantum mechanics. Non-Newtonian linear Schrödinger equation can be Newtonian nonlinear. In such nonlinear quantum mechanics, the superposition principle remains the same as in the linear theory; only the meaning of “plus” and “times” is different. The same type of duality was introduced by Maslov to optimization theory [15], with the key idea that something very difficult in a nonlinear framework can become easy if we rewrite the problem in new arithmetic.

Speaking of a non-Newtonian paradigm, one typically has in mind a non-Newtonian theory of gravity (hence general relativity) or non-Newtonian mechanics (hence quantum mechanics). In the new paradigm that looms on the horizon, the term non-Newtonian may be understood in a much broader sense.

14. Conclusions

Bell overlooked the fundamental possibility of probabilistic theories based on generalized calculi and arithmetics. Both structures can be formulated at different levels, leading to a hierarchy of models related to one another by a new type of relativity principle. The violation of Bell's inequality by quantum probabilities has the same status as the paradoxes from special or general theories of relativity. They all disappear if we correctly apply appropriate relativity principles. The new framework of generalized arithmetics and calculi creates theoretical possibilities that are comparable only to those opened by the discovery of non-Euclidean geometries.

Acknowledgments

I am indebted to Kamil Nalikowski for the discussions. Calculations were carried out at the Academic Computer Center in Gdańsk. The work was supported by the CI TASK grant “Non-Newtonian calculus with interdisciplinary applications”.

References

[1] J.S. Bell, *Physics* **1**, 195 (1964).
 [2] D. Dubois, H. Prade, *Fuzzy Sets Syst.* **8**, 1 (1982).
 [3] D. Dubois, H. Prade, *Fuzzy Sets Syst.* **8**, 105 (1982).

[4] D. Dubois, H. Prade, *Fuzzy Sets Syst.* **8**, 225 (1982).
 [5] D. Schmeidler, *Econometria* **57**, 571 (1989).
 [6] R. Mesiar, *J. Math. Anal. App.* **194**, 477 (1995).
 [7] H.-J. Zimmermann, *Fuzzy Set Theory — and its Applications*, 3rd Ed., Kluwer, Boston (MA) 1996.
 [8] M. Marinacci, *Decisions Econ. Finan.* **20**, 153 (1997).
 [9] E. Pap, in: *Handbook of Measure Theory*, Vol. II, Ed. E. Pap, Elsevier, 2002, p. 1403.
 [10] M. Grabisch, J.-L. Marichal, R. Mesiar, E. Pap, *Aggregation Functions*, Cambridge University Press, Cambridge 2009.
 [11] D. Aerts, M. Czachor, M. Kuna, *Chaos Solitons Fractals* **91**, 461 (2016).
 [12] M. Burgin, M. Czachor, *Non-Diophantine Arithmetics in Mathematics, Physics, and Psychology*, World Scientific, Singapore 2020.
 [13] Q. Lei, Z. Xu, *Knowledge-Based Sys.* **76**, 1 (2015).
 [14] D. Aerts, M. Czachor, M. Kuna, *Rep. Math. Phys.* **81**, 357 (2018).
 [15] V.P. Maslov, *Uspekhi Mat. Nauk* **42**, 43 (1987) (in Russian).
 [16] V.N. Kolokoltsov, V.P. Maslov, *Idempotent Analysis and Applications*, Kluwer, Dordrecht 1997.
 [17] M. Gondran, M. Minoux, *Graphs, Dioids, and Semirings. New Models and Algorithms*, Springer, 2008.
 [18] G.L. Litvinov, V.P. Maslov, G.B. Shpiz, *Math. Notes* **69**, 696 (2001).
 [19] A. Einstein, B. Podolsky, N. Rosen, *Phys. Rev.* **47**, 777 (1935).
 [20] D. Bohm, *Phys. Rev.* **85**, 166 (1952).
 [21] D. Bohm, *Phys. Rev.* **85**, 180 (1952).
 [22] J.F. Clauser, M.A. Horne, *Phys. Rev. D* **10**, 526 (1974).
 [23] B.S. Cirel'son, *Lett. Math. Phys.* **4**, 93 (1980).
 [24] D.M. Greenberger, M.A. Horne, A. Shimony, A. Zeilinger, *Am. J. Phys.* **58**, 1131 (1990).
 [25] D. Bouwmeester, J.-W. Pan, M. Daniell, H. Weinfurter, A. Zeilinger, *Phys. Rev. Lett.* **82**, 1345 (1999).
 [26] M. Czachor, *Phys. Rev. A* **49**, 2231 (1994).
 [27] M. Czachor, *Quantum Stud. Math. Found.* **3**, 123 (2016).
 [28] M. Czachor, *Acta Phys. Pol. A* **139**, 70 (2021).
 [29] M. Czachor, *Entropy* **22**, 1180 (2020).

- [30] M. Burgin, [arXiv:1010.3287](https://arxiv.org/abs/1010.3287), 2010.
- [31] M. Czachor, *Found. Sci.* **25**, 971 (2020).
- [32] M. Grossman, R. Katz, *Non-Newtonian Calculus*, Lee Press, Pigeon Cove (MA) 1972.
- [33] M. Grossman, *The First Nonlinear System of Differential and Integral Calculus*, Mathco, Rockport (ME) 1979.
- [34] M. Grossman, *Bigeometric Calculus: A System with Scale-Free Derivative*, Archimedes Foundation, Rockport (ME) 1983.
- [35] E. Pap, *Zb. Rad. Prirod.-Mat. Fak. Ser. Mat.* **23**, 145 (1993).
- [36] E. Pap, *Int. J. Approx. Reasoning* **47**, 368 (2008).
- [37] A. Parvate, A.D. Gangal, *Fractals* **17**, 53 (2009).
- [38] A. Parvate, A.D. Gangal, *Fractals* **19**, 271 (2011).
- [39] A.K. Golmankhaneh, *Fractal Calculus and its Applications: F^α -calculus*, World Scientific, Singapore (2022).
- [40] M. Czachor, *Acta Phys. Pol. B* **50**, 813 (2019).
- [41] M. Czachor, *Int. J. Theor. Phys.* **56**, 1364 (2017).
- [42] M. Czachor, *Found. Sci.* **26**, 75 (2021).
- [43] M. Czachor, K. Nalikowski, *Found. Sci.* **138** (2022).
- [44] M.P. Piłat, *Found. Sci.* (2022).
- [45] G. Boole, *Phil. Trans. R. Soc.* **152**, 225 (1862).
- [46] I. Pitowsky, *British J. Phil. Sci.* **45**, 95 (1994).
- [47] C.H. Bennett, G. Brassard, in: *Proc. of the IEEE Int. Conf. on Computers, Systems, and Signal Processing, Bangalore*, 1984, p. 175.
- [48] A. Ekert, *Phys. Rev. Lett.* **67**, 661 (1991).
- [49] C.H. Bennett, G. Brassard, N.D. Mermin, *Phys. Rev. Lett.* **68**, 557 (1992).
- [50] N. Gisin, G. Ribordy, W. Tittel, H. Zbinden, *Rev. Mod. Phys.* **74**, 145 (2002).
- [51] D. Aerts, M. Czachor, M. Pawłowski, *Phys. Rev. A* **73**, 034303 (2006); Erratum: *Phys. Rev. A* **73**, 059901(E) (2006).
- [52] I. Białyński-Birula, J. Mycielski, *Ann. Phys.* **100**, 62 (1976).
- [53] R. Gähler, A.G. Klein, A. Zeilinger, *Phys. Rev. A* **23**, 1611 (1981).
- [54] S. Weinberg, *Ann. Phys.* **194**, 336 (1989).
- [55] H.D. Doebner, G.A. Goldin, *Phys. Rev. A* **54**, 3764 (1996).
- [56] N. Gisin, *Helv. Phys. Acta* **62**, 363 (1989).
- [57] J. Polchinski, *Phys. Rev. Lett.* **66**, 397 (1991).
- [58] M. Czachor, *Found. Phys. Lett.* **4**, 351 (1991).
- [59] M. Czachor, H.-D. Doebner, *Phys. Lett. A* **301**, 139 (2002).
- [60] P. Benioff, *Found. Phys.* **32**, 989 (2002).
- [61] P. Benioff, *Found. Phys.* **35**, 1825 (2005).
- [62] P. Benioff, *Quant. Stud. Math. Found.* **2**, 289 (2015).
- [63] P. Benioff, *Quant. Inf. Proc.* **15**, 1081 (2016).
- [64] P. Benioff, *Quant. Inf. Proc.* **15**, 3005 (2016).
- [65] G. Etesi, *Found. Sci.* **25**, 327 (2020).
- [66] M. Czachor, [arXiv:2209.06070](https://arxiv.org/abs/2209.06070), 2022.
- [67] A.N. Kolmogorov, *Atti. Acad. Naz. Lincei. Rend.* **12**, 388 (1930), reprinted in: *Selected Works of A.N. Kolmogorov, Vol. 1, Mathematics and Mechanics*, Ed. V.M. Tikhomirov, Kluwer, Dordrecht 1991.
- [68] M. Nagumo, *Japan J. Math.* **7**, 71 (1930), reprinted in: *Mitio Nagumo Collected Papers*, Eds. M. Yamaguti, L. Nirenberg, S. Mizohata, Y. Sibuya, Springer, Tokyo 1993.
- [69] A. Rényi, *MTA III. Oszt. Közl.* **10**, 251 (1960), reprinted in: *Selected Papers of Alfréd Rényi*, Ed. P. Turán, Akadémiai Kiadó, Budapest 1976.
- [70] G. Kaniadakis, *Physica A* **296**, 405 (2001).
- [71] G. Kaniadakis, *Phys. Lett. A* **288**, 283 (2001).
- [72] G. Kaniadakis, *Phys. Rev. E* **66**, 056125 (2002).
- [73] G. Kaniadakis, *Entropy* **15**, 3983 (2013).
- [74] A.V. Oppenheim, R.W. Schaffer, *IEEE Signal Process. Mag.* **21**, 95 (2004).
- [75] M.S. Burgin, *Uspekhi Mat. Nauk* **32**, 209 (1977) (in Russian).
- [76] Y.D. Sergeyev, *EMS Surv. Math. Sci.* **4**, 219 (2017).
- [77] C.S. Calude, M. Dumitrescu, *SN Comput. Sci.* **1**, 36 (2020).
- [78] D. Aerts, S. Sozzo, J. Tapia, in: *Quantum Interaction*, Lecture Notes in Computer Science, vol. 7620, 2012 p. 48.
- [79] J.R. Busemeyer, P.D. Bruza, *Quantum Models of Cognition and Decision*, Cambridge University Press, Cambridge 2014.