

## COMMUNICATION

## Side-on phosphinoboryl platinum(II) complexes

Anna Ordyszewska,\*<sup>a</sup> Antoni Czaplewski,<sup>a</sup> Tomasz Wojnowski,<sup>a</sup> Iwona Anusiewicz,<sup>b</sup> Jarosław Chojnacki,<sup>a</sup> and Rafał Grubba\*<sup>a</sup>Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

**In this work, we report the synthesis of the first phosphinoboryl transition metal complexes, featuring a direct covalent bond between boron and phosphorus atoms. The oxidative addition of bromo(phosphino)boranes to platinum(0) compounds enabled the formation of platinum(II) complexes with unprecedented side-on coordination of the boryl ligand.**

Transition metal boryl complexes have been extensively studied computationally and experimentally due to the strong  $\sigma$ -donors of Lewis acidic boryl ligands with minimal  $\pi$ -backbonding to the metal center<sup>1–13</sup>. Many groundbreaking discoveries have been made since the first mention of boryls in the 1960s<sup>14</sup>. These complexes are well established and have been used as key intermediates in numerous reactions such as diboration<sup>15–23</sup>, hydroboration<sup>24,25</sup>, the activation of CO<sub>2</sub> and CS<sub>2</sub><sup>26</sup>, and C–H borylation<sup>27–29</sup>.


Several methods have been used for the preparation of mono-, bis-, and tris-transition-metal boryl complexes. A method widely used to obtain mono- and bis-boryl complexes is the oxidative addition of diboron<sup>22,23,30–33</sup> (B–B) or borane<sup>34–37</sup> (B–X, X = H or halogen) to low-valent metal centers to provide bis(boryl) and boryl complexes, respectively. Braunschweig reported iron, ruthenium, and platinum complexes bearing ferrocenyl(bromo)boryl ligands<sup>38</sup> as well as cationic platinum(II) boryl complexes<sup>39,40</sup> demonstrating the high *trans* influence of the boryl group and applied *trans*-[(Cy<sub>3</sub>P)<sub>2</sub>Pt(Br){B(Mes)Br}] (Cy = cyclohexyl, Mes = mesityl) in the synthesis of base-free platinum(II) borylene complexes<sup>41</sup>. Furthermore, the trisboryl complexes [(Me<sub>3</sub>P)<sub>3</sub>Co(Bcat)<sub>3</sub>]<sup>42</sup> and [xant(P<sup>i</sup>Pr<sub>2</sub>)<sub>2</sub>Ir(Bcat)<sub>3</sub>] [Bcat = catecholboryl; xant(P<sup>i</sup>Pr<sub>2</sub>)<sub>2</sub> = 9,9-dimethyl-4,5-bis(diisopropylphosphino)xanthene]<sup>43</sup> have also been investigated. Notably, complexes bearing PBP polydentate borane<sup>44</sup> and boryl ligands<sup>45–52</sup> have found application in

hydrogen activation<sup>53</sup>, as catalysts in the dehydrogenation of dimethylamine<sup>54</sup> and the hydrosilylation of aldehydes<sup>55</sup>, and as pre-catalysts in ethylene polymerization<sup>56</sup> (Chart 1A). Interestingly, the cobalt(I) diphosphinoboryl complex underwent reversible double H<sub>2</sub> activation<sup>57</sup>.

While platinum complexes bearing directly bonded low-valent P–B ligands have been previously described by Bourissou<sup>58</sup> with phosphinoborane (Chart 1B) and by Drover<sup>59</sup> and our group<sup>60</sup> with diphosphinoboranes, no transition metal boryl complexes with directly bonded boron and phosphorus centers have been reported to the best of our knowledge.

In this work, we synthesized and characterized the first platinum(II) complexes with a side-on bonded phosphinoboryl ligand featuring a direct B–P bond (Chart 1C). Recently, we used bromo(phosphino)boranes as substrates to synthesize phosphinoborinium<sup>61,62</sup> and phosphinoborenium<sup>63,64</sup> cations via bromide anion abstraction. The presence of reactive B–Br and B–P bonds in these boranes encouraged us to explore them as precursors for boryl ligands. Furthermore, we supposed that the introduction of a donor phosphanyl group to the boron atom will facilitate the side-on coordination of the boryl ligand to the transition metal center.


First, we tested the reactivity of *i*Pr<sub>2</sub>N(Br)BPtBu<sub>2</sub> (**a**) with [(C<sub>2</sub>H<sub>4</sub>)Pt(PPh<sub>3</sub>)<sub>2</sub>] (**1**) in dichloromethane (Scheme 1). The reaction was initiated at –50 °C, after which the reaction mixture was gradually warmed to room temperature and stirred for 72 h. The reaction progress was monitored by <sup>31</sup>P and <sup>11</sup>B NMR spectroscopy, with the near complete consumption of the starting materials upon reaction completion.



**Chart 1.** Structures of selected transition metal complexes with boron–phosphorus ligands.

<sup>a</sup> Department of Inorganic Chemistry, Faculty of Chemistry and Advanced Materials Center, Gdańsk University of Technology, Narutowicza 11/12, 80-233 Gdańsk, Poland. E-mail: rafal.grubba@pg.edu.pl

<sup>b</sup> Laboratory of Quantum Chemistry, Department of Theoretical Chemistry, Faculty of Chemistry, University of Gdańsk, Wita Stwosza 63, 80-308 Gdańsk, Poland. Electronic Supplementary Information (ESI) available: Experimental, crystallographic, spectroscopic and computational details. See DOI: 10.1039/x0xx00000x



Scheme 1. Syntheses of platinum(II) complexes **1a** and **2a–2d**.

The desired phosphinoboryl Pt(II) complex (**1a**) was isolated as yellow crystals by diffusion crystallization from a toluene/pentane system at room temperature. The  $^{31}\text{P}$ -NMR spectrum of **1a** displayed two doublets at 42.5 and  $-67.2$  ppm with a  $^2J_{\text{PP}}$  coupling constant of 265 Hz, which were respectively attributed to the phosphine ligand and  $t\text{Bu}_2\text{P}$  fragment of the phosphinoboryl ligand. The large  $^2J_{\text{PP}}$  coupling constant indicated that the phosphine ligand and the phosphorus atom of the phosphinoboryl group were positioned trans to each other. Both resonances exhibit platinum satellites, with coupling constants of 3314 Hz ( $\text{PPh}_3$ ) and 1168 Hz ( $i\text{Pr}_2\text{NB-PtBu}_2$ ), confirming the coordination of the phosphorus atoms to the platinum(II) center. Notably, the  $^{31}\text{P}$  NMR resonance of the phosphinoboryl ligand is significantly upfield shifted relative to the phosphorus resonance of the parent borane ( $-11.2$  ppm). The  $^{11}\text{B}$  NMR spectrum of **1a** showed a single broad resonance at 40.4 ppm, similar to the corresponding resonance of the borane (39.5 ppm), suggesting the presence of a tricoordinate boron atom. Furthermore, the oxidative addition of the B–Br bond of the parent bromo(phosphino)borane to the Pt(0) center was unambiguously confirmed by the X-ray diffraction (XRD) analysis of the crystals of **1a** (for the molecular structure of this complex, see Fig. S31).

Despite the successful isolation of the target complex **1a**, NMR analysis of the reaction mixture indicated the formation of several unidentified byproducts, explaining the low isolation yield of **1a** (18%). This finding encouraged us to search for more effective alternative metal precursors for side-on phosphinoboryl complexes. We turned our attention to  $\text{Pt}(\text{PCy}_3)_2$  (**2**), which had been previously used as a substrate to synthesize numerous platinum boryl complexes.<sup>33–35,38–41</sup> Unlike the reaction involving **1**, precursor **2** cleanly reacted with  $i\text{Pr}_2\text{N}(\text{Br})\text{BPtBu}_2$  (**a**) in toluene.


Based on  $^{31}\text{P}$  NMR spectroscopy, the reaction mixture contained only the target side-on phosphinoboryl complex **2a** and free  $\text{PCy}_3$  as products (Fig. S13). This observation aligned with the proposed stoichiometry for the reaction of **2** with **a** (Scheme 1). The NMR spectroscopic features of **2a** were similar to those observed for **1a**, displaying a broad singlet in the  $^{11}\text{B}$  NMR spectrum at 39.5 ppm and two coupled doublets with platinum satellites in the  $^{31}\text{P}$  spectrum:  $\delta = 40.3$  ppm (d,  $^2J_{\text{PP}} = 239$  Hz,  $^1J_{\text{PtP}} = 3148$  Hz,  $\text{PCy}_3$ );  $\delta = -75.9$  ppm (bd,  $^2J_{\text{PP}} = 239$  Hz,  $^1J_{\text{PtP}} = 1080$  Hz,  $t\text{Bu}_2\text{P}$ ). Evaporating the solvent, dissolving the residue in

petroleum ether, and then storing the solution at  $-20$  °C provided **2a** as yellow crystals with a 61% yield. The formation of **2a** was further confirmed by single-crystal XRD (Fig. 1).

To study the influence of the steric and electronic features of parent bromo(phosphino)boranes on the reaction outcomes of **2**, we conducted a series of analogous experiments with  $i\text{Pr}_2\text{N}(\text{Br})\text{BPCy}_2$  (**b**),  $\text{Cy}_2\text{N}(\text{Br})\text{BPtBuPh}$  (**c**), and  $\text{Cy}_2\text{N}(\text{Br})\text{BPPH}_2$  (**d**). The boranes **b** and **c** reacted with **2** in a manner similar to **a**, producing only the desired side-on phosphinoboryl complexes (**2b**, **2c**) and free  $\text{PCy}_3$ . The isolation yields of complexes **2b** and **2c** were 60% and 74%, respectively. NMR spectroscopic data and XRD analyses of **2b** and **2c** fully supported the formation of side-on phosphinoboryl complexes (see ESI for details, Figures S32 and S33).

In contrast to experiments with compounds **a–c**, phosphinoborane **d**, with less bulky and electron-withdrawing phenyl substituents on the phosphorus atom, reacted with **2** in toluene through the oxidative addition of the B–P bond instead of the B–Br bond to the metal fragment (Scheme 1). Evaporation of the solvent and crystallization from petroleum ether at  $-20$  °C yielded **2d** as red crystals along with byproducts such as  $\text{PCy}_3$  and the hydrido complex  $[\text{Br}(\text{H})\text{Pt}(\text{PCy}_3)_2]$  in a molar ratio of 2:4:1. The complex **2d** exhibited solubility in hydrocarbons similar to that of the byproducts, which prevented its purification via crystallization. X-ray analysis revealed that **2d** contained two Pt(II) centers connected by two bridging  $\text{PPh}_2$  phosphido ligands (Fig. S34). Additionally, two  $\text{Cy}_2\text{NBBr}$  boryl ligands were coordinated to the first platinum atom, while one  $\text{PCy}_3$  ligand completed the coordination sphere of the second platinum atom. The XRD results aligned with the  $^{31}\text{P}\{^1\text{H}\}$  NMR data for **2d**, which displayed two coupled doublets ( $^2J_{\text{PP}} = 80$  Hz): a highly deshielded resonance at 235.3 ppm with two sets of platinum satellites ( $^1J_{\text{Pt1P}} = 979$  Hz,  $^1J_{\text{Pt2P}} = 2780$  Hz) attributed to the bridging phosphido ligands, and a second resonance at 61.4 ppm for the  $\text{PCy}_3$  ligand coupled only to one platinum center ( $^1J_{\text{Pt2P}} = 2780$  Hz). Unlike **1a** and **2a–2c**, **2d** did not show any resonance in the  $^{11}\text{B}$  NMR spectra despite the presence of two boryl ligands. The electron-withdrawing phenyl substituents on the phosphorus atom in the parent phosphinoborane facilitated the heterolytic cleavage of the B–P bond after the nucleophilic substitution of the electron-rich Pt(0) center onto the electrophilic boron atom and additionally stabilized the negative charge of the departing phosphido ligand. To our knowledge, compound **2d** is the first example of a transition metal complex containing both phosphido and boryl ligands.

All title phosphinoboryl complexes exhibited very similar structural features; therefore, they will be discussed in detail using the representative compound **2a** as an example (Fig. 1). The phosphinoboryl ligand coordinated to the Pt(II) center in a side-on manner *via* B1 and P1 atoms, forming a Pt1–B1–P1 triangle. Additionally, the phosphine and bromido ligands coordinated to the metal and were respectively positioned trans to the P1 and B1 atoms of the phosphinoboryl group. The geometry around the Pt1 atom was a distorted square planar, with the P2 and Br1 atoms deflected from the Pt1–B1–P1 plane in opposite directions by 0.411 and 0.321 Å, respectively.



**Figure 1.** X-ray structure of complex **2a**. H-atoms are omitted for clarity. Thermal ellipsoids are shown at the 50% probability level.

Due to the side-on coordination of the phosphinoboryl ligand, the B1–Pt1–P1 angle was acute, measuring  $52.55(7)^\circ$ , while the other angles around the Pt1 atom were obtuse, ranging from  $93.75(2)^\circ$  to  $112.10(7)^\circ$ . The Pt1–B1 bond distance of  $2.020 \text{ \AA}$  was very close to analogous distances reported for aminoboryl platinum complexes.<sup>34</sup> The Pt1–P1 bond ( $2.286 \text{ \AA}$ ) was slightly shorter than the distance between Pt1 and the P2 atom of the PCy<sub>3</sub> ligand ( $2.323 \text{ \AA}$ ). Moreover, the B–P bond within the phosphinoboryl ligand ( $1.921(3) \text{ \AA}$ ) was somewhat shorter than the expected length for a single covalent bond ( $1.96 \text{ \AA}$ ).<sup>65</sup> The B1 and N1 atoms exhibited planar trigonal geometries, with the sums of the angles around these atoms close to  $360^\circ$ . This is consistent with the significantly shortened B1–N1 bond distance ( $1.385(3) \text{ \AA}$ ) and suggests  $\pi$ -donation from the N1 atom to the B1 atom. Furthermore, the Pt1–Br bond distance of  $2.608 \text{ \AA}$  was notably longer than the typical Pt–Br bond length ( $2.37 \text{ \AA}$ ),<sup>65</sup> which can be attributed to the strong trans effect of the phosphinoboryl ligand.

The electronic structure of **2a** was investigated through theoretical methods. According to our findings, the HOMO of **2a** was delocalized mainly at the Pt1–B1–P1 triangle in a bonding manner, with some contribution from the lone electron pair on the Br1 atom (of nonbonding character) (Fig. S36). The LUMO was also based on the Pt1–B1–P1 region (in a bonding manner) but with antibonding contributions between the Pt1–B1–P1 triangle and neighboring atoms (Fig. S36). According to natural bond orbital analysis, the largest positive charges were located on the P2 (+1.08), P1 (+0.57), and B1 (+0.50) atoms, whereas the platinum center carried only a slight positive charge (+0.13). The largest negative charges were found on the N1 (–0.81) and Br1 (–0.78) atoms. The phosphinoboryl ligand was stabilized by  $\pi$ -donation from the N1 atom to a formally empty, unhybridized p orbital of boron, consistent with the planar trigonal geometry around these atoms. The lone pair on the P1 atom interacted not only with the metal center, forming a dative bond but also, to a lesser extent, with the antibonding  $\pi^*(\text{B1–N1})$  orbital, shortening the B1–P1 bond. The calculated 57% contribution of Pt to the  $\sigma(\text{Pt1–B1})$  bond confirmed the covalent character of this bond. The ratio of boron p- to s-orbital contributions in the  $\sigma(\text{Pt1–B1})$  bond was 1:1.82, indicating the strong  $\sigma$ -donor

properties of the phosphinoboryl ligand. These properties led to the strong trans effect of this ligand,<sup>66</sup> which manifested in a significant elongation of the Pt1–Br1 bond.

In conclusion, we developed a simple synthetic route for side-on phosphinoboryl complexes. The presence of reactive Pt–B and B–P bonds in these structures makes them promising candidates for applications in small molecule activation and catalysis.

Financial support of these studies from Gdańsk University of Technology by the DEC-2/2021/IDUB/V.6/Si grant under the SILICIUM - ‘Excellence Initiative - Research University’ program and Minigrant No. 037741 is gratefully acknowledged. The calculations have been carried out using resources provided by Wroclaw Centre for Networking and Supercomputing (<http://wccs.pl>) grant No. 435.

## Conflicts of interest

There are no conflicts to declare

## Data availability

The data supporting this article have been included as part of the ESI. Crystallographic data has been deposited at the CCDC under 2402129 (for **1a**), 2402130 (for **2**), 2402131 (for **2a**), 2402132 (for **2b**), 2402133 (for **2c**), 2402134 (for **2d**), and 2402135 (for **2H**).

## Notes and references

- U. Kaur, K. Saha, S. Gayen and S. Ghosh, *Coord Chem Rev*, 2021, **446**, 214106.
- S. Aldridge and D. L. Coombs, *Coord Chem Rev*, 2004, **248**, 535–559.
- Kays D. L. and S. Aldridge, *Contemporary Metal Boron Chemistry I: Borylenes, Boryls, Borane  $\sigma$ -Complexes, and Borohydrides*, Springer Berlin Heidelberg, 2008.
- G. J. Irvine, M. J. G. Lesley, T. B. Marder, N. C. Norman, C. R. Rice, E. G. Robins, W. R. Roper, G. R. Whittell and L. J. Wright, *Chem Rev*, 1998, **98**, 2685–2722.
- H. Braunschweig, R. D. Dewhurst and A. Schneider, *Chem Rev*, 2010, **110**, 3924–3957.
- Y. Li, J. Liu, X. Huang, L. B. Qu, C. Zhao, R. Langer and Z. Ke, *Chemistry - A European Journal*, 2019, **25**, 13785–13798.
- X. Guo and Z. Lin, *Chem Sci*, 2024, **15**, 3060–3070.
- T. R. Cundari and Y. Zhao, *Inorganica Chim Acta*, 2003, **345**, 70–80.
- W. Drescher and C. Kleeberg, *Dalton Transactions*, 2022, **51**, 4749–4759.
- S. Sakaki and T. Kikuno, *Inorg. Chem.*, 1997, **36**, 226–229.
- J. Zhu, Z. Lin and T. B. Marder, *Inorg Chem*, 2005, **44**, 9384–9390.
- H. Braunschweig and M. Colling, *Coord Chem Rev*, 2001, **223**, 1–51.
- L. Dang, Z. Lin and T. B. Marder, *Chemical Communications*, 2009, 3987–3995.
- G. Schmid and H. Noth, *Chem. Ber.*, 1968, **101**, 2502–2505.
- T. B. Marder and N. C. Norman, *Top Catal*, 1998, **5**, 63–73.

- 16 T. Ishiyama, T. Kitano and N. Miyaoura, *Tetrahedron Lett*, 1998, **39**, 2357–2360.
- 17 W. Clegg, T. R. F. Johann, T. B. Marder, N. C. Norman, A. G. Orpen, T. M. Peakman, M. J. Quayle, R. Rice and A. J. Scott, 1998, 1431–1438.
- 18 Y. G. Lawson, M. J. G. Lesley, T. B. Marder, N. C. Norman and C. R. Rice, *Chemical Communications*, 1997, 2051–2052.
- 19 K. M. Anderson, M. J. G. Lesley, N. C. Norman, A. G. Orpen and J. Starbuck, *New Journal of Chemistry*, 1999, **23**, 1053–1055.
- 20 T. Ishiyama, N. Matsuda, N. Miyaoura and A. Suzuki, *J. Am. Chem. Soc.*, 1993, **115**, 11018–11019.
- 21 C. N. Iverson and M. R. Smith III, *Organometallics*, 1997, **16**, 5450–5452.
- 22 C. N. Iverson and M. R. Smith, *J Am Chem Soc*, 1995, **117**, 4403–4404.
- 23 G. Lesley, P. Nguyen, N. J. Taylor, T. B. Marder, A. J. Scott, W. Clegg and N. C. Norman, *Organometallics*, 1996, **15**, 5137–5154.
- 24 S. A. Westcott, T. B. Marder, R. T. Baker, R. L. Harlow, J. C. Calabrese, K. C. Lam and Z. Lin, *Polyhedron*, 2004, **23**, 2665–2677.
- 25 K. Burgess and M. J. Ohlmeyer, *Chem Rev*, 1991, **91**, 1179–1191.
- 26 S. I. Källäne, T. Braun, M. Teltewskoi, B. Braun, R. Herrmann and R. Laubenstein, *Chemical Communications*, 2015, **51**, 14613–14616.
- 27 S. Ding, L. Wang, Z. Miao and P. Li, *Molecules*, 2019, **24**, 1–14.
- 28 G. Wang, L. Xu and P. Li, *J Am Chem Soc*, 2015, **137**, 8058–8061.
- 29 G. Wang, L. Liu, H. Wang, Y. S. Ding, J. Zhou, S. Mao and P. Li, *J Am Chem Soc*, 2017, **139**, 91–94.
- 30 D. Curtis, M. J. G. Lesley, N. C. Norman, A. G. Orpen and J. Starbuck, *J. Chem. Soc. { } Dalton Trans.*, 1999, **2**, 1687–1694.
- 31 W. Drescher, C. Borner, D. J. Tindall and C. Kleeberg, *RSC Adv*, 2019, **9**, 3900–3911.
- 32 C. Borner and C. Kleeberg, *Eur J Inorg Chem*, 2014, **2**, 2486–2489.
- 33 J. H. Muessig, D. Prieschl, A. Deißberger, R. D. Dewhurst, M. Dietz, J. O. C. Jiménez-Halla, A. Trumpp, S. R. Wang, C. Brunecker, A. Haefner, A. Gärtner, T. Thiess, J. Böhnke, K. Radacki, R. Bertermann, T. B. Marder and H. Braunschweig, *J Am Chem Soc*, 2018, **140**, 13056–13063.
- 34 H. Braunschweig, P. Brenner, A. Müller, K. Radacki, D. Rais and K. Uttinger, *Chemistry - A European Journal*, 2007, **13**, 7171–7176.
- 35 N. Arnold, H. Braunschweig, P. Brenner, J. O. C. Jimenez-Halla, T. Kupfer and K. Radacki, *Organometallics*, 2012, **31**, 1897–1907.
- 36 J. P. H. Charmant, C. Fan, N. C. Norman and P. G. Pringle, *Dalton Transactions*, 2007, **2**, 114–123.
- 37 J. F. Hartwig and S. Huber, 1993, **2**, 4908–4909.
- 38 H. Braunschweig, K. Radacki, D. Rais and F. Seeler, *Organometallics*, 2004, **23**, 5545–5549.
- 39 H. Braunschweig, K. Radacki and K. Uttinger, *Chemistry - A European Journal*, 2008, **14**, 7858–7866.
- 40 H. Braunschweig, K. Radacki, D. Rais and D. Scheschkewitz, *Angewandte Chemie - International Edition*, 2005, **44**, 5651–5654.
- 41 H. Braunschweig, K. Radacki and K. Uttinger, *Angewandte Chemie - International Edition*, 2007, **46**, 3979–3982.
- 42 W. Drescher, D. Schmitt-Monreal, C. R. Jacob and C. Kleeberg, *Organometallics*, 2020, **39**, 538–543.
- 43 M. A. Esteruelas, I. Fernández, A. Martínez, M. Oliván, E. Oñate and A. Vélez, *Inorg Chem*, 2019, **58**, 4712–4717.
- 44 R. C. Handford, L. Zhong and T. D. Tilley, *Organometallics*, 2022, **41**, 2631–2637.
- 45 Y. Segawa, M. Yamashita and K. Nozaki, *J Am Chem Soc*, 2009, **131**, 9201–9203.
- 46 H. Ogawa and M. Yamashita, *Dalton Transactions*, 2013, **42**, 625–629.
- 47 H. Schubert, W. Leis, H. A. Mayer and L. Wesemann, *Chemical Communications*, 2014, **50**, 2738–2740.
- 48 Y. Ding, Q. Q. Ma, J. Kang, J. Zhang, S. Li and X. Chen, *Dalton Transactions*, 2019, **48**, 17633–17643.
- 49 F. Fang, M. M. Xue, M. Ding, J. Zhang, S. Li and X. Chen, *Chem Asian J*, 2021, **16**, 2489–2494.
- 50 P. M. Rutz, J. Grunenberg and C. Kleeberg, *Molecules*, DOI:10.3390/molecules28176191.
- 51 P. Ríos, A. Rodríguez and J. López-Serrano, *ACS Catal*, 2016, **6**, 5715–5723.
- 52 P. M. Rutz, J. Grunenberg and C. Kleeberg, *Organometallics*, 2022, **41**, 3044–3054.
- 53 P. Ríos, J. Borge, F. Fernández de Córdoba, G. Sciortino, A. Lledós and A. Rodríguez, *Chem Sci*, 2021, **12**, 2540–2548.
- 54 E. H. Kwan, H. Ogawa and M. Yamashita, *ChemCatChem*, 2017, **9**, 2457–2462.
- 55 M. M. Xue, J. Chang, J. Zhang and X. Chen, *Dalton Transactions*, 2022, **51**, 2304–2312.
- 56 F. W. Seidel and K. Nozaki, *Angewandte Chemie International Edition*, 2022, **61**, e202111691.
- 57 T.-P. Lin and J. C. Peters, *J Am Chem Soc*, 2013, **135**, 15310–15313.
- 58 A. Amgoune, S. Ladiera, K. Miqueu and D. Bourissou, *J Am Chem Soc*, 2012, **134**, 6560–6563.
- 59 M. W. Drover and J. C. Peters, *Dalton Transactions*, 2018, **47**, 3733–3738.
- 60 A. Ordyszewska, N. Szykiewicz, E. Perzanowski, J. Chojnacki, A. Wiśniewska and R. Grubba, *Dalton Transactions*, 2019, **48**, 12482–12495.
- 61 K. Kaniewska-Laskowska, K. Klimsiak, N. Szykiewicz, J. Chojnacki and R. Grubba, *Chemical Communications*, 2022, **58**, 10068–10071.
- 62 K. Kaniewska-Laskowska, M. Czapla, J. Chojnacki and R. Grubba, *Dalton Transactions*, 2023, **52**, 8311–8315.
- 63 K. Kaniewska-Laskowska, A. Ordyszewska, T. Wojnowski, H. Halenka, M. Czapla, J. Chojnacki and R. Grubba, *Dalton Transactions*, 2023, **52**, 16061–16066.
- 64 T. Wojnowski, A. Ordyszewska, H. Halenka, I. Anusiewicz, J. Chojnacki, K. Kaniewska-Laskowska and R. Grubba, *Dalton Transactions*, 2024, DOI:10.1039/D4DT02656J.
- 65 P. Pyykkö and M. Atsumi, *Chemistry - A European Journal*, 2009, **15**, 186–197.
- 66 J. Zhu, Z. Lin and T. B. Marder, *Inorg Chem*, 2005, **44**, 9384–9390.