



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Discrete Applied Mathematics 129 (2003) 361–369

DISCRETE
APPLIED
MATHEMATICS

www.elsevier.com/locate/dam

The complexity of the T -coloring problem for graphs with small degree

Krzysztof Giaro, Robert Janczewski^{*,1}, Michał Małafiejski

*Foundations of Informatics Department, Faculty of Electronics, Telecommunications and Informatics,
Technical University of Gdańsk, ul. Narutowicza 11/12, Gdańsk, Poland*

Received 18 September 2000; received in revised form 10 July 2002; accepted 29 July 2002

Abstract

In the paper we consider a generalized vertex coloring model, namely T -coloring. For a given finite set T of nonnegative integers including 0, a proper vertex coloring is called a T -coloring if the distance of the colors of adjacent vertices is not an element of T . This problem is a generalization of the classic vertex coloring and appeared as a model of the frequency assignment problem. We present new results concerning the complexity of T -coloring with the smallest span on graphs with small degree Δ . We distinguish between the cases that appear to be polynomial or NP-complete. More specifically, we show that our problem is polynomial on graphs with $\Delta \leq 2$ and in the case of k -regular graphs it becomes NP-hard even for every fixed T and every $k > 3$. Also, the case of graphs with $\Delta = 3$ is under consideration. Our results are based on the complexity properties of the homomorphism of graphs.

© 2003 Published by Elsevier B.V.

Keywords: Vertex coloring; T -coloring; T -span; Homomorphism; NP-completeness

1. Introduction

We consider the T -coloring problem, as a generalized classical vertex coloring problem, which is one of the variants of the channel assignment problem in broadcast networks [8,16]. In this problem one wishes to assign to each transmitter $x_i \in \{x_1, \dots, x_n\}$, located in a region, a frequency $f(x_i)$ avoiding interference between transmitters, i.e.

* Corresponding author.

E-mail addresses: giaro@eti.pg.gda.pl (K. Giaro), skalar@eti.pg.gda.pl (R. Janczewski), mima@eti.pg.gda.pl (M. Małafiejski).

¹ Supported by FNP.

two interfering transmitters (because of proximity, meteorological or other reasons) must be assigned frequencies so that the distance between them does not belong to the forbidden set T of nonnegative integers including 0. The most common objective is to minimize the span of a frequency band. For more about applications of this problem the reader is referred to [2,3,14,15].

Let $G = (V, E)$ be a simple loopless graph with vertex set $V = V(G)$ and edge set $E = E(G)$. By $\Delta(G)$ we mean the maximum degree $\rho(v)$ over all vertices v of graph G , by $\chi(G)$ and $\omega(G)$ we denote the chromatic number and the clique number of graph G , respectively. Let $G(W)$ denote the subgraph of graph G induced by $W \subset V$.

Definition 1. Let T be a finite set of nonnegative integers satisfying $0 \in T$. By a T -coloring of graph G we mean a vertex coloring $c: V \rightarrow \mathbb{N}$ satisfying $|c(v) - c(w)| \notin T$, whenever $\{v, w\} \in E$. The T -span is defined as $\text{sp}_T(G) = \min_c \text{sp}_T(G, c)$, where $\text{sp}_T(G, c) = \max c(V) - \min c(V)$ and c is a proper vertex T -coloring of graph G . A T -coloring c is said to be optimal if $\text{sp}_T(G, c) = \text{sp}_T(G)$.

Following [13] we introduce the notion of T -graphs.

Definition 2. For a given set T , we define an infinite T -graph G_T , with vertex set $V(G_T) = \mathbb{N} \cup \{0\}$ and edge set $E(G_T) = \{\{x, y\} : |x - y| \notin T\}$. By G_T^{d+1} we mean the subgraph of G_T induced by $\{0, \dots, d\}$.

Given a graph G , set T and positive integer k , the problem of verifying the inequality $\text{sp}_T(G) \leq k$ we call the T -SPAN PROBLEM. This differs from the T -COLORING PROBLEM, which requires an optimal T -coloring as its output. The notion of a T -coloring was introduced in [8]. The problem has been studied extensively (see [3,4,12,13–18]). The majority of results concern lower and upper bounds on $\text{sp}_T(G)$, see [3,11,17]. The first complexity result comes independently from [6,12], where the authors showed NP-completeness in the strong sense of the T -SPAN PROBLEM on complete graphs (so even a pseudopolynomial algorithm for the T -SPAN PROBLEM cannot exist unless $P=NP$). We call the above problems FIXED T -SPAN PROBLEM and FIXED T -COLORING PROBLEM if set T is fixed. Furthermore, in [7] the authors have developed a linear algorithm for solving the FIXED T -COLORING PROBLEM on complete graphs (but exponential with respect to $\max T$). So far, the problem on graphs with “small” degree has been still open. Therefore, in Sections 2 and 3 we deal with some new properties of homomorphisms and in Section 5 we show NP-completeness of the FIXED T -SPAN PROBLEM on subcubic graphs (i.e. with $\Delta \leq 3$), and r -regular graphs (i.e. with all vertices of degree r) with $r \geq 3$. In Section 4 we show a polynomial time algorithm for the T -COLORING PROBLEM on graphs with $\Delta \leq 2$.

2. Simple properties of graph homomorphisms

The idea of graph homomorphism is a generalization of vertex coloring. Moreover, it generalizes the T -coloring problem as well.



Definition 3. For two simple graphs G and H a *graph homomorphism* is a function $h: V(G) \rightarrow V(H)$ such that $\{h(v), h(w)\} \in E(H)$, whenever, $\{v, w\} \in E(G)$ for all $v, w \in V(G)$.

We write $G \rightarrow H$ if there exists a homomorphism from G to H . Furthermore, if the homomorphism is onto, then it is called an *epimorphism*. In addition, if there exists h^{-1} and it is a homomorphism from H to G , then we call it an *isomorphism* and graphs G and H are said to be isomorphic, in symbols $G \simeq H$. We write $H \tilde{\subset} G$ if H is isomorphic to any subgraph of G .

There is a straightforward equivalence between the properties of T -span and the existence of homomorphism from G to G_T^{d+1} (see [13]).

Proposition 4. Given a graph G , any set T and a nonnegative integer d we have $\text{sp}_T(G) \leq d$ if and only if $G \rightarrow G_T^{d+1}$.

Let us note that if $T = \{0\}$, then the T -coloring problem reduces to the well-known vertex coloring problem, and moreover $G_T^{d+1} \simeq K_{d+1}$. Thus we get

Corollary 5. Given a graph G and a positive integer d we have $\chi(G) \leq d$ if and only if $G \rightarrow K_d$.

The composition of graph homomorphisms is still a graph homomorphism. Moreover, an image of a complete graph under a homomorphism is a complete graph with the same number of vertices so

Corollary 6. If $K_n \rightarrow G$ then $K_n \tilde{\subset} G$.

And

Proposition 7. If $h: V(G) \rightarrow V(H)$ is a homomorphism then $\psi(G) \leq \psi(H(h(V(G))))$, where ψ is any of the functions from the list $\{\chi, \omega, \text{sp}_T\}$.

From the above is easy to see that if $G \rightarrow H$ and H is bipartite, then graph G is bipartite. Concluding this section note an important upper bound proved in [17].

Theorem 8 (Tesman [17]). For any given graph G and set T the following inequality holds

$$\text{sp}_T(G) \leq |T| \cdot (\chi(G) - 1).$$

Let us also recall that

Theorem 9 (Brooks). If G is a connected graph that is neither a complete graph nor an odd cycle, then $\chi(G) \leq \Delta(G)$.




Fig. 1. Graph A_v^k replacing the vertex v .

3. Homomorphisms into odd cycles

The problem of graph homomorphism is considered in [1,5]. Let H be a fixed graph, the decision problem of the existence of a homomorphism from G to H will be denoted $\text{HOM}(H)$, where G is any graph from the specified family. The most important result comes from [9].

Theorem 10 (Hell and Nešetřil [9]). *The problem $\text{HOM}(H)$ on arbitrary graphs is polynomial, whenever H is bipartite, otherwise it is NP-complete.*

In this section we prove that the problem $\text{HOM}(C_{2k+1})$ on subcubic graphs is NP-complete for every positive integer $k \geq 2$, in contrast to the problem $\text{HOM}(C_3)$, which is polynomial. Moreover, we prove analogous result for 3-regular graphs and NP-completeness of the problem $\text{HOM}(C_{2k+1})$ on r -regular graphs, for every $r \geq 4$ and $k \geq 1$.

We start with a general construction. Let G be an arbitrary graph and k be any positive integer greater than 1. We replace each vertex $v \in V(G)$ of degree $\rho(v)$ with the graph A_v^k shown in Fig. 1 (the dotted vertical lines in Fig. 1 mean path P_k). We replace also every edge $\{v, w\} \in E(G)$ with the edge $\{v_i, w_j\}$ such that no two inserted edges are incident. Let G'_k be the graph constructed from G as above. It is easy to see that G'_k is always a subcubic graph.

Theorem 11. *The problem $\text{HOM}(C_{2k+1})$, $k \geq 2$ is NP-complete on subcubic graphs.*

Proof. By Theorem 10 it suffices to show $G \rightarrow C_{2k+1}$ iff $G'_k \rightarrow C_{2k+1}$. First, observe that $A_v^k \rightarrow C_{2k+1}$ and moreover for every homomorphism $h_v: V(A_v^k) \rightarrow V(C_{2k+1})$ we have $|h_v(\{v_1, \dots, v_{\rho(v)}\})| = 1$. Otherwise, we have $h_v(v_i) \neq h_v(v_{i+1})$ for some $i \in \{1, \dots, \rho(v) - 1\}$, hence $h_v(v_i) = h_v(x)$, where $\{v_i, s\}, \{v_{i+1}, s\}, \{s, x\} \in E(A_v^k)$ and $x \notin \{v_1, \dots, v_{\rho(v)}\}$. Thus C_{2l-1} is subgraph of $C_{2k+1}(h(V(A_v^k)))$ for some $l < k$, which is impossible. So, constructing a homomorphism $g: V(G) \rightarrow V(C_{2k+1})$ from a homomorphism $g': V(G'_k) \rightarrow V(C_{2k+1})$ is straightforward.

Conversely, let $g: V(G) \rightarrow V(C_{2k+1})$ be a homomorphism, then we let $g'(v_i) = g(v)$ and for $w \in V(A_v^k) \setminus \{v_1, \dots, v_{\rho(v)}\}$ $g'(w) = \tau_v \circ h_v(w)$, where $h_v: V(A_v^k) \rightarrow V(C_{2k+1})$ is a






Fig. 2. A graph G' .

homomorphism and τ_v is any automorphism of C_{2k+1} such that $\tau_v(h_v(v_i)) = g(v)$. One can check that $g' : V(G'_k) \rightarrow V(C_{2k+1})$ is a homomorphism. \square

Theorem 12. *The problem $\text{HOM}(C_{2k+1})$, $k \geq 2$ is NP-complete on 3-regular graphs.*

Proof. It suffices to show the equivalence $G \rightarrow C_{2k+1}$ iff $G' \rightarrow C_{2k+1}$ for any subcubic connected graph G , where $k \geq 2$ and G' is a cubic graph defined as follows. Let α_i be an isomorphism from graph G to its i th isomorphic copy G_i , for $i = 1, 2, 3$, which are vertex disjoint. Let $V_j \subset V(G)$ be the set of vertices of degree j . We define $V(G') = \bigcup_{i=1}^3 V(G_i) \cup \bigcup_{v \in V_1} \{x_v^1, x_v^2\} \cup \bigcup_{u \in V_2} \{y_u\}$ and $E(G') = \bigcup_{i=1}^3 E(G_i) \cup \bigcup_{v \in V_1} \bigcup_{i=1}^3 \{\{x_v^1, \alpha_i(v)\}, \{x_v^2, \alpha_i(v)\}\} \cup \bigcup_{u \in V_2} \bigcup_{i=1}^3 \{\{y_u, \alpha_i(u)\}\}$ (see Fig. 2). Assuming that x_v^j and y_u are different vertices for $j = 1, 2$ and $v, u \in V(G)$, it is obvious that G' is a cubic graph.

Now, suppose $g : V(G) \rightarrow V(C_{2k+1})$ is a homomorphism. Let $g' : V(G') \rightarrow V(C_{2k+1})$ be defined $g'(w) = g(v)$ for $w \in \{\alpha_1(v), \alpha_2(v), \alpha_3(v)\}$ and $v \in V(G)$, $g'(x_v^i) = g(z)$ for $\{z, v\} \in E(G)$, $g'(y_u) = g(z)$ for any z adjacent to u . Thus g' is a well-defined homomorphism. Conversely, if g' is a homomorphism from G' to C_{2k+1} then $g = g' \circ \alpha_1$ is a homomorphism from G to C_{2k+1} . \square

Theorem 13. *The problem $\text{HOM}(C_{2k+1})$ is NP-complete on r -regular graphs for every fixed integer $k \geq 1$ and $r \geq 4$.*

Proof. By induction on $r \geq 4$, consider $r + 1$ isomorphic copies of any r regular graph. Using the analogous method as that in Theorem 12 we can show that the problem $\text{HOM}(C_{2k+1})$ is NP-complete for any $k \geq 2$ and for all $r \geq 4$. In [10] the author proved NP-completeness of edge 3-chromaticity of 3-regular graphs. Since line

graphs of 3-regular graphs are 4-regular, the problem of 3-chromaticity of 4-regular graphs is NP-complete. The construction from Theorem 12 is carried over to the case $\text{Hom}(C_3)$ on r -regular graphs with $r \geq 4$. \square

4. Polynomial algorithm for cycles

We show a polynomial-time algorithm for graphs with $\Delta \leq 2$.

Theorem 14. *The T -COLORING PROBLEM on graphs with degree not exceeding 2 can be solved in time $O(n|T|^2 \log|T|)$.*

Proof. Bipartite graphs can be optimally colored with 1 and $\min \mathbb{N} \setminus T + 1$, thus all we need is considering odd cycles. Let T be any set and a be an arbitrary integer. We ask if $\text{sp}_T(C_{2k+1}) \leq a - 1$. By Theorem 8 we have $\text{sp}_T(C_{2k+1}) \leq 2|T|$. Thus using the standard bisection method we need only check $1 + \log_2|T|$ inequalities to find $\text{sp}_T(C_{2k+1})$.

In the following, we sketch the idea of the algorithm. Let $\text{TAB}(v_i)[1 \dots a]$ be a table of logical values associated with vertex v_i and defined as follows: $\text{TAB}(v_i)[j] = \text{TRUE}$ if and only if there exists a T -coloring of path v_1, \dots, v_i using colors not greater than a such that v_1 is colored with 1 and v_i is colored with j . So, $\text{TAB}(v_1)$ has value TRUE only on its first position and $\text{TAB}(v_{i+1})[y] = \text{TRUE}$ if and only if there exists $z \in \{1, \dots, a\}$ such that $|z - y| \notin T$ and $\text{TAB}(v_i)[z] = \text{TRUE}$. We see that there exists a T -coloring iff $\text{TAB}(v_{2k+1})[j] = \text{TRUE}$ for some $j - 1 \notin T$, so constructing the T -coloring is straightforward. It is obvious that the complexity of the above algorithm is $O(k|T|^2 \log|T|)$. \square

5. Main results

Based on Theorem 11 we can prove the main result of this paper. Before doing this, we introduce the following notion.

Definition 15. For a given set T , by d_T we mean the number such that $G_T^{d_T}$ is bipartite and $G_T^{d_T+1}$ is not bipartite.

Lemma 16. *For any set T the following inequality holds:*

$$d_T \leq \text{sp}_T(K_3)$$

and, moreover, d_T can be determined in polynomial time.

Proof. Let us notice that $\chi(G_T^{d_T+1}) = \chi(G_T^{d_T}) + 1 = 3$. Thus from Corollary 5 it follows $G_T^{d_T+1} \rightarrow K_3$, hence by Proposition 7 $\text{sp}_T(G_T^{d_T+1}) \leq \text{sp}_T(K_3)$. By Proposition 4 $\text{sp}_T(G_T^{d_T+1}) \leq d_T$. Assuming $\text{sp}_T(G_T^{d_T+1}) \leq d_T - 1$ we get at once $G_T^{d_T+1} \rightarrow G_T^{d_T}$ but this contradicts the definition of d_T . So, we get $d_T = \text{sp}_T(G_T^{d_T+1}) \leq \text{sp}_T(K_3)$. By






Fig. 3. A graph $G_T^{d_T+1}$ (left) and a cycle C_{2k+1} (right).

Theorem 8 $sp_T(K_3) \leq 2|T|$, hence using the bisection method we can determine the greatest d_T such that $G_T^{d_T}$ is bipartite. This can be done in time $O(|T|^2 \log|T|)$. \square

Lemma 17. Given any set T , we have $d_T = sp_T(K_3)$ if and only if $K_3 \tilde{C} G_T^{d_T+1}$.

Proof. By Corollary 6 $K_3 \tilde{C} G_T^{d_T+1}$ is equivalent to $K_3 \rightarrow G_T^{d_T+1}$. Assume $K_3 \tilde{C} G_T^{d_T+1}$, then by Proposition 4 $sp_T(K_3) \leq d_T$, hence from Lemma 16 it follows that $d_T = sp_T(K_3)$. The converse implication is straightforward by Proposition 4. \square

Let us denote by C_T the shortest odd-length cycle in graph $G_T^{d_T+1}$.

Lemma 18. There exists a homomorphism $h: V(G_T^{d_T+1}) \rightarrow V(C_T)$.

Proof. We only have to construct a homomorphism on the vertices of the connected component of $G_T^{d_T+1}$ containing vertex d_T , because the other components are bipartite. So let V_1 and V_2 be a bipartition of a bipartite graph obtained from this component by removing d_T and let W_i^j , $i = 1, 2$ and $j \geq 1$, be the vertex subset of V_i of distance j from vertex d_T in the graph $G_T^{d_T+1}$. Finally, let $W_1^0 = W_2^0 = \{d_T\}$. Let $C_{2k+1} = (\{d, a_1, b_1, \dots, a_k, b_k\}, \{\{d, a_1\}, \{d, b_1\}, \{a_1, b_2\}, \{b_1, a_2\}, \dots, \{a_{k-1}, b_k\}, \{b_{k-1}, a_k\}, \{a_k, b_k\}\})$ be any cycle isomorphic to C_T . Let us define $h(d_T) = d$, $h(W_1^j) = \{a_j\}$ and $h(W_2^j) = \{b_j\}$ for $j = 1, \dots, k$ and $h(W_i^j) = h(W_i^k)$ for $j > k$, $i = 1, 2$ (see Fig. 3).

The construction of h is correct because any vertex from W_i^j , $j > 0$, can have neighbours only in the sets $W_{3-i}^{j\pm 1}$ and W_{3-i}^j , and the latter case is impossible for $j < k$. \square

Lemma 19. For any graph G the following equivalence holds: $G \rightarrow G_T^{d_T+1}$ if and only if $G \rightarrow C_T$.

Proof. Let $G \rightarrow G_T^{d_T+1}$, hence from Lemma 18 it follows $G \rightarrow C_T$. Conversely, assume that $G \rightarrow C_T$. By definition $C_T \tilde{C} G_T^{d_T+1}$, thus we get $G \rightarrow G_T^{d_T+1}$. \square

Theorem 20. *The T -SPAN PROBLEM can be solved in polynomial time on subcubic graphs for all sets T satisfying $K_3 \tilde{C} G_T^{d_T+1}$. The FIXED T -SPAN PROBLEM is NP-complete on cubic graphs for all sets T not satisfying $K_3 \tilde{C} G_T^{d_T+1}$.*

Proof. Let T be a fixed set and k be any positive integer. By Theorem 8 the case $G = K_4$ is polynomial and can be solved in $O(|T|^3)$ time (by Proposition 4 it reduces to the problem of finding the smallest d such that $K_4 \tilde{C} G_T^d$; by Theorem 8 $K_4 \tilde{C} G_T^{3|T|+1}$ and the fact that 0 is a vertex of a maximal clique of G_T^d , it reduces to searching all the triples of vertices of $G_T^{3|T|+1}$). For any subcubic graph $G \neq K_4$ we ask if $\text{sp}_T(G) \leq k$.

Suppose that $K_3 \tilde{C} G_T^{d_T+1}$. Brooks' theorem implies $G \rightarrow K_3$, thus by Lemma 17 and Proposition 7 $\text{sp}_T(G) \leq d_T$. According to Proposition 4 we have $\text{sp}_T(G) < d_T$ iff G is bipartite, hence to solve T -SPAN PROBLEM for graph G we only need to check if G is bipartite ($O(n+m)$ time) and if it is so then $\text{sp}_T(G)$ equals the smallest positive integer not belonging to T (which we can find in $O(|T|)$ time). Otherwise, $\text{sp}_T(G) = d_T$, computable in time $O(|T|^2 \log |T|)$.

Now assume that K_3 is not isomorphic to any subgraph of $G_T^{d_T+1}$ and let $k = d_T$. From Proposition 4 we have $\text{sp}_T(G) \leq k$ iff $G \rightarrow G_T^{d_T+1}$. By Lemma 19 we get $\text{sp}_T(G) \leq k$ iff $G \rightarrow C_T$ and, moreover, C_T is an odd cycle of length greater than 4. By Theorem 12 the problem $\text{HOM}(C_T)$ on cubic graphs is NP-complete and so is the FIXED T -SPAN PROBLEM. \square

Corollary 21. *The T -SPAN PROBLEM is NP-complete in the strong sense on 3-regular graphs.*

Proof. By Theorem 20 and Lemma 17 it suffices to verify that for $T = \{0, 2, 3\}$ we have $d_T = 4 < \text{sp}_T(K_3) = 5$. \square

It is worth observing that if for some set T we put $k = d_T$, then by Lemma 19 for any graph G the question if $\text{sp}_T(G) \leq k$ is equivalent to $G \rightarrow C_T$. So, if for every $k \geq 1$ the problem $\text{HOM}(C_{2k+1})$ is NP-complete on a class \mathcal{G} , then the FIXED T -SPAN PROBLEM on the class \mathcal{G} is NP-complete as well. Thus from Theorem 13 we have the following:

Theorem 22. *For every set T and integer $r \geq 4$ the FIXED T -SPAN PROBLEM is NP-complete on r -regular graphs.*

Corollary 23. *The T -SPAN PROBLEM is NP-complete in the strong sense on r -regular graphs for any $r \geq 3$.*

Table 1 Now we sum up all the above results in the following table. Recall that the numbers appearing in the third column are polynomially computable functions of T .



Table 1

The complexity of the T -SPAN PROBLEM and T -COLORING PROBLEM on graphs with bounded degree

| Graph | Problem | Property of T | Complexity | Reference |
|-----------------------------|-------------------------|------------------------------|---------------------|------------|
| $\Delta \leq 2$ | T -COLORING PROBLEM | any | $O(n T ^2 \log T)$ | Theorem 14 |
| $\Delta \leq 3$ | T -COLORING PROBLEM | $\omega(G_T^{d_T+1}) \geq 3$ | $O(n^2 + T ^3)$ | Theorem 20 |
| 3-regular | FIXED T -SPAN PROBLEM | $\omega(G_T^{d_T+1}) \leq 2$ | NPC | Theorem 20 |
| r -regular, $r \geq 4$ | FIXED T -SPAN PROBLEM | any | NPC | Theorem 22 |

Acknowledgements

The authors are grateful to the two anonymous referees and prof. Kubale for valuable comments.

References

- [1] M.O. Albertson, Generalized Colorings, Academic Press, New York, 1987, pp. 35–49.
- [2] M. Bellare, O. Goldreich, M. Sudan, Free bits, PCP and non-approximability towards tight results, Proceedings of the 36th IEEE Symposium on Foundations of Computer Science, Los Alamos, 1995, pp. 422–431.
- [3] M.B. Cozzens, F.S. Roberts, T -colorings of graphs and the channel assignment problem, Congr. Numer. 35 (1982) 191–208.
- [4] M.B. Cozzens, F.S. Roberts, Greedy algorithms for T -colorings of complete graphs and the meaningfulness of conclusions about them, J. Combin. Inform. System Sci. 16 (1991) 286–299.
- [5] A.M.H. Gerards, Homomorphisms of graphs into odd cycles, J. Graph Theory 12 (1988) 73–83.
- [6] A. Gräf, Distance graphs and the T -coloring problem, Discrete Math. 196 (1999) 153–166.
- [7] J.R. Griggs, D.D.-F. Liu, The channel assignment problem for mutually adjacent sites, J. Combin. Theory Ser. A 68 (1994) 169–183.
- [8] W.K. Hale, Frequency assignment: theory and applications, Proceedings IEEE 68 (1980) 1497–1514.
- [9] P. Hell, J. Nešetřil, On the complexity of H -coloring, J. Combin. Theory Ser. B 48 (1990) 92–110.
- [10] I. Holyer, The NP-completeness of edge-coloring, SIAM J. Comput. 10 (1981) 718–720.
- [11] R. Janczewski, A note on divisibility and T -span of graphs, Discrete Math. 234 (2001) 171–179.
- [12] K. Jansen, A rainbow about T -colorings for complete graphs, Discrete Math. 154 (1996) 129–139.
- [13] D.D.-F. Liu, T -colorings of graphs, Discrete Math. 101 (1992) 202–212.
- [14] D.D.-F. Liu, T -graphs and the channel assignment problem, Discrete Math. 161 (1996) 197–205.
- [15] A. Raychaudhuri, Further results on T -colorings and frequency assignment problem, Discrete Math. 7 (1994) 605–613.
- [16] F.S. Roberts, T -coloring of graphs: recent results and open problems, Discrete Math. 93 (1991) 229–245.
- [17] B. Tesman, T -colorings, list T -colorings and set T -colorings of graphs, Ph.D. Thesis, Department of Math. Rutgers University, New Brunswick, NJ, 1989.
- [18] B. Tesman, Applications of forbidden difference graphs to T -colorings, Congressus Numerantium 74 (1990), 15–24.

