

First-order impulsive ordinary differential equations with advanced arguments

Tadeusz Jankowski

*Gdansk University of Technology, Department of Differential Equations, 11/12 G. Narutowicz Str.,
80–952 Gdańsk, Poland*

Received 7 March 2006

Available online 15 September 2006

Submitted by Steven G. Krantz

Abstract

This paper deals with impulsive advanced ordinary differential equations with boundary conditions. We investigate the existence of solutions and quasolutions for advanced impulsive differential equations. To obtain such results we apply Schauder's fixed point theorem. Corresponding results are also formulated for differential inequalities.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Impulsive differential equations; Equations with advanced arguments; Nonlinear boundary conditions; Existence results

1. Introduction

For $J = [0, T]$, $T > 0$, let $0 = t_0 < t_1 < \dots < t_m < t_{m+1} = T$. Put $J' = J \setminus \{t_1, t_2, \dots, t_m\}$.

In this paper, we investigate first-order impulsive advanced differential equations of type

$$\begin{cases} x'(t) = f(t, x(t), x(\alpha(t))) \equiv Fx(t), & t \in J', \\ \Delta x(t_k) = I_k(x(t_k)), & k = 1, 2, \dots, m, \\ 0 = g(x(0), x(T)), \end{cases} \quad (1)$$

where as usual $\Delta x(t_k) = x(t_k^+) - x(t_k^-)$, $x(t_k^+)$ and $x(t_k^-)$ denote the right and left limits of x at t_k , respectively, and

E-mail address: tjank@mif.pg.gda.pl.

(H₁) $f \in C(J \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$, $\alpha \in C(J, J)$, $t \leq \alpha(t) \leq T$, $t \in J$, $I_k \in C(\mathbb{R}, \mathbb{R})$ for $k = 1, 2, \dots, m$, $g \in C(\mathbb{R} \times \mathbb{R}, \mathbb{R})$ and if there exists a point $\bar{t} \in J$ such that $\alpha(\bar{t}) \in \{t_1, t_2, \dots, t_m\}$, then $\bar{t} \in \{t_1, t_2, \dots, t_m\}$.

Put $J_0 = [0, t_1]$, $J_k = (t_k, t_{k+1}]$, $k = 1, 2, \dots, m$. Let us introduce the spaces:

$$PC(J) = PC(J, \mathbb{R}) = \left\{ \begin{array}{l} x: J \rightarrow \mathbb{R}, x|_{J_k} \in C(J_k, \mathbb{R}), k = 0, 1, \dots, m, \\ \text{and there exist } x(t_k^+) \text{ for } k = 1, 2, \dots, m \end{array} \right\}$$

and

$$PC^1(J) = PC^1(J, \mathbb{R}) = \left\{ \begin{array}{l} x \in PC(J), x|_{J_k} \in C^1(J_k, \mathbb{R}), k = 0, 1, \dots, m, \\ \text{and there exist } x'(t_k^+) \text{ for } k = 1, 2, \dots, m \end{array} \right\}.$$

Indeed, $PC(J)$ and $PC^1(J)$ are Banach spaces with the respective norms:

$$\|x\|_{PC} = \sup_{t \in J} \|x(t)\|, \quad \|x\|_{PC^1} = \|x\|_{PC} + \|x'\|_{PC}.$$

By a solution of problem (1) we mean a function $x \in PC^1(J)$ which satisfies

- the differential equation in (1) for every $t \in J'$,
- the boundary condition in problem (1) and
- at every t_k , $k = 1, 2, \dots, m$, the function x satisfies the second condition in problem (1).

Throughout this paper we assume that $\alpha(t) \neq t$, $t \in J$.

An interesting and fruitful technique for proving existence results for nonlinear differential problems is the monotone iterative method, for details, see, for example, [11]. There exists a vast literature devoted to the applications of this method to differential equations with initial and boundary conditions. This technique can also be applied to impulsive differential equations, for details, see, for example, [12]. However, only a few papers have appeared where the monotone iterative technique is applied to delay impulsive differential problems, see, for example, [2,3,6,14]. Usually, it is assumed that the function f satisfies a one-sided Lipschitz condition with corresponding Lipschitz constants. For problems with deviating arguments, it is better to assume that the above constants are replaced by corresponding Lipschitz functions. I know only a few papers where such assumptions appeared for differential equations without impulsive, see [7–10]. I do not know any paper where it is done for impulsive problems with deviating arguments. Just in this paper the function f from problem (1) satisfies a one-sided Lipschitz condition (with respect to the last two variables) with functional coefficients and argument α being of advanced type. Note that impulsive differential equations are also discussed in papers [1,4,5,13].

The plan of this paper is as follows. In Section 2, we formulate sufficient conditions which guarantee that problem (1) has a solution. To prove Theorem 2 we apply Schauder's theorem. It is assumed that a lower solution of (1) is bigger than an upper solution. Indeed, first impulsive differential inequalities are investigated. In Section 3, we discuss existence of quasisolutions of problem (1). Given are two examples to show that the assumptions of this paper are satisfied.

2. Lower and upper solutions of problem (1)

Let us introduce the following definition.

We say that $u \in PC^1(J)$ is a lower solution of (1) if

$$\begin{cases} u'(t) \leq Fu(t), & t \in J', \\ \Delta u(t_k) \leq I_k(u(t_k)), & k = 1, 2, \dots, m, \\ g(u(0), u(T)) \leq 0, \end{cases} \quad (2)$$

and it is an upper solution of (1) if the above inequalities are reversed.

We assume that $z_0(t) \leq y_0(t)$, $t \in J$, and define the sector

$$[z_0, y_0]_* = \{v \in PC^1(J, \mathbb{R}): z_0(t) \leq v(t) \leq y_0(t), t \in J\}.$$

Lemma 1. Assume that $K \in C(J, \mathbb{R})$, $\alpha \in C(J, J)$, $t \leq \alpha(t) \leq T$, $t \in J$, and $L_k \geq 0$, $k = 1, 2, \dots, m$. Let $p \in PC^1(J)$ and

$$\begin{cases} p'(t) \geq K(t)p(t) + M(t)p(\alpha(t)), & t \in J', \\ \Delta p(t_k) \geq L_k p(t_k), & k = 1, 2, \dots, m, \\ p(T) \leq 0, \end{cases} \quad (3)$$

where M is nonnegative and $M \in PC(J)$.

In addition assume that

$$\int_0^T M^*(t) dt \left(\prod_{i=1}^m (1 + L_i) \right) \leq 1 \quad \text{with } M^*(t) = M(t) e^{\int_t^{\alpha(t)} K(s) ds}. \quad (4)$$

Then $p(t) \leq 0$, $t \in J$.

Proof. Put

$$q(t) = e^{\int_t^T K(s) ds} p(t), \quad t \in J.$$

Then $q(T) = p(T) \leq 0$, $\Delta q(t_k) \geq L_k q(t_k)$, $k = 1, 2, \dots, m$, and

$$q'(t) = e^{\int_t^T K(s) ds} \{-K(t)p(t) + p'(t)\} \geq e^{\int_t^T K(s) ds} M(t)p(\alpha(t)), \quad t \in J'.$$

Then system (3) takes the form

$$\begin{cases} q'(t) \geq M^*(t)q(\alpha(t)), & t \in J', \\ q(t_k^+) \geq (1 + L_k)q(t_k), & k = 1, 2, \dots, m, \\ q(T) \leq 0. \end{cases} \quad (5)$$

Note that if $q(t) \leq 0$, $t \in J$, then also $p(t) \leq 0$ on J .

We need to prove that $q(t) \leq 0$, $t \in J$. Suppose that the inequality $q(t) \leq 0$, $t \in J$, is not true. It means that we can find $t_1^* \in [0, T)$ such that $q(t_1^*) > 0$. Then $\inf_{[t_1^*, T]} q(t) = -\rho$. Indeed, $\rho \geq 0$ and there exists $t_0^* \in J_p$ for some fixed p such that $q(t_0^*) = -\rho$ or $q(t_p^+) = -\rho$. Below we discuss only the situation when $q(t_0^*) = -\rho$ because in the case when $q(t_p^+) = -\rho$, the proof is similar.

Let $t_1^* \in J_r$ for some r . Indeed, $t_1^* < t_0^*$, so $r \leq p$.

Now, for $\sigma \in PC(J)$, we consider the following inequalities

$$\begin{cases} q'(t) \geq \sigma(t), & t \in [t_1^*, T] \setminus \{t_{r+1}, \dots, t_m\}, \\ q(t_k^+) \geq (1 + L_k)q(t_k), & k = r + 1, \dots, m. \end{cases}$$

Then



$$q(t) \geq q(t_1^*) \prod_{i=r+1}^k (1 + L_i) + \sum_{i=r+1}^k \int_{\bar{t}_{i-1}}^{\bar{t}_i} \sigma(s) ds \left(\prod_{j=i}^k (1 + L_j) \right) + \int_{\bar{t}_k}^t \sigma(s) ds \tag{6}$$

for $t \in \bar{J}_k$, $k = r, r + 1, \dots, m$. Here $\bar{J}_r = [\bar{t}_r, \bar{t}_{r+1}]$, $\bar{t}_r = t_1^*$, $\bar{t}_k = t_k$, $\bar{J}_k = (\bar{t}_k, \bar{t}_{k+1}]$ for $k = r + 1, \dots, m$, and $\sum_{i=a}^b \dots = 0$, $\prod_{i=a}^b \dots = 1$ if $a > b$.

Let $\sigma(t) = M^*(t)q(\alpha(t))$. It yields $\sigma(t) \geq -\rho M^*(t)$, $t \in [t_1^*, T]$. Put $t = t_0^*$. Then

$$\begin{aligned} q(t_0^*) &\geq q(t_1^*) \prod_{i=r+1}^p (1 + L_i) + \sum_{i=r+1}^p \int_{\bar{t}_{i-1}}^{\bar{t}_i} \sigma(s) ds \left(\prod_{j=i}^p (1 + L_j) \right) + \int_{\bar{t}_p}^{t_0^*} \sigma(s) ds \\ &> \sum_{i=r+1}^p \int_{\bar{t}_{i-1}}^{\bar{t}_i} \sigma(s) ds \left(\prod_{j=i}^p (1 + L_j) \right) + \int_{\bar{t}_p}^{t_0^*} \sigma(s) ds \\ &\geq -\rho \left\{ \sum_{i=r+1}^p \int_{\bar{t}_{i-1}}^{\bar{t}_i} M^*(s) ds \left(\prod_{j=i}^p (1 + L_j) \right) + \int_{\bar{t}_p}^{t_0^*} M^*(s) ds \right\}. \end{aligned}$$

Hence, if $\rho > 0$, we have

$$\begin{aligned} 1 &< \sum_{i=r+1}^p \int_{\bar{t}_{i-1}}^{\bar{t}_i} M^*(s) ds \left(\prod_{j=i}^p (1 + L_j) \right) + \int_{\bar{t}_p}^{t_0^*} M^*(s) ds \\ &\leq \int_0^T M^*(s) ds \left(\prod_{i=1}^m (1 + L_i) \right). \end{aligned}$$

It contradicts (4).

If $\rho = 0$, then

$$0 \geq q(t_1^*) \prod_{i=r+1}^p (1 + L_i) > 0.$$

It is a contradiction too. The proof is complete. \square

Remark 1. If $M(t) = 0$, $t \in J$, then Lemma 1 reduces to Lemma 4 of [4]. Note that condition (4) is satisfied if $K(t) \geq 0$, $t \in J$, and

$$\int_0^T M(t) e^{\int_t^T K(s) ds} dt \left(\prod_{i=1}^m (1 + L_i) \right) \leq 1. \tag{7}$$

We see that condition (7) does not depend on the advanced argument α . If we extra assume that $K(t) = K > 0$, $M(t) = M > 0$, then condition (4) holds if

$$M(e^{KT} - 1) \prod_{i=1}^m (1 + L_i) \leq K.$$

For example, if we take $m = 2$, $L_1 = \frac{1}{3}$, $L_2 = \frac{1}{2}$, $T = \frac{2}{3}$, $K = \frac{3}{2}$, then from the last condition we have

$$M \leq \frac{3}{4(e-1)} \approx 0.43648.$$

Theorem 1. Assume that $K \in C(J, \mathbb{R})$, $\eta \in PC(J)$, M is nonnegative, $M(t) \neq 0$, and $M \in PC(J)$. Moreover, let $\alpha \in C(J, J)$, $t \leq \alpha(t) \leq T$, $\gamma_k, L_k \in \mathbb{R}$ and $L_k \geq 0$ for $k = 1, 2, \dots, m$. In addition, assume that

$$\delta \equiv \int_0^T M^*(s) ds + \sum_{i=1}^m L_i < 1, \quad (8)$$

where M^* is defined as in Lemma 1. Then the impulsive problem

$$\begin{cases} v'(t) = K(t)v(t) + M(t)v(\alpha(t)) + \eta(t), & t \in J', \\ v(t_k^+) = (1 + L_k)v(t_k) + \gamma_k, & k = 1, 2, \dots, m, \\ v(T) = k_0 \end{cases} \quad (9)$$

has a unique solution $v \in PC^1(J)$.

Proof. Put

$$z(t) = e^{\int_t^T K(s) ds} v(t), \quad t \in J.$$

Then problem (9) takes the form

$$\begin{cases} z'(t) = M(t)e^{\int_t^{\alpha(t)} K(s) ds} z(\alpha(t)) + \eta(t)e^{\int_t^T K(s) ds} \equiv \mathcal{F}z(t), & t \in J', \\ z(t_k^+) = (1 + L_k)z(t_k) + \gamma_k e^{\int_t^T K(s) ds} \equiv z(t_k) + \mathcal{P}_k z(t_k), & k = 1, 2, \dots, m, \\ z(T) = k_0. \end{cases} \quad (10)$$

Note that z is the solution of the following impulsive integral equation

$$z(t) = k_0 - \int_t^T \mathcal{F}z(s) ds - \sum_{i=k+1}^m \mathcal{P}_i z(t_i) \equiv \mathcal{A}z(t), \quad t \in J_k, \quad (11)$$

for $k = 0, 1, \dots, m$.

To find a solution of problem (11) is equivalent to get a fixed point of the operator $\mathcal{A}: PC(J) \rightarrow PC(J)$. Let $x, y \in PC(J)$. Then

$$\begin{aligned} \|\mathcal{A}x - \mathcal{A}y\| &= \sup_{t \in J} |\mathcal{A}x(t) - \mathcal{A}y(t)| \\ &\leq \sup_{t \in J} \left[\int_t^T |\mathcal{F}x(s) - \mathcal{F}y(s)| ds \right] + \sum_{i=1}^m |\mathcal{P}_i x(t_i) - \mathcal{P}_i y(t_i)| \\ &\leq \sup_{t \in J} \int_t^T M^*(s) |x(\alpha(s)) - y(\alpha(s))| ds + \sum_{i=1}^m L_i |x(t_i) - y(t_i)| \\ &= \delta \|x - y\|. \end{aligned}$$



Problem (11) has a unique solution, by the Banach fixed point theorem. It means that also problem (9) has a unique solution. This ends the proof. \square

Remark 2. If $M(t) = 0$ on J , then condition (8) is superfluous. Note that in this case system (10) takes the form

$$\begin{cases} z'(t) = \bar{\eta}(t), & t \in J', \\ z(t_k^+) = (1 + L_k)z(t_k) + \bar{\gamma}_k, & k = 1, 2, \dots, m, \\ x(T) = k_0 \end{cases} \quad (12)$$

with

$$\bar{\eta}(t) = \eta(t)e^{\int_t^T K(s) ds}, \quad \bar{\gamma}_k = \gamma_k e^{\int_{t_k}^T K(s) ds}.$$

It is easy to verify that the solution z of problem (12) has now the form

$$\begin{aligned} z(t) = z(0) \prod_{i=1}^k (1 + L_i) + \sum_{j=1}^k \left\{ \int_{t_{j-1}}^{t_j} \bar{\eta}(s) ds \prod_{i=j}^k (1 + L_i) + \bar{\gamma}_j \prod_{i=j+1}^k (1 + L_i) \right\} \\ + \int_{t_k}^t \bar{\eta}(s) ds, \quad t \in J_k, \text{ for } k = 0, 1, \dots, m, \end{aligned}$$

where

$$z(0) = \frac{1}{V} \left\{ k_0 - \sum_{j=1}^m \left[\int_{t_{j-1}}^{t_j} \bar{\eta}(s) ds \prod_{i=j}^m (1 + L_i) + \bar{\gamma}_j \prod_{i=j+1}^m (1 + L_i) \right] - \int_{t_m}^T \bar{\eta}(s) ds \right\}$$

with

$$V = \prod_{i=1}^m (1 + L_i).$$

Now we give sufficient conditions when problem (1) has a solution.

Theorem 2. Let assumption (H_1) hold. Moreover, assume that

(H_2) $y_0, z_0 \in PC^1(J)$ are lower and upper solutions of problem (1), respectively, and $z_0(t) \leq y_0(t)$ on J ,

(H_3) there exist functions $K, M \in C(J, \mathbb{R})$, M is nonnegative and such that

$$f(t, u, v) - f(t, \bar{u}, \bar{v}) \geq -K(t)(\bar{u} - u) - M(t)(\bar{v} - v)$$

for $z_0(t) \leq u \leq \bar{u} \leq y_0(t)$, $z_0(\alpha(t)) \leq v \leq \bar{v} \leq y_0(\alpha(t))$, $t \in J$,

(H_4) there exist constants $L_k \in [0, 1)$, $k = 1, 2, \dots, m$, such that

$$I_k(w(t_k)) - I_k(\bar{w}(t_k)) \geq -L_k[\bar{w}(t_k) - w(t_k)], \quad k = 1, 2, \dots, m,$$

for any w, \bar{w} with $z_0(t_k) \leq w(t_k) \leq \bar{w}(t_k) \leq y_0(t_k)$, $k = 1, 2, \dots, m$,

(H_5) conditions (4) and (8) hold,

(H₆) there exists $\gamma > 0$ such that for any $u, \bar{u} \in [z_0(0), y_0(0)]$ with $u \leq \bar{u}$ and $v, \bar{v} \in [z_0(T), y_0(T)]$ with $v \leq \bar{v}$ we have

$$g(u, v) \leq g(\bar{u}, v), \quad (13)$$

$$g(u, v) - g(u, \bar{v}) \leq \gamma(\bar{v} - v). \quad (14)$$

Then there exist solutions $v, w \in [z_0, y_0]_*$ of problem (1).

Proof. Some ideas are taken from paper [5]. Let $\eta, \xi \in [z_0, y_0]$, where

$$[z_0, y_0] = \{u \in PC(J, \mathbb{R}) : z_0(t) \leq u \leq y_0(t), t \in J\}.$$

Put $\varphi(t) = \sup[\eta(t), \xi(t)]$, $\Phi(t) = \inf[\eta(t), \xi(t)]$. Consider the initial value problems

$$\begin{cases} v'(t) = F\varphi(t) + K(t)[v(t) - \varphi(t)] + M(t)[v(\alpha(t)) - \varphi(\alpha(t))], & t \in J', \\ \Delta v(t_k) = I_k(\varphi(t_k)) + L_k[v(t_k) - \varphi(t_k)], & k = 1, 2, \dots, m, \\ v(T) = \varphi(T) + \frac{1}{\gamma}g(\varphi(0), \varphi(T)), \end{cases} \quad (15)$$

$$\begin{cases} w'(t) = F\Phi(t) + K(t)[w(t) - \Phi(t)] + M(t)[w(\alpha(t)) - \Phi(\alpha(t))], & t \in J', \\ \Delta w(t_k) = I_k(\Phi(t_k)) + L_k[w(t_k) - \Phi(t_k)], & k = 1, 2, \dots, m, \\ w(T) = \Phi(T) + \frac{1}{\gamma}g(\Phi(0), \Phi(T)). \end{cases} \quad (16)$$

By Theorem 1, problems (15), (16) have a unique solution. Therefore, we can define the operator

$$B : \bar{\Omega} \rightarrow PC(J) \times PC(J), \quad [z_0, y_0] \subset PC(J), \quad B(\eta, \xi) = (v, w), \quad (17)$$

where v, w are solutions of (15), (16), $\bar{\Omega} = [z_0, y_0] \times [z_0, y_0]$.

Now, we want to show that

$$z_0(t) \leq w(t) \leq v(t) \leq y_0(t), \quad t \in J. \quad (18)$$

Put $p = z_0 - w$. Then

$$\begin{aligned} p'(t) &\geq Fz_0(t) - F\Phi(t) - K(t)[w(t) - \Phi(t)] - M(t)[w(\alpha(t)) - \Phi(\alpha(t))] \\ &\geq -K(t)[\Phi(t) - z_0(t)] - M(t)[\Phi(\alpha(t)) - z_0(\alpha(t))] - K(t)[w(t) - \Phi(t)] \\ &\quad - M(t)[w(\alpha(t)) - \Phi(\alpha(t))] \\ &= K(t)p(t) + M(t)p(\alpha(t)). \end{aligned}$$

Moreover,

$$\begin{aligned} p(T) &= z_0(T) - \Phi(T) - \frac{1}{\gamma}g(\Phi(0), \Phi(T)) \\ &\leq z_0(T) - \Phi(T) - \frac{1}{\gamma}g(z_0(0), \Phi(T)) \\ &= z_0(T) - \Phi(T) + \frac{1}{\gamma}[g(z_0(0), z_0(T)) - g(z_0(0), \Phi(T))] - \frac{1}{\gamma}g(z_0(0), z_0(T)) \\ &\leq z_0(T) - \Phi(T) + \frac{1}{\gamma}\gamma[\Phi(T) - z_0(T)] = 0, \end{aligned}$$



and

$$\Delta p(t_k) \geq I_k(z_0(t_k)) - I_k(\Phi(t_k)) - L_k[w(t_k) - \Phi(t_k)] \geq L_k p(t_k), \quad k = 1, 2, \dots, m.$$

This and Lemma 1 show that $z_0(t) \leq w(t), t \in J$. Similarly we can show that $v(t) \leq y_0(t), t \in J$. To show that $w(t) \leq v(t), t \in J$, we put $p = w - v$. Then

$$\begin{aligned} p'(t) &= F\Phi(t) - F\varphi(t) + K(t)[w(t) - \Phi(t) - v(t) + \varphi(t)] \\ &\quad + M(t)[w(\alpha(t)) - \Phi(\alpha(t)) - v(\alpha(t)) + \varphi(\alpha(t))] \\ &\geq -K(t)[\varphi(t) - \Phi(t)] - M(t)[\varphi(\alpha(t)) - \Phi(\alpha(t))] \\ &\quad + K(t)[w(t) - \Phi(t) - v(t) + \varphi(t)] \\ &\quad + M(t)[w(\alpha(t)) - \Phi(\alpha(t)) - v(\alpha(t)) + \varphi(\alpha(t))] \\ &= K(t)p(t) + M(t)p(\alpha(t)). \end{aligned}$$

Moreover,

$$\begin{aligned} p(T) &= \Phi(T) - \varphi(T) + \frac{1}{\gamma} [g(\Phi(0), \Phi(T)) - g(\varphi(0), \varphi(T))] \\ &\leq \Phi(T) - \varphi(T) + \frac{1}{\gamma} \gamma [\varphi(T) - \Phi(T)] = 0, \end{aligned}$$

and, for $k = 1, 2, \dots, m$, we have

$$\Delta p(t_k) = I_k(\Phi(t_k)) - I_k(\varphi(t_k)) + L_k[w(t_k) - \Phi(t_k) - v(t_k) + \varphi(t_k)] \geq L_k p(t_k).$$

This and Lemma 1 show that $w(t) \leq v(t), t \in J$, so (18) holds.

Hence $B : \bar{\Omega} \rightarrow \bar{\Omega}$. In order to apply Schauder’s fixed point theorem we need to show that the operator B is continuous and compact. Let $(v_n, w_n) \in \bar{\Omega}$, and $v_n \rightarrow v, w_n \rightarrow w$ in $PC(J)$. Put

$$\begin{aligned} \mathcal{D}_0 v(t) &= F\varphi(t) + K(t)[v(t) - \varphi(t)] + M(t)[v(\alpha(t)) - \varphi(\alpha(t))], \\ \mathcal{D}_k v(t_k) &= I_k(\varphi(t_k)) + L_k[v(t_k) - \varphi(t_k)], \quad k = 1, 2, \dots, m, \\ \bar{k}_0 &= \varphi(T) + \frac{1}{\gamma} g(\varphi(0), \varphi(T)). \end{aligned}$$

Then problem (15) takes the form

$$\begin{cases} v'(t) = \mathcal{D}_0 v(t), & t \in J', \\ \Delta v(t_k) = \mathcal{D}_k v(t_k), & k = 1, 2, \dots, m, \\ v(T) = \bar{k}_0. \end{cases}$$

Similarly as in the proof of Theorem 1, v is the solution of the following impulsive integral equation

$$v(t) = \bar{k}_0 - \int_t^T \mathcal{D}_0 v(s) ds - \sum_{i=k+1}^m \mathcal{D}_i v(t_i) \equiv \mathcal{D}v(t), \quad t \in J_k, \quad k = 0, 1, \dots, m.$$

Then, for $t \in J$, we have

$$\begin{aligned}
|\mathcal{D}v_n(t) - \mathcal{D}v(t)| &\leq \int_t^T |\mathcal{D}_0v_n(s) - \mathcal{D}_0v(s)| ds + \sum_{i=1}^m |\mathcal{D}_i v_n(t_i) - \mathcal{D}_i v(t_i)| \\
&\leq \int_0^T [K(s)|v_n(s) - v(s)| + M(s)|v_n(\alpha(s)) - v(\alpha(s))|] ds \\
&\quad + \sum_{i=1}^m L_i |v_n(t_i) - v(t_i)|.
\end{aligned}$$

Thus the Lebesgue dominated convergence theorem implies

$$\sup_{t \in J} |\mathcal{D}v_n(t) - \mathcal{D}v(t)| \rightarrow 0 \quad \text{if } n \rightarrow \infty,$$

so operator \mathcal{D} is continuous. Similar property holds for $w_n \rightarrow w$ too. As a result $B: \bar{\Omega} \rightarrow \bar{\Omega}$ is continuous. In view of (18), the operator $B: \bar{\Omega} \rightarrow \bar{\Omega}$ is bounded too.

Now we need to show that the operator $B: \bar{\Omega} \rightarrow \bar{\Omega}$ is compact. Note that

$$|\mathcal{D}v(t_1) - \mathcal{D}v(t_2)| \leq \left| \int_{t_1}^{t_2} \mathcal{D}_0v(s) ds \right|.$$

Similar property also holds for the solution w . It proves that the operator $B: \bar{\Omega} \rightarrow \bar{\Omega}$ is equicontinuous on J . The Arzela–Ascoli theorem guarantees that B is compact. Hence, by Schauder's fixed point theorem, operator B has a fixed point, i.e. there exist $(v, w) \in \bar{\Omega}$ such that $B(v, w) = (v, w)$ and $v \leq w$.

Now, by (17), we see that v, w satisfy the following relations

$$\begin{cases} v'(t) = Fv(t) + K(t)[v(t) - v(t)] + M(t)[v(\alpha(t)) - v(\alpha(t))], & t \in J', \\ \Delta v(t_k) = I_k(v(t_k)) + L_k[v(t_k) - v(t_k)], & k = 1, 2, \dots, m, \\ v(T) = v(T) + \frac{1}{\gamma}g(v(0), v(T)), \end{cases}$$

$$\begin{cases} w'(t) = Fw(t) + K(t)[w(t) - w(t)] + M(t)[w(\alpha(t)) - w(\alpha(t))], & t \in J', \\ \Delta w(t_k) = I_k(w(t_k)) + L_k[w(t_k) - w(t_k)], & k = 1, 2, \dots, m, \\ w(T) = w(T) + \frac{1}{\gamma}g(w(0), w(T)). \end{cases}$$

It shows that $v, w \in PC^1(J)$ are solutions of problem (1). This ends the proof. \square

Example 1. For $J = [0, T]$, we consider the problem

$$\begin{cases} x'(t) = \lambda_1(t)e^{x(t)} + \lambda_2(t) \sin(x(\alpha(t))) - \lambda_1(t), & t \in J \setminus \{t_1\}, \\ \Delta x(t_1) = Lx(t_1), \\ 0 = 2x^2(0) + x(T) - k, \end{cases} \quad (19)$$

where $\lambda_1, \lambda_2 \in C(J, \mathbb{R}_+)$, $\mathbb{R}_+ = [0, \infty)$, $\alpha \in C(J, J)$, $t \leq \alpha(t) \leq T$, $t \in J$, $0 < t_1 < T$, $L \geq 0$, $0 \leq k \leq 1$.

Take $y_0(t) = 0$, $z_0(t) = -1$, $t \in J$. Indeed, $z_0(t) < y_0(t)$ on J , and



$$\begin{aligned}
 Fy_0(t) &= \lambda_1(t) - \lambda_1(t) = 0 = y_0'(t), \\
 Fz_0(t) &= \lambda_1(t)[e^{-1} - 1] - \lambda_2(t) \sin 1 \leq 0 = z_0'(t), \\
 \Delta y_0(t_1) &= L \cdot 0 = I_1(y_0(t_1)), \\
 \Delta z_0(t_1) &= 0 \geq L(-1) = I_1(z_0(t_1)), \\
 g(y_0(0), y_0(T)) &= g(0, 0) = -k \leq 0, \\
 g(z_0(0), z_0(T)) &= g(-1, -1) = 1 - k \geq 0.
 \end{aligned}$$

It proves that y_0, z_0 are lower and upper solutions of problem (19), respectively. Moreover, $K(t) = \lambda_1(t)$, $M(t) = \lambda_2(t)$, $L_1 = L$, so assumptions (H₃), (H₄), (H₆) are satisfied. If we extra assume that

$$\int_0^T \lambda_2(t) e^{\int_t^T \lambda_1(s) ds} dt + L < 1, \tag{20}$$

then problem (19) has solutions in the segment $[-1, 0]_*$, by Theorem 2. Note that condition (20) guaranties that condition (4) is satisfied too.

For example, if we take $L = \frac{1}{2}$, $T = \pi$, $\lambda_1(t) = \lambda > 0$, $\lambda_2(t) = \beta e^{\lambda(t-T)} \sin t$ for $t \in J$, then condition (20) holds if $0 < \beta < \frac{1}{4}$.

3. Coupled lower and upper solutions of problem (1)

Let us introduce the following definition.

We say that $u, w \in PC^1(J, \mathbb{R})$ are coupled lower and upper solutions of (1) if

$$\begin{cases}
 u'(t) \leq Fu(t), & t \in J', \\
 \Delta u(t_k) \leq I_k(u(t_k)), & k = 1, 2, \dots, m, \\
 g(u(0), w(T)) \leq 0, \\
 \\
 w'(t) \geq Fw(t), & t \in J', \\
 \Delta w(t_k) \geq I_k(w(t_k)), & k = 1, 2, \dots, m, \\
 g(w(0), u(T)) \geq 0.
 \end{cases}$$

The next result deals with the case when problem (1) has quasisolutions.

Theorem 3. Assume that assumptions (H₁), (H₃)–(H₅) hold, where y_0, z_0 are coupled lower and upper solutions of problem (1) and $z_0(t) \leq y_0(t)$ on J . In addition, we assume that

(H'₆) there exists $\gamma > 0$ such that for any $u, \bar{u} \in [z_0(0), y_0(0)]$ with $u \leq \bar{u}$ and $v, \bar{v} \in [z_0(T), y_0(T)]$ with $v \leq \bar{v}$ we have

$$\begin{aligned}
 g(u, v) &\leq g(\bar{u}, v), \\
 g(u, v) - g(u, \bar{v}) &\geq -\gamma(\bar{v} - v).
 \end{aligned}$$

Then there exist $u, v \in [z_0, y_0]_*$ coupled quasisolutions of problem (1), i.e. the pair (v, w) is a solution of the system:

$$\left\{ \begin{array}{l} \left\{ \begin{array}{l} v'(t) = Fv(t), \quad t \in J', \\ \Delta v(t_k) = I_k(v(t_k)), \quad k = 1, 2, \dots, m, \\ g(v(0), w(T)) = 0, \end{array} \right. \\ \left\{ \begin{array}{l} w'(t) = Fw(t), \quad t \in J', \\ \Delta w(t_k) = I_k(w(t_k)), \quad k = 1, 2, \dots, m, \\ g(w(0), v(T)) = 0. \end{array} \right. \end{array} \right.$$

Proof. Consider the initial value problems

$$\left\{ \begin{array}{l} \left\{ \begin{array}{l} v'(t) = F\varphi(t) + K(t)[v(t) - \varphi(t)] + M(t)[v(\alpha(t)) - \varphi(\alpha(t))], \quad t \in J', \\ \Delta v(t_k) = I_k(\varphi(t_k)) + L_k[v(t_k) - \varphi(t_k)], \quad k = 1, 2, \dots, m, \\ v(T) = \varphi(T) - \frac{1}{\gamma}g(\varphi(0), \varphi(T)), \end{array} \right. \\ \left\{ \begin{array}{l} w'(t) = F\Phi(t) + K(t)[w(t) - \Phi(t)] + M(t)[w(\alpha(t)) - \Phi(\alpha(t))], \quad t \in J', \\ \Delta w(t_k) = I_k(\Phi(t_k)) + L_k[w(t_k) - \Phi(t_k)], \quad k = 1, 2, \dots, m, \\ w(T) = \Phi(T) - \frac{1}{\gamma}g(\varphi(0), \Phi(T)), \end{array} \right. \end{array} \right.$$

where Φ and φ are defined as in the proof of Theorem 2. The proof is similar to the proof of Theorem 2 and therefore it is omitted. \square

Example 2. Now we consider the problem

$$\left\{ \begin{array}{l} x'(t) = bx(t) - bx^2(\alpha(t)) + (a-1)(b-5) \equiv Fx(t), \quad t \in J = [0, T], \\ \Delta x(t_i) = L_i x(t_i), \quad i = 1, 2, \dots, m, \text{ with } 0 < t_1 < t_2 < \dots < t_m < T, \\ 0 = \lambda[x(0) + x^2(0)] - x(T) - a, \end{array} \right. \quad (21)$$

where $a > 1$, $b \geq 5$, $L_i \geq 0$, $i = 1, 2, \dots, m$, $\lambda > 0$, $\alpha \in C(J, J)$, $t \leq \alpha(t) \leq T$ and

$$a[\lambda(a-1) - 1] \geq 0. \quad (22)$$

In addition, we assume that

$$\sum_{i=1}^m L_i < 1. \quad (23)$$

Put $y_0(t) = 0$, $z_0(t) = -a$, $t \in J$. Then

$$\begin{aligned} Fy_0(t) &= (a-1)(b-5) \geq 0 = y_0'(t), \\ Fz_0(t) &= -ab - ba^2 + (a-1)(b-5) < 0 = z_0'(t), \\ \Delta y_0(t_i) &= 0 = 0L_i = I_i(y_0(t_i)), \quad i = 1, 2, \dots, m, \\ \Delta z_0(t_i) &= 0 \geq -L_i a = I_i(z_0(t_i)), \quad i = 1, 2, \dots, m, \\ g(y_0(0), z_0(T)) &= g(0, -a) = a - a = 0, \\ g(z_0(0), y_0(T)) &= g(-a, 0) = a[\lambda(a-1) - 1] \geq 0, \end{aligned}$$

by (22). It shows that y_0, z_0 are weakly coupled lower and upper solutions of (21). Note that $K(t) = b$, $M(t) = 0$, $t \in J$, so assumption (H_3) holds. Assumptions (H_4) , (H_5) , (H'_6) are also satisfied. By Theorem 3, problem (21) has, in the sector $[z_0, y_0]_*$, coupled quasisolutions.



References

- [1] R.P. Agarwal, D. Franco, D. O'Regan, Singular boundary value problems for first and second order impulsive differential equations, *Aequationes Math.* 69 (2005) 83–96.
- [2] W. Ding, M. Han, J. Mi, Periodic boundary value problem for second-order impulsive functional differential equations, *Comput. Math. Appl.* 50 (2005) 491–507.
- [3] W. Ding, J. Mi, M. Han, Periodic boundary value problems for the first order impulsive functional differential equations, *Appl. Math. Comput.* 165 (2005) 433–446.
- [4] D. Franco, J.J. Nieto, First-order impulsive ordinary differential equations with anti-periodic and nonlinear boundary conditions, *Nonlinear Anal.* 42 (2000) 163–173.
- [5] D. Franco, J.J. Nieto, D. O'Regan, Existence of solutions for first order ordinary differential equations with nonlinear boundary conditions, *Appl. Math. Comput.* 153 (2004) 793–802.
- [6] Z. He, J. Yu, Periodic boundary value problem for first-order impulsive functional differential equations, *J. Comput. Appl. Math.* 138 (2002) 205–217.
- [7] T. Jankowski, On delay differential equations with nonlinear boundary conditions, *Bound. Value Probl.* 2005 (2005) 201–214.
- [8] T. Jankowski, Advanced differential equations with nonlinear boundary conditions, *J. Math. Anal. Appl.* 304 (2005) 490–503.
- [9] T. Jankowski, Solvability of three point boundary value problems for second order differential equations with deviating arguments, *J. Math. Anal. Appl.* 312 (2005) 620–636.
- [10] T. Jankowski, Boundary value problems for first order differential equations of mixed type, *Nonlinear Anal.* 64 (2006) 1984–1997.
- [11] G.S. Ladde, V. Lakshmikantham, A.S. Vatsala, *Monotone Iterative Techniques for Nonlinear Differential Equations*, Pitman, Boston, 1985.
- [12] V. Lakshmikantham, D.D. Bainov, P.S. Simeonov, *Theory of Impulsive Ordinary Differential Equations*, World Scientific, Singapore, 1989.
- [13] A.M. Samoilenko, N.A. Perestyuk, *Impulsive Differential Equations*, World Scientific, Singapore, 1995.
- [14] F. Zhang, Z. Ma, J. Yan, Boundary value problems for first order impulsive delay differential equations with a parameter, *J. Math. Anal. Appl.* 290 (2004) 213–223.

