

2,2,6,6-Tetramethylpiperidinium triisopropoxysilanethiolate

Katarzyna Baranowska,* Paweł Roman and Justyna Socha

Department of Inorganic Chemistry, Faculty of Chemistry, Gdańsk University of Technology, 11/12 G. Narutowicz St., 80233 - PL Gdańsk, Poland
Correspondence e-mail: kasiab29@wp.pl


Received 3 September 2009; accepted 19 October 2009

Key indicators: single-crystal X-ray study; $T = 120\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.049; wR factor = 0.122; data-to-parameter ratio = 17.9.

The crystal of the title compound, $\text{C}_9\text{H}_{20}\text{N}^+\cdot\text{C}_9\text{H}_{21}\text{O}_3\text{SSi}^-$, is built of aggregates, each made up of two 2,2,6,6-tetramethylpiperidinium cations and two triisopropoxysilanethiolate anions. The aggregates are linked by four $\text{N}-\text{H}\cdots\text{S}$ bonds and correspond to an $R_4^2(8)$ graph-set motif.

Related literature

For the structures of similar compounds and comparison of bond distances, see Baranowska, Chojnacki, Gosiewska & Wojnowski (2006); Baranowska, Chojnacki, Konitz *et al.* (2006); Baranowska & Piwowarska (2008); Becker *et al.* (2004). For the graph-set description of hydrogen-bonding patterns, see Bernstein *et al.* (1995); Etter (1990).



Experimental

Crystal data

$\text{C}_9\text{H}_{20}\text{N}^+\cdot\text{C}_9\text{H}_{21}\text{O}_3\text{SSi}^-$
 $M_r = 379.67$
Triclinic, $P\bar{1}$
 $a = 9.2433 (6)\text{ \AA}$
 $b = 11.5545 (8)\text{ \AA}$
 $c = 11.7593 (7)\text{ \AA}$
 $\alpha = 85.955 (5)^\circ$
 $\beta = 77.190 (6)^\circ$

$\gamma = 67.620 (6)^\circ$
 $V = 1132.28 (13)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.21\text{ mm}^{-1}$
 $T = 120\text{ K}$
 $0.42 \times 0.39 \times 0.33\text{ mm}$

Data collection

Oxford Diffraction KM4/Xcalibur diffractometer with Sapphire2 detector
Absorption correction: analytical (*CrysAlis RED*; Oxford)
Diffraction, 2006)
 $T_{\min} = 0.908$, $T_{\max} = 0.934$
6937 measured reflections
4209 independent reflections
3561 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.122$
 $S = 1.10$
4209 reflections
235 parameters
1 restraint
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.45\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.43\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A \cdots S1	0.966 (17)	2.352 (17)	3.3166 (14)	177 (2)
N1—H1B \cdots S1 ⁱ	0.927 (17)	2.338 (17)	3.2354 (14)	163.0 (17)

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2006); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2006); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* publication routines (Farrugia, 1999).

The authors thank Dr Agnieszka Pladzyk and Dr Jarosław Chojnacki for helpful discussions during the preparation of the manuscript.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YA2106).

References

- Baranowska, K., Chojnacki, J., Gosiewska, M. & Wojnowski, W. (2006). *Z. Anorg. Allg. Chem.* **632**, 1086–1090.
- Baranowska, K., Chojnacki, J., Konitz, A., Wojnowski, W. & Becker, B. (2006). *Polyhedron*, **25**, 1555–1560.
- Baranowska, K. & Piwowarska, N. (2008). *Acta Cryst. E* **64**, o1781.
- Becker, B., Baranowska, K., Chojnacki, J. & Wojnowski, W. (2004). *Chem. Commun.*, pp. 620–621.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Oxford Diffraction (2006). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2009). E65, o2825 [https://doi.org/10.1107/S1600536809042962]

2,2,6,6-Tetramethylpiperidinium triisopropoxysilanethiolate

Katarzyna Baranowska, Paweł Roman and Justyna Socha

S1. Comment

Hydrogen bonding is arguably the most prominent interaction in selfassembly of molecules in crystals and plays an important role in determining structure of chemical and biological systems. Hydrogen bonds of the $(^+)N\cdots H-S^-$ -type have gained relatively little attention, as these bonds are mostly weak and quite seldom lead to the proton transfer. One notable exception are silanethiolates, where ionization of the SH group is facilitated by the neighbouring silicon atom. The salts of these anions with primary amines as counter-ions often feature tetrameric aggregates with a cubane-like hydrogen bonded core (Becker *et al.*, 2004). Secondary amines give derivatives with discrete dimeric units in the solid state (Baranowska, Chojnacki, Konitz *et al.* 2006).

We present here the crystal structure of the title compound, which was obtained by the reaction of tri-*iso*-propoxysilane-thiol with 2,2,6,6-tetramethylpiperidine.


The crystal structure is built of aggregates, made up of two tri-*iso*-propoxysilanethiolate anions and two piperidinium cations (Fig. 1). The aggregates contain eight-membered ring built due to the formation of four charge-assisted $(^+)N-H\cdots S^-$ -hydrogen bonds (graph theory motif $R^2_4(8)$ according to Etter, 1990; Bernstein *et al.*, 1995). The N \cdots S distances (Table 1) lie in the range comparable with the values observed in aromatic thiolates (Baranowska & Piwowarska, 2008) or silanethiolates (Baranowska, Chojnacki, Gosiewska, Wojnowski, 2006).

S2. Experimental

Tri-*iso*-propoxysilanethiol (2 mmol) was dissolved in 8 ml of propanol-2 and 2,2,6,6-tetramethylpiperidine (0.338 ml, 2 mmol) was added. The solvent was then added gradually until a white deposit formed was completely dissolved. The solution was left to stand at 4 °C for a few days for crystallization. The obtained colourless crystals were suitable for X-ray diffraction analysis. The product is hygroscopic and slowly oxidizes in the air, therefore all operations were carried out using a vacuum-nitrogen line and Schlenk techniques.

S3. Refinement

Hydrogen atoms were placed in geometrically calculated positions (C—H 0.98 Å for methyl, 0.99 Å for methylene and 1.00 Å for methine H atoms) and refined as riding on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for methylene and methine and $1.5U_{\text{eq}}(\text{C})$ for methyl groups. Hydrogen atoms of ammonium group were found in the difference map and refined in isotropic approximation constrained to produce N—H bonds equal within 0.04 Å (SADI instruction of *SHELXL97*; Sheldrick, 2008)

**Figure 1**

Dimeric aggregates $[\text{C}_9\text{H}_{20}\text{N}^+\text{C}_9\text{H}_{21}\text{O}_3\text{SiS}^-]_2$ in the structure of the title compound. The unlabeled atoms are derived from the reference atoms by the $(1 - x, 1 - y, 1 - z)$ symmetry transformation. Displacement ellipsoids are drawn at the 50% probability level. C-bound H atoms have been omitted for clarity.

2,2,6,6-Tetramethylpiperidinium triisopropoxysilanethiolate

Crystal data


 $M_r = 379.67$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 9.2433 (6) \text{ \AA}$
 $b = 11.5545 (8) \text{ \AA}$
 $c = 11.7593 (7) \text{ \AA}$
 $\alpha = 85.955 (5)^\circ$
 $\beta = 77.190 (6)^\circ$
 $\gamma = 67.620 (6)^\circ$
 $V = 1132.28 (13) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 420$
 $D_x = 1.114 \text{ Mg m}^{-3}$
 $\text{Mo } K\alpha \text{ radiation, } \lambda = 0.71073 \text{ \AA}$

Cell parameters from 5191 reflections

 $\theta = 2.4\text{--}28.8^\circ$
 $\mu = 0.21 \text{ mm}^{-1}$
 $T = 120 \text{ K}$

Prism, colourless

 $0.42 \times 0.39 \times 0.33 \text{ mm}$

Data collection

Oxford Diffraction KM4/Xcalibur
diffractometer with Sapphire2 detector

6937 measured reflections

4209 independent reflections

Graphite monochromator

3561 reflections with $I > 2\sigma(I)$

Detector resolution: 8.1883 pixels mm^{-1}
 $R_{\text{int}} = 0.015$
 ω scans

 $\theta_{\text{max}} = 25.5^\circ, \theta_{\text{min}} = 2.4^\circ$

Absorption correction: analytical

 $h = -10 \rightarrow 11$

(*CrysAlis RED*; Oxford Diffraction, 2006)

 $k = -13 \rightarrow 13$
 $T_{\text{min}} = 0.908, T_{\text{max}} = 0.934$
 $l = -7 \rightarrow 14$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.049$$

$$wR(F^2) = 0.122$$

$$S = 1.10$$

4209 reflections

235 parameters

1 restraint

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0855P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.45 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.43 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^* / U_{\text{eq}}$
C1	0.7770 (2)	0.42248 (16)	-0.00766 (13)	0.0205 (4)
H1	0.7182	0.3662	-0.0104	0.025*
C2	0.6713 (2)	0.55565 (17)	-0.02787 (15)	0.0272 (4)
H2A	0.5712	0.5799	0.031	0.041*
H2B	0.6476	0.5616	-0.1058	0.041*
H2C	0.7266	0.6117	-0.022	0.041*
C3	0.9338 (2)	0.37720 (18)	-0.09727 (15)	0.0302 (4)
H3A	0.9935	0.4305	-0.0932	0.045*
H3B	0.9115	0.3814	-0.1755	0.045*
H3C	0.9975	0.2905	-0.0809	0.045*
C4	1.0748 (2)	0.28839 (16)	0.23790 (15)	0.0236 (4)
H4	1.0715	0.3583	0.182	0.028*
C5	1.1457 (3)	0.3044 (3)	0.33692 (19)	0.0506 (6)
H5A	1.0774	0.3838	0.3781	0.076*
H5B	1.253	0.3049	0.3057	0.076*
H5C	1.1527	0.235	0.3912	0.076*
C6	1.1706 (3)	0.1663 (2)	0.1731 (2)	0.0481 (6)
H6A	1.1779	0.0969	0.2274	0.072*
H6B	1.2783	0.1635	0.1377	0.072*
H6C	1.1179	0.1587	0.1119	0.072*
C7	0.7501 (2)	0.09780 (16)	0.18044 (15)	0.0246 (4)
H7	0.6694	0.1273	0.255	0.03*
C8	0.6687 (3)	0.0770 (2)	0.0905 (2)	0.0460 (6)
H8A	0.5784	0.1542	0.0818	0.069*

H8B	0.6295	0.0096	0.1156	0.069*
H8C	0.7452	0.0538	0.0156	0.069*
C9	0.8863 (3)	-0.01829 (18)	0.2035 (2)	0.0416 (5)
H9A	0.9654	-0.0482	0.1306	0.062*
H9B	0.8453	-0.0834	0.2343	0.062*
H9C	0.9369	0.0013	0.2605	0.062*
C10	0.3378 (2)	0.77040 (17)	0.24261 (14)	0.0256 (4)
H10A	0.3714	0.6889	0.2026	0.031*
H10B	0.2393	0.828	0.2184	0.031*
C11	0.4694 (2)	0.82307 (17)	0.20331 (15)	0.0280 (4)
H11A	0.4912	0.8302	0.1173	0.034*
H11B	0.4333	0.9078	0.2376	0.034*
C12	0.6221 (2)	0.73804 (16)	0.24148 (14)	0.0233 (4)
H12A	0.705	0.7745	0.216	0.028*
H12B	0.6617	0.6555	0.202	0.028*
C13	0.5987 (2)	0.71909 (15)	0.37349 (14)	0.0197 (4)
C14	0.5719 (2)	0.83674 (16)	0.44152 (16)	0.0290 (4)
H14A	0.5284	0.8278	0.5242	0.044*
H14B	0.496	0.9102	0.4105	0.044*
H14C	0.6739	0.8473	0.4333	0.044*
C15	0.7431 (2)	0.61143 (15)	0.40297 (15)	0.0234 (4)
H15A	0.8393	0.6312	0.3766	0.035*
H15B	0.7574	0.5346	0.3637	0.035*
H15C	0.7255	0.5993	0.4875	0.035*
C16	0.2995 (2)	0.75210 (16)	0.37445 (14)	0.0218 (4)
C17	0.2107 (2)	0.87568 (17)	0.44395 (16)	0.0319 (4)
H17A	0.2099	0.8596	0.527	0.048*
H17B	0.1004	0.9126	0.4327	0.048*
H17C	0.265	0.9337	0.4166	0.048*
C18	0.1976 (2)	0.67189 (17)	0.40504 (15)	0.0255 (4)
H18A	0.2555	0.5901	0.3656	0.038*
H18B	0.0963	0.7139	0.3795	0.038*
H18C	0.1757	0.6603	0.4896	0.038*
O1	0.81444 (14)	0.41560 (10)	0.10576 (9)	0.0198 (3)
O2	0.91262 (14)	0.29862 (11)	0.28713 (10)	0.0212 (3)
O3	0.81113 (14)	0.19313 (10)	0.13885 (10)	0.0233 (3)
Si1	0.77654 (5)	0.32207 (4)	0.20952 (4)	0.01653 (15)
S1	0.55226 (5)	0.38899 (4)	0.31590 (3)	0.01896 (14)
N1	0.45724 (16)	0.67745 (12)	0.41129 (12)	0.0182 (3)
H1A	0.489 (3)	0.5932 (17)	0.3823 (19)	0.046 (6)*
H1B	0.437 (2)	0.6748 (18)	0.4920 (15)	0.033 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0237 (9)	0.0253 (9)	0.0159 (8)	-0.0128 (8)	-0.0048 (7)	0.0011 (6)
C2	0.0229 (9)	0.0336 (10)	0.0226 (9)	-0.0078 (8)	-0.0055 (7)	0.0023 (7)
C3	0.0304 (11)	0.0329 (10)	0.0224 (9)	-0.0099 (9)	0.0024 (8)	-0.0050 (7)

C4	0.0152 (9)	0.0275 (9)	0.0286 (9)	-0.0100 (8)	-0.0036 (7)	0.0046 (7)
C5	0.0249 (11)	0.0914 (19)	0.0412 (13)	-0.0269 (13)	-0.0081 (9)	-0.0038 (12)
C6	0.0200 (10)	0.0409 (12)	0.0749 (17)	-0.0051 (10)	-0.0006 (11)	-0.0118 (11)
C7	0.0254 (10)	0.0226 (9)	0.0265 (9)	-0.0123 (8)	0.0003 (7)	-0.0025 (7)
C8	0.0505 (15)	0.0505 (13)	0.0541 (14)	-0.0334 (12)	-0.0207 (12)	0.0073 (11)
C9	0.0400 (13)	0.0258 (10)	0.0625 (14)	-0.0147 (10)	-0.0153 (11)	0.0079 (9)
C10	0.0215 (9)	0.0296 (9)	0.0209 (9)	-0.0043 (8)	-0.0050 (7)	0.0022 (7)
C11	0.0295 (10)	0.0269 (9)	0.0220 (9)	-0.0068 (8)	-0.0033 (8)	0.0055 (7)
C12	0.0238 (9)	0.0233 (9)	0.0223 (9)	-0.0107 (8)	-0.0007 (7)	0.0015 (7)
C13	0.0200 (9)	0.0202 (8)	0.0199 (8)	-0.0094 (7)	-0.0023 (7)	-0.0015 (6)
C14	0.0344 (11)	0.0262 (9)	0.0288 (10)	-0.0140 (9)	-0.0044 (8)	-0.0067 (7)
C15	0.0199 (9)	0.0249 (9)	0.0272 (9)	-0.0096 (8)	-0.0059 (7)	-0.0013 (7)
C16	0.0168 (8)	0.0223 (9)	0.0205 (8)	-0.0010 (7)	-0.0036 (7)	-0.0007 (7)
C17	0.0279 (10)	0.0267 (10)	0.0301 (10)	0.0005 (8)	-0.0022 (8)	-0.0041 (8)
C18	0.0171 (9)	0.0312 (10)	0.0252 (9)	-0.0061 (8)	-0.0038 (7)	-0.0003 (7)
O1	0.0239 (6)	0.0233 (6)	0.0161 (6)	-0.0124 (5)	-0.0057 (5)	0.0009 (5)
O2	0.0149 (6)	0.0279 (6)	0.0208 (6)	-0.0084 (5)	-0.0036 (5)	0.0028 (5)
O3	0.0258 (7)	0.0200 (6)	0.0230 (6)	-0.0114 (5)	0.0034 (5)	-0.0040 (5)
Si1	0.0155 (3)	0.0168 (2)	0.0167 (2)	-0.00606 (19)	-0.00172 (18)	-0.00110 (17)
S1	0.0151 (2)	0.0219 (2)	0.0187 (2)	-0.00605 (18)	-0.00163 (16)	-0.00349 (16)
N1	0.0169 (7)	0.0187 (7)	0.0178 (7)	-0.0053 (6)	-0.0033 (6)	-0.0006 (6)

Geometric parameters (\AA , $^{\circ}$)

C1—O1	1.4399 (18)	C10—H10A	0.99
C1—C2	1.512 (2)	C10—H10B	0.99
C1—C3	1.519 (2)	C11—C12	1.523 (2)
C1—H1	1.00	C11—H11A	0.99
C2—H2A	0.98	C11—H11B	0.99
C2—H2B	0.98	C12—C13	1.531 (2)
C2—H2C	0.98	C12—H12A	0.99
C3—H3A	0.98	C12—H12B	0.99
C3—H3B	0.98	C13—C15	1.526 (2)
C3—H3C	0.98	C13—N1	1.527 (2)
C4—O2	1.444 (2)	C13—C14	1.536 (2)
C4—C6	1.502 (3)	C14—H14A	0.98
C4—C5	1.510 (2)	C14—H14B	0.98
C4—H4	1.00	C14—H14C	0.98
C5—H5A	0.98	C15—H15A	0.98
C5—H5B	0.98	C15—H15B	0.98
C5—H5C	0.98	C15—H15C	0.98
C6—H6A	0.98	C16—N1	1.525 (2)
C6—H6B	0.98	C16—C18	1.531 (2)
C6—H6C	0.98	C16—C17	1.535 (2)
C7—O3	1.430 (2)	C17—H17A	0.98
C7—C8	1.506 (3)	C17—H17B	0.98
C7—C9	1.507 (3)	C17—H17C	0.98
C7—H7	1.00	C18—H18A	0.98

C8—H8A	0.98	C18—H18B	0.98
C8—H8B	0.98	C18—H18C	0.98
C8—H8C	0.98	O1—Si1	1.6408 (11)
C9—H9A	0.98	O2—Si1	1.6441 (11)
C9—H9B	0.98	O3—Si1	1.6452 (11)
C9—H9C	0.98	Si1—S1	2.0558 (6)
C10—C11	1.530 (3)	N1—H1A	0.966 (17)
C10—C16	1.530 (2)	N1—H1B	0.927 (17)
O1—C1—C2	108.93 (13)	C10—C11—H11A	109.6
O1—C1—C3	107.87 (14)	C12—C11—H11B	109.6
C2—C1—C3	112.55 (14)	C10—C11—H11B	109.6
O1—C1—H1	109.1	H11A—C11—H11B	108.1
C2—C1—H1	109.1	C11—C12—C13	113.24 (15)
C3—C1—H1	109.1	C11—C12—H12A	108.9
C1—C2—H2A	109.5	C13—C12—H12A	108.9
C1—C2—H2B	109.5	C11—C12—H12B	108.9
H2A—C2—H2B	109.5	C13—C12—H12B	108.9
C1—C2—H2C	109.5	H12A—C12—H12B	107.7
H2A—C2—H2C	109.5	C15—C13—N1	105.89 (13)
H2B—C2—H2C	109.5	C15—C13—C12	110.41 (14)
C1—C3—H3A	109.5	N1—C13—C12	107.30 (12)
C1—C3—H3B	109.5	C15—C13—C14	108.82 (13)
H3A—C3—H3B	109.5	N1—C13—C14	111.15 (14)
C1—C3—H3C	109.5	C12—C13—C14	113.03 (14)
H3A—C3—H3C	109.5	C13—C14—H14A	109.5
H3B—C3—H3C	109.5	C13—C14—H14B	109.5
O2—C4—C6	111.26 (14)	H14A—C14—H14B	109.5
O2—C4—C5	107.29 (15)	C13—C14—H14C	109.5
C6—C4—C5	112.44 (17)	H14A—C14—H14C	109.5
O2—C4—H4	108.6	H14B—C14—H14C	109.5
C6—C4—H4	108.6	C13—C15—H15A	109.5
C5—C4—H4	108.6	C13—C15—H15B	109.5
C4—C5—H5A	109.5	H15A—C15—H15B	109.5
C4—C5—H5B	109.5	C13—C15—H15C	109.5
H5A—C5—H5B	109.5	H15A—C15—H15C	109.5
C4—C5—H5C	109.5	H15B—C15—H15C	109.5
H5A—C5—H5C	109.5	N1—C16—C10	107.55 (13)
H5B—C5—H5C	109.5	N1—C16—C18	106.11 (13)
C4—C6—H6A	109.5	C10—C16—C18	110.72 (14)
C4—C6—H6B	109.5	N1—C16—C17	110.86 (13)
H6A—C6—H6B	109.5	C10—C16—C17	113.25 (14)
C4—C6—H6C	109.5	C18—C16—C17	108.14 (15)
H6A—C6—H6C	109.5	C16—C17—H17A	109.5
H6B—C6—H6C	109.5	C16—C17—H17B	109.5
O3—C7—C8	107.94 (15)	H17A—C17—H17B	109.5
O3—C7—C9	108.78 (15)	C16—C17—H17C	109.5
C8—C7—C9	113.02 (17)	H17A—C17—H17C	109.5

O3—C7—H7	109	H17B—C17—H17C	109.5
C8—C7—H7	109	C16—C18—H18A	109.5
C9—C7—H7	109	C16—C18—H18B	109.5
C7—C8—H8A	109.5	H18A—C18—H18B	109.5
C7—C8—H8B	109.5	C16—C18—H18C	109.5
H8A—C8—H8B	109.5	H18A—C18—H18C	109.5
C7—C8—H8C	109.5	H18B—C18—H18C	109.5
H8A—C8—H8C	109.5	C1—O1—Si1	124.60 (10)
H8B—C8—H8C	109.5	C4—O2—Si1	123.87 (10)
C7—C9—H9A	109.5	C7—O3—Si1	126.27 (11)
C7—C9—H9B	109.5	O1—Si1—O2	104.18 (6)
H9A—C9—H9B	109.5	O1—Si1—O3	103.66 (6)
C7—C9—H9C	109.5	O2—Si1—O3	110.48 (6)
H9A—C9—H9C	109.5	O1—Si1—S1	116.03 (5)
H9B—C9—H9C	109.5	O2—Si1—S1	109.64 (5)
C11—C10—C16	113.34 (14)	O3—Si1—S1	112.43 (5)
C11—C10—H10A	108.9	C16—N1—C13	119.89 (13)
C16—C10—H10A	108.9	C16—N1—H1A	105.6 (13)
C11—C10—H10B	108.9	C13—N1—H1A	108.5 (14)
C16—C10—H10B	108.9	C16—N1—H1B	108.0 (13)
H10A—C10—H10B	107.7	C13—N1—H1B	106.7 (13)
C12—C11—C10	110.38 (14)	H1A—N1—H1B	107.6 (18)
C12—C11—H11A	109.6		
C16—C10—C11—C12	57.4 (2)	C1—O1—Si1—O3	37.28 (14)
C10—C11—C12—C13	-57.91 (19)	C1—O1—Si1—S1	-86.47 (13)
C11—C12—C13—C15	167.08 (14)	C4—O2—Si1—O1	-35.98 (13)
C11—C12—C13—N1	52.13 (18)	C4—O2—Si1—O3	74.77 (13)
C11—C12—C13—C14	-70.76 (19)	C4—O2—Si1—S1	-160.78 (11)
C11—C10—C16—N1	-51.19 (19)	C7—O3—Si1—O1	-158.72 (12)
C11—C10—C16—C18	-166.70 (14)	C7—O3—Si1—O2	90.20 (13)
C11—C10—C16—C17	71.65 (19)	C7—O3—Si1—S1	-32.64 (14)
C2—C1—O1—Si1	123.01 (13)	C10—C16—N1—C13	50.41 (18)
C3—C1—O1—Si1	-114.55 (14)	C18—C16—N1—C13	168.93 (13)
C6—C4—O2—Si1	-72.47 (18)	C17—C16—N1—C13	-73.89 (18)
C5—C4—O2—Si1	164.15 (14)	C15—C13—N1—C16	-168.72 (13)
C8—C7—O3—Si1	126.51 (15)	C12—C13—N1—C16	-50.79 (18)
C9—C7—O3—Si1	-110.51 (16)	C14—C13—N1—C16	73.25 (17)
C1—O1—Si1—O2	152.92 (12)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···Si1	0.97 (2)	2.35 (2)	3.3166 (14)	177 (2)
N1—H1B···Si1 ⁱ	0.93 (2)	2.34 (2)	3.2354 (14)	163 (2)

Symmetry code: (i) $-x+1, -y+1, -z+1$.