

## Genetic Features of Clinical *Pseudomonas aeruginosa* Strains

KATARZYNA WOLSKA<sup>1\*</sup> and PIOTR SZWEDA<sup>2</sup>

<sup>1</sup>Department of Microbiology, Natural Faculty, University of Podlasie, Siedlce, Poland

<sup>2</sup>Department of Food Chemistry, Technology and Biotechnology, Chemistry Faculty,  
Gdansk University of Technology, Gdańsk, Poland

Received 18 February 2009, revised 20 July 2009, accepted 31 July 2009

### Abstract

The genetic features of each isolate were determined by enterobacterial repetitive intergenic consensus (ERIC) primer sequences used in PCR and by searching for six virulence genes (*alg D*, *las B*, *tox A*, *plc H*, *plc N*, *exo S*). 49 (79%) of the isolates were distributed in three ERIC PCR subgroups and showed 62% of similarity. The remaining 13 strains generated unique patterns. The first subgroup was primarily composed of isolates from faeces, these strains indicated over 70% relationship with the next subgroup, and primarily contained strains isolated from wounds and bronchial washings and the last subgroup contained strains isolated from wounds and urine. The unique strains were isolated mainly from urine. Statistical analysis indicated that variations in distribution of virulence genes in *P. aeruginosa* isolates with respect to strain origin and genomic subgroups were not significant. In the group of 49 strains, 100% gave a positive reaction to *alg D*, *las B* and *plc H* genes, 91.8% to *tox A* and *plc N* genes and 83.7% to *exo S* gene. Among the strains that generated unique (ERIC-PCR) patterns, 69.2% gave a positive reaction to *alg D* gene, 84.6% to *las B* gene, 76.9% to *tox A*, *plc N* and *plc H* genes, and 46.15% to *exo S* gene.

---

Key words: *Pseudomonas aeruginosa*, ERIC-PCR, virulence genes

---

### Introduction

*Pseudomonas aeruginosa* is an opportunistic pathogen that is responsible for a wide range of infections. It is a common hospital-acquired pathogen and is responsible for ventilator-associated pneumonia in patients with underlying immune defects (Carratala *et al.*, 1998), as well as wound and catheter-associated infections. Also intestinal colonization by *P. aeruginosa* could be a reservoir for invasive infections caused by this bacterium and for its dissemination (Zaborina *et al.*, 2006). In healthy individuals, it is a leading cause of contact lens keratitis (Cheng *et al.*, 1999), otitis externa (swimmer's ear), and hot-tub folliculitis (Speert, 2002). Additionally, *P. aeruginosa* is the most common pathogen in lung infections affecting cystic fibrosis (CF) patients and is the leading cause of morbidity and mortality in this patient group (Govan and Deretic, 1996). The ability of *P. aeruginosa* to cause infection is further exacerbated by a high level of resistance to antibiotics, which makes *Pseudomonas* infections difficult to treat (Hancock and Speert, 2000).

*P. aeruginosa* produces several virulent factors to colonize the cells of its host. Many of these factors are

controlled by regulatory systems involving cell-to-cell signaling (Van Delden and Iglewski, 1998). Among these are: exotoxin A, exoenzyme S, las B elastase, phospholipases C and alginate. Exotoxin A, encoded by the *tox A* gene, inhibits protein biosynthesis by transferring an ADP-ribosyl moiety to elongation factor 2 of eukaryotic cells. Exoenzyme S, encoded by the *exo S* gene, is also an ADP-ribosyltransferase that is secreted by a type-III secretion system directly into the cytosol of epithelial cells (Rumbaugh *et al.*, 1999a). Las B elastase, a zinc metalloprotease encoded by the *las B* gene, attacks eukaryotic proteins such as collagen and elastin, and destroys the structural proteins of the cell (Toder *et al.*, 1994). While phospholipids are hydrolyzed by two phospholipases C encoded by *plc H* and *plc N* genes (PLC-H and PLC N, respectively). Alginate, encoded by *alg D* gene, protects the bacterium from the host's immune response and from antibiotics (Ostroff *et al.*, 1990; Storey *et al.*, 1997).

Genetic methods have been used to explore the genetic diversity of various populations of *P. aeruginosa* strains. Various methods, such as pulsed-field gel electrophoresis, arbitrarily primed polymerase chain reaction, and ribotyping are currently available for genotyping of *P. aeruginosa* (Bennekov *et al.*, 1996).

---

\* Corresponding author: K. Wolska, Department of Microbiology, Natural Faculty, University of Podlasie in Siedlce, Prusa 12, 08-110 Siedlce, Poland; e-mail: kwolska@ap.siedlce.pl

On the other hand, the enterobacterial repetitive intergenic consensus (ERIC) sequences are known to be dispersed throughout the prokaryotic genome, and polymerase chain reaction (PCR) studies of eubacteria have revealed that inter-ERIC distances and patterns are highly specific to the individual group of the same bacterial species (Versalovic *et al.*, 1991).

In the present study we aimed to determine the genetic features of clinical *P. aeruginosa* isolates by ERIC – based PCR (ERIC-PCR) and by assessing variations in the prevalence of six virulence genes (*alg D*, *las B*, *tox A*, *plc H*, *plc N*, *exo S*).

## Experimental

### Materials and Methods

**Source of isolates and identification.** A total of 62 strains of *P. aeruginosa*, were originally isolated from a variety of clinical specimens: faeces (26), urine (11), blood (1), bronchial washings (9), sputum (1), wound swab (9), throat swab (2), ulceration swab (1), swab from skin round tracheotomy (1) and from ear (1). The bacteria were obtained from 62 patients from different wards of the municipal hospital, main hospital and outpatients' department in Siedlce (Poland), between December 2005 and March 2006. The strains were identified as *P. aeruginosa* according to biochemical patterns in the Api 20NE system (bio Mérieux). The reference strain NCTC 6749 was also examined. Stock cultures were stored in TSB (tripticase soy broth, Difco) containing 20% glycerol at  $-80^{\circ}\text{C}$ .

**DNA extraction.** Isolates were grown in trypticase-soya broth at  $37^{\circ}\text{C}$  for 24 h and DNA was extracted by using the Genomic DNA Pre Plus (A&A Biotechnology, Gdańsk, Poland).

**Detection of virulence genes by PCR.** The prevalence of virulence genes encoding alginate (*alg D*), Las B elastase (*las B*), haemolytic phospholipase C (*plc H*), non-haemolytic phospholipase C (*plc N*), exoenzyme S (*exo S*) and exotoxin A (*tox A*) was determined by PCR. The genes were amplified with primers selected on the basis of the published PAO1 sequence (Stover *et al.*, 2000). The PCR mixture contained PCR buffer (10 mM Tris/HCl, 50 mM KCl, 1.5 mM  $\text{MgCl}_2$ , pH 8.3), 200  $\mu\text{M}$  of each dNTP (Boehringer), 12.5 pmol of each primer, DMSO at a final concentration of 4%, 1 U Ampli Tag DNA polymerase (Perkin Elmer) and 25 ng DNA template. The DNA was amplified in PTC-100 Programmable Thermo Controller (MJ Research) using the following protocol:  $94^{\circ}\text{C}$  for 3 min, 30 cycles of  $94^{\circ}\text{C}$  for 30 s,  $55^{\circ}\text{C}$  for 1 min and  $72^{\circ}\text{C}$  for 1 min 30 s, and  $72^{\circ}\text{C}$  for 5 min. Each gene was amplified separately. PCR products were separated in a 1% agarose gel for 1 h at 100 V, stained with ethidium bromide and detected by UV transillumination. Amplified genes were identified on the basis of fragment size (Table I).

**Enterobacterial Repetitive Intergenic Consensus (ERIC-PCR).** ERIC primer sequences were used in PCR to detect differences in the number and distribution of these bacterial repetitive sequences in the bacterial genome. ERIC PCR was carried out using the primer sequences ERIC-1R, 5'-CACTTAGGGGTCC TCGAATGTA-3' and ERIC-2, 5'-AAGTAAGTGACT GGGGTGAGCG-3' to amplify the regions in the bacterial genome placed between the ERIC sequences. Dice coefficient was calculated and compared to evaluate similarity among strains through the use of BIOGENE software. The method was described in the previous study (Wolska and Szwedka, 2008).

**Statistical methods.** The distribution of virulence genes with respect to genomic groups or strain origin was compared using chi-square ( $\chi^2$ ) test.

Table I  
Primers used for PCR amplification of virulence factors

Gene	Primer sequence (5'-3')	Product (bp)	Number of pair G+C
<i>alg D</i>	alg DF – CGTCTGCCGCGAGATCGGCT	313	14
	alg DR – GACCTCGACGGTCTTGCGGA		13
<i>las B</i>	las BF – GGAATGAACGAAGCGTTCTCCGAC	284	13
	las BR – TTGGCGTCGACGAACACCTCG		13
<i>tox A</i>	tox AF – CTGCGCGGGTCTATGTGCC	270	13
	tox AR – GATGCTGGACGGGTGCGAG		12
<i>plc H</i>	plc HF – GCACGTGGTCATCCTGATGC	608	14
	plc HR – TCCGTAGGCGTCGACGTAC		12
<i>plc N</i>	plc NF – TCCGTTATCGCAACCAGCCCTACG	481	14
	plc NR – TCGCTGTGAGCAGGTGCGAAC		13
<i>exo S</i>	exo SF – CGTCGTGTTCAAGCAGATGGTGCTG	444	14
	exo SR – CCGAACCGCTTACCAGGC		13

## Results

We used PCR to assess the prevalence of six virulence genes. PCR detected *alg D*, *las B*, *tox A*, *plc H*, *plc N* and *exo S* in 58 (93.55%), 60 (96.8%), 55 (88.7%), 59 (95.2%), 55 (88.7%) and 47 (75.8%) isolates, respectively. 40 (64.5%) isolates gave positive PCR results for all studied genes. In this group, 17 (65.4%) isolates were obtained from faeces, 5 (45.45%) isolates were obtained from urine, 7 (77.8%) from wound, 8 (88.9%) from bronchial washings and individual strains were isolated from sputum, skin and blood. The remaining strains (22–35.5%) gave a negative result for one or more genes. All studied genes were not detected in two strains; the first strain (number 2) was isolated from urine and the second one (42) was obtained from throat swab. The strain isolated from faeces (59) gave a negative PCR reaction to *alg D*, *tox A*, *plc N* and *exo S* genes. The next strain isolated from the same source (53) gave a negative PCR reaction to *alg D*, *plc N* and *exo S* genes, and the last strain of this origin (8) gave a negative PCR reaction to *plc N* and *exo S* genes. The other strains gave a negative result to one of the three following genes: *exo S*, *tox A* and *plc N*. *Exo S* gene was not presented in strains isolated from urine (4 isolates), faeces (1), throat swab (1), wound (2), bronchial washings (1) and ear (1). *Tox A* gene and *plc N* were not detected

in 2 and 3 strains isolated from faeces. The variations in distribution of virulence genes in *P. aeruginosa* isolates with respect to origin were not significant by the use the chi-square analysis ( $\lambda^2 = 4.01$ ) (Table II).

Nearly all strains, with the exception of 2, 42 and 59 which gave negative reactions to nearly all studied virulence genes, showed 50% of similarity according to Dice coefficient values (Fig. 1). 41 (66.1% of all) isolates revealed a high similarity, over 70%. Dendrogram analysis enabled the division of strains into groups (50–62% of similarity), then subgroups (64–80%), genotypes (80–86%) and lastly subtypes (86–100%). 13 (21%) strains generated unique ERIC-PCR patterns. 49 (79%) isolates and reference strain of *P. aeruginosa* creating group 1 showed 62% of similarity. The chi-square analysis indicated that variations in distribution of *P. aeruginosa* isolates in genomic subgroups with respect to ecological origin were significant ( $\lambda^2 = 80.186$ ) (Table III). The most numerous clonal subgroup of 1, which was characterized by nearly 80% of similarity, consisted of 2 genotypes. This subgroup contained 29 isolates, among which, 24 (92.3%) were isolated from faeces and the individual strains in this group were isolated from wound, skin, bronchial washings, urine and throat swab. The mentioned subgroup demonstrated over 70% of similarity with the next clonal subgroup, which included 12 isolates situated in 2 types. Among

Table II  
Distribution of virulence genes in *Pseudomonas aeruginosa* isolates with respect to origin

Origin (No) of strains	Virulence genes					
	<i>alg D</i>	<i>las B</i>	<i>tox A</i>	<i>plc H</i>	<i>plc N</i>	<i>exo S</i>
Faeces (26)	24 (92.3%)	26 (100%)	23 (88.5%)	25 (96.1%)	21 (80.8%)	22 (84.6%)
Urine (11)	10 (90.9%)	10 (90.9%)	9 (81.8%)	10 (90.9%)	10 (90.9%)	6 (54.5%)
Wound (9)	9 (100%)	9 (100%)	9 (100%)	9 (100%)	9 (100%)	7 (77.8%)
Bronchial washings (9)	9 (100%)	9 (100%)	9 (100%)	9 (100%)	9 (100%)	8 (88.9%)
Throat (2)	1	1	1	1	1	–
Ulceration (1)	1	1	–	1	1	1
Ear (1)	1	1	1	1	1	–
Blood (1)	1	1	1	1	1	1
Skin (1)	1	1	1	1	1	1
Sputum (1)	1	1	1	1	1	1
Reference (1)	1	1	1	1	1	1

Table III  
Distribution of the 63 *Pseudomonas aeruginosa* isolates in genomic subgroups with respect to origin

Genomic subgroups (No of strains)	Faecies (n = 26)	Urine (n = 11)	Wound (n = 9)	Bronchial washings (n = 9)	Throat (n = 2)	Ulceration (n = 1)	Ear (n = 1)	Blood (n = 1)	Skin (n = 1)	Sputum (n = 1)	Reference (n = 1)
1 (29)	24 (92.3%)	1 (9.1%)	1 (11.1%)	1 (11.1%)	1 (50%)	–	–	–	1	–	–
2 (12)	–	1 (8.3%)	5 (55.5%)	4 (44.4%)	–	1	1	–	–	–	–
3 (9)	–	3 (27.3%)	3 (33.3%)	1 (11.1%)	–	–	–	–	–	1	1
The unique patterns (13)	2 (7.7%)	6 (54.5%)	–	3 (33.3%)	1 (50%)	–	–	1	–	–	–




Fig. 1. Dendrogram showing genetic relationships between 62 *P. aeruginosa* isolates. The results were obtained by ERIC-PCR analysis using two primers.

Table IV  
Distribution of virulence genes in *Pseudomonas aeruginosa* isolates with respect to genomic subgroups

Genomic subgroups (No of strains)	Virulence genes					
	<i>alg D</i>	<i>las B</i>	<i>tox A</i>	<i>plc H</i>	<i>plc N</i>	<i>exo S</i>
1 (29)	29 (100%)	29 (100%)	27 (93.1%)	29 (100%)	25 (86.2%)	24 (82.7%)
2 (12)	12 (100%)	12 (100%)	11 (91.7%)	12 (100%)	12 (100%)	11 (91.7%)
3 (9)	9 (100%)	9 (100%)	8 (88.9%)	9 (100%)	9 (100%)	7 (77.8%)
The unique patterns (13)	9 (69.2%)	11 (84.6%)	10 (76.9%)	10 (76.9%)	10 (76.9%)	6 (46.15)

these 12 isolates, 5 (55.6%) strains were obtained from wound, 4 (44.4%) from bronchial washings, 1 (9.1%) from urine and 1 from ulceration. Among 41 strains forming subgroups 1 and 2, 29 (70.7%) gave a positive reaction to all six genes. 11 (26.8%) strains gave a positive reaction to five genes and only 1 (2.4%) strain gave a positive reaction to four genes. All isolates of these subgroups gave positive reactions to *alg D*, *las B* and *plc H* genes. The prevalence of *tox A* gene in isolates of subgroup 1 was similar to isolates of subgroup 2. And the prevalence of *exo S* gene was higher in the isolates of group 2 than in group 1. The last 9 strains forming group 1, showed 62% of similarity with subgroups 1 and 2. Among which, there were isolates from urine (3 (27.3%) strains), wound (3 (33.3%)), sputum (1), bronchial washings (1 (11.1%)) and the reference strain of *P. aeruginosa* NCTC 6749. These strains gave a positive reaction to nearly all virulence genes with the exception of two strains isolated from urine. One gave a negative result for *exo S* gene and the second one gave a negative result for *tox A* gene. The remaining strains which were collected (13–21%) generated unique (ERIC-PCR) patterns, showing the genetic diversity of these bacteria. They were isolated from urine (6 (54.5%)), bronchial washings (3 (33.3%)), blood (1), faeces (2 (7.7%)) and throat swab (1). Among these strains, 6 (46.15%) gave positive reactions to all virulence genes, but 7 (53.85%) gave negative results to one or more (from 3 to 6) virulence gene. The statistical analysis showed that variations in distribution of virulence genes in *P. aeruginosa* isolates with respect to genomic subgroups were not significant ( $\lambda^2 = 4.01$ ) (Table IV).

### Discussion

The relationship between isolates was calculated by numerical analysis of genetic features determined by ERIC PCR. 79% of isolates were distributed in three ERIC PCR subgroups and showed 62% of similarity. The remaining strains generated unique patterns. The first subgroup was primarily composed of isolates from faeces that were strictly related with each other. These strains indicated over 70% relationship with the

next subgroup, and primarily contained strains isolated from wound and bronchial washings. Above all, the last subgroup contained strains isolated from wound and urine. The unique strains were isolated mainly from urine. Virulence genes were detected more frequently in strains isolated from bronchial washings and wound rather than in strains isolated from urine and faeces. However, the statistical analysis indicated that variations in distribution of virulence genes in *P. aeruginosa* isolates with respect to strain origin and genomic subgroups were not significant. This result is in agreement with the studies of Hamood *et al.* (1996) and Rumaugh *et al.* (1999b). They suggest that elastase, phospholipase C, exotoxin A and exoenzyme S are produced by *P. aeruginosa* isolates from the different sites of infection. The studies also demonstrated that the production of higher levels of elastase and phospholipase C is important in all types of infections. In some bacterial infections, complex systems of coordinate regulation control the expression of virulence genes (Mecalanos, 1992). Storey *et al.* (1997) found a positive correlation between *alg D* transcript accumulation and both *las B* and *las A* transcript accumulation levels. This indicates a common regulatory element in a cascade of regulators or a common environmental cue that triggers transcription. Numerous authors (Joly *et al.*, 2005; Rietsch *et al.*, 2005; Yahr *et al.*, 1995) have documented the presence of a type III secretion system in *P. aeruginosa* that appears to play a major role in the virulence of this organism. Endimiani *et al.* (2006) showed that 100% of *P. aeruginosa* isolates from bloodstream infections were positive for the following genes: *exo T*, *exo U*, *las B*, *plc H*, *plc N*, *tox A* and *nan 2*. *Exo S*, *exo Y* and *nan 1* genes were detected in 78.9%, 73.7% and 57.9% of isolates. Our data demonstrated that the studied genes: *alg D*, *las B*, *tox A*, *plc H*, *plc N* and *exo S* were present in 93.55%, 96.8%, 88.7%, 95.2%, 88.7% and 75.8% *P. aeruginosa* isolates, respectively. Feltman *et al.* (2001) and Ferguson *et al.* (2001) indicated that over 90% of clinical *P. aeruginosa* strains contain *exo T* and *exo Y* genes, but clinical isolates from urine frequently have the *exo Y* gene present at a relatively lower level-about 70%. Similarly, in our study *exo S* gene was rarely detected in *P. aeruginosa* strains isolated from urine. According



to Lin *et al.* (2006), *exo U* gene of *P. aeruginosa* is the major contributor to cytotoxicity against mammalian cells. Others (Rumbaugh *et al.*, 1999a) revealed that *P. aeruginosa* isolates recovered from patients suffering from urinary tract infections produced significant levels of exotoxin A. Over 80% of studied isolates from urine contained the *tox A* gene. In all probability, this toxin can play a significant role as a virulence factor of *P. aeruginosa* within catheter-associated urinary tract infections (Goldsworthy, 2008). Lanotte *et al.* (2004) demonstrated that 80% of all pulmonary isolates harboured *exo S*, indicating that this gene plays a major role in pulmonary infections. In our study also, strains isolated from bronchial washings showed the prevalence of *exo S*. *Exo S* gene was frequently associated with multi-drug resistant (MDR) strains which suggests that *P. aeruginosa* isolates showing MDR traits may be more virulent than susceptible strains (Endimiani *et al.*, 2006). Results of Zaborina *et al.* (2006) showed that the multi-drug resistant *P. aeruginosa* clinical strains that disrupted the barrier function of cultured intestinal epithelial cells are characterized by the *exo U* gene. The presence of virulent strains of *P. aeruginosa* within the intestinal tract could be the major source of systemic sepsis and death among immuno-compromised patients.

The above presented data has been founded in Poland. So far nobody has applied the ERIC-PCR method in discriminating clinical *P. aeruginosa* strains and evaluated the occurrence of such numerous groups of virulence genes.

## Literature

- Bennekov T., H. Colding, B. Ojeniyi, M.W. Bentzon and N. Hoiby. 1996. Comparison of ribotyping of *Pseudomonas aeruginosa* isolates from cystic fibrosis patients. *J. Clin. Microbiol.* 34: 202–204.
- Carratala J., B. Roson, A. Fernandez-Sevilla, F. Alcaide and F. Gudiol. 1998. Bacteremic pneumonia in neutropenic patients with cancer. *Arch. Intern. Med.* 158: 868–872.
- Cheng K.H., S.L. Leung, H.W. Hoekman, W.H. Beekhuis, P.G. Mulder, A.J. Geerards and A. Kijlstra. 1999. Incidence of contact-lens-associated microbial keratitis and its related morbidity. *Lancet* 354: 181–185.
- Endimiani A., B. Pini, A. Baj, F. Luzzaro and A. Toniolo. 2006. Bloodstream infections due to *Pseudomonas aeruginosa*: clinical outcome associated with pathogenesis-related genes. Abstracts 16<sup>th</sup> European Congress Clinical Microbiology and Infectious Disease 2006. Nice, France p. 1164.
- Feltman H., G. Schlert, S. Khan, M. Jain, L. Peterson and A.R. Hauser. 2001. Prevalence of type III secretion genes in clinical and environmental isolates of *Pseudomonas aeruginosa*. *Microbiology* 147: 2659–2669.
- Ferguson M.W., J.A. Maxwell, T.S. Vincent, J. da Silva and J.C. Olson. 2001. Comparison of the *exo S* gene and protein expression in soil and clinical isolates of *Pseudomonas aeruginosa*. *Infect. Immun.* 69: 2198–2210.
- Goldsworthy M.J.H. 2008. Gene expression of *Pseudomonas aeruginosa* and MRSA within a catheter-associated urinary tract infection biofilm model. *Bioscience Horizons* 1: 28–37.
- Govan J.R. and V. Deretic. 1996. Microbial pathogenesis in cystic fibrosis: mucoid *Pseudomonas aeruginosa* and *Burkholderia cepacia*. *Microbiol. Rev.* 60: 539–574.
- Hamood A.N., J.A. Griswold and C.M. Duhan. 1996. Production of extracellular virulence factors by *Pseudomonas aeruginosa* isolates obtained from tracheal, urinary tract and wound infections. *J. Surg. Research* 61: 425–432.
- Hancock R.E. and D.P. Speert. 2000. Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and impact on treatment. *Drug. Resist. Updat.* 4: 247–255.
- Joly B., M. Pierre, S. Auvin, F. Colin, F.B. Guery and M.O. Husson. 2005. Relative expression of *Pseudomonas aeruginosa* virulence genes analyzed by a real time RT-PCR method during lung infection in rats. *FEMS Microbiol. Lett.* 243: 271–278.
- Lanotte P., S. Watt, L. Mereghetti, N. Dartiguelongue, A. Raster-Lari, A. Goudeau and R. Quentin. 2004. Genetic features of *Pseudomonas aeruginosa* isolates from cystic fibrosis patients compared with those of isolates from other origins. *J. Med. Microbiol.* 53: 73–81.
- Lin H., S. Huang, H. Teng, D. Ji, Y. Chen and Y. Chen. 2006. Presence of the *exo U* gene of *Pseudomonas aeruginosa* is correlated with cytotoxicity in MDCK cells but not with colonization in BALB/c mice. *J. Clin. Microbiol.* 44: 4596–4597.
- Mecalanos J.J. 1992. Environmental signals controlling expression of virulence determinants in bacteria. *J. Bacteriol.* 174: 1–7.
- Ostroff R.M., A.I. Vasil and M.L. Vasil. 1990. Molecular comparison of nonhemolytic and a hemolytic phospholipase C from *Pseudomonas aeruginosa*. *J. Bacteriol.* 172: 5915–5923.
- Rietsch A., I. Vallet-Gely, S.L. Dove and J.J. Mekalanos. 2005. ExsE, a secreted regulator of type III secretion genes in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci.* 102: 9006–8011.
- Rumbaugh K.P., B.S. Abdul, N. Hamood and J.A. Griswold. 1999. Analysis of *Pseudomonas aeruginosa* clinical isolates for possible variations within the virulence genes exotoxin A and exoenzyme S. *J. Surg. Research* 82: 95–105.
- Rumbaugh K.P., J.A. Griswold and A.N. Hamood. 1999. *Pseudomonas aeruginosa* strains obtained from patients with tracheal, urinary tract and wound infection: variations in virulence factors and virulence genes. *J. Hosp. Infect.* 43: 211–218.
- Speert D.P. 2002. Molecular epidemiology of *Pseudomonas aeruginosa*. *Front. Biosci.* 7: e354–e361.
- Storey D.G., E.E. Ujack, J. Mitchell and H.R. Rabin. 1997. Positive correlation of *algD* transcription to *las B* and *las A* transcription by populations of *Pseudomonas aeruginosa* in the lungs of patients with cystic fibrosis. *Infect. Immun.* 65: 4061–4067.
- Stover C.K., X.Q. Pham, A.L. Erwin, S.D. Mizoguchi, P. Warren, M.J. Hickey, F.S.L. Brinkman, W.O. Hufnagle and D.J. Kowalik. 2000. Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an opportunistic pathogen. *Nature* 406: 959–964.
- Toder D.S., S.J. Ferrell, J.L. Nezezon, L. Rust and B.H. Iglewski. 1994. *Las A* and *las B* genes of *Pseudomonas aeruginosa*: Analysis transcription and gene product activity. *Infect. Immun.* 62: 1320–1327.
- Van Delden C. and B.H. Iglewski 1998. Cell-to-cell signaling and *Pseudomonas aeruginosa* infections. *Emerg. Infect. Dis.* 4: 551–60.
- Versalovic J., T. Koeuth and J.R. Lupski. 1991. Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Res.* 19: 6823–6831.
- Wolska K. and P. Szwedra. 2008. A comparative evaluation of PCR ribotyping and ERIC PCR for determining the diversity of clinical *Pseudomonas aeruginosa* isolates. *Pol. J. Microbiol.* 57: 157–163.
- Yahr T.L., A.K. Hovey, S.M. Kulich and D.W. Frank. 1995. Transcriptional analysis of the *Pseudomonas aeruginosa* exoenzyme S structural gene. *J. Bacteriol.* 177: 1169–1178.
- Zaborina O., J.E. Kohler, Y. Wang, C. Bethel, O. Shevchenko, L. Wu, J.R. Turner and J.C. Alverdy. 2006. Identification of multi-drug resistant *Pseudomonas aeruginosa* clinical isolates that highly disruptive to the intestinal epithelial barrier. *An. Clin. Microbiol. Antimicrob.* 5: 1–14.