

Trichloridobis(ethyldiphenylphosphine)- (tetrahydrofuran)molybdenum(III)

Tomasz Kruczyński, Jerzy Pikies and Łukasz Ponikiewski*

Chemical Faculty, Gdańsk University of Technology, Narutowicza 11/12, Gdańsk PL 80233, Poland

Correspondence e-mail: lukasz_ponikiewski@vp.pl


Received 10 May 2010; accepted 7 June 2010

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; disorder in main residue; R factor = 0.044; wR factor = 0.115; data-to-parameter ratio = 18.2.

In the mononuclear title compound, $[\text{MoCl}_3(\text{C}_4\text{H}_8\text{O})(\text{C}_{14}\text{H}_{15}\text{P})_2]$, obtained by the reaction of trichlorotris(tetrahydrofuran)molybdenum(III) and ethyldiphenylphosphine in tetrahydrofuran (THF) solution, the Mo^{III} atom is six-coordinated by one O atom of a THF molecule, two P atoms from two ethyldiphenylphosphine ligands and three Cl atoms in a distorted octahedral geometry. The C atoms of the THF molecule are disordered over two positions in a 0.55 (2):0.45 (2) ratio.

Related literature

For the structures of similar molybdenum complexes and for bond-length data, see: Cotton & Jianrui (1996); Cotton & Vidyasagar (1995); Hofacker *et al.* (1989); Borgmann *et al.* (1997). For the synthesis, see: Anker *et al.* (1975).



Experimental

Crystal data

$[\text{MoCl}_3(\text{C}_4\text{H}_8\text{O})(\text{C}_{14}\text{H}_{15}\text{P})_2]$
 $M_r = 702.85$
 Monoclinic, $P2_1/c$
 $a = 15.437$ Å
 $b = 13.356$ Å
 $c = 20.229$ Å
 $\beta = 128.87^\circ$

$V = 3247.1$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.77$ mm⁻¹
 $T = 120$ K
 $0.13 \times 0.10 \times 0.04$ mm

Data collection

Oxford Diffraction KM-4/Xcalibur diffractometer with a Sapphire2 detector
 Absorption correction: analytical [*CrysAlis RED* (Oxford Diffraction, 2006)], based on expressions derived by Clark &

Reid (1995)
 $T_{\min} = 0.937$, $T_{\max} = 0.970$
 25240 measured reflections
 7022 independent reflections
 2985 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.089$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.115$
 $S = 0.81$
 7022 reflections

385 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.70$ e Å⁻³
 $\Delta\rho_{\min} = -0.54$ e Å⁻³

Table 1

Selected bond lengths (Å).

Mo1—O1	2.206 (3)	Mo1—Cl3	2.4126 (13)
Mo1—Cl1	2.3871 (13)	Mo1—P1	2.5964 (14)
Mo1—Cl2	2.3822 (13)	Mo1—P2	2.5974 (13)

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2006); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2006); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The work was undertaken with financial support from the Polish State Committee, grant No. NN204271535.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZZ211).

References

- Anker, M. W., Chatt, J., Leigh, G. J. & Wedd, A. G. (1975). *J. Chem. Soc. Dalton Trans.* pp. 2639–2645.
 Borgmann, C., Limberg, C. & Driess, A. (1997). *J. Organomet. Chem.* **541**, 367–375.
 Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
 Cotton, F. A. & Jianrui, S. (1996). *Inorg. Chim. Acta*, **251**, 101–104.
 Cotton, F. A. & Vidyasagar, K. (1995). *Polyhedron*, **14**, 3077–3085.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Hofacker, P., Friebel, C., Dehnicke, K., Bauml, P., Hiller, W. & Strähle, J. (1989). *Z. Naturforsch. Teil B*, **44**, 1161–1166.
 Oxford Diffraction (2006). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2010). E66, m786 [doi:10.1107/S1600536810021690]

Trichloridobis(ethyldiphenylphosphine)(tetrahydrofuran)molybdenum(III)**Tomasz Kruczyński, Jerzy Pikies and Łukasz Ponikiewski****S1. Comment**

The title molecule [MoCl₃(PEtPh₂)(THF)] was prepared as a potential adduct for synthesis with lithium phosphanides of the formula R₂P—P(SiMe₃)Li (*R* = ^tBu, ⁱPr, Et₂N, ⁱPr₂N).

The Mo^{III} atom resides in a distorted MoCl₃OP₂ octahedral environment. The equatorial positions are occupied by three Cl atoms and one O atom from the THF, while the axial positions are occupied by P atoms from two ethyldiphenylphosphine residues. The Mo—Cl bond length [2.3871 (13) Å, 2.3822 (13) Å, 2.4126 (13) Å], the Mo—P bond length [2.5964 (14) Å, 2.5974 (13) Å] and the Mo—O bond length [2.206 (3) Å] are very similar to the previously reported molybdenum complexes (Cotton & Vidyasagar, 1995; Cotton & Jianrui, 1996; Hofacker *et al.*, 1989; Borgmann *et al.*, 1997).

Atoms C29, C30, C31, C32 from the THF molecule were disordered over two positions. During the refinement process the disorder models were refined with occupancies of 0.55 (2) and 0.45 (2).

S2. Experimental

The title compound was prepared according to the previously published method (Anker *et al.*, 1975)

S3. Refinement

Atoms C29, C30, C31, C32 were disordered over two positions. During the refinement process the disorder models were refined with occupancies of 0.55 (2) and 0.45 (2). H atoms bonded to C were included in calculated positions and refined as riding on their parent C atom with C—H = 0.95 Å, *U*_{iso}(H) = 1.2 *U*_{eq}(C) for aromatic; C—H = 0.99 Å, *U*_{iso}(H) = 1.2 *U*_{eq}(C) for methylene; and C—H = 0.98 Å, *U*_{iso}(H) = 1.5 *U*_{eq}(C) for methyl atoms.




Figure 1

The molecular structure of the title compound showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. For clarity, all H atoms and the minor component of the disorder have been omitted.

Trichloridobis(ethylidiphenylphosphine)(tetrahydrofuran)molybdenum(III)

Crystal data

$[\text{MoCl}_3(\text{C}_4\text{H}_8\text{O})(\text{C}_{14}\text{H}_{15}\text{P})_2]$

$M_r = 702.85$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 15.437\ \text{\AA}$

$b = 13.356\ \text{\AA}$

$c = 20.229\ \text{\AA}$

$\beta = 128.87^\circ$

$V = 3247.1\ \text{\AA}^3$

$Z = 4$

$F(000) = 1444$

$D_x = 1.438\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 5220 reflections

$\theta = 2.0\text{--}28.9^\circ$

$\mu = 0.77\ \text{mm}^{-1}$

$T = 120\ \text{K}$

Block, orange

$0.13 \times 0.1 \times 0.04\ \text{mm}$

Data collection

Oxford Diffraction KM-4/Xcalibur
diffractometer with a Sapphire2 (large Be
window) detector

Graphite monochromator

Detector resolution: $8.1883\ \text{pixels mm}^{-1}$

ω scans

Absorption correction: analytical

[*CrysAlis RED* (Oxford Diffraction, 2006).
based on expressions derived by Clark & Reid
(1995)]

$T_{\min} = 0.937$, $T_{\max} = 0.97$

25240 measured reflections

7022 independent reflections

2985 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.089$

$\theta_{\max} = 27^\circ$, $\theta_{\min} = 2^\circ$

$h = -15 \rightarrow 19$

$k = -16 \rightarrow 17$

$l = -25 \rightarrow 22$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H-atom parameters constrained
$wR(F^2) = 0.115$	$w = 1/[\sigma^2(F_o^2) + (0.0551P)^2]$
$S = 0.81$	where $P = (F_o^2 + 2F_c^2)/3$
7022 reflections	$(\Delta/\sigma)_{\max} = 0.001$
385 parameters	$\Delta\rho_{\max} = 0.70 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.54 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Experimental. CrysAlis RED (Oxford Diffraction, 2006) Analytical numeric absorption correction using a multifaceted crystal model based on expressions derived by Clark & Reid (1995)

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Mo1	0.57979 (3)	0.74957 (4)	0.63166 (2)	0.03356 (12)	
Cl1	0.53421 (13)	0.92350 (10)	0.61183 (10)	0.0612 (4)	
Cl2	0.62891 (9)	0.74919 (11)	0.76928 (7)	0.0511 (3)	
Cl3	0.62090 (10)	0.57325 (9)	0.64269 (9)	0.0488 (3)	
P1	0.78469 (10)	0.78721 (10)	0.69759 (9)	0.0415 (3)	
P2	0.37606 (10)	0.70228 (10)	0.56343 (8)	0.0370 (3)	
O1	0.5361 (2)	0.7484 (3)	0.50482 (18)	0.0411 (7)	
C1	0.8589 (4)	0.8883 (4)	0.7753 (3)	0.0495 (14)	
C2	0.8104 (5)	0.9421 (5)	0.8018 (4)	0.075 (2)	
H2A	0.7358	0.9277	0.7785	0.09*	
C3	0.8682 (6)	1.0174 (6)	0.8621 (5)	0.099 (3)	
H3A	0.8334	1.0547	0.8797	0.118*	
C4	0.9750 (6)	1.0370 (5)	0.8957 (4)	0.084 (2)	
H4A	1.0161	1.0871	0.938	0.101*	
C5	1.0232 (5)	0.9849 (5)	0.8687 (4)	0.0711 (19)	
H5A	1.0976	1	0.8918	0.085*	
C6	0.9674 (4)	0.9118 (4)	0.8095 (4)	0.0645 (17)	
H6A	1.0026	0.8765	0.7913	0.077*	
C7	0.8016 (4)	0.8198 (4)	0.6187 (3)	0.0462 (13)	
C8	0.8262 (6)	0.7500 (6)	0.5828 (4)	0.0873 (15)	
H8A	0.8389	0.6826	0.6018	0.105*	
C9	0.8329 (5)	0.7744 (6)	0.5206 (4)	0.084 (2)	
H9A	0.8507	0.7242	0.4977	0.101*	

C10	0.8145 (5)	0.8694 (6)	0.4914 (4)	0.080 (2)	
H10A	0.8208	0.8866	0.449	0.096*	
C11	0.7871 (6)	0.9387 (5)	0.5234 (4)	0.0873 (15)	
H11A	0.7712	1.0052	0.5021	0.105*	
C12	0.7820 (7)	0.9141 (5)	0.5872 (5)	0.098 (2)	
H12A	0.7641	0.9649	0.6097	0.117*	
C13	0.8800 (4)	0.6808 (4)	0.7548 (4)	0.0563 (16)	
H13A	0.85	0.6221	0.7166	0.068*	
H13B	0.9529	0.6977	0.7698	0.068*	
C14	0.8976 (5)	0.6530 (5)	0.8342 (4)	0.084 (2)	
H14A	0.9562	0.6022	0.8654	0.126*	
H14B	0.8283	0.6261	0.8194	0.126*	
H14C	0.9198	0.7126	0.87	0.126*	
C15	0.2980 (4)	0.7756 (3)	0.5873 (3)	0.0414 (13)	
C16	0.1881 (4)	0.7565 (4)	0.5442 (3)	0.0572 (14)	
H16A	0.1526	0.705	0.5026	0.069*	
C17	0.1273 (5)	0.8118 (5)	0.5608 (4)	0.0656 (17)	
H17A	0.051	0.7972	0.5318	0.079*	
C18	0.1780 (5)	0.8869 (5)	0.6187 (4)	0.0639 (17)	
H18A	0.1364	0.9264	0.6291	0.077*	
C19	0.2870 (5)	0.9059 (5)	0.6615 (4)	0.0675 (18)	
H19A	0.3217	0.9584	0.7021	0.081*	
C20	0.3483 (4)	0.8502 (4)	0.6471 (3)	0.0524 (14)	
H20A	0.4255	0.8633	0.6785	0.063*	
C21	0.2853 (4)	0.7107 (4)	0.4475 (3)	0.0416 (13)	
C22	0.2670 (4)	0.6302 (4)	0.3982 (3)	0.0497 (14)	
H22A	0.2983	0.567	0.4241	0.06*	
C23	0.2041 (4)	0.6398 (5)	0.3121 (3)	0.0594 (16)	
H23A	0.1927	0.5835	0.2787	0.071*	
C24	0.1579 (4)	0.7294 (5)	0.2740 (3)	0.0630 (18)	
H24A	0.1137	0.7357	0.2141	0.076*	
C25	0.1750 (4)	0.8081 (5)	0.3209 (4)	0.0621 (16)	
H25A	0.1427	0.8707	0.2942	0.075*	
C26	0.2390 (4)	0.8001 (4)	0.4079 (4)	0.0599 (16)	
H26A	0.2509	0.8574	0.4405	0.072*	
C27	0.3571 (4)	0.5747 (4)	0.5836 (3)	0.0466 (13)	
H27A	0.2765	0.5598	0.5468	0.056*	
H27B	0.3913	0.5281	0.5678	0.056*	
C28	0.4074 (4)	0.5557 (4)	0.6752 (3)	0.0624 (16)	
H28A	0.3979	0.485	0.6824	0.094*	
H28B	0.3699	0.5977	0.6901	0.094*	
H28C	0.4868	0.5721	0.7123	0.094*	
C29	0.542 (4)	0.6602 (15)	0.4672 (19)	0.064 (3)	0.55 (2)
H29A	0.4819	0.6128	0.4502	0.077*	0.55 (2)
H29B	0.6146	0.6263	0.5077	0.077*	0.55 (2)
C30	0.5284 (16)	0.6963 (11)	0.3911 (10)	0.066 (3)	0.55 (2)
H30A	0.4919	0.6449	0.3458	0.079*	0.55 (2)
H30B	0.6012	0.7139	0.4062	0.079*	0.55 (2)

C31	0.4556 (16)	0.7877 (16)	0.3635 (13)	0.073 (5)	0.55 (2)
H31A	0.4697	0.8363	0.3343	0.088*	0.55 (2)
H31B	0.3758	0.7691	0.325	0.088*	0.55 (2)
C32	0.488 (5)	0.8294 (17)	0.4431 (19)	0.068 (7)	0.55 (2)
H32A	0.5427	0.8838	0.4634	0.081*	0.55 (2)
H32B	0.4218	0.857	0.4344	0.081*	0.55 (2)
C29A	0.541 (5)	0.6650 (18)	0.461 (2)	0.056 (3)	0.45 (2)
H29C	0.5087	0.604	0.4656	0.067*	0.45 (2)
H29D	0.6191	0.6509	0.4857	0.067*	0.45 (2)
C30A	0.474 (2)	0.6981 (14)	0.3695 (13)	0.075 (4)	0.45 (2)
H30C	0.4962	0.6617	0.3397	0.091*	0.45 (2)
H30D	0.393	0.6895	0.3382	0.091*	0.45 (2)
C31A	0.506 (2)	0.8069 (16)	0.3820 (17)	0.063 (4)	0.45 (2)
H31C	0.4502	0.8462	0.3304	0.076*	0.45 (2)
H31D	0.5797	0.8153	0.3959	0.076*	0.45 (2)
C32A	0.510 (6)	0.839 (2)	0.455 (3)	0.064 (7)	0.45 (2)
H32C	0.5683	0.8906	0.4901	0.077*	0.45 (2)
H32D	0.4375	0.867	0.4346	0.077*	0.45 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mo1	0.0421 (2)	0.0281 (2)	0.0318 (2)	0.00186 (18)	0.02382 (17)	0.0012 (2)
Cl1	0.0978 (11)	0.0283 (8)	0.0685 (10)	0.0094 (7)	0.0575 (9)	0.0041 (7)
Cl2	0.0565 (7)	0.0651 (8)	0.0343 (6)	-0.0004 (7)	0.0297 (6)	0.0013 (8)
Cl3	0.0583 (8)	0.0320 (7)	0.0577 (9)	0.0070 (6)	0.0372 (7)	0.0059 (6)
P1	0.0426 (8)	0.0390 (8)	0.0446 (8)	-0.0019 (5)	0.0281 (7)	-0.0007 (6)
P2	0.0405 (7)	0.0363 (8)	0.0323 (8)	0.0025 (5)	0.0220 (6)	-0.0010 (6)
O1	0.0568 (19)	0.0352 (18)	0.0328 (16)	0.0028 (17)	0.0288 (15)	0.0022 (19)
C1	0.057 (4)	0.048 (4)	0.045 (3)	-0.008 (2)	0.032 (3)	-0.005 (3)
C2	0.076 (4)	0.081 (5)	0.086 (5)	-0.033 (3)	0.060 (4)	-0.042 (4)
C3	0.109 (6)	0.114 (6)	0.105 (6)	-0.055 (5)	0.083 (5)	-0.069 (5)
C4	0.088 (5)	0.074 (5)	0.069 (5)	-0.040 (4)	0.039 (4)	-0.031 (4)
C5	0.055 (4)	0.062 (5)	0.075 (5)	-0.018 (3)	0.031 (4)	-0.012 (4)
C6	0.060 (4)	0.057 (4)	0.080 (5)	-0.013 (3)	0.046 (4)	-0.011 (3)
C7	0.049 (3)	0.046 (4)	0.053 (3)	-0.004 (2)	0.036 (3)	-0.004 (3)
C8	0.154 (5)	0.060 (3)	0.093 (4)	0.013 (3)	0.099 (4)	0.013 (3)
C9	0.100 (5)	0.096 (6)	0.086 (5)	0.038 (4)	0.073 (4)	0.009 (4)
C10	0.114 (5)	0.085 (6)	0.089 (5)	0.006 (4)	0.087 (5)	0.007 (4)
C11	0.154 (5)	0.060 (3)	0.093 (4)	0.013 (3)	0.099 (4)	0.013 (3)
C12	0.204 (8)	0.041 (4)	0.116 (6)	0.005 (4)	0.133 (6)	0.006 (4)
C13	0.048 (3)	0.049 (4)	0.066 (4)	0.003 (2)	0.033 (3)	0.014 (3)
C14	0.074 (4)	0.084 (5)	0.067 (5)	0.013 (3)	0.031 (4)	0.029 (4)
C15	0.046 (3)	0.042 (4)	0.033 (3)	0.007 (2)	0.024 (3)	0.002 (2)
C16	0.051 (3)	0.067 (4)	0.057 (3)	0.003 (3)	0.036 (3)	-0.008 (3)
C17	0.058 (4)	0.086 (5)	0.064 (4)	0.018 (3)	0.044 (4)	0.009 (4)
C18	0.072 (4)	0.069 (5)	0.061 (4)	0.028 (3)	0.047 (4)	0.010 (3)
C19	0.079 (5)	0.070 (5)	0.056 (4)	0.011 (3)	0.044 (4)	-0.011 (3)

C20	0.051 (3)	0.061 (4)	0.042 (3)	0.004 (3)	0.028 (3)	-0.005 (3)
C21	0.040 (3)	0.044 (3)	0.036 (3)	0.002 (2)	0.022 (3)	-0.006 (2)
C22	0.054 (3)	0.050 (4)	0.048 (3)	0.005 (2)	0.033 (3)	0.001 (3)
C23	0.060 (4)	0.072 (5)	0.035 (3)	0.003 (3)	0.024 (3)	-0.012 (3)
C24	0.056 (4)	0.089 (5)	0.030 (3)	0.013 (3)	0.020 (3)	-0.005 (3)
C25	0.067 (4)	0.061 (4)	0.047 (4)	0.022 (3)	0.030 (3)	0.013 (3)
C26	0.076 (4)	0.047 (4)	0.047 (4)	0.014 (3)	0.034 (3)	0.002 (3)
C27	0.051 (3)	0.040 (3)	0.052 (3)	-0.004 (2)	0.034 (3)	0.001 (3)
C28	0.077 (4)	0.059 (4)	0.058 (4)	-0.004 (3)	0.045 (3)	0.014 (3)
C29	0.089 (5)	0.066 (4)	0.053 (5)	-0.004 (4)	0.052 (5)	-0.013 (4)
C30	0.084 (6)	0.076 (5)	0.053 (6)	-0.004 (5)	0.051 (6)	-0.011 (4)
C31	0.086 (14)	0.083 (10)	0.040 (9)	-0.014 (9)	0.035 (11)	0.004 (7)
C32	0.094 (19)	0.057 (6)	0.049 (8)	0.010 (8)	0.043 (12)	0.020 (6)
C29A	0.086 (5)	0.055 (5)	0.052 (5)	-0.005 (4)	0.056 (5)	-0.012 (4)
C30A	0.094 (7)	0.077 (6)	0.058 (6)	-0.003 (6)	0.049 (6)	-0.014 (5)
C31A	0.076 (14)	0.085 (10)	0.036 (10)	-0.011 (9)	0.039 (12)	0.003 (7)
C32A	0.092 (19)	0.055 (6)	0.047 (8)	0.008 (8)	0.044 (12)	0.020 (6)

Geometric parameters (Å, °)

Mo1—O1	2.206 (3)	C16—H16A	0.95
Mo1—Cl1	2.3871 (13)	C17—C18	1.356 (8)
Mo1—Cl2	2.3822 (13)	C17—H17A	0.95
Mo1—Cl3	2.4126 (13)	C18—C19	1.348 (7)
Mo1—P1	2.5964 (14)	C18—H18A	0.95
Mo1—P2	2.5974 (13)	C19—C20	1.371 (7)
P1—C7	1.824 (6)	C19—H19A	0.95
P1—C1	1.830 (5)	C20—H20A	0.95
P1—C13	1.840 (5)	C21—C26	1.364 (7)
P2—C27	1.818 (5)	C21—C22	1.370 (7)
P2—C21	1.830 (5)	C22—C23	1.368 (7)
P2—C15	1.838 (5)	C22—H22A	0.95
O1—C29	1.434 (15)	C23—C24	1.358 (8)
O1—C29A	1.453 (16)	C23—H23A	0.95
O1—C32	1.455 (15)	C24—C25	1.326 (7)
O1—C32A	1.459 (17)	C24—H24A	0.95
C1—C2	1.366 (7)	C25—C26	1.380 (7)
C1—C6	1.389 (7)	C25—H25A	0.95
C2—C3	1.390 (8)	C26—H26A	0.95
C2—H2A	0.95	C27—C28	1.514 (7)
C3—C4	1.358 (9)	C27—H27A	0.99
C3—H3A	0.95	C27—H27B	0.99
C4—C5	1.359 (9)	C28—H28A	0.98
C4—H4A	0.95	C28—H28B	0.98
C5—C6	1.355 (8)	C28—H28C	0.98
C5—H5A	0.95	C29—C30	1.499 (17)
C6—H6A	0.95	C29—H29A	0.99
C7—C12	1.358 (8)	C29—H29B	0.99

C7—C8	1.373 (8)	C30—C31	1.510 (16)
C8—C9	1.364 (9)	C30—H30A	0.99
C8—H8A	0.95	C30—H30B	0.99
C9—C10	1.352 (9)	C31—C32	1.464 (16)
C9—H9A	0.95	C31—H31A	0.99
C10—C11	1.341 (8)	C31—H31B	0.99
C10—H10A	0.95	C32—H32A	0.99
C11—C12	1.381 (9)	C32—H32B	0.99
C11—H11A	0.95	C29A—C30A	1.514 (18)
C12—H12A	0.95	C29A—H29C	0.99
C13—C14	1.498 (8)	C29A—H29D	0.99
C13—H13A	0.99	C30A—C31A	1.506 (17)
C13—H13B	0.99	C30A—H30C	0.99
C14—H14A	0.98	C30A—H30D	0.99
C14—H14B	0.98	C31A—C32A	1.509 (18)
C14—H14C	0.98	C31A—H31C	0.99
C15—C16	1.359 (7)	C31A—H31D	0.99
C15—C20	1.371 (7)	C32A—H32C	0.99
C16—C17	1.391 (7)	C32A—H32D	0.99
O1—Mo1—C12	179.21 (10)	C18—C17—H17A	120.3
O1—Mo1—C11	88.22 (10)	C16—C17—H17A	120.3
C12—Mo1—C11	92.44 (5)	C19—C18—C17	120.3 (5)
O1—Mo1—C13	88.34 (10)	C19—C18—H18A	119.8
C12—Mo1—C13	91.01 (5)	C17—C18—H18A	119.8
C11—Mo1—C13	176.50 (6)	C18—C19—C20	120.8 (6)
O1—Mo1—P1	89.05 (9)	C18—C19—H19A	119.6
C12—Mo1—P1	90.49 (5)	C20—C19—H19A	119.6
C11—Mo1—P1	91.92 (5)	C19—C20—C15	119.8 (5)
C13—Mo1—P1	88.66 (4)	C19—C20—H20A	120.1
O1—Mo1—P2	89.76 (8)	C15—C20—H20A	120.1
C12—Mo1—P2	90.67 (4)	C26—C21—C22	117.8 (5)
C11—Mo1—P2	90.94 (5)	C26—C21—P2	120.3 (4)
C13—Mo1—P2	88.41 (4)	C22—C21—P2	121.8 (4)
P1—Mo1—P2	176.87 (5)	C23—C22—C21	120.8 (5)
C7—P1—C1	102.7 (2)	C23—C22—H22A	119.6
C7—P1—C13	103.8 (3)	C21—C22—H22A	119.6
C1—P1—C13	101.9 (3)	C24—C23—C22	120.4 (6)
C7—P1—Mo1	113.20 (16)	C24—C23—H23A	119.8
C1—P1—Mo1	119.23 (19)	C22—C23—H23A	119.8
C13—P1—Mo1	114.15 (18)	C25—C24—C23	119.6 (6)
C27—P2—C21	104.1 (2)	C25—C24—H24A	120.2
C27—P2—C15	102.5 (2)	C23—C24—H24A	120.2
C21—P2—C15	101.7 (2)	C24—C25—C26	120.9 (6)
C27—P2—Mo1	114.84 (16)	C24—C25—H25A	119.6
C21—P2—Mo1	111.36 (16)	C26—C25—H25A	119.6
C15—P2—Mo1	120.40 (17)	C21—C26—C25	120.6 (6)
C29—O1—C32	108.9 (15)	C21—C26—H26A	119.7

C29A—O1—C32	104.0 (14)	C25—C26—H26A	119.7
C29—O1—C32A	113.6 (15)	C28—C27—P2	113.6 (4)
C29A—O1—C32A	108.5 (16)	C28—C27—H27A	108.8
C29—O1—Mo1	123.2 (9)	P2—C27—H27A	108.8
C29A—O1—Mo1	128.2 (10)	C28—C27—H27B	108.8
C32—O1—Mo1	127.6 (11)	P2—C27—H27B	108.8
C32A—O1—Mo1	122.9 (12)	H27A—C27—H27B	107.7
C2—C1—C6	118.3 (5)	C27—C28—H28A	109.5
C2—C1—P1	121.5 (4)	C27—C28—H28B	109.5
C6—C1—P1	120.2 (4)	H28A—C28—H28B	109.5
C1—C2—C3	121.1 (6)	C27—C28—H28C	109.5
C1—C2—H2A	119.5	H28A—C28—H28C	109.5
C3—C2—H2A	119.5	H28B—C28—H28C	109.5
C4—C3—C2	119.2 (7)	O1—C29—C30	105.5 (14)
C4—C3—H3A	120.4	O1—C29—H29A	110.6
C2—C3—H3A	120.4	C30—C29—H29A	110.6
C5—C4—C3	120.0 (6)	O1—C29—H29B	110.6
C5—C4—H4A	120	C30—C29—H29B	110.6
C3—C4—H4A	120	H29A—C29—H29B	108.8
C6—C5—C4	121.3 (6)	C29—C30—C31	103.3 (18)
C6—C5—H5A	119.3	C29—C30—H30A	111.1
C4—C5—H5A	119.3	C31—C30—H30A	111.1
C5—C6—C1	120.1 (6)	C29—C30—H30B	111.1
C5—C6—H6A	120	C31—C30—H30B	111.1
C1—C6—H6A	120	H30A—C30—H30B	109.1
C12—C7—C8	115.8 (6)	C32—C31—C30	104.0 (19)
C12—C7—P1	121.3 (5)	C32—C31—H31A	111
C8—C7—P1	122.7 (4)	C30—C31—H31A	111
C9—C8—C7	122.1 (7)	C32—C31—H31B	111
C9—C8—H8A	118.9	C30—C31—H31B	111
C7—C8—H8A	118.9	H31A—C31—H31B	109
C10—C9—C8	120.7 (7)	O1—C32—C31	107.4 (17)
C10—C9—H9A	119.7	O1—C32—H32A	110.2
C8—C9—H9A	119.7	C31—C32—H32A	110.2
C11—C10—C9	118.7 (7)	O1—C32—H32B	110.2
C11—C10—H10A	120.7	C31—C32—H32B	110.2
C9—C10—H10A	120.7	H32A—C32—H32B	108.5
C10—C11—C12	120.4 (6)	O1—C29A—C30A	105.3 (18)
C10—C11—H11A	119.8	O1—C29A—H29C	110.7
C12—C11—H11A	119.8	C30A—C29A—H29C	110.7
C7—C12—C11	122.2 (7)	O1—C29A—H29D	110.7
C7—C12—H12A	118.9	C30A—C29A—H29D	110.7
C11—C12—H12A	118.9	H29C—C29A—H29D	108.8
C14—C13—P1	113.7 (4)	C31A—C30A—C29A	100 (2)
C14—C13—H13A	108.8	C31A—C30A—H30C	111.8
P1—C13—H13A	108.8	C29A—C30A—H30C	111.8
C14—C13—H13B	108.8	C31A—C30A—H30D	111.8
P1—C13—H13B	108.8	C29A—C30A—H30D	111.8

H13A—C13—H13B	107.7	H30C—C30A—H30D	109.5
C13—C14—H14A	109.5	C30A—C31A—C32A	104 (2)
C13—C14—H14B	109.5	C30A—C31A—H31C	110.9
H14A—C14—H14B	109.5	C32A—C31A—H31C	110.9
C13—C14—H14C	109.5	C30A—C31A—H31D	110.9
H14A—C14—H14C	109.5	C32A—C31A—H31D	110.9
H14B—C14—H14C	109.5	H31C—C31A—H31D	108.9
C16—C15—C20	119.4 (5)	O1—C32A—C31A	105 (2)
C16—C15—P2	119.5 (4)	O1—C32A—H32C	110.7
C20—C15—P2	121.1 (4)	C31A—C32A—H32C	110.7
C15—C16—C17	120.3 (6)	O1—C32A—H32D	110.7
C15—C16—H16A	119.8	C31A—C32A—H32D	110.7
C17—C16—H16A	119.8	H32C—C32A—H32D	108.8
C18—C17—C16	119.4 (6)		
O1—Mo1—P1—C7	11.0 (2)	P1—C7—C8—C9	176.5 (5)
Cl2—Mo1—P1—C7	-169.7 (2)	C7—C8—C9—C10	-0.5 (12)
Cl1—Mo1—P1—C7	-77.2 (2)	C8—C9—C10—C11	-1.4 (11)
Cl3—Mo1—P1—C7	99.3 (2)	C9—C10—C11—C12	2.3 (11)
O1—Mo1—P1—C1	131.9 (2)	C8—C7—C12—C11	-0.5 (11)
Cl2—Mo1—P1—C1	-48.8 (2)	P1—C7—C12—C11	-175.6 (6)
Cl1—Mo1—P1—C1	43.7 (2)	C10—C11—C12—C7	-1.4 (12)
Cl3—Mo1—P1—C1	-139.8 (2)	C7—P1—C13—C14	168.8 (4)
O1—Mo1—P1—C13	-107.4 (2)	C1—P1—C13—C14	62.4 (5)
Cl2—Mo1—P1—C13	71.9 (2)	Mo1—P1—C13—C14	-67.6 (5)
Cl1—Mo1—P1—C13	164.4 (2)	C27—P2—C15—C16	-58.1 (5)
Cl3—Mo1—P1—C13	-19.1 (2)	C21—P2—C15—C16	49.4 (5)
O1—Mo1—P2—C27	108.1 (2)	Mo1—P2—C15—C16	173.0 (4)
Cl2—Mo1—P2—C27	-71.3 (2)	C27—P2—C15—C20	122.9 (4)
Cl1—Mo1—P2—C27	-163.7 (2)	C21—P2—C15—C20	-129.7 (4)
Cl3—Mo1—P2—C27	19.7 (2)	Mo1—P2—C15—C20	-6.1 (5)
O1—Mo1—P2—C21	-9.9 (2)	C20—C15—C16—C17	-0.2 (8)
Cl2—Mo1—P2—C21	170.76 (18)	P2—C15—C16—C17	-179.3 (4)
Cl1—Mo1—P2—C21	78.31 (18)	C15—C16—C17—C18	1.8 (9)
Cl3—Mo1—P2—C21	-98.26 (18)	C16—C17—C18—C19	-1.9 (9)
O1—Mo1—P2—C15	-128.8 (2)	C17—C18—C19—C20	0.3 (9)
Cl2—Mo1—P2—C15	51.90 (18)	C18—C19—C20—C15	1.3 (9)
Cl1—Mo1—P2—C15	-40.55 (18)	C16—C15—C20—C19	-1.3 (8)
Cl3—Mo1—P2—C15	142.89 (18)	P2—C15—C20—C19	177.8 (4)
Cl1—Mo1—O1—C29	-180 (2)	C27—P2—C21—C26	150.8 (5)
Cl3—Mo1—O1—C29	0 (2)	C15—P2—C21—C26	44.6 (5)
P1—Mo1—O1—C29	89 (2)	Mo1—P2—C21—C26	-84.9 (5)
P2—Mo1—O1—C29	-89 (2)	C27—P2—C21—C22	-33.3 (5)
Cl1—Mo1—O1—C29A	179 (3)	C15—P2—C21—C22	-139.5 (4)
Cl3—Mo1—O1—C29A	-2 (3)	Mo1—P2—C21—C22	91.0 (4)
P1—Mo1—O1—C29A	87 (3)	C26—C21—C22—C23	-0.1 (8)
P2—Mo1—O1—C29A	-90 (3)	P2—C21—C22—C23	-176.1 (4)
Cl1—Mo1—O1—C32	-6 (3)	C21—C22—C23—C24	-0.6 (9)

C13—Mo1—O1—C32	173 (3)	C22—C23—C24—C25	0.7 (9)
P1—Mo1—O1—C32	-98 (3)	C23—C24—C25—C26	-0.1 (9)
P2—Mo1—O1—C32	85 (3)	C22—C21—C26—C25	0.7 (8)
C11—Mo1—O1—C32A	7 (4)	P2—C21—C26—C25	176.7 (4)
C13—Mo1—O1—C32A	-173 (4)	C24—C25—C26—C21	-0.6 (9)
P1—Mo1—O1—C32A	-85 (4)	C21—P2—C27—C28	-169.0 (4)
P2—Mo1—O1—C32A	98 (4)	C15—P2—C27—C28	-63.3 (4)
C7—P1—C1—C2	128.5 (5)	Mo1—P2—C27—C28	69.0 (4)
C13—P1—C1—C2	-124.3 (5)	C32—O1—C29—C30	17 (4)
Mo1—P1—C1—C2	2.4 (6)	C32A—O1—C29—C30	5 (5)
C7—P1—C1—C6	-52.0 (5)	Mo1—O1—C29—C30	-168.8 (15)
C13—P1—C1—C6	55.3 (5)	O1—C29—C30—C31	-30 (4)
Mo1—P1—C1—C6	-178.0 (4)	C29—C30—C31—C32	33 (4)
C6—C1—C2—C3	-0.9 (10)	C29—O1—C32—C31	4 (5)
P1—C1—C2—C3	178.7 (6)	C29A—O1—C32—C31	6 (5)
C1—C2—C3—C4	-0.6 (12)	C32A—O1—C32—C31	120 (18)
C2—C3—C4—C5	1.6 (12)	Mo1—O1—C32—C31	-170.0 (17)
C3—C4—C5—C6	-1.2 (11)	C30—C31—C32—O1	-23 (4)
C4—C5—C6—C1	-0.3 (10)	C32—O1—C29A—C30A	-11 (4)
C2—C1—C6—C5	1.3 (9)	C32A—O1—C29A—C30A	-22 (5)
P1—C1—C6—C5	-178.2 (5)	Mo1—O1—C29A—C30A	165.2 (15)
C1—P1—C7—C12	-47.7 (6)	O1—C29A—C30A—C31A	38 (4)
C13—P1—C7—C12	-153.6 (5)	C29A—C30A—C31A—C32A	-39 (4)
Mo1—P1—C7—C12	82.1 (5)	C29—O1—C32A—C31A	-4 (6)
C1—P1—C7—C8	137.5 (5)	C29A—O1—C32A—C31A	-3 (6)
C13—P1—C7—C8	31.6 (6)	C32—O1—C32A—C31A	-73 (13)
Mo1—P1—C7—C8	-92.7 (5)	Mo1—O1—C32A—C31A	170 (2)
C12—C7—C8—C9	1.4 (10)	C30A—C31A—C32A—O1	27 (5)