


This is an Accepted Manuscript version of the following article, accepted for publication in **Supramolecular Chemistry**.  
Postprint of: Pomećko R., Bocheńska M., Hubscher-Bruder V., Zouhair A., Arnaud-Neu F., Anion recognition by phosphonium calix[4]arenes: synthesis and physico-chemical studies, *Supramolecular Chemistry*, Vol. 22, Iss. 5 (2010), pp. 275-288, DOI: [10.1080/10610270903437051](https://doi.org/10.1080/10610270903437051)  
It is deposited under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Anion recognition by phosphonium calix[4]arenes: synthesis and physico-chemical studies

Journal:	<i>Supramolecular Chemistry</i>
Manuscript ID:	GSCH-2009-0077.R3
Manuscript Type:	Full Paper
Date Submitted by the Author:	20-Oct-2009
Complete List of Authors:	Pomećko, Radosław; Gdansk University of Technology, Chemical technology Asfari, Zouhair; IPHC, CNRS, UDS, DSA Hubscher-Bruder, Véronique; IPHC, CNRS, UDS, DSA Bochenska, Maria; Gdansk University, Chemical technology Arnaud-Neu, Françoise; IPHC, CNRS, UDS, DSA
Keywords:	Phosphonium calix[4]arenes , anions, binding properties, microcalorimetry, ion selective electrodes
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
scheme 1.cdx scheme 2.cdx	



## Index abstract



Radoslaw Pomecko, Zouhair Asfari,  
Véronique Hubscher-Bruder, Maria  
Bochenska and Françoise Arnaud-Neu

Anion recognition by phosphonium  
calix[4]arenes: synthesis and physico-  
chemical studies

## Anion recognition by phosphonium calix[4]arenes : synthesis and physico-chemical studies

Radoslaw Pomecko,<sup>a,b</sup> Zouhair Asfari,<sup>a</sup> Véronique Hubscher-Bruder,<sup>a</sup> Maria Bochenska<sup>b\*</sup> and Françoise Arnaud-Neu<sup>a\*</sup>

<sup>a</sup>*IPHC-DISA, ULP, CNRS, ECPM, 25 rue Becquerel, 67087 Strasbourg Cedex 2, France; E-mail: farnaud@chimie.u-strasbg.fr*

<sup>b</sup>*Department of Chemical Technology, Chemical Faculty, Gdansk University of Technology, ul. Narutowicza 11/12, 80-264 Gdansk, Poland. E-mail: marboch@chem.pg.gda.pl*

*p*-tert-Butylcalix[4]arenes, in the cone conformation, di- and tetrasubstituted at the narrow rim with charged phosphonium groups, have been synthesized and characterized. Their interactions with a wide range of anions have been investigated in solution in chloroform and acetonitrile by means of <sup>1</sup>H and <sup>31</sup>P NMR and microcalorimetry (ITC). These compounds have also been incorporated as sensing material in PVC ion selective electrodes (ISE). The results showed that they interact strongly with the more lipophilic anions ClO<sub>4</sub><sup>-</sup>, SCN<sup>-</sup> and I<sup>-</sup>, in solution as in the electrode membranes. The origin of this selectivity is discussed and, in particular, the role of the salt counterion is examined.

**Keywords:** Phosphonium calix[4]arenes; anion binding properties; microcalorimetry; ion selective electrodes.

### INTRODUCTION

Anions play an important role in many biological processes, such as regulation of cell activity, synthesis of proteins, transport of hormones [1,2,3]. They are also frequently used in many industrial technologies, which very often generate an increase of the concentration of anions in the environment or even introduce anionic species which were unknown so far in ecosystems. The presence of these anions is crucial in environmental and medical concerns, as they are pollutants and may have harmful effects on living organisms and human health [4,5,6]. Therefore there is a need of fast and selective anion detection methods allowing real-time monitoring of anion concentration changes and of efficient clean up processes. Design of anion receptors for such applications remains a great challenge for chemists because they have to take into consideration the specific

1  
2  
3 anion properties, such as a large range of shapes and geometries, small electric charges vs.  
4 sizes, high free energies of solvation, and in some cases multiple oxidation states of the  
5 central atoms in oxoanions or pH dependence. In many artificial anion hosts, noncovalent  
6 interactions are responsible for host-guest recognition. They include electrostatic  
7 interactions, hydrogen bonding, hydrophobic effects, coordination to a metal ion or  
8 combinations of these interactions. The hosts can be neutral, containing urea [7,8,9,10],  
9 thiourea [11,12] or amide functions [13]. They can also be positively charged, containing  
10 pyridinium [14], polyammonium [15] or quaternary ammonium [16] binding sites.  
11 Calix[4]arenes [17,18] and porphyrins [19] are often used as scaffolds onto which these  
12 functional groups can be grafted. Calixpyrroles are also known as efficient anion receptors  
13 [20,21].

22  
23 Recently, we described the synthesis and characterization of a new calix[4]arene  
24 derivative (**5**) bearing four positively charged triphenyl phosphonium groups [22]. The  
25 presence of these highly polarisable moieties, where the charge is spread over the three  
26 aromatic rings, was expected to favour the interaction with lipophilic anions. Preliminary  
27 binding studies showed that this ligand interacted selectively with some anions, namely  
28  $\text{ClO}_4^-$  and  $\text{SCN}^-$ . This compound was also incorporated, as ionophore-sensing material, in  
29 ion selective electrodes (ISEs) which exhibited a selectivity order similar to the  
30 Hofmeister series. The disubstituted derivative **1** was also synthesised as the  
31 hexafluorophosphate [23].

32  
33 In order to get more information on the mechanisms involved in the recognition  
34 process and to optimize its selectivity, we have now extended this study to new di- and  
35 tetrasubstituted phosphonium calix[4]arene derivatives (compounds **2–4**, **6** and **7**) and re-  
36 examined the properties of **1** and **5** (Fig. 1). In some of these compounds, one of the  
37 phenyl rings on the phosphonium groups has been replaced by a methyl radical or a  
38 hydrogen atom. The presence of such small substituents is expected to increase the charge  
39 density on the phosphorus atoms and their accessibility [24]. Moreover, the presence of  
40 hydrogen atoms may induce the formation of hydrogen bonds with anions. The possibility  
41 of tuning the charge density on the phosphorus atoms by changing the nature of the their  
42 substituents could allow the design of receptors able to distinguish lipophilic anions like  
43  $\text{ClO}_4^-$ ,  $\text{SCN}^-$ ,  $\text{I}^-$  or  $\text{NO}_3^-$  from other anions and between them. The binding properties of  
44 these compounds towards a variety of anions have been followed by  $^1\text{H}$  and  $^{31}\text{P}$  NMR and  
45 by titration microcalorimetry (ITC). In particular the role of the salt counterion was  
46 examined using these techniques.

Fig. 1

## RESULTS AND DISCUSSION

### Synthesis

The ligands synthesised in this work (Fig. 1) are based on a tetrakis-*p-tert*-butylcalix[4]arene platform which presents several advantages. It has a well defined size and is readily amenable to substitution at its lower rim where ligating groups can be attached and in some extent preorganized. The amphiphilicity of such derivatives should allow their introduction into the membranes of ion selective electrodes.

Diphosphonium ligands were synthesised in two or three steps according to the known procedure [23]. The first step was the selective bromoalkylation of the tetrakis-*p-tert*-butylcalix[4]arene (**S**<sub>1</sub>), leading to the intermediate molecules **S**<sub>2</sub> or **S**<sub>3</sub> in 71 % and 55 % yield, respectively (Scheme 1). 1,3-bis-(4-triphenylphosphonium-butoxy)-*p-tert*-butylcalix[4]arene dibromide (**1**) was obtained by the reaction of **S**<sub>2</sub> with 10 equivalents of triphenylphosphine. After 6 days under reflux in chloroform, the product was precipitated from a dichloromethane/hexane mixture in 76% yield. 1,3-bis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-*p-tert*-butylcalix[4]arene dibromide (**2**) was obtained during the reaction of **S**<sub>2</sub> with 10 equivalents of diphenylmethylphosphine in the same conditions in 69% yield.

#### Scheme 1

The more lipophilic molecules **3** and **4**, where the free phenolic protons are substituted with propyl groups, were prepared in order to increase their stability in the lipophilic membrane of ion selective electrodes. The reaction of **S**<sub>3</sub> with 10 equivalents of 1,4-dibromobutyl, refluxed for 4 days in dimethylformamide in the presence of 7 equivalents of NaH gave the 1,3-bis-(propoxy)-2,4-bis-(butoxy-4-bromide)-tetrakis-*p-tert*-butylcalix[4]arene (**S**<sub>4</sub>) in 57% yield. Compounds **3** and **4** were obtained by the reaction of **S**<sub>4</sub> with 10 equivalents of triphenylphosphine and 10 equivalents of diphenylmethylphosphine in chloroform, in 76% and 64% yield, respectively (Scheme 1).

Tetraphosphonium ligands **6** and **7** were synthesised according to the procedure already described for **5** [22] from the intermediate molecule tetrakis-(butoxy-4-bromide)-*p-tert*-butylcalix[4]arene (**S**<sub>5</sub>) by substitution of the bromine atoms of the alkyl chains (Scheme 2). The tetrakis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-*p-tert*-butylcalix[4]arene tetrabromide (**6**) was obtained from the reaction of **S**<sub>5</sub> with 20 equivalents of diphenylmethylphosphine in chloroform with 59% yield.

## Scheme 2

Ligand **7** was synthesised in three steps. The reaction of **S<sub>5</sub>** with 5 equivalents of  $\text{KPPH}_2$  gave the product **S<sub>6</sub>** in 40% yield [25,26,27]. **S<sub>6</sub>** was then protonated with an excess of HBr giving ligand **7** in 52 % yield.

In order to study the influence of the ligand counterion on the ligand – anion interactions, the tetra substituted phosphonium ligands were also synthesised as perchlorates (**5a**, **6a**) and hexafluorophosphates (**5b**, **6b**). These compounds were obtained by reacting ligands **5** and **6** with the appropriate silver salts.

For comparison purpose, the monomeric subunit of ligand **5**, the triphenylphosphonium-butoxy-*p-tert*-butylphenol bromide (**8**) was synthesized as previously described [22].

The cone conformation of the di-substituted calix[4]arenes was indicated by the presence in their  $^1\text{H}$  NMR spectra of two singlets for the *tert*-butyl protons (at 0.99 and 1.28 ppm for **1** and 0.96 and 1.29 ppm for **2**) and a AB system for the methylene protons (at 3.19 and 3.90 ppm for **1** and at 3.23 and 4.02 ppm for **2**). The  $^1\text{H}$  NMR spectra of ligands **5** and **6** are characteristic of tetrasubstituted derivatives of calix[4]arenes in the cone conformation. For instance in the case of **6**, it is indicated by the presence of the singlet corresponding to the protons of the *tert*-butyl groups at 1.03 ppm, the AB system of the methylene protons of the calixarene at 4.26 and 3.03 ppm. Two well defined multiplets for the protons of the aromatic phosphonium groups can be observed as well as one doublet for the methyl protons in direct neighbourhood of the phosphorus atoms.

The spectrum of the protonated phosphonium ligand **7** presents several broad peaks corresponding to multiplets in coalescence. Only two singlets could be clearly observed, one for the *tert*-butyl groups and one for the aromatic protons of the calix[4]arene, suggesting also the cone conformation of this molecule.

## Binding studies

### $^1\text{H}$ and $^{31}\text{P}$ NMR studies in chloroform

The interactions between the phosphonium calixarenes and the following anions:  $\text{NO}_3^-$ ,  $\text{ClO}_4^-$ ,  $\text{I}^-$ ,  $\text{SCN}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{HCO}_3^-$ ,  $\text{Cr}_2\text{O}_7^{2-}$  provided as sodium salts were studied by  $^1\text{H}$  and  $^{31}\text{P}$  NMR in  $\text{CDCl}_3$ . Among these anions, only perchlorate, thiocyanate and iodide salts induced changes in the  $^1\text{H}$  NMR spectra of the ligands. The changes observed in the case of the disubstituted ligand **1** upon addition of sodium perchlorate are illustrated in Fig. 2. The main signals affected were those of the  $\text{CH}_2$  protons (i, j) directly bound to the carbon

atoms next to the phosphorus atoms and the aromatic protons (k) of the phosphonium groups (see Table S1, Supplementary Online Material). The shifts of these signals due to changes in the charge density on the phosphorus atoms reflect interactions with these anions.

*Figure 2*

Similar but smaller changes were induced in the spectrum of ligand **2** by the presence of the same anions (see Table S2, Supplementary Online Material). As with ligand **1**, no change is observed for the aromatic protons (c, d) and for the protons (e, f) of the methylene bridge, indicating that the conformation of the calixarene unit is not disturbed. This may be explained by the high rigidity of these disubstituted derivatives due to hydrogen bonds involving the free phenolic groups.

In the  $^{31}\text{P}$  NMR spectra of the free ligands **1** and **2**, and of these ligands in the presence of sodium iodide, thiocyanate and perchlorate, the phosphorus atoms appear as a singlet indicating that the two phosphonium groups are chemically equivalent and participate in the anion-ligand interaction. The most important changes in chemical shifts are observed for perchlorate:  $\Delta\delta = 0.602$  and  $0.620$  ppm with **1** and **2**, respectively (Table 1).

*Table 1*

With the tetrasubstituted calix[4]arene **5** previously studied [22] and  $\text{ClO}_4^-$ ,  $\text{SCN}^-$  and  $\text{I}^-$  the most important changes of chemical shifts corresponded to the signals of the methylene bridge protons and the aromatic protons of the calixarene, indicating changes in the conformational tensions of the calixarene scaffold. The signals corresponding to protons of the butyl chains were also shifted as well as the aromatic protons of the phosphonium. On the contrary no change was observed with the monomer **8** [22].

The spectrum of ligand **6** was also modified in the presence of these anions (see Table S3, Supplementary Online Material). In particular the signals corresponding to the protons of the calix[4]arene scaffold are moved in a comparable way for the three anions. The changes observed in the spectrum of this ligand in the presence of  $\text{ClO}_4^-$  are illustrated in Fig. 3.

*Figure 3*

With this anion, the multiplet corresponding to the protons (g) and (j) from 3.87 to 3.70 ppm gives two multiplets from 4.36 to 4.17 ppm for (g) and from 3.28 to 3.11 ppm for (j). The multiplet from 8.17 to 8.02 ppm for protons (k) is shifted upfield and gives one multiplet from 7.90 to 7.70 ppm, whereas the multiplet from 7.78 to 7.54 ppm of protons (l, m) does not move significantly. The doublet of the  $\text{CH}_3$  protons (o) adjacent to the phosphorus atoms at 2.91 ppm is strongly shifted to 2.50 ppm.

1  
2  
3 The addition of the anions studied to **7** did not induce any change in its  $^1\text{H}$  NMR  
4 spectrum. Especially the signals of the protons of the phosphonium moieties, expected to be  
5 involved in hydrogen bond formation, were not shifted.  
6  
7

8  
9 With ligand **6** as with **5**, only one singlet for the phosphorus atoms was detected in its  $^{31}\text{P}$   
10 NMR spectrum indicating all the four phosphonium groups being chemically equivalent. In  
11 the presence of sodium perchlorate and thiocyanate, the changes in chemical shifts are larger  
12 than those observed with the disubstituted derivatives and suggest stronger interactions (Table  
13 1). With all ligands the most important values were observed for perchlorate.  
14  
15  
16  
17

### 18 19 $^1\text{H}$ NMR studies in acetonitrile

20  
21  $^1\text{H}$  NMR experiments were repeated with **5** and sodium perchlorate in deuterated  
22 acetonitrile, a more dissociating solvent, in which association phenomena are not as  
23 important as in chloroform [22]. The spectra of the ligand are very similar in both  
24 solvents. In acetonitrile the addition of  $\text{NaClO}_4$  induces shifts of the same signals as in  
25 chloroform. Moreover this study showed the influence of the counterion of the salts on the  
26 shifts observed in the spectra (See Table S4, Supplementary Online Material) . The signal  
27 of the methylene protons (g) adjacent to the phenolic oxygen atoms was shifted only with  
28  $\text{NaClO}_4$  and  $\text{LiClO}_4$ . With both salts the shifts corresponding to the signals of the protons  
29 (j) next to the phosphorus atoms (which are supposed to interact with perchlorate) were  
30 similar, whereas with  $\text{Et}_4\text{NClO}_4$  and especially with  $\text{CsClO}_4$ , the values were very small.  
31 The changes observed suggested that ligand – anion interaction was connected with the  
32 nature of the counterion and its affinity for the ligand.  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 In order to observe the influence of the ligand counterion, bromide anions were  
43 replaced by the more lipophilic perchlorate (**5a** and **6a**) or hexafluorophosphate (**5b** and  
44 **6b**) anions. The chemical shifts ( $\delta$ ) of selected protons, given in Table 2, show only slight  
45 differences for protons (c, e, f) and (g) close to the calix[4]arene scaffold ( $\Delta\delta$  in the range  
46 0.02 – 0.07 ppm). In contrast, the signals of the protons (j) next to the charged phosphorus  
47 atoms are greatly shifted ( $\Delta\delta = -0.64$  ppm for **5a**,  $-0.75$  ppm for **5b** and  $\Delta\delta = -0.73$  ppm  
48 for **6a**,  $-0.83$  ppm for **6b**). With ligands **6a** and **6b**, the signals of the protons (o) of the  
49 methyl substituents are also displaced ( $\Delta\delta = -0.41$  ppm for **6a** and  $-0.46$  ppm for **6b**).  
50 These results show that the chemical shifts of the protons next to the phosphonium groups  
51 are influenced by the ligand counterion.  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 2



1  
2  
3 On the other hand, it was also shown that when the lipophilic  $\text{PF}_6^-$  anions were  
4 replacing the  $\text{Br}^-$  counterions of the ligand, there was still a significant shift of the protons  
5 (j) close to the phosphorus atoms for ligand **5b** in the presence of  $\text{ClO}_4^-$  (Table S4) [22].  
6  
7 This result suggested complexation of perchlorate with this ligand, where no exchange is  
8 normally possible.  
9  
10  
11

### 12 13 14 **Microcalorimetric studies in acetonitrile**

15 In order to get more information on the influence of the counterion of the salt,  
16 microcalorimetric titrations were carried out with ligands **5** and **5b** against  $\text{NaClO}_4$ ,  
17  $\text{LiClO}_4$  and  $\text{Et}_4\text{NClO}_4$  in acetonitrile. The thermograms recorded during the titration of  
18 these ligands with  $\text{NaClO}_4$  and  $\text{LiClO}_4$  showed significant exothermic heat effects,  
19 whereas their titration with  $\text{Et}_4\text{NClO}_4$  led to no thermal effect (see Figure S1,  
20 Supplementary Online Material). For comparison purpose, the titration of the monomer **8**  
21 with  $\text{NaClO}_4$  was also carried out showing no significant heat effect.  
22  
23  
24  
25  
26  
27

28 If only the  $\text{ClO}_4^-$  anion were involved in the complexation, a similar heat effect  
29 should be observed in all the titrations. The fact that an effect is only observed for  $\text{NaClO}_4$   
30 and  $\text{LiClO}_4$  suggests that it is not only related to the anion interaction (complexation or  
31 anion exchange). Assuming that the large tetraethylammonium cation cannot be  
32 complexed with a calix[4]arene, the heat effects observed during the titration with  $\text{NaClO}_4$   
33 and  $\text{LiClO}_4$  would rather be due to the complexation of the cations with the two ligands.  
34 This is supported by the fact that no heat effect was detected with the monomer **8**,  
35 supposed to be unable to complex these cations.  
36  
37  
38  
39  
40  
41

42 Calorimetric data obtained with  $\text{NaClO}_4$  and  $\text{LiClO}_4$  were interpreted assuming  
43 different cation complexation models. With sodium and both ligands, the best fit was  
44 obtained by considering the presence of  $\text{ML}$  and  $\text{ML}_2$  complexes. The same species were  
45 found with  $\text{LiClO}_4$  and **5b**, whereas only a 1:1 complex was formed with **5**. The formation  
46 of  $\text{ML}_2$  species could be explained by the complexation of the ion pair  $\text{Na}^+\text{ClO}_4^-$  by two  
47 ligands. The stability constants of these complexes are given in Table S5 (see  
48 Supplementary Online Material).  
49  
50  
51  
52  
53  
54

55 An important heat effect was observed during the titration of ligands **5** and **5b**  
56 against  $\text{LiBr}$ , which should be related directly to cation complexation as no anion  
57 exchange is possible with these ligands. The data interpretation led to species of the same  
58 stoichiometry as with  $\text{LiClO}_4$  (Table S5). The values of the stability constants of 1:1  
59 species formed in the presence of  $\text{LiClO}_4$  and  $\text{LiBr}$  are comparable (with **5**,  $\log \beta = 3.24$   
60

1  
2  
3 and 3.15, respectively, and with **5b**,  $\log \beta = 4.29$  and 4.32, respectively). They are lower  
4 than that of the complex formed by the tetra-methylated *p-tert*-butylcalix[4]arene with  
5 lithium ( $\log \beta = 5.10$  in acetonitrile [28]).  
6  
7  
8  
9

10 All that considered, the UV spectrophotometric titrations of ligand **5** against  $\text{ClO}_4^-$   
11 previously performed may certainly be interpreted in terms of cation rather than anion  
12 complexation [22]. The values of the stability constants of the 1:1 complexes with sodium  
13 perchlorate ( $\log \beta = 3.81 \pm 0.02$ ) and with lithium perchlorate ( $\log \beta = 3.71 \pm 0.04$ ) [22] are  
14 of the same order of magnitude as those obtained from microcalorimetric measurements.  
15  
16  
17  
18  
19

### 20 21 22 **Potentiometric studies** 23

24 Only few ligands containing phosphorus atoms have been studied so far as active material in  
25 ion selective membrane electrodes [29,30,31]. They showed a selective response for  $\text{ClO}_4^-$ ,  
26 with, however, little discrimination with respect to  $\text{I}^-$  and  $\text{SCN}^-$ . The results suggested a  
27 particular affinity of ligands containing phosphorus atoms for  $\text{ClO}_4^-$ ,  $\text{SCN}^-$  and  $\text{I}^-$  anions. By  
28 attaching phosphonium moieties to a calix[4]arene scaffold and taking advantage of the  
29 preorganization of the ligand, it was expected to enhance the selectivity for tetrahedral or  
30 spherical anions. Such selectivity (especially for  $\text{ClO}_4^-$  over  $\text{I}^-$ ) is hard to obtain with other  
31 kinds of receptors.  
32  
33  
34  
35  
36  
37

38 Disubstituted phosphonium ligands **1** and **2** were tested as ionophores in the  
39 membrane electrodes. The electrodes were sensitive to perchlorate, thiocyanate, iodide  
40 and nitrate, showing fast, near-Nernstian responses (Figure 4 and Table S6 of  
41 Supplementary Online Material). However, their characteristics changed with time.  
42 Attempts to optimize the composition of the conditioning solutions as well as the  
43 conditioning time did not improve the situation which might be due to slow leakage of the  
44 ionophores from the membrane. These ligands had also the tendency to crystallize in the  
45 membrane phase. Crystallization of our ligands within the membranes depends strongly  
46 on the kind of plasticizer used. The ligands in the membranes based on bis-(2-  
47 ethylhexyl)sebacate (BEHS) has crystallized strongly, which decreased their stability.  
48 This phenomenon originates from the higher lipophilicity of this plasticizer ( $\log P = 10.1$ )  
49 as compared to 2-nitrophenyloctylether (*o*-NPOE) ( $\log P = 5.9$ ) [32], which is more  
50 suitable for charged ligands. Electrodes with membranes based on *o*-NPOE had the best  
51 lifetime and response characteristics and were chosen for further studies.  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Figure 4*

The more lipophilic ligands **3** and **4**, in which n-propyl chains replace the two phenolic OH groups, were also synthesised. The lifetime of electrodes incorporating these ligands for perchlorate was increased to at least three weeks. The repeatability of the measurements was also good, but their detection limits increased (see Table S7, Supplementary Online Material).

The membranes of electrodes incorporating the tetrasubstituted ligand **6** showed rather quick (within 15-20 s), stable and fully reversible responses (Figure 5 and Table S8 of Supplementary Online Material). The repeatability of the measurements was also good and their lifetime was more than three weeks. They showed close to Nernstian response to  $\text{ClO}_4^-$ ,  $\text{I}^-$ ,  $\text{NO}_3^-$  and  $\text{SCN}^-$  and no significant response for  $\text{SO}_4^{2-}$ ,  $\text{CO}_3^{2-}$ ,  $\text{HPO}_4^{2-}$ ,  $\text{PO}_4^{3-}$ . A similar behaviour was already observed with electrodes incorporating ligand **5** (Table S8). The highest selectivity was obtained for  $\text{ClO}_4^-$  ions in buffered solution (pH=5.5) and in water (pH=6.5). The over-Nernstian slope of the electrode response for  $\text{Cr}_2\text{O}_7^{2-}$  could indicate a mechanism where both processes, anion complexation and anion exchange, play an important role. It can also be explained as the presence in the sample of different forms of chromates.

*Figure 5*

While the addition of lipophilic anionic sites (KTCIPB) to the membranes in the case of ligand **5** did not change much the properties of the electrodes, it affected the properties of the electrodes containing **6** (Table 3). Without salt, the slope of the electrode is -54.2 mV, and slightly decreases to -51.8 and -49.9 mV, respectively, in the presence of salt. According to the literature data [29,31], such results suggest that none of the ligands works as a neutral carrier because the addition of the lipophilic anion to the membrane does not induce a cationic response of the potentiometric cell. Ligand **6** seems to work in the membrane as a typical anion-exchanger, whereas ligand **5** could be considered as a charged ligand despite the small influence of KTCIPB.

*Table 3*

The influence of the ligand counterions ( $\text{Br}^-$ ,  $\text{ClO}_4^-$ ,  $\text{PF}_6^-$ ) on the properties of the membrane electrodes was also studied. Table 4 compares the responses to perchlorate of electrodes based on ligands **5a**, **6a** (perchlorates) and on ligands **5b** and **6b** (hexafluorophosphates) to corresponding electrodes based on ligands **5** and **6** (bromides). The less good slope of the electrodes containing **6a** and **6b** as compared to that of electrodes containing **6** suggests rather the anion-exchange nature of the latter ligand. In

1  
2  
3 this case the presence of more lipophilic anions (perchlorate or hexafluorophosphate)  
4 slows down the anion exchange process. In contrast, the properties of electrodes with the  
5 bromide ligand **5** and with the perchlorate ligand **5a** are comparable. The presence of  
6 highly lipophilic perchlorate anions does not disturb the electrode response, while the  
7 presence of hexafluorophosphate anions in the membrane phase (electrode containing  
8 ligand **5b**) decreases the slope and the linearity range.  
9

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
*Table 4*

The poorer properties of the electrode containing **5b** and **6b** could be explained by the higher lipophilicity of hexafluorophosphate anions which hinders the process of anion exchange. The complexation of  $\text{ClO}_4^-$  by ligand **5** could explain the good response of electrodes based on **5** and **5a** to perchlorate. Such interpretation is consistent with the results of previous experiments (Table 4) and indicates that ligand **5** behaves more like a charged carrier for  $\text{ClO}_4^-$ , while ligand **6** behaves more as an anion-exchanger.

The order of selectivity observed with all phosphonium ligands **1-6** and **8** follows the Hofmeister series:



The highest selectivity is observed for perchlorate (Table 5).

*Table 5*

The electrodes based on tetraphosphonium derivatives show higher selectivities for perchlorate over thiocyanate, iodide and nitrate than those based on their di-phosphonium counterparts. These selectivities are also better than the selectivity of electrodes based on the monomer **8**.

The replacement of one phenyl substituent on the phosphorus atoms by one methyl group does not change significantly the selectivity pattern and the values of the selectivity coefficients of electrodes containing either di- or tetraphosphonium ligands. The only exception is the selectivity of electrodes based on compound **5** against  $\text{Cr}_2\text{O}_7^{2-}$  which increases from 2.6 to 3.2 log units.

Electrodes based on alkylated compounds **3** and **4** are also selective for perchlorate but the selectivity over iodide and thiocyanate is decreased (Table 5).

Ligands **5** and **6** display better potentiometric properties than the protonated cyclam [33] or its copper complex [34] and than a phosphodithia macrocycle [35]. For instance the detection limit is  $2.5 \times 10^{-7}$  M with ligand **5** when compared to  $4.2 \times 10^{-6}$  M for cyclam and  $8 \times 10^{-7}$  M for the phosphodithiamacrocycle. Ligands **6** as **5** presents generally higher selectivities than TDMACl [36], the protonated cyclam and  $[\text{Cu}(\text{cyclam})]^{2+}$ .

## CONCLUDING REMARKS

The different techniques used to assess the binding properties of phosphonium derivatives showed strong interactions with  $\text{SCN}^-$ ,  $\text{I}^-$  and especially  $\text{ClO}_4^-$  and pointed out the important role played by the salt counterion ( $\text{Na}^+$  or  $\text{Li}^+$ ), which may be complexed by the calix[4]arene. Incorporated in PVC membrane electrodes, these molecules are efficient sensing material for anions with a selectivity order following the Hofmeister series generally observed for ion exchangers. However, the electrodes based on tetraphosphonium derivatives showed better selectivities than those based on the diphosphonium analogues or on the monomeric unit, indicating the importance of the ligand preorganisation which should not be observed in the case of simple anion-exchangers.

A question which must be addressed concerns the nature of the interaction between the ligands and the anions, e.g.  $\text{ClO}_4^-$ . Is it a simple ion-exchange between the bromides of the ligand and this more lipophilic anion, or is it complexation within the charged phosphonium groups? What is the exact role of the salt counterion?

If NMR gives some indications on the changes in the molecule, suggesting interactions, it does not tell if there is complexation or anion exchange, since the nature of the counterion of the ligand has been shown to influence the chemical shifts of the protons near the charged atoms. In favour of ion-exchange is the fact that the most important shifts are observed with the more lipophilic anions  $\text{ClO}_4^-$ ,  $\text{SCN}^-$  and  $\text{I}^-$ . The behaviour in selective electrodes is consistent also with this assumption. However, the fact that no change occurred in the spectrum of the monomer, where only exchange is possible, is against this hypothesis. In favour of complexation is the fact that, in the presence of  $\text{NaClO}_4$ , strong shifts are observed for the signal of the protons next to the phosphonium groups in the spectrum of the hexafluorophosphate ligand where no exchange is possible. On the other hand  $^1\text{H}$  NMR and microcalorimetry emphasized the importance of the cation which can be complexed in the cavity of the calixarene. With perchlorate, the best interaction takes place with  $\text{Na}^+$  and  $\text{Li}^+$ , whereas little or no interaction occurs with the larger  $\text{Et}_4\text{N}^+$  and  $\text{Cs}^+$ . It can be noted also that the anion plays also a role in the complexation of the cation, since the spectrum of the calixarene part is not affected in the presence of  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{HCO}_3^-$  and  $\text{Cr}_2\text{O}_7^{2-}$ , i.e. the less lipophilic ones.

## EXPERIMENTAL

FAB mass spectra were obtained on a VG analytical ZAB HF instrument. All reagents and solvents were commercial and used without further purification.

Chromatography columns were prepared from Kieselgel Merck Si 60(40-63  $\mu\text{m}$ ). TLC was performed on 250  $\mu\text{m}$  silica gel plates Merck containing a fluorescent indicator.

### Synthesis of intermediate compounds

#### *1,3-bis-(butoxy-4-bromide)-p-tert-butylcalix[4]arene (S<sub>2</sub>)*

Into a 250  $\text{cm}^3$  flask containing (3.244 g, 5.00 mmol) of tetrakis-*p-tert*-butylcalix[4]arene **S<sub>1</sub>** and acetone (50  $\text{cm}^3$ ), (1.383 g, 10.00 mmol) of  $\text{K}_2\text{CO}_3$  were added. The mixture was stirred at room temperature for 2h. Then (3.236 g, 15.00 mmol) of 1,4-butyl-dibromide in acetone (50  $\text{cm}^3$ ) were added. The mixture was left for 4 days under reflux. After four days 5  $\text{cm}^3$  of methanol were added. The solvents were evaporated, and the reaction mixture was dissolved in 100  $\text{cm}^3$  of dichloromethane. After extraction with 150  $\text{cm}^3$  of water the organic phase was dried with  $\text{Na}_2\text{SO}_4$  and evaporated. The crude product was purified by crystallization from a 1/10 dichloromethane/methanol mixture giving (3.252 g, 3.54 mmol) of compound **S<sub>2</sub>** in 71 % yield. Mp > 280 °C.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 0.98 (s, 18H, C-( $\text{CH}_3$ )<sub>3</sub>), 1.30 (s, 18H, C-( $\text{CH}_3$ )<sub>3</sub>), 2.15 (qn, 4H,  $J = 4.4$ ,  $\text{CH}_2$ - $\text{CH}_2$ -O), 2.32 (qn, 4H,  $J = 4.2$ ,  $\text{CH}_2$ - $\text{CH}_2$ -Br), 3.32 (d, 4H,  $J = 13.0$ , Ar- $\text{CH}_2$ -Ar), 3.65 (t, 4H,  $J = 6.6$ ,  $\text{CH}_2$ - $\text{CH}_2$ -Br), 4.01 (t, 4H,  $J = 5.8$ ,  $\text{CH}_2$ - $\text{CH}_2$ -O), 4.25 (d, 4H,  $J = 13.0$ , Ar- $\text{CH}_2$ -Ar), 6.78 (s, 4H, Ar- $H$ ), 7.08 (s, 4H, Ar- $H$ ), 7.40 (s, 2H, OH). Anal. Calcd. for  $\text{C}_{52}\text{H}_{70}\text{O}_4\text{Br}_2$ : C, 67.97; H, 7.68. Found: C, 68.21; H, 7.94.

#### *1,3-bis-(propoxy)-p-tert-butylcalix[4]arene (S<sub>3</sub>)*

Into a 250  $\text{cm}^3$  flask containing (3.244g, 5.00 mmol) of tetrakis-*p-tert*-butylcalix[4]arene **S<sub>1</sub>** and acetone (50  $\text{cm}^3$ ), (1.383g, 10.00 mmol) of  $\text{K}_2\text{CO}_3$  were added. The mixture was stirred at room temperature for 2h. Then (1.845 g, 15.00 mmol) of bromopropane in acetone (40  $\text{cm}^3$ ) were added. The mixture was left for 4 days under reflux. After four days 5  $\text{cm}^3$  of methanol were added. The solvents were evaporated, and the reaction mixture was dissolved in 100  $\text{cm}^3$  dichloromethane. After extraction with 150  $\text{cm}^3$  of water the organic phase was dried with  $\text{Na}_2\text{SO}_4$  and evaporated. The crude product was purified by crystallization from a 1/9 acetone/methanol mixture giving (2.016 g, 2.75 mmol) of pure compound **S<sub>3</sub>** in 55 % yield. Mp > 280 °C.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.02 (s, 18H, C-( $\text{CH}_3$ )<sub>3</sub>), 1.27 (t, 6H,  $J = 5.0$ ,  $\text{CH}_3$ - $\text{CH}_2$ -), 1.28 (s, 18H, C-( $\text{CH}_3$ )<sub>3</sub>),

2.06 (sx, 4H,  $J = 4.9$ ,  $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-O}$ ), 3.32 (d, 4H,  $J = 12.8$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 3.96 (t, 4H,  $J = 5.9$ ,  $\text{O-CH}_2\text{-CH}_2$ ), 4.31 (d, 4H,  $J = 12.8$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 6.86 (s, 4H,  $\text{Ar-H}$ ), 7.05 (s, 4H,  $\text{Ar-H}$ ), 7.89 (s, 2H,  $\text{OH}$ ). Anal. Calcd for  $\text{C}_{50}\text{H}_{68}\text{O}_4$ : C, 81.92; H, 9.35. Found: C, 82.05; H, 9.40.

### 1,3-bis-(propoxy)-2,4-bis-(butoxy-4-bromide)-tetrakis-*p*-tert-butylcalix[4]arene (**S<sub>4</sub>**)

Into a 250 cm<sup>3</sup> flask containing (2.016 g, 2.75 mmol) of **S<sub>3</sub>** and DMF (50 cm<sup>3</sup>), (0.480 g, 20.00 mmol) of NaH were added. NaH was washed twice with hexane before addition. The mixture was stirred at room temperature for 4h. After this time, (5.940 g, 27.5 mmol) of 1,4-dibromobutyl in DMF (40 cm<sup>3</sup>) were added. The mixture was left for 2 days at 80-90 °C. After 2 days 30 cm<sup>3</sup> of methanol were added. The solvents were evaporated and the reaction mixture was dissolved in 100 cm<sup>3</sup> of dichloromethane. After extraction with 250 cm<sup>3</sup> of water the organic phase was dried with  $\text{Na}_2\text{SO}_4$  and evaporated. The crude product was purified by crystallization from a 1/10 dichloromethane/methanol mixture giving (1.584 g, 1.57 mmol) of compound **S<sub>4</sub>** in 57 % yield. Mp 170 °C. <sup>1</sup>H NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.01 (t, 6H,  $J = 5.2$ ,  $\text{CH}_3\text{-CH}_2\text{-}$ ), 1.05 (s, 18H,  $\text{C-(CH}_3)_3$ ), 1.12 (s, 18H,  $\text{C-(CH}_3)_3$ ), 1.99-2.05 (m, 4H,  $\text{CH}_3\text{-CH}_2\text{-}$ ), 2.07-2.25 (m, 8H,  $\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-Br}$ ), 3.13 (d, 4H,  $J = 13.0$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 3.51 (t, 4H,  $J = 6.6$ ,  $\text{-CH}_2\text{-Br}$ ), 3.81 (t, 4H,  $J = 5.8$ ,  $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-O}$ ), 3.91 (t, 4H,  $J = 5.8$ ,  $\text{Br-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-O}$ ), 4.39 (d, 4H,  $J = 13.0$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 6.74 (s, 4H,  $\text{Ar-H}$ ), 6.83 (s, 4H,  $\text{Ar-H}$ ). Anal. Calcd for  $\text{C}_{58}\text{H}_{82}\text{O}_4\text{Br}_2$ : C, 69.45; H, 8.24. Found: C, 69.65; H, 8.30.

### Tetrakis-(butoxy-4-bromide)-tetrakis-*p*-tert-butylcalix[4]arene (**S<sub>5</sub>**)

The suspension of *p*-tert-butylcalix[4]arene **S<sub>1</sub>** (1.947 g, 3.00 mmol) and NaH in oil washed three times with hexane (0.700 g, 29.17 mmol) were stirred at room temperature in DMF (50 cm<sup>3</sup>) for 1h. Then 1,4-dibromobutane (12.947 g, 59.06 mmol) was added and the mixture was heated to 80 °C. After 4 days of heating, the mixture was cooled and MeOH (20 cm<sup>3</sup>) was added. After removing of the solvent, the residue was dissolved in dichloromethane and water and acidified with 1M HCl. The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and evaporated. After precipitation from methanol the pure compound **S<sub>5</sub>** (1.520 g, 1.28 mmol) was obtained in 43% yield. Mp 180 °C. <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.09 (s, 36H,  $\text{-C(CH}_3)_3$ ), 1.98-2.09 (m, 8H,  $\text{-CH}_2\text{-}$ ), 2.16-2.23 (m, 8H,  $\text{-CH}_2\text{-}$ ), 3.15 (d, 4H,  $J = 13.0$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 3.53 (t, 8H,  $J = 6.9$ ,  $\text{-CH}_2\text{-Br}$ ), 3.91 (t, 8H,  $J = 6.9$ , -

1  
2  
3  $CH_2-O-$ ), 4.36 (d, 4H,  $J = 13.0$ , Ar- $CH_2$ -Ar), 6.79, (s, 8H, Ar- $H$ ). Anal. Calcd. for  
4  $C_{60}H_{84}O_4Br_4$ : C, 60.61; H, 7.12; Found: C, 60.87; H, 7.32.  
5  
6  
7

### 8 **Tetrakis-(4-(diphenylphosphine)-butoxy)-*p*-*tert*-butylcalix[4]arene ( $S_6$ )**

9  
10 Into a 100 cm<sup>3</sup> flask containing compound  $S_5$  (1.510 g, 1.27 mmol) of freshly distilled  
11 THF (10 cm<sup>3</sup>), (1.282 g, 5.72 mmol) of KP(Ph)<sub>2</sub> in THF (15 cm<sup>3</sup>) were added via a  
12 syringe. The mixture was stirred for 2 hours at room temperature, during which the colour  
13 of the reaction mixture changed from red to dark yellow. The mixture was then evaporated  
14 and extracted twice with 30 cm<sup>3</sup> of dichloromethane. Purification of the crude product on  
15 silica column with a 3/7 dichloromethane/hexane mixture as eluent gave compound  $S_6$   
16 (0.409 g, 0.26 mmol) in 20 % yield. Mp 110 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  [ppm]:  
17 1.08 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.42-1.63 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.00-2.18 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-  
18 O), 2.00-2.18 (m, 8H, -CH<sub>2</sub>-P), 3.05 (d, 4H,  $J = 12.9$ , Ar- $CH_2$ -Ar), 3.81 (t, 8H,  $J = 5.4$ , -  
19 CH<sub>2</sub>-CH<sub>2</sub>-O), 4.29 (d, 4H,  $J = 12.9$ , Ar- $CH_2$ -Ar), 6.75 (s, 8H, Ar- $H$ ), 7.20-7.46 (m, 40H,  
20 P-Ar- $H$ ). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  [ppm]: -14.91 [P].  $m/z$  (MALDI) 1610.88 (M +  
21 H)<sup>+</sup>. Anal. Calcd. for C<sub>108</sub>H<sub>124</sub>O<sub>4</sub>P<sub>4</sub>: C, 80.57; H, 7.76; Found: C, 80.73; H, 7.87.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 **Synthesis of phosphonium ligands**

#### 34 **1,3-bis-(4-triphenylphosphonium-butoxy)-*p*-*tert*-butylcalix[4]arene dibromide (1)**

35  
36 Into a 100 cm<sup>3</sup> flask containing  $S_2$  (1.184 g, 2.00 mmol) in chloroform (30 cm<sup>3</sup>), (5.248 g,  
37 20.00 mmol) of triphenylphosphine in chloroform (20 cm<sup>3</sup>) were added. After 6 days  
38 under reflux, the mixture was cooled and the solvent evaporated. The residue was  
39 dissolved in dichloromethane. The excess of triphenylphosphine was precipitated from  
40 methanol and filtered out. The filtrate was evaporated. The pure product **1** (2.179 g, 1.51  
41 mmol) was obtained by precipitation from a 1/9 dichloromethane/hexane mixture, as a  
42 white-light green powder in 76 % yield: Mp > 280 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$   
43 [ppm]: 0.99 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.28 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 2.05-2.29 (m, 8H, CH<sub>2</sub>-CH<sub>2</sub>-  
44 CH<sub>2</sub>-CH<sub>2</sub>-O), 3.19 (d, 4H,  $J = 12.8$ , Ar- $CH_2$ -Ar), 3.80-4.00 (m, 4H, -CH<sub>2</sub>-O), 3.90 (d, 4H,  
45  $J = 12.8$ , Ar- $CH_2$ -Ar), 3.90-4.08 (m, 4H, -CH<sub>2</sub>-P), 6.79 (s, 4H, Ar- $H$ ), 6.99 (s, 4H, Ar- $H$ ),  
46 7.49 (s, 2H, OH), 7.54 – 7.64 (m, 12H, P-Ar- $H$  meta), 7.65-7.74 (m, 6H, P-Ar- $H$  para),  
47 7.80-7.93 (m, 12H, P-Ar- $H$  ortho). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  [ppm]: 25.80.  $m/z$   
48 (FAB<sup>+</sup>) 721.7 (M + 2H)<sup>2+</sup>;  $m/z$  (MALDI) 1361.7 (M - Br)<sup>+</sup>. Anal. Calcd for  
49 C<sub>88</sub>H<sub>100</sub>O<sub>4</sub>P<sub>2</sub>Br<sub>2</sub>: C, 73.22; H, 6.98. Found: C, 73.46; H, 7.20.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5 **1,3-bis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-p-tert-butylcalix[4]arene**  
6 **dibromide (2)**  
7

8  
9 Compound **2** was prepared following the same procedure as for compound **1** with **S**<sub>2</sub>  
10 (2.753 g, 3.00 mmol) and diphenylmethylphosphine (6.006 g, 30.00 mmol) in 69 % yield.  
11 Mp > 280 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 0.96 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.29 (s, 18H,  
12 -C(CH<sub>3</sub>)<sub>3</sub>), 1.91-2.10 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.11-2.27 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.99 (d, 6H,  
13 J = 13.8, CH<sub>3</sub>-P), 3.23 (d, 4H, J = 13.5, Ar-CH<sub>2</sub>-Ar), 3.58-3.75 (m, 4H, -CH<sub>2</sub>-P), 3.90 (t,  
14 4H, J = 5.3, -CH<sub>2</sub>-O), 4.02 (d, 4H, J = 13.5, Ar-CH<sub>2</sub>-Ar), 6.76 (s, 4H, Ar-H), 7.01 (s, 4H,  
15 Ar-H), 7.31 (s, 2H, OH), 7.48-7.59 (m, 8 H, P-Ar-H meta), 7.61-7.70 (m, 4H, P-Ar-H  
16 para), 7.90-8.06 (m, 8H, P-Ar-H ortho). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 25.96. m/z  
17 (MALDI) 1239.58 (M - Br)<sup>+</sup>. Anal. Calcd. for C<sub>78</sub>H<sub>96</sub>O<sub>4</sub>P<sub>2</sub>Br<sub>2</sub>: C, 71.01; H, 7.33. Found:  
18 C, 71.21; H, 7.52.  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 **1,3-bis-(4-triphenylphosphonium-butoxy)-2,4-bis-propoxy-p-tert-butyl-calix[4]arene**  
29 **dibromide (3)**  
30

31 Into a 100 cm<sup>3</sup> flask containing **S**<sub>4</sub> (2.012 g, 2.00 mmol) in chloroform (30 cm<sup>3</sup>), 5.248 g  
32 of triphenylphosphine (20.00 mmol) in chloroform (20 cm<sup>3</sup>) were added and left for 6  
33 days under reflux. After that time the mixture was cooled and the solvent evaporated. The  
34 residue was dissolved in dichloromethane. The excess of triphenylphosphine was  
35 precipitated from methanol and filtered off. The filtrate was evaporated. The pure product  
36 **3** (2.179g, 1.51 mmol) was obtained by precipitation from a 1/9 dichloromethane/hexane  
37 mixture, as a white-light green powder in 76% yield. Mp 120 °C. <sup>1</sup>H NMR (300 MHz,  
38 CDCl<sub>3</sub>) δ [ppm]: 0.86 (t, 6H, J = 5.3, CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 0.99 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.11 (s,  
39 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.65-1.87 (m, 4H, CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 1.77-1.94 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-P),  
40 2.28-2.43 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.95 (d, 4H, J = 13.0, Ar-CH<sub>2</sub>-Ar), 3.63 (t, 4H, J = 5.9,  
41 CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 3.85-4.00 (m, 4H, -CH<sub>2</sub>-P), 3.85-4.00 (m, 4H, -CH<sub>2</sub>-O), 4.18 (d, 4H, J  
42 = 13.0, Ar-CH<sub>2</sub>-Ar), 6.61 (s, 4H, Ar-H), 6.76 (s, 4H, Ar-H), 7.60-7.92 (m, 30H, P-Ar-H  
43 ortho, meta, para). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 25.84. m/z (MALDI) 1447.6 (M  
44 - Br)<sup>+</sup>. Anal. Calcd. for C<sub>94</sub>H<sub>112</sub>O<sub>4</sub>P<sub>2</sub>Br<sub>2</sub>: C, 73.91; H, 7.39. Found: C, 73.67; H, 7.66.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

58 **1,3-bis-propoxy-2,4-bis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-p-tert-**  
59 **butylcalix[4]arene dibromide (4)**  
60

Compound **4** was obtained according to the same procedure as for **3** with **S**<sub>4</sub> (3.050 g, 3.04

mmol) and diphenylmethylphosphine (6.006 g, 30.00 mmol) a white powder in 64 % yield. Mp 148 °C.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 0.79 (s, 18H,  $-\text{C}(\text{CH}_3)_3$ ), 1.03 (t, 6H,  $J = 6.9$ ,  $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-O}$ ), 1.31 (s, 18H,  $-\text{C}(\text{CH}_3)_3$ ), 1.55-1.75 (m, 4H,  $-\text{CH}_2\text{-CH}_2\text{-P}$ ), 1.88-2.04 (m, 4H,  $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-O}$ ), 2.40-2.58 (m, 4H,  $\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-O}$ ), 3.05 (d, 4H,  $J = 12.5$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 3.23 (d, 6H,  $J = 14.3$ ,  $\text{CH}_3\text{-P}$ ), 3.62-3.80 (m, 8H,  $-\text{CH}_2\text{-P}$  and  $-\text{CH}_2\text{-O}$ ), 3.88 (t, 4H,  $J = 5.9$ ,  $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-O}$ ), 4.32 (d, 4H,  $J = 12.5$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 6.43 (s, 4H,  $\text{Ar-H}$ ), 7.08 (s, 4H,  $\text{Ar-H}$ ), 7.60-7.80 (m, 12H,  $\text{P-Ar-H meta, para}$ ), 8.01-8.13 (m, 8H,  $\text{P-Ar-H ortho}$ ).  $^{31}\text{P}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 25.98.  $m/z$  (MALDI) 1323.7 ( $\text{M} - \text{Br}$ ) $^+$ . Anal. Calcd. for  $\text{C}_{84}\text{H}_{108}\text{O}_4\text{P}_2\text{Br}_2$ : C, 71.88; H, 7.76. Found: C, 71.99; H, 7.85.

### **Tetrakis-(4-triphenylphosphonium-butoxy)-tetrakis-*p*-tert-butylcalix[4]arene tetrabromide (5)**

Compound **S<sub>5</sub>** (1.184 g, 1.00 mmol) was dissolved in chloroform (30  $\text{cm}^3$ ). After a few minutes of stirring triphenylphosphine (5.248 g, 20.00 mmol) and chloroform (20  $\text{cm}^3$ ) were added. After 6 days of refluxing the mixture was cooled and solvent was evaporated. The residue was dissolved in dichloromethane. The excess of triphenylphosphine was precipitated from methanol and filtered. The organic layer was evaporated. Chromatography on a silica column with 90:10 dichloromethane: methanol mixture as eluent gave compound **5** (0.67 g, 0.30 mmol) in 30% yield. Mp 132 °C.  $^1\text{H}$  NMR (300 MHz;  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.02 (s, 36H,  $-\text{C}(\text{CH}_3)_3$ ), 1.56-1.72 (m, 8H,  $-\text{CH}_2-$ ), 2.24-2.41 (m, 8H,  $-\text{CH}_2-$ ), 2.91 (d, 4H,  $J = 13.0$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 3.78-4.01 (m, 16H,  $-\text{CH}_2\text{-P}$  and  $\text{Ar-O-CH}_2$ ), 4.23 (d, 4H,  $J = 13.0$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 6.63 (s, 8H,  $\text{Ar-H}$ ), 7.59-7.71- (m, 36H,  $\text{P-Ar-H, meta, para}$ ), 7.76-7.88 (m, 24H,  $\text{P-Ar-H, ortho}$ ).  $^{31}\text{P}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 25.76.  $m/z$  ( $\text{FAB}^+$ ) 479.5 ( $\text{M}$ ) $^{4+}$ ;  $m/z$  (MALDI) 2157.7 ( $\text{M} - \text{Br}$ ) $^+$ . Anal. Calcd. for  $\text{C}_{132}\text{H}_{144}\text{O}_4\text{P}_4\text{Br}_4$ : C, 70.84; H, 6.49. Found: C, 70.97; H, 6.69.

### **Tetrakis-(4-triphenylphosphonium-butoxy)-*p*-tert-butylcalix[4]arene tetraperchlorate (5a)**

In a 10  $\text{cm}^3$  flask **5** (0.100 g, 0.045 mmol) was dissolved in acetonitrile (1  $\text{cm}^3$ ). (0.050 g, 0.241 mmol) of  $\text{AgClO}_4$  in acetonitrile (1  $\text{cm}^3$ ) were added dropwise to the ligand solution. After 24 hours of stirring at room temperature, the precipitate of  $\text{AgBr}$  was filtered off. The filtrate was evaporated to give compound **5a** (0.088 g, 0.038 mmol) in 84 % yield. Mp > 120 °C decomposition.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.07 (s, 36H,  $-\text{C}(\text{CH}_3)_3$ ), 1.70-1.85 (m, 8H,  $-\text{CH}_2\text{-CH}_2\text{-P}$ ), 2.45-2.53 (m, 8H,  $-\text{CH}_2\text{-CH}_2\text{-O}$ ), 2.95 (d, 4H,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

$J = 13.0$ , Ar- $\text{CH}_2$ -Ar), 3.43-3.60 (m, 8H, - $\text{CH}_2$ -P), 4.20-4.30 (m, 8H, - $\text{CH}_2$ -O), 4.44 (d, 4H,  $J = 13.0$ , Ar- $\text{CH}_2$ -Ar), 6.97 (s, 8H, Ar- $H$ ), 7.56-7.90 (m, 60H, P-Ar- $H$ , *ortho*, *meta*, *para*),  $^{31}\text{P}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 24.54.  $m/z$  (MALDI) 2214.2 ( $\text{M} - \text{ClO}_4$ ) $^+$ . Anal. Calcd: for  $\text{C}_{132}\text{H}_{144}\text{O}_4\text{P}_4(\text{ClO}_4)_4$ : C, 68.45; H, 6.27. Found: C, 68.70; H, 6.45.

**Tetrakis-(4-triphenylphosphonium-butoxy)-tetrakis-*p*-tert-butylcalix[4]arene tetrahexafluorophosphate (5b)**

Compound **5** (0.100 g, 0.045 mmoles) was dissolved in acetonitrile (1  $\text{cm}^3$ ). Then (0.062 g, 0.245 mmol)  $\text{AgPF}_6$  were dissolved in acetonitrile and added dropwise to the ligand solution. After 24h the precipitate of NaBr was removed and the solution was evaporated. Compound **5b** (0.090 g, 0.036 mmole) was obtained in 80 % yield. Mp 128  $^\circ\text{C}$ .  $^1\text{H}$  NMR (500 MHz;  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.12 (s, 36H, - $\text{C}(\text{CH}_3)_3$ ), 1.65-1.81 (m, 8H, - $\text{CH}_2$ -), 2.38-2.49 (m, 8H, - $\text{CH}_2$ -), 3.25-3.38 (m, 8H, - $\text{CH}_2$ -P), 3.47 (d, 4H,  $J = 13.0$ , Ar- $\text{CH}_2$ -Ar), 4.15-4.24 (m, 8H, Ar-O- $\text{CH}_2$ ), 4.47 (d, 4H,  $J = 13.0$ , Ar- $\text{CH}_2$ -Ar), 7.04 (s, 8H, Ar- $H$ ), 7.66-7.70 (m, 60H, P-Ar- $H$ ).  $^{31}\text{P}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 24.72 [ $\text{P}^+$ ], -143.39 [ $\text{PF}_6$ ].  $m/z$  (MALDI) 1447.60 ( $\text{M} - \text{PF}_6$ ) $^+$ . Anal. Calcd. for  $\text{C}_{132}\text{H}_{144}\text{O}_4\text{P}_4(\text{PF}_6)_4$ : C, 63.46; H, 5.81. Found: C, 63.30; H, 5.78.

**Tetrakis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-*p*-tert-butylcalix[4]arene tetrabromide (6)**

Into a 100  $\text{cm}^3$  flask containing **S5** (1.184 g, 1.00 mmol) dissolved in chloroform (30  $\text{cm}^3$ ), (4.004 g, 20.00 mmol) diphenylmethylphosphine in chloroform (20  $\text{cm}^3$ ) were added. After 6 days under reflux the mixture was cooled and the solvent was evaporated. The product was purified by precipitation from a 1/9 dichloromethane/hexane mixture to give compound **6** (1.190 g, 0.59 mmol) in 59 % yield. Mp 160  $^\circ\text{C}$ .  $^1\text{H}$  NMR (300 MHz;  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.03 (s, 36H, - $\text{C}(\text{CH}_3)_3$ ), 1.55-1.75 (m, 8H, - $\text{CH}_2$ - $\text{CH}_2$ -P), 2.30-2.48 (m, 8H, - $\text{CH}_2$ - $\text{CH}_2$ -O), 2.91 (d, 12H,  $J = 13.5$ ,  $\text{CH}_3$ -P), 3.03 (d, 4H,  $J = 12.8$ , Ar- $\text{CH}_2$ -Ar), 3.70-3.87 (m, 16H, - $\text{CH}_2$ -P and  $\text{CH}_2$ -O), 4.26 (d, 4H,  $J = 12.8$ , Ar- $\text{CH}_2$ -Ar), 6.71 (s, 8H, Ar- $H$ ), 7.54-7.78 (m, 24H, P-Ar- $H$  *meta*, *para*), 8.02-8.17 (m, 16H, P-Ar- $H$  *ortho*).  $^{31}\text{P}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 25.98.  $m/z$  (MALDI) 1909.68 ( $\text{M} - \text{Br}$ ) $^+$ . Anal. Calcd. for  $\text{C}_{112}\text{H}_{136}\text{O}_4\text{P}_4\text{Br}_4$ : C, 67.61; H, 6.89. Found: C, 68.59; H, 7.10.

**Tetrakis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-*p*-tert-butylcalix[4]arene**

**tetraperchlorate (6a)**

Compound **6a** was obtained according to the same procedure as **5a** with **6** (0.100 g, 0.05 mmol) and AgClO<sub>4</sub> (0.050 g, 0.24 mmol) in 90 % yield. Mp > 120 °C decomposition. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 1.09 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.65-1.80 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.35-2.50 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.50 (d, 12H, *J* = 12.0, CH<sub>3</sub>-P), 3.18-3.28 (m, 8H, -CH<sub>2</sub>-P), 3.38 (d, 4H, *J* = 12.8, Ar-CH<sub>2</sub>-Ar), 4.15-4.28 (m, 8H, -CH<sub>2</sub>-O), 4.45 (d, 4H, *J* = 12.8, Ar-CH<sub>2</sub>-Ar), 6.97 (s, 8H, Ar-*H*), 7.60-7.87 (m, 40H, P-Ar-*H*, *ortho*, *meta*, *para*). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 24.77. *m/z* (MALDI) 1966.24 (M - ClO<sub>4</sub>)<sup>+</sup>. Anal. Calcd. for C<sub>112</sub>H<sub>136</sub>O<sub>4</sub>P<sub>4</sub>(ClO<sub>4</sub>)<sub>4</sub>: C, 65.05; H, 6.63. Found: C, 65.13; H, 6.72.

**Tetrakis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-p-tert-butylcalix[4]arene tetrahexafluorophosphate (6b)**

Compound **6b** was obtained according to the same procedure as for **5b** with **6** (0.100 g, 0.05 mmol) and AgPF<sub>6</sub> (0.069 g, 0.27 mmol) in 80 % yield. Mp 155 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 1.11 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.55-1.70 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.24-2.38 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.42 (d, 12H, *J* = 12.0, CH<sub>3</sub>-P), 2.95-3.10 (m, 8H, -CH<sub>2</sub>-P), 3.39 (d, 4H, *J* = 12.8, Ar-CH<sub>2</sub>-Ar), 4.05-4.20 (m, 8H, -CH<sub>2</sub>-O), 4.42 (d, 4H, *J* = 12.8, Ar-CH<sub>2</sub>-Ar), 7.00 (s, 8H, Ar-*H*), 7.58-7.80 (m, 40H, P-Ar-*H*, *ortho*, *meta*, *para*). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 24.50 [P<sup>+</sup>], -143.07 [PF<sub>6</sub>], *m/z* (MALDI) 2104.80 (M - PF<sub>6</sub>)<sup>+</sup>. Anal. Calcd. for C<sub>112</sub>H<sub>136</sub>O<sub>4</sub>P<sub>4</sub>(PF<sub>6</sub>)<sub>4</sub>: C, 59.79; H, 6.09. Found: C, 59.87; H, 6.03.

**Protonated tetrakis-(4-(P,P-diphenyl-phosphine)-butoxy)-p-tert-butylcalix[4]arene tetrabromide (7)**

Into a 50 cm<sup>3</sup> flask containing **S<sub>6</sub>** (1.510 g, 0.94 mmol) in dichloromethane (10 cm<sup>3</sup>), 5 cm<sup>3</sup> of HBr (solution of 33 % wt. in glacial acetic acid) in dichloromethane (15 cm<sup>3</sup>) were added. The mixture was left stirred for 24 hours at room temperature. The mixture was then evaporated to give the pure product **7** (0.954 g, 0.49 mmol) in 52 % yield. Mp 175 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 1.22 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.80-2.15 (m, 16H, O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-P), 2.92-3.10 (m, 4H, Ar-CH<sub>2</sub>-Ar), 3.18-3.30 (m, 8H, CH<sub>2</sub>-O), 3.40-3.55 (m, 8H, -CH<sub>2</sub>-P), 4.05-4.27 (m, 4H, Ar-CH<sub>2</sub>-Ar), 7.05 (s, 8H, Ar-*H*), 7.60-8.05 (m, 40H, P-Ar-*H*, *ortho*, *meta*, *para*), 10.30-10.40 (m, 4H, H-P). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 46.32 [P<sup>+</sup>]. *m/z* (MALDI) 1849.04 (M - Br)<sup>+</sup>. Anal. Calcd. for C<sub>108</sub>H<sub>128</sub>O<sub>4</sub>P<sub>4</sub>Br<sub>4</sub>: C, 67.08; H, 6.67. Found: C, 66.84; H, 6.92.

### NMR studies

5 mg of ligand were introduced with 6 equivalents of the solid alkali metal salts ( $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{Cs}^+$ ) or tetraethylammonium perchlorate in a glass vessel and dissolved in a small volume of deuterated chloroform or acetonitrile. After manual shaking for a few minutes, the mixture was left in contact for 24 hours before filtration of the excess of salt if necessary. The solution (or the filtrate) was then sampled in a NMR tube and its spectrum recorded on Bruker SY300 MHz or SY400 MHz spectrometers equipped for  $^1\text{H}$  or  $^{31}\text{P}$  resonances.

### Microcalorimetric studies


Microcalorimetric titrations were performed using a 2277 Thermal Activity Monitor Microcalorimeter (Thermometric). Titration were carried out at 25 °C on 2.7 cm<sup>3</sup> of 10<sup>-5</sup> to 5×10<sup>-4</sup> M solutions of the ligand in acetonitrile using a glass cell of 4 cm<sup>3</sup>. The heat changes were measured after injection of 15×15 μL of 10<sup>-3</sup> and 10<sup>-2</sup> M  $\text{LiClO}_4$ ,  $\text{LiBr}$ ,  $\text{NaClO}_4$ ,  $\text{NaPF}_6$ , or  $\text{Et}_4\text{NClO}_4$  solutions in the same solvent. Chemical calibration was made by determination of the complexation enthalpy of  $\text{Ba}^{2+}$  with 18C6 in water or of  $\text{Rb}^+$  with 18C6 in methanol, as recommended [37]. Values of the stability constants ( $\beta$ ) and of the enthalpies of complexation ( $\Delta H$ ) were refined simultaneously from these data using the ligand binding analysis program DIGITAM version 4.1 [38] and after correction for the heat of dilution determined in separate experiments by adding the salt solutions to 2.7 cm<sup>3</sup> of pure solvent. The values of the corresponding entropies of complexation ( $\Delta S$ ) were then derived from the expressions  $\Delta G = -RT\ln\beta$  and  $\Delta G = \Delta H - T\Delta S$ .

### Ion selective electrodes

THF was dried and freshly distilled before used for the preparation of the ion selective membranes. PVC (high molecular weight poly(vinyl chloride), 2-nitrophenyl octyl ether (*o*-NPOE), bis-(2-ethylhexyl)sebacate (BEHS), (2-morpholino)ethanesulfonic acid monohydrate (MES) were from Fluka Selectophore. The  $\text{LiClO}_4$ ,  $\text{CsClO}_4$  and sodium salts:  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$ ,  $\text{ClO}_4^-$ ,  $\text{SCN}^-$ ,  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{CO}_3^{2-}$ ,  $\text{HPO}_4^{2-}$ ,  $\text{PO}_4^{3-}$ ,  $\text{Cr}_2\text{O}_7^{2-}$ , citrate, acetate, benzoate, and oxalate were of p.a. grade. All aqueous salt solutions were prepared with demineralised water (conductivity < 1.0 μS/cm).

The membranes were composed of 4 mg of ionophores **1** – **6** and **8**, 60 mg of PVC and 120 mg of plasticizer. All the components were dissolved in 1.5 cm<sup>3</sup> of dried, freshly distilled

1  
2  
3 THF and the solutions were poured into glass rings of 24 mm in diameter. The solutions were  
4 left for 24 h for slow solvent evaporation giving the mother membranes of thickness about  
5 0.1mm. Several membranes of 7 mm diameter were cut from each mother membrane and  
6  
7 were incorporated into the Ag/AgCl electrodes bodies of IS 561 type (Moeller S.A., Zurich).  
8  
9 The two plasticizers BEHS and *o*-NPOE were used for the preparation of the membranes.  
10  
11 However, electrodes with membranes based on NPOE had the best lifetime and response  
12 characteristics. The EMF measurements were carried out at zero current conditions using a  
13  
14 Lawson Lab 16 EMF station (multi-channel millivoltmeter) or a Metrohm 654 millivoltmeter.  
15  
16 A double-junction reference Radelkis 0P0820P electrode with a 1M CH<sub>3</sub>COOLi solution in  
17  
18 the bridge cell was used. The measurements were carried out using cells of the type:



26  
27 At least three identical electrodes of the same membrane composition and containing the  
28 same inner electrolyte were prepared [39]. The studies were repeated several times over the  
29  
30 period of one month.

31  
32 To reduce the pH changes during the titrations solutions were prepared with 0.05 M  
33  
34 MES/NaOH buffer of pH = 5.5 ((2-morpholino)ethanesulfonic acid monohydrate (MES)). All  
35  
36 salt solutions contained 10<sup>-2</sup> M NaCl as supporting electrolyte [22].

37  
38 The selectivity coefficients  $K_{A,B}^{pot}$  of the electrodes were determined by the separate  
39  
40 solution method (SSM) and in some cases by the fixed interference method (FIM) [40,41,42].  
41  
42 The calibration curves were obtained by addition of standard solutions of different anions to  
43  
44 50 cm<sup>3</sup> of 0.01 M NaCl in 0.05 M MES/NaOH buffer solution of pH = 5.5. The concentration  
45  
46 of the primary anion [A] was increased from 10<sup>-7</sup> to 10<sup>-2</sup> M. They were also measured by  
47  
48 successive dilution of initial 5×10<sup>-2</sup> M salt solutions until further dilution resulted in no  
49  
50 potential change.

### 51 52 53 **Acknowledgements**

54  
55 R.P. gratefully acknowledges the Gdansk University of Technology, Poland, and the  
56  
57 Ministère de la Recherche et des Nouvelles Technologies, France (Cotutelles de thèses),  
58  
59 for financial support.  
60

## References

- [1] Bianchi, A.; Bowman-James, K.; Garcia-Espana, E. Eds., *Supramolecular Chemistry of Anions*; John Wiley-VCH, New York, 1997, ch. 3, pp. 63.
- [2] Kanyo, Z. F.; Christianson, D. W. *J. Biol. Chem.*, **1991**, 266, 4264.
- [3] Pardee, A. P. *J. Biol. Chem.*, **1966**, 241, 5887.
- [4] Puljak L.; Kilic, G. *Biochim. Biophys. Acta*, **2006**, 1762, 404.
- [5] Weiner, M. L.; Salminen, W. F.; Larson, P. R.; Barter, R. A.; Kranetz J. L.; Simon, G. *S. Food and Chemical Toxicology*, **2001**, 39, 759.
- [6] Matzer, W. E. *Environmental Forensic*, **2001**, 2, 301.
- [7] Bondy, C. R.; Gale P. A.; Loeb, S. J. *J. Am. Chem. Soc.*, **2004**, 126, 5030.
- [8] Amendola, V.; Boiocchi, M.; Esteban-Gómez, D.; Fabbri L.; Monzani, E. *Org. Biomol. Chem.*, **2005**, 3, 2632.
- [9] Brooks, S. J.; Edwards, P. R.; Gale P. A.; Light, M. E. *New J. Chem.*, **2006**, 30, 65.
- [10] Turner, D. R.; Paterson M. J.; Steed, J. W. *J. Org. Chem.*, **2006**, 71, 1598.
- [11] Boerrigter, H.; Grave, L.; M. Nissink, J. W.; Chrisstoffels, L. A. J.; van der Maas, J. H.; Verboom, W.; de Jong F.; Reinhoudt, D. N.; *J. Org. Chem.*, **1998**, 63, 4174.
- [12] Lee K. H.; Hong, J. I. *Tetrahedron Lett.*, **2000**, 41, 6083.
- [13] Chmielewski M.; Jurczak, J. *Tetrahedron Lett.*, **2004**, 45, 6007.
- [14] Beer, P. D.; Drew M. G. B.; Gradwell, . K. *J. Chem. Soc., Perkin Trans. 2*, **2000**, 511.
- [15] Dietrich, B. *Pure & Appl. Chem.*, **1993**, 65, 1457.
- [16] Egorov, V. V.; Rakhman'ko, E. M.; Okaev E. B.; Pomelenok E. V.; Nazarov, V. A.; *Talanta*, **2004**, 63, 119.
- [17] Matthews S. E.; Beer, P. D. *Calixarenes 2001*; Asfari, Z.; Böhmer, V.; Harrowfield, J.; Vicens J. Eds., Kluwer Academic Publishers, Dordrecht, 2001, ch. 23, p. 421.
- [18] Miao, R.; Zheng, Q. Y.; Chen C. F.; Huang Z. T. *Tetrahedron Lett.*, **2004**, 45, 4959.

- 1  
2  
3  
4  
5 [19] Jagessar, R. C.; Shang, M.; Scheidt, W. R.; Burns; D. H. *J. Am. Chem. Soc.*, **1998**,  
6 *120*, 11684.  
7  
8 [20] Gale, P. A.; Sessler, J. L.; Kral, V.; Lynch, V. *J. Am. Chem. Soc.*, **1996**, *118*, 5140.  
9  
10 [21] Gale, P. A.; Sessler, J. L.; Allen, W. E.; Tvermoes, N. A.; Lynch, V. *Chem. Commun.*,  
11 **1997**, 665.  
12  
13 [22] Pomecko, R.; Asfari, Z.; Hubscher-Bruder, V.; Bochenska M.; Arnaud-Neu, F.  
14 *Supramol. Chem.*, **2007**, *19*, 459.  
15  
16 [23] Hamdi, A.; Nam, K. C.; Ryu, B.J.; Kim, J. S.; Vicens, J. *Tetrahedron Letters* **2004**, *45*,  
17 4689.  
18  
19 [24] Broder, C. K.; Davidson, M. G.; Forsyth, V. T.; Howard, J. A. K.; Lamb S.; Mason, S.  
20 *A. Crys. Grow. Des.*, **2002**, *3*, 163.  
21  
22 [25] Issleib K.; Muller, D.-W. *Chem. Berichte*, **1959**, *92*, 3175.  
23  
24 [26] Clark, P. W. *Org. Prep. and Proc. Int.*, **1979**, *11*, 103.  
25  
26 [27] Mohr, W.; Horn, C. R.; Stahl J.; Gladysz, J. A. *Synthesis*, **2003**, *8*, 1279.  
27  
28 [28] Baklouti, L.; Cherif J.; Abidi, R.; Arnaud-Neu, F.; Harrowfield J.; Vicens, J. *Org.*  
29 *Biomol. Chem.*, **2004**, *2*, 2786.  
30  
31 [29] Savage, P. B.; Holmgren S. K.; Gellman, S. H. *J. Am. Chem. Soc.*, **1994**, *116*, 4069.  
32  
33 [30] Casabo, J.; Munoz J. A.; Errachid, A. *Sens. Actuators, B*, **1997**, *43*, 206.  
34  
35 [31] Shamsipur, M.; Soleymannpour, A.; Akhond, M.; Sharghi H.; Hasaninejad, A. R. *Sens.*  
36 *Actuators, B*, **2003**, *89*, 9.  
37  
38 [32] Dinten, O.; Spichiger, U. E.; Chaniotakis, N.; Gehrig, P.; Rusterholz, B.; Morf, W. E.;  
39 Simon, W. *Anal. Chem.*, **1991**, *63*, 596.  
40  
41 [33] Lizondo-Sabater, J.; Segui, M. J.; Loris, J. M.; Martinez-Manez, R.; Pardo, T.;  
42 Sancenon, F.; Soto, J. *Sens. Actuators B*, **2004**, *101*, 20.  
43  
44 [34] Segui, M. J.; Lizondo-Sabater, J.; Martinez-Manez, R.; Soto, J. *Analyst*, **2002**, *127*,  
45 387.  
46  
47 [35] Casabo, J.; Escriche, L.; Perez-Jimenez, C.; Munoz, J. A.; Teixidor, F.; Bausells, J.;  
48 Errachid, A. *Anal. Chim. Acta*, **1996**, *320*, 63.  
49  
50 [36] Schaller, U.; Bakker, E.; Spichiger, U. E.; Pretsch, E. *Anal. Chem.*, **1994**, *66*, 391.  
51  
52 [37] Arnaud-Neu, F.; Delgado, R.; Chaves, S. *Pure Appl. Chem.*, **2003**, *75*, 71.  
53  
54 [38] Hallen, D. *Pure Appl. Chem.*, **1993**, *65*, 1527.  
55  
56 [39] Bochenska, M.; Zielenska, A.; Pomecko, R.; Hubscher-Bruder, V.; Arnaud-Neu, F. *J.*  
57 *Incl. Phenom.*, **2005**, *52*, 129.  
58  
59  
60



- 
- 1  
2  
3  
4  
5 [40] Buck, R. P.; Linder, E. *Pure Appl. Chem.*, **1994**, *66*, 2527.  
6  
7 [41] Umezawa, Y.; Buhlmann, P.; Umezawa, K.; Tohda, K.; Amemiya, S. *Pure Appl.*  
8 *Chem.* **2000**, *72*, 1851.  
9  
10 [42] Lindner, E.; Umezawa, Y. *Pure Appl. Chem.* **2008**, *80*, 85.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review Only

**Table 1.** Changes ( $\Delta\delta$  in ppm) in the  $^{31}\text{P}$  NMR spectra of phosphonium ligands in the presence of sodium iodide, thiocyanate and perchlorate in  $\text{CDCl}_3$

	<b>1</b>	<b>2</b>	<b>5</b>	<b>6</b>
Ligand ( $\delta$ )	25.797	25.962	25.756	25.982
Ligand + $\text{I}^-$ ( $\Delta\delta$ )	0.215	0.203	0.210	0.175
Ligand + $\text{SCN}^-$ ( $\Delta\delta$ )	0.470	0.530	0.767	0.965
Ligand + $\text{ClO}_4^-$ ( $\Delta\delta$ )	0.602	0.620	0.992	1.170

**Table 2.** Differences ( $\Delta\delta$ ) in the proton chemical shifts ( $\delta$ )[ppm] in the spectra of ligands **5** and **6**, **5a** and **6a**, **5b** and **6b** in CD<sub>3</sub>CN

	<b>c</b>	<b>e</b>	<b>f</b>	<b>g</b>	<b>j</b>	<b>o</b>
$\delta$ ( <b>5</b> )	6.87	2.93	4.15	3.74	3.81	-
$\delta$ ( <b>5a</b> )	6.83	2.91	4.11	3.76	3.17	-
$\Delta\delta$ ( <b>5a-5</b> )	-0.04	-0.02	-0.04	0.02	-0.64	-
$\delta$ ( <b>5b</b> )	6.80	2.87	4.09	3.77	3.06	-
$\Delta\delta$ ( <b>5b-5</b> )	-0.07	-0.06	-0.06	0.03	-0.75	-
$\delta$ ( <b>6</b> )	6.92	3.02	4.19	3.70	3.58	2.78
$\delta$ ( <b>6a</b> )	6.89	2.98	4.15	3.73	2.85	2.37
$\Delta\delta$ ( <b>6a-6</b> )	-0.03	-0.04	-0.04	0.03	-0.73	-0.41
$\delta$ ( <b>6b</b> )	6.88	2.97	4.12	3.75	2.75	2.32
$\Delta\delta$ ( <b>6b-6</b> )	-0.04	-0.05	-0.07	0.05	-0.83	-0.46

**Table 3.** Characteristics of potentiometric responds for perchlorate of PVC/NPOE electrodes containing ligand **5** and **6** and different amount of KTCIPB. (Inner and conditioning electrolyte MES/NaOH pH=5.5/10<sup>-2</sup>M NaCl)

Ligand	KTCIPB (mol %)	S (mV/decade)	LR (log[A])
<b>5</b>	0	-55.6	-6.0
<b>5</b>	40	-54.6	-6.0
<b>5</b>	120	-55.9	-5.7
<b>6</b>	0	-54.2	-6.0
<b>6</b>	40	-51.8	-6.0
<b>6</b>	120	-49.9	-5.7

**Table 4.** Characteristics of potentiometric responds for perchlorate of PVC/NPOE electrodes containing tetrasubstituted phosphonium ligands with different counterions (Inner and conditioning electrolyte MES/NaOH pH=5.5/10<sup>-2</sup>M NaCl).

Ligand	Counterion	S (mV/decade)	LR (log[A])
<b>5</b>	Br <sup>-</sup>	-56.5	-6.0
<b>5a</b>	ClO <sub>4</sub> <sup>-</sup>	-56.1	-6.0
<b>5b</b>	PF <sub>6</sub> <sup>-</sup>	-38.3	-5.5
<b>6</b>	Br <sup>-</sup>	-54.4	-5.7
<b>6a</b>	ClO <sub>4</sub> <sup>-</sup>	-40.4	-6.0
<b>6b</b>	PF <sub>6</sub> <sup>-</sup>	-36.2	-6.0

**Table 5.** Selectivity coefficients as  $\log K_{\text{ClO}_4^-,X}^{\text{pot}}$  of the PVC/NPOE membrane electrodes based on phosphonium calixarenes **1 - 6** and the monomer **8**

Anion X	$\log K_{\text{ClO}_4^-,X}^{\text{pot}}$						
	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>8</b>
$\text{ClO}_4^-$	0	0	0	0	0	0	0
$\text{SCN}^-$	-1.2	-1.4	-1.1	-0.6	-1.3	-1.4	-1.1
$\text{I}^-$	-1.7	-1.7	-1.3	-0.9	-2.0	-2.0	-1.6
$\text{NO}_3^-$	-2.6	-2.6	-2.4	-1.9	-2.9	-3.0	-2.5
$\text{HCO}_3^-$	-4.6	-4.5	-4.6	-4.4	-4.6	-4.4	nd
$\text{Cr}_2\text{O}_7^{2-}$	-2.2	-2.4	-2.6	-1.7	-2.6	-3.2	-1.9
$\text{HPO}_4^{2-}$	-4.5	-4.5	-4.5	-4.4	-4.5	-4.4	nd
$\text{SO}_4^{2-}$	-4.9	-4.7	-4.7	-4.7	-4.7	-4.5	nd

nd: not determined

**Figure captions**

- Figure 1.** Chemical structures of the ligands under study
- Figure 2.**  $^1\text{H}$  NMR spectra of ligand **1** alone and in the presence of  $\text{ClO}_4^-$  in  $\text{CDCl}_3$
- Figure 3.**  $^1\text{H}$  NMR spectra of ligand **6** alone and in the presence of  $\text{ClO}_4^-$  in  $\text{CDCl}_3$
- Figure 4.** Potentiometric anion responses of electrodes with PVC/NPOE membrane containing ligand **1** in MES buffer at pH 5.5.
- Figure 5.** Potentiometric anion responses of electrodes with PVC/NPOE membrane containing ligand **6** in MES buffer at pH 5.5.

## Anion recognition by phosphonium calix[4]arenes : synthesis and physico-chemical studies

Radoslaw Pomecko,<sup>a,b</sup> Zouhair Asfari,<sup>a</sup> Véronique Hubscher-Bruder,<sup>a</sup> Maria Bochenska<sup>b\*</sup> and Françoise Arnaud-Neu<sup>a\*</sup>

<sup>a</sup>*IPHC-DISA, ULP, CNRS, ECPM, 25 rue Becquerel, 67087 Strasbourg Cedex 2, France; E-mail: farnaud@chimie.u-strasbg.fr*

<sup>b</sup>*Department of Chemical Technology, Chemical Faculty, Gdansk University of Technology, ul. Narutowicza 11/12, 80-264 Gdansk, Poland. E-mail: marboch@chem.pg.gda.pl*

*p*-tert-Butylcalix[4]arenes, in the cone conformation, di- and tetrasubstituted at the narrow rim with charged phosphonium groups, have been synthesized and characterized. Their interactions with a wide range of anions have been investigated in solution in chloroform and acetonitrile by means of <sup>1</sup>H and <sup>31</sup>P NMR and microcalorimetry (ITC). These compounds have also been incorporated as sensing material in PVC ion selective electrodes (ISE). The results showed that they interact strongly with the more lipophilic anions ClO<sub>4</sub><sup>-</sup>, SCN<sup>-</sup> and I<sup>-</sup>, in solution as in the electrode membranes. The origin of this selectivity is discussed and, in particular, the role of the salt counterion is examined.

**Keywords:** Phosphonium calix[4]arenes; anion binding properties; microcalorimetry; ion selective electrodes.

### INTRODUCTION

Anions play an important role in many biological processes, such as regulation of cell activity, synthesis of proteins, transport of hormones [1,2,3]. They are also frequently used in many industrial technologies, which very often generate an increase of the concentration of anions in the environment or even introduce anionic species which were unknown so far in ecosystems. The presence of these anions is crucial in environmental and medical concerns, as they are pollutants and may have harmful effects on living organisms and human health [4,5,6]. Therefore there is a need of fast and selective anion detection methods allowing real-time monitoring of anion concentration changes and of efficient clean up processes. Design of anion receptors for such applications remains a great challenge for chemists because they have to take into consideration the specific



1  
2  
3 anion properties, such as a large range of shapes and geometries, small electric charges vs.  
4 sizes, high free energies of solvation, and in some cases multiple oxidation states of the  
5 central atoms in oxoanions or pH dependence. In many artificial anion hosts, noncovalent  
6 interactions are responsible for host-guest recognition. They include electrostatic  
7 interactions, hydrogen bonding, hydrophobic effects, coordination to a metal ion or  
8 combinations of these interactions. The hosts can be neutral, containing urea [7,8,9,10],  
9 thiourea [11,12] or amide functions [13]. They can also be positively charged, containing  
10 pyridinium [14], polyammonium [15] or quaternary ammonium [16] binding sites.  
11 Calix[4]arenes [17,18] and porphyrins [19] are often used as scaffolds onto which these  
12 functional groups can be grafted. Calixpyrroles are also known as efficient anion receptors  
13 [20,21].

22  
23 Recently, we described the synthesis and characterization of a new calix[4]arene  
24 derivative (**5**) bearing four positively charged triphenyl phosphonium groups [22]. The  
25 presence of these highly polarisable moieties, where the charge is spread over the three  
26 aromatic rings, was expected to favour the interaction with lipophilic anions. Preliminary  
27 binding studies showed that this ligand interacted selectively with some anions, namely  
28  $\text{ClO}_4^-$  and  $\text{SCN}^-$ . This compound was also incorporated, as ionophore-sensing material, in  
29 ion selective electrodes (ISEs) which exhibited a selectivity order similar to the  
30 Hofmeister series. The disubstituted derivative **1** was also synthesised as the  
31 hexafluorophosphate [23].

32  
33 In order to get more information on the mechanisms involved in the recognition  
34 process and to optimize its selectivity, we have now extended this study to new di- and  
35 tetrasubstituted phosphonium calix[4]arene derivatives (compounds **2–4**, **6** and **7**) and re-  
36 examined the properties of **1** and **5** (Fig. 1). In some of these compounds, one of the  
37 phenyl rings on the phosphonium groups has been replaced by a methyl radical or a  
38 hydrogen atom. The presence of such small substituents is expected to increase the charge  
39 density on the phosphorus atoms and their accessibility [24]. Moreover, the presence of  
40 hydrogen atoms may induce the formation of hydrogen bonds with anions. The possibility  
41 of tuning the charge density on the phosphorus atoms by changing the nature of the their  
42 substituents could allow the design of receptors able to distinguish lipophilic anions like  
43  $\text{ClO}_4^-$ ,  $\text{SCN}^-$ ,  $\text{I}^-$  or  $\text{NO}_3^-$  from other anions and between them. The binding properties of  
44 these compounds towards a variety of anions have been followed by  $^1\text{H}$  and  $^{31}\text{P}$  NMR and  
45 by titration microcalorimetry (ITC). In particular the role of the salt counterion was  
46 examined using these techniques.

Fig. 1

## RESULTS AND DISCUSSION

### Synthesis

The ligands synthesised in this work (Fig. 1) are based on a tetrakis-*p-tert*-butylcalix[4]arene platform which presents several advantages. It has a well defined size and is readily amenable to substitution at its lower rim where ligating groups can be attached and in some extent preorganized. The amphiphilicity of such derivatives should allow their introduction into the membranes of ion selective electrodes.

Diphosphonium ligands were synthesised in two or three steps according to the known procedure [23]. The first step was the selective bromoalkylation of the tetrakis-*p-tert*-butylcalix[4]arene (**S**<sub>1</sub>), leading to the intermediate molecules **S**<sub>2</sub> or **S**<sub>3</sub> in 71 % and 55 % yield, respectively (Scheme 1). 1,3-bis-(4-triphenylphosphonium-butoxy)-*p-tert*-butylcalix[4]arene dibromide (**1**) was obtained by the reaction of **S**<sub>2</sub> with 10 equivalents of triphenylphosphine. After 6 days under reflux in chloroform, the product was precipitated from a dichloromethane/hexane mixture in 76% yield. 1,3-bis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-*p-tert*-butylcalix[4]arene dibromide (**2**) was obtained during the reaction of **S**<sub>2</sub> with 10 equivalents of diphenylmethylphosphine in the same conditions in 69% yield.

### Scheme 1

The more lipophilic molecules **3** and **4**, where the free phenolic protons are substituted with propyl groups, were prepared in order to increase their stability in the lipophilic membrane of ion selective electrodes. The reaction of **S**<sub>3</sub> with 10 equivalents of 1,4-dibromobutyl, refluxed for 4 days in dimethylformamide in the presence of 7 equivalents of NaH gave the 1,3-bis-(propoxy)-2,4-bis-(butoxy-4-bromide)-tetrakis-*p-tert*-butylcalix[4]arene (**S**<sub>4</sub>) in 57% yield. Compounds **3** and **4** were obtained by the reaction of **S**<sub>4</sub> with 10 equivalents of triphenylphosphine and 10 equivalents of diphenylmethylphosphine in chloroform, in 76% and 64% yield, respectively (Scheme 1).

Tetraphosphonium ligands **6** and **7** were synthesised according to the procedure already described for **5** [22] from the intermediate molecule tetrakis-(butoxy-4-bromide)-*p-tert*-butylcalix[4]arene (**S**<sub>5</sub>) by substitution of the bromine atoms of the alkyl chains (Scheme 2). The tetrakis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-*p-tert*-butylcalix[4]arene tetrabromide (**6**) was obtained from the reaction of **S**<sub>5</sub> with 20 equivalents of diphenylmethylphosphine in chloroform with 59% yield.

*Scheme 2*

Ligand **7** was synthesised in three steps. The reaction of **S<sub>5</sub>** with 5 equivalents of KPPH<sub>2</sub> gave the product **S<sub>6</sub>** in 40% yield [25,26,27]. **S<sub>6</sub>** was then protonated with an excess of HBr giving ligand **7** in 52 % yield.

In order to study the influence of the ligand counterion on the ligand – anion interactions, the tetra substituted phosphonium ligands were also synthesised as perchlorates (**5a**, **6a**) and hexafluorophosphates (**5b**, **6b**). These compounds were obtained by reacting ligands **5** and **6** with the appropriate silver salts.

For comparison purpose, the monomeric subunit of ligand **5**, the triphenylphosphonium-butoxy-*p-tert*-butylphenol bromide (**8**) was synthesized as previously described [22].

The cone conformation of the di-substituted calix[4]arenes was indicated by the presence in their <sup>1</sup>H NMR spectra of two singlets for the *tert*-butyl protons (at 0.99 and 1.28 ppm for **1** and 0.96 and 1.29 ppm for **2**) and a AB system for the methylene protons (at 3.19 and 3.90 ppm for **1** and at 3.23 and 4.02 ppm for **2**). The <sup>1</sup>H NMR spectra of ligands **5** and **6** are characteristic of tetrasubstituted derivatives of calix[4]arenes in the cone conformation. For instance in the case of **6**, it is indicated by the presence of the singlet corresponding to the protons of the *tert*-butyl groups at 1.03 ppm, the AB system of the methylene protons of the calixarene at 4.26 and 3.03 ppm. Two well defined multiplets for the protons of the aromatic phosphonium groups can be observed as well as one doublet for the methyl protons in direct neighbourhood of the phosphorus atoms.

The spectrum of the protonated phosphonium ligand **7** presents several broad peaks corresponding to multiplets in coalescence. Only two singlets could be clearly observed, one for the *tert*-butyl groups and one for the aromatic protons of the calix[4]arene, suggesting also the cone conformation of this molecule.

**Binding studies****<sup>1</sup>H and <sup>31</sup>P NMR studies in chloroform**

The interactions between the phosphonium calixarenes and the following anions: NO<sub>3</sub><sup>-</sup>, ClO<sub>4</sub><sup>-</sup>, I<sup>-</sup>, SCN<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, HCO<sub>3</sub><sup>-</sup>, Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> provided as sodium salts were studied by <sup>1</sup>H and <sup>31</sup>P NMR in CDCl<sub>3</sub>. Among these anions, only perchlorate, thiocyanate and iodide salts induced changes in the <sup>1</sup>H NMR spectra of the ligands. The changes observed in the case of the disubstituted ligand **1** upon addition of sodium perchlorate are illustrated in Fig. 2. The main signals affected were those of the CH<sub>2</sub> protons (i, j) directly bound to the carbon

atoms next to the phosphorus atoms and the aromatic protons (k) of the phosphonium groups (see Table S1, Supplementary Online Material). The shifts of these signals due to changes in the charge density on the phosphorus atoms reflect interactions with these anions.

*Figure 2*

Similar but smaller changes were induced in the spectrum of ligand **2** by the presence of the same anions (see Table S2, Supplementary Online Material). As with ligand **1**, no change is observed for the aromatic protons (c, d) and for the protons (e, f) of the methylene bridge, indicating that the conformation of the calixarene unit is not disturbed. This may be explained by the high rigidity of these disubstituted derivatives due to hydrogen bonds involving the free phenolic groups.

In the  $^{31}\text{P}$  NMR spectra of the free ligands **1** and **2**, and of these ligands in the presence of sodium iodide, thiocyanate and perchlorate, the phosphorus atoms appear as a singlet indicating that the two phosphonium groups are chemically equivalent and participate in the anion-ligand interaction. The most important changes in chemical shifts are observed for perchlorate:  $\Delta\delta = 0.602$  and  $0.620$  ppm with **1** and **2**, respectively (Table 1).

*Table 1*

With the tetrasubstituted calix[4]arene **5** previously studied [22] and  $\text{ClO}_4^-$ ,  $\text{SCN}^-$  and  $\text{I}^-$  the most important changes of chemical shifts corresponded to the signals of the methylene bridge protons and the aromatic protons of the calixarene, indicating changes in the conformational tensions of the calixarene scaffold. The signals corresponding to protons of the butyl chains were also shifted as well as the aromatic protons of the phosphonium. On the contrary no change was observed with the monomer **8** [22].

The spectrum of ligand **6** was also modified in the presence of these anions (see Table S3, Supplementary Online Material). In particular the signals corresponding to the protons of the calix[4]arene scaffold are moved in a comparable way for the three anions. The changes observed in the spectrum of this ligand in the presence of  $\text{ClO}_4^-$  are illustrated in Fig. 3.

*Figure 3*

With this anion, the multiplet corresponding to the protons (g) and (j) from 3.87 to 3.70 ppm gives two multiplets from 4.36 to 4.17 ppm for (g) and from 3.28 to 3.11 ppm for (j). The multiplet from 8.17 to 8.02 ppm for protons (k) is shifted upfield and gives one multiplet from 7.90 to 7.70 ppm, whereas the multiplet from 7.78 to 7.54 ppm of protons (l, m) does not move significantly. The doublet of the  $\text{CH}_3$  protons (o) adjacent to the phosphorus atoms at 2.91 ppm is strongly shifted to 2.50 ppm.

1  
2  
3 The addition of the anions studied to **7** did not induce any change in its  $^1\text{H}$  NMR  
4 spectrum. Especially the signals of the protons of the phosphonium moieties, expected to be  
5 involved in hydrogen bond formation, were not shifted.  
6  
7

8  
9 With ligand **6** as with **5**, only one singlet for the phosphorus atoms was detected in its  $^{31}\text{P}$   
10 NMR spectrum indicating all the four phosphonium groups being chemically equivalent. In  
11 the presence of sodium perchlorate and thiocyanate, the changes in chemical shifts are larger  
12 than those observed with the disubstituted derivatives and suggest stronger interactions (Table  
13 1). With all ligands the most important values were observed for perchlorate.  
14  
15  
16  
17

### 18 19 $^1\text{H}$ NMR studies in acetonitrile

20  
21  $^1\text{H}$  NMR experiments were repeated with **5** and sodium perchlorate in deuterated  
22 acetonitrile, a more dissociating solvent, in which association phenomena are not as  
23 important as in chloroform [22]. The spectra of the ligand are very similar in both  
24 solvents. In acetonitrile the addition of  $\text{NaClO}_4$  induces shifts of the same signals as in  
25 chloroform. Moreover this study showed the influence of the counterion of the salts on the  
26 shifts observed in the spectra (See Table S4, Supplementary Online Material) . The signal  
27 of the methylene protons (g) adjacent to the phenolic oxygen atoms was shifted only with  
28  $\text{NaClO}_4$  and  $\text{LiClO}_4$ . With both salts the shifts corresponding to the signals of the protons  
29 (j) next to the phosphorus atoms (which are supposed to interact with perchlorate) were  
30 similar, whereas with  $\text{Et}_4\text{NClO}_4$  and especially with  $\text{CsClO}_4$ , the values were very small.  
31 The changes observed suggested that ligand – anion interaction was connected with the  
32 nature of the counterion and its affinity for the ligand.  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 In order to observe the influence of the ligand counterion, bromide anions were  
43 replaced by the more lipophilic perchlorate (**5a** and **6a**) or hexafluorophosphate (**5b** and  
44 **6b**) anions. The chemical shifts ( $\delta$ ) of selected protons, given in Table 2, show only slight  
45 differences for protons (c, e, f) and (g) close to the calix[4]arene scaffold ( $\Delta\delta$  in the range  
46 0.02 – 0.07 ppm). In contrast, the signals of the protons (j) next to the charged phosphorus  
47 atoms are greatly shifted ( $\Delta\delta = -0.64$  ppm for **5a**,  $-0.75$  ppm for **5b** and  $\Delta\delta = -0.73$  ppm  
48 for **6a**,  $-0.83$  ppm for **6b**). With ligands **6a** and **6b**, the signals of the protons (o) of the  
49 methyl substituents are also displaced ( $\Delta\delta = -0.41$  ppm for **6a** and  $-0.46$  ppm for **6b**).  
50 These results show that the chemical shifts of the protons next to the phosphonium groups  
51 are influenced by the ligand counterion.  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 2

1  
2  
3 On the other hand, it was also shown that when the lipophilic  $\text{PF}_6^-$  anions were  
4 replacing the  $\text{Br}^-$  counterions of the ligand, there was still a significant shift of the protons  
5 (j) close to the phosphorus atoms for ligand **5b** in the presence of  $\text{ClO}_4^-$  (Table S4) [22].  
6  
7 This result suggested complexation of perchlorate with this ligand, where no exchange is  
8 normally possible.  
9  
10  
11

### 12 13 14 **Microcalorimetric studies in acetonitrile**

15 In order to get more information on the influence of the counterion of the salt,  
16 microcalorimetric titrations were carried out with ligands **5** and **5b** against  $\text{NaClO}_4$ ,  
17  $\text{LiClO}_4$  and  $\text{Et}_4\text{NClO}_4$  in acetonitrile. The thermograms recorded during the titration of  
18 these ligands with  $\text{NaClO}_4$  and  $\text{LiClO}_4$  showed significant exothermic heat effects,  
19 whereas their titration with  $\text{Et}_4\text{NClO}_4$  led to no thermal effect (see Figure S1,  
20 Supplementary Online Material). For comparison purpose, the titration of the monomer **8**  
21 with  $\text{NaClO}_4$  was also carried out showing no significant heat effect.  
22  
23  
24  
25  
26  
27

28 If only the  $\text{ClO}_4^-$  anion were involved in the complexation, a similar heat effect  
29 should be observed in all the titrations. The fact that an effect is only observed for  $\text{NaClO}_4$   
30 and  $\text{LiClO}_4$  suggests that it is not only related to the anion interaction (complexation or  
31 anion exchange). Assuming that the large tetraethylammonium cation cannot be  
32 complexed with a calix[4]arene, the heat effects observed during the titration with  $\text{NaClO}_4$   
33 and  $\text{LiClO}_4$  would rather be due to the complexation of the cations with the two ligands.  
34 This is supported by the fact that no heat effect was detected with the monomer **8**,  
35 supposed to be unable to complex these cations.  
36  
37  
38  
39  
40  
41

42 Calorimetric data obtained with  $\text{NaClO}_4$  and  $\text{LiClO}_4$  were interpreted assuming  
43 different cation complexation models. With sodium and both ligands, the best fit was  
44 obtained by considering the presence of  $\text{ML}$  and  $\text{ML}_2$  complexes. The same species were  
45 found with  $\text{LiClO}_4$  and **5b**, whereas only a 1:1 complex was formed with **5**. The formation  
46 of  $\text{ML}_2$  species could be explained by the complexation of the ion pair  $\text{Na}^+\text{ClO}_4^-$  by two  
47 ligands. The stability constants of these complexes are given in Table S5 (see  
48 Supplementary Online Material).  
49  
50  
51  
52  
53  
54

55 An important heat effect was observed during the titration of ligands **5** and **5b**  
56 against  $\text{LiBr}$ , which should be related directly to cation complexation as no anion  
57 exchange is possible with these ligands. The data interpretation led to species of the same  
58 stoichiometry as with  $\text{LiClO}_4$  (Table S5). The values of the stability constants of 1:1  
59 species formed in the presence of  $\text{LiClO}_4$  and  $\text{LiBr}$  are comparable (with **5**,  $\log \beta = 3.24$   
60

1  
2  
3 and 3.15, respectively, and with **5b**,  $\log \beta = 4.29$  and 4.32, respectively). They are lower  
4  
5 than that of the complex formed by the tetra-methylated *p-tert*-butylcalix[4]arene with  
6  
7 lithium ( $\log \beta = 5.10$  in acetonitrile [28]).  
8  
9

10 All that considered, the UV spectrophotometric titrations of ligand **5** against  $\text{ClO}_4^-$   
11 previously performed may certainly be interpreted in terms of cation rather than anion  
12 complexation [22]. The values of the stability constants of the 1:1 complexes with sodium  
13 perchlorate ( $\log \beta = 3.81 \pm 0.02$ ) and with lithium perchlorate ( $\log \beta = 3.71 \pm 0.04$ ) [22] are  
14  
15 of the same order of magnitude as those obtained from microcalorimetric measurements.  
16  
17  
18  
19

### 20 21 22 **Potentiometric studies** 23

24 Only few ligands containing phosphorus atoms have been studied so far as active material in  
25 ion selective membrane electrodes [29,30,31]. They showed a selective response for  $\text{ClO}_4^-$ ,  
26 with, however, little discrimination with respect to  $\text{I}^-$  and  $\text{SCN}^-$ . The results suggested a  
27 particular affinity of ligands containing phosphorus atoms for  $\text{ClO}_4^-$ ,  $\text{SCN}^-$  and  $\text{I}^-$  anions. By  
28 attaching phosphonium moieties to a calix[4]arene scaffold and taking advantage of the  
29 preorganization of the ligand, it was expected to enhance the selectivity for tetrahedral or  
30 spherical anions. Such selectivity (especially for  $\text{ClO}_4^-$  over  $\text{I}^-$ ) is hard to obtain with other  
31 kinds of receptors.  
32  
33  
34  
35  
36  
37

38 Disubstituted phosphonium ligands **1** and **2** were tested as ionophores in the  
39 membrane electrodes. The electrodes were sensitive to perchlorate, thiocyanate, iodide  
40 and nitrate, showing fast, near-Nernstian responses (Figure 4 and Table S6 of  
41 Supplementary Online Material). However, their characteristics changed with time.  
42 Attempts to optimize the composition of the conditioning solutions as well as the  
43 conditioning time did not improve the situation which might be due to slow leakage of the  
44 ionophores from the membrane. These ligands had also the tendency to crystallize in the  
45 membrane phase. Crystallization of our ligands within the membranes depends strongly  
46 on the kind of plasticizer used. The ligands in the membranes based on bis-(2-  
47 ethylhexyl)sebacate (BEHS) has crystallized strongly, which decreased their stability.  
48 This phenomenon originates from the higher lipophilicity of this plasticizer ( $\log P = 10.1$ )  
49 as compared to 2-nitrophenyloctylether (*o*-NPOE) ( $\log P = 5.9$ ) [32], which is more  
50 suitable for charged ligands. Electrodes with membranes based on *o*-NPOE had the best  
51 lifetime and response characteristics and were chosen for further studies.  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Figure 4*

The more lipophilic ligands **3** and **4**, in which n-propyl chains replace the two phenolic OH groups, were also synthesised. The lifetime of electrodes incorporating these ligands for perchlorate was increased to at least three weeks. The repeatability of the measurements was also good, but their detection limits increased (see Table S7, Supplementary Online Material).

The membranes of electrodes incorporating the tetrasubstituted ligand **6** showed rather quick (within 15-20 s), stable and fully reversible responses (Figure 5 and Table S8 of Supplementary Online Material). The repeatability of the measurements was also good and their lifetime was more than three weeks. They showed close to Nernstian response to  $\text{ClO}_4^-$ ,  $\text{I}^-$ ,  $\text{NO}_3^-$  and  $\text{SCN}^-$  and no significant response for  $\text{SO}_4^{2-}$ ,  $\text{CO}_3^{2-}$ ,  $\text{HPO}_4^{2-}$ ,  $\text{PO}_4^{3-}$ . A similar behaviour was already observed with electrodes incorporating ligand **5** (Table S8). The highest selectivity was obtained for  $\text{ClO}_4^-$  ions in buffered solution (pH=5.5) and in water (pH=6.5). The over-Nernstian slope of the electrode response for  $\text{Cr}_2\text{O}_7^{2-}$  could indicate a mechanism where both processes, anion complexation and anion exchange, play an important role. It can also be explained as the presence in the sample of different forms of chromates.

*Figure 5*

While the addition of lipophilic anionic sites (KTCIPB) to the membranes in the case of ligand **5** did not change much the properties of the electrodes, it affected the properties of the electrodes containing **6** (Table 3). Without salt, the slope of the electrode is -54.2 mV, and slightly decreases to -51.8 and -49.9 mV, respectively, in the presence of salt. According to the literature data [29,31], such results suggest that none of the ligands works as a neutral carrier because the addition of the lipophilic anion to the membrane does not induce a cationic response of the potentiometric cell. Ligand **6** seems to work in the membrane as a typical anion-exchanger, whereas ligand **5** could be considered as a charged ligand despite the small influence of KTCIPB.

*Table 3*

The influence of the ligand counterions ( $\text{Br}^-$ ,  $\text{ClO}_4^-$ ,  $\text{PF}_6^-$ ) on the properties of the membrane electrodes was also studied. Table 4 compares the responses to perchlorate of electrodes based on ligands **5a**, **6a** (perchlorates) and on ligands **5b** and **6b** (hexafluorophosphates) to corresponding electrodes based on ligands **5** and **6** (bromides). The less good slope of the electrodes containing **6a** and **6b** as compared to that of electrodes containing **6** suggests rather the anion-exchange nature of the latter ligand. In



1  
2  
3 this case the presence of more lipophilic anions (perchlorate or hexafluorophosphate)  
4 slows down the anion exchange process. In contrast, the properties of electrodes with the  
5 bromide ligand **5** and with the perchlorate ligand **5a** are comparable. The presence of  
6 highly lipophilic perchlorate anions does not disturb the electrode response, while the  
7 presence of hexafluorophosphate anions in the membrane phase (electrode containing  
8 ligand **5b**) decreases the slope and the linearity range.  
9

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
*Table 4*

The poorer properties of the electrode containing **5b** and **6b** could be explained by the higher lipophilicity of hexafluorophosphate anions which hinders the process of anion exchange. The complexation of  $\text{ClO}_4^-$  by ligand **5** could explain the good response of electrodes based on **5** and **5a** to perchlorate. Such interpretation is consistent with the results of previous experiments (Table 4) and indicates that ligand **5** behaves more like a charged carrier for  $\text{ClO}_4^-$ , while ligand **6** behaves more as an anion-exchanger.

The order of selectivity observed with all phosphonium ligands **1-6** and **8** follows the Hofmeister series:



The highest selectivity is observed for perchlorate (Table 5).

*Table 5*

The electrodes based on tetraphosphonium derivatives show higher selectivities for perchlorate over thiocyanate, iodide and nitrate than those based on their di-phosphonium counterparts. These selectivities are also better than the selectivity of electrodes based on the monomer **8**.

The replacement of one phenyl substituent on the phosphorus atoms by one methyl group does not change significantly the selectivity pattern and the values of the selectivity coefficients of electrodes containing either di- or tetraphosphonium ligands. The only exception is the selectivity of electrodes based on compound **5** against  $\text{Cr}_2\text{O}_7^{2-}$  which increases from 2.6 to 3.2 log units.

Electrodes based on alkylated compounds **3** and **4** are also selective for perchlorate but the selectivity over iodide and thiocyanate is decreased (Table 5).

Ligands **5** and **6** display better potentiometric properties than the protonated cyclam [33] or its copper complex [34] and than a phosphodithia macrocycle [35]. For instance the detection limit is  $2.5 \times 10^{-7}$  M with ligand **5** when compared to  $4.2 \times 10^{-6}$  M for cyclam and  $8 \times 10^{-7}$  M for the phosphodithiamacrocycle. Ligands **6** as **5** presents generally higher selectivities than TDMACl [36], the protonated cyclam and  $[\text{Cu}(\text{cyclam})]^{2+}$ .

## CONCLUDING REMARKS

The different techniques used to assess the binding properties of phosphonium derivatives showed strong interactions with  $\text{SCN}^-$ ,  $\text{I}^-$  and especially  $\text{ClO}_4^-$  and pointed out the important role played by the salt counterion ( $\text{Na}^+$  or  $\text{Li}^+$ ), which may be complexed by the calix[4]arene. Incorporated in PVC membrane electrodes, these molecules are efficient sensing material for anions with a selectivity order following the Hofmeister series generally observed for ion exchangers. However, the electrodes based on tetraphosphonium derivatives showed better selectivities than those based on the diphosphonium analogues or on the monomeric unit, indicating the importance of the ligand preorganisation which should not be observed in the case of simple anion-exchangers.

A question which must be addressed concerns the nature of the interaction between the ligands and the anions, e.g.  $\text{ClO}_4^-$ . Is it a simple ion-exchange between the bromides of the ligand and this more lipophilic anion, or is it complexation within the charged phosphonium groups? What is the exact role of the salt counterion?

If NMR gives some indications on the changes in the molecule, suggesting interactions, it does not tell if there is complexation or anion exchange, since the nature of the counterion of the ligand has been shown to influence the chemical shifts of the protons near the charged atoms. In favour of ion-exchange is the fact that the most important shifts are observed with the more lipophilic anions  $\text{ClO}_4^-$ ,  $\text{SCN}^-$  and  $\text{I}^-$ . The behaviour in selective electrodes is consistent also with this assumption. However, the fact that no change occurred in the spectrum of the monomer, where only exchange is possible, is against this hypothesis. In favour of complexation is the fact that, in the presence of  $\text{NaClO}_4$ , strong shifts are observed for the signal of the protons next to the phosphonium groups in the spectrum of the hexafluorophosphate ligand where no exchange is possible. On the other hand  $^1\text{H}$  NMR and microcalorimetry emphasized the importance of the cation which can be complexed in the cavity of the calixarene. With perchlorate, the best interaction takes place with  $\text{Na}^+$  and  $\text{Li}^+$ , whereas little or no interaction occurs with the larger  $\text{Et}_4\text{N}^+$  and  $\text{Cs}^+$ . It can be noted also that the anion plays also a role in the complexation of the cation, since the spectrum of the calixarene part is not affected in the presence of  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{HCO}_3^-$  and  $\text{Cr}_2\text{O}_7^{2-}$ , i.e. the less lipophilic ones.

## EXPERIMENTAL

FAB mass spectra were obtained on a VG analytical ZAB HF instrument. All reagents and solvents were commercial and used without further purification.

Chromatography columns were prepared from Kieselgel Merck Si 60(40-63  $\mu\text{m}$ ). TLC was performed on 250  $\mu\text{m}$  silica gel plates Merck containing a fluorescent indicator.

### Synthesis of intermediate compounds

#### *1,3-bis-(butoxy-4-bromide)-p-tert-butylcalix[4]arene (S<sub>2</sub>)*

Into a 250  $\text{cm}^3$  flask containing (3.244 g, 5.00 mmol) of tetrakis-*p-tert*-butylcalix[4]arene **S<sub>1</sub>** and acetone (50  $\text{cm}^3$ ), (1.383 g, 10.00 mmol) of  $\text{K}_2\text{CO}_3$  were added. The mixture was stirred at room temperature for 2h. Then (3.236 g, 15.00 mmol) of 1,4-butyl-dibromide in acetone (50  $\text{cm}^3$ ) were added. The mixture was left for 4 days under reflux. After four days 5  $\text{cm}^3$  of methanol were added. The solvents were evaporated, and the reaction mixture was dissolved in 100  $\text{cm}^3$  of dichloromethane. After extraction with 150  $\text{cm}^3$  of water the organic phase was dried with  $\text{Na}_2\text{SO}_4$  and evaporated. The crude product was purified by crystallization from a 1/10 dichloromethane/methanol mixture giving (3.252 g, 3.54 mmol) of compound **S<sub>2</sub>** in 71 % yield. Mp > 280 °C.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 0.98 (s, 18H, C-( $\text{CH}_3$ )<sub>3</sub>), 1.30 (s, 18H, C-( $\text{CH}_3$ )<sub>3</sub>), 2.15 (qn, 4H,  $J = 4.4$ ,  $\text{CH}_2$ - $\text{CH}_2$ -O), 2.32 (qn, 4H,  $J = 4.2$ ,  $\text{CH}_2$ - $\text{CH}_2$ -Br), 3.32 (d, 4H,  $J = 13.0$ , Ar- $\text{CH}_2$ -Ar), 3.65 (t, 4H,  $J = 6.6$ ,  $\text{CH}_2$ - $\text{CH}_2$ -Br), 4.01 (t, 4H,  $J = 5.8$ ,  $\text{CH}_2$ - $\text{CH}_2$ -O), 4.25 (d, 4H,  $J = 13.0$ , Ar- $\text{CH}_2$ -Ar), 6.78 (s, 4H, Ar- $H$ ), 7.08 (s, 4H, Ar- $H$ ), 7.40 (s, 2H, OH). Anal. Calcd. for  $\text{C}_{52}\text{H}_{70}\text{O}_4\text{Br}_2$ : C, 67.97; H, 7.68. Found: C, 68.21; H, 7.94.

#### *1,3-bis-(propoxy)-p-tert-butylcalix[4]arene (S<sub>3</sub>)*

Into a 250  $\text{cm}^3$  flask containing (3.244g, 5.00 mmol) of tetrakis-*p-tert*-butylcalix[4]arene **S<sub>1</sub>** and acetone (50  $\text{cm}^3$ ), (1.383g, 10.00 mmol) of  $\text{K}_2\text{CO}_3$  were added. The mixture was stirred at room temperature for 2h. Then (1.845 g, 15.00 mmol) of bromopropane in acetone (40  $\text{cm}^3$ ) were added. The mixture was left for 4 days under reflux. After four days 5  $\text{cm}^3$  of methanol were added. The solvents were evaporated, and the reaction mixture was dissolved in 100  $\text{cm}^3$  dichloromethane. After extraction with 150  $\text{cm}^3$  of water the organic phase was dried with  $\text{Na}_2\text{SO}_4$  and evaporated. The crude product was purified by crystallization from a 1/9 acetone/methanol mixture giving (2.016 g, 2.75 mmol) of pure compound **S<sub>3</sub>** in 55 % yield. Mp > 280 °C.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.02 (s, 18H, C-( $\text{CH}_3$ )<sub>3</sub>), 1.27 (t, 6H,  $J = 5.0$ ,  $\text{CH}_3$ - $\text{CH}_2$ -), 1.28 (s, 18H, C-( $\text{CH}_3$ )<sub>3</sub>),

2.06 (sx, 4H,  $J = 4.9$ ,  $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-O}$ ), 3.32 (d, 4H,  $J = 12.8$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 3.96 (t, 4H,  $J = 5.9$ ,  $\text{O-CH}_2\text{-CH}_2$ ), 4.31 (d, 4H,  $J = 12.8$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 6.86 (s, 4H,  $\text{Ar-H}$ ), 7.05 (s, 4H,  $\text{Ar-H}$ ), 7.89 (s, 2H,  $\text{OH}$ ). Anal. Calcd for  $\text{C}_{50}\text{H}_{68}\text{O}_4$ : C, 81.92; H, 9.35. Found: C, 82.05; H, 9.40.

### 1,3-bis-(propoxy)-2,4-bis-(butoxy-4-bromide)-tetrakis-*p*-tert-butylcalix[4]arene (**S**<sub>4</sub>)

Into a 250 cm<sup>3</sup> flask containing (2.016 g, 2.75 mmol) of **S**<sub>3</sub> and DMF (50 cm<sup>3</sup>), (0.480 g, 20.00 mmol) of NaH were added. NaH was washed twice with hexane before addition. The mixture was stirred at room temperature for 4h. After this time, (5.940 g, 27.5 mmol) of 1,4-dibromobutyl in DMF (40 cm<sup>3</sup>) were added. The mixture was left for 2 days at 80-90 °C. After 2 days 30 cm<sup>3</sup> of methanol were added. The solvents were evaporated and the reaction mixture was dissolved in 100 cm<sup>3</sup> of dichloromethane. After extraction with 250 cm<sup>3</sup> of water the organic phase was dried with  $\text{Na}_2\text{SO}_4$  and evaporated. The crude product was purified by crystallization from a 1/10 dichloromethane/methanol mixture giving (1.584 g, 1.57 mmol) of compound **S**<sub>4</sub> in 57 % yield. Mp 170 °C. <sup>1</sup>H NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.01 (t, 6H,  $J = 5.2$ ,  $\text{CH}_3\text{-CH}_2\text{-}$ ), 1.05 (s, 18H,  $\text{C-(CH}_3)_3$ ), 1.12 (s, 18H,  $\text{C-(CH}_3)_3$ ), 1.99-2.05 (m, 4H,  $\text{CH}_3\text{-CH}_2\text{-}$ ), 2.07-2.25 (m, 8H,  $\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-Br}$ ), 3.13 (d, 4H,  $J = 13.0$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 3.51 (t, 4H,  $J = 6.6$ ,  $\text{-CH}_2\text{-Br}$ ), 3.81 (t, 4H,  $J = 5.8$ ,  $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{-O}$ ), 3.91 (t, 4H,  $J = 5.8$ ,  $\text{Br-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-O}$ ), 4.39 (d, 4H,  $J = 13.0$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 6.74 (s, 4H,  $\text{Ar-H}$ ), 6.83 (s, 4H,  $\text{Ar-H}$ ). Anal. Calcd for  $\text{C}_{58}\text{H}_{82}\text{O}_4\text{Br}_2$ : C, 69.45; H, 8.24. Found: C, 69.65; H, 8.30.

### Tetrakis-(butoxy-4-bromide)-tetrakis-*p*-tert-butylcalix[4]arene (**S**<sub>5</sub>)

The suspension of *p*-tert-butylcalix[4]arene **S**<sub>1</sub> (1.947 g, 3.00 mmol) and NaH in oil washed three times with hexane (0.700 g, 29.17 mmol) were stirred at room temperature in DMF (50 cm<sup>3</sup>) for 1h. Then 1,4-dibromobutane (12.947 g, 59.06 mmol) was added and the mixture was heated to 80 °C. After 4 days of heating, the mixture was cooled and MeOH (20 cm<sup>3</sup>) was added. After removing of the solvent, the residue was dissolved in dichloromethane and water and acidified with 1M HCl. The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and evaporated. After precipitation from methanol the pure compound **S**<sub>5</sub> (1.520 g, 1.28 mmol) was obtained in 43% yield. Mp 180 °C. <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  [ppm]: 1.09 (s, 36H,  $\text{-C(CH}_3)_3$ ), 1.98-2.09 (m, 8H,  $\text{-CH}_2\text{-}$ ), 2.16-2.23 (m, 8H,  $\text{-CH}_2\text{-}$ ), 3.15 (d, 4H,  $J = 13.0$ ,  $\text{Ar-CH}_2\text{-Ar}$ ), 3.53 (t, 8H,  $J = 6.9$ ,  $\text{-CH}_2\text{-Br}$ ), 3.91 (t, 8H,  $J = 6.9$ , -

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

$CH_2-O$ ), 4.36 (d, 4H,  $J = 13.0$ , Ar- $CH_2$ -Ar), 6.79, (s, 8H, Ar- $H$ ). Anal. Calcd. for  $C_{60}H_{84}O_4Br_4$ : C, 60.61; H, 7.12; Found: C, 60.87; H, 7.32.

### Tetrakis-(4-(diphenylphosphine)-butoxy)-*p*-*tert*-butylcalix[4]arene ( $S_6$ )

Into a 100 cm<sup>3</sup> flask containing compound  $S_5$  (1.510 g, 1.27 mmol) of freshly distilled THF (10 cm<sup>3</sup>), (1.282 g, 5.72 mmol) of KP(Ph)<sub>2</sub> in THF (15 cm<sup>3</sup>) were added via a syringe. The mixture was stirred for 2 hours at room temperature, during which the colour of the reaction mixture changed from red to dark yellow. The mixture was then evaporated and extracted twice with 30 cm<sup>3</sup> of dichloromethane. Purification of the crude product on silica column with a 3/7 dichloromethane/hexane mixture as eluent gave compound  $S_6$  (0.409 g, 0.26 mmol) in 20 % yield. Mp 110 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  [ppm]: 1.08 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.42-1.63 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.00-2.18 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.00-2.18 (m, 8H, -CH<sub>2</sub>-P), 3.05 (d, 4H,  $J = 12.9$ , Ar- $CH_2$ -Ar), 3.81 (t, 8H,  $J = 5.4$ , -CH<sub>2</sub>-CH<sub>2</sub>-O), 4.29 (d, 4H,  $J = 12.9$ , Ar- $CH_2$ -Ar), 6.75 (s, 8H, Ar- $H$ ), 7.20-7.46 (m, 40H, P-Ar- $H$ ). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  [ppm]: -14.91 [P].  $m/z$  (MALDI) 1610.88 (M + H)<sup>+</sup>. Anal. Calcd. for  $C_{108}H_{124}O_4P_4$ : C, 80.57; H, 7.76; Found: C, 80.73; H, 7.87.

### Synthesis of phosphonium ligands

#### 1,3-bis-(4-triphenylphosphonium-butoxy)-*p*-*tert*-butylcalix[4]arene dibromide (**1**)

Into a 100 cm<sup>3</sup> flask containing  $S_2$  (1.184 g, 2.00 mmol) in chloroform (30 cm<sup>3</sup>), (5.248 g, 20.00 mmol) of triphenylphosphine in chloroform (20 cm<sup>3</sup>) were added. After 6 days under reflux, the mixture was cooled and the solvent evaporated. The residue was dissolved in dichloromethane. The excess of triphenylphosphine was precipitated from methanol and filtered out. The filtrate was evaporated. The pure product **1** (2.179 g, 1.51 mmol) was obtained by precipitation from a 1/9 dichloromethane/hexane mixture, as a white-light green powder in 76 % yield: Mp > 280 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  [ppm]: 0.99 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.28 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 2.05-2.29 (m, 8H, CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 3.19 (d, 4H,  $J = 12.8$ , Ar- $CH_2$ -Ar), 3.80-4.00 (m, 4H, -CH<sub>2</sub>-O), 3.90 (d, 4H,  $J = 12.8$ , Ar- $CH_2$ -Ar), 3.90-4.08 (m, 4H, -CH<sub>2</sub>-P), 6.79 (s, 4H, Ar- $H$ ), 6.99 (s, 4H, Ar- $H$ ), 7.49 (s, 2H, OH), 7.54 – 7.64 (m, 12H, P-Ar- $H$  meta), 7.65-7.74 (m, 6H, P-Ar- $H$  para), 7.80-7.93 (m, 12H, P-Ar- $H$  ortho). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  [ppm]: 25.80.  $m/z$  (FAB<sup>+</sup>) 721.7 (M + 2H)<sup>2+</sup>;  $m/z$  (MALDI) 1361.7 (M - Br)<sup>+</sup>. Anal. Calcd for  $C_{88}H_{100}O_4P_2Br_2$ : C, 73.22; H, 6.98. Found: C, 73.46; H, 7.20.

1  
2  
3  
4  
5 **1,3-bis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-p-tert-butylcalix[4]arene**  
6 **dibromide (2)**  
7

8  
9 Compound **2** was prepared following the same procedure as for compound **1** with **S**<sub>2</sub>  
10 (2.753 g, 3.00 mmol) and diphenylmethylphosphine (6.006 g, 30.00 mmol) in 69 % yield.  
11  
12 Mp > 280 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 0.96 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.29 (s, 18H,  
13 -C(CH<sub>3</sub>)<sub>3</sub>), 1.91-2.10 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.11-2.27 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.99 (d, 6H,  
14 J = 13.8, CH<sub>3</sub>-P), 3.23 (d, 4H, J = 13.5, Ar-CH<sub>2</sub>-Ar), 3.58-3.75 (m, 4H, -CH<sub>2</sub>-P), 3.90 (t,  
15 4H, J = 5.3, -CH<sub>2</sub>-O), 4.02 (d, 4H, J = 13.5, Ar-CH<sub>2</sub>-Ar), 6.76 (s, 4H, Ar-H), 7.01 (s, 4H,  
16 Ar-H), 7.31 (s, 2H, OH), 7.48-7.59 (m, 8 H, P-Ar-H meta), 7.61-7.70 (m, 4H, P-Ar-H  
17 para), 7.90-8.06 (m, 8H, P-Ar-H ortho). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 25.96. m/z  
18 (MALDI) 1239.58 (M - Br)<sup>+</sup>. Anal. Calcd. for C<sub>78</sub>H<sub>96</sub>O<sub>4</sub>P<sub>2</sub>Br<sub>2</sub>: C, 71.01; H, 7.33. Found:  
19 C, 71.21; H, 7.52.  
20  
21  
22  
23  
24  
25  
26  
27

28 **1,3-bis-(4-triphenylphosphonium-butoxy)-2,4-bis-propoxy-p-tert-butyl-calix[4]arene**  
29 **dibromide (3)**  
30

31 Into a 100 cm<sup>3</sup> flask containing **S**<sub>4</sub> (2.012 g, 2.00 mmol) in chloroform (30 cm<sup>3</sup>), 5.248 g  
32 of triphenylphosphine (20.00 mmol) in chloroform (20 cm<sup>3</sup>) were added and left for 6  
33 days under reflux. After that time the mixture was cooled and the solvent evaporated. The  
34 residue was dissolved in dichloromethane. The excess of triphenylphosphine was  
35 precipitated from methanol and filtered off. The filtrate was evaporated. The pure product  
36 **3** (2.179g, 1.51 mmol) was obtained by precipitation from a 1/9 dichloromethane/hexane  
37 mixture, as a white-light green powder in 76% yield. Mp 120 °C. <sup>1</sup>H NMR (300 MHz,  
38 CDCl<sub>3</sub>) δ [ppm]: 0.86 (t, 6H, J = 5.3, CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 0.99 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.11 (s,  
39 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.65-1.87 (m, 4H, CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 1.77-1.94 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-P),  
40 2.28-2.43 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.95 (d, 4H, J = 13.0, Ar-CH<sub>2</sub>-Ar), 3.63 (t, 4H, J = 5.9,  
41 CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 3.85-4.00 (m, 4H, -CH<sub>2</sub>-P), 3.85-4.00 (m, 4H, -CH<sub>2</sub>-O), 4.18 (d, 4H, J  
42 = 13.0, Ar-CH<sub>2</sub>-Ar), 6.61 (s, 4H, Ar-H), 6.76 (s, 4H, Ar-H), 7.60-7.92 (m, 30H, P-Ar-H  
43 ortho, meta, para). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 25.84. m/z (MALDI) 1447.6 (M  
44 - Br)<sup>+</sup>. Anal. Calcd. for C<sub>94</sub>H<sub>112</sub>O<sub>4</sub>P<sub>2</sub>Br<sub>2</sub>: C, 73.91; H, 7.39. Found: C, 73.67; H, 7.66.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

58 **1,3-bis-propoxy-2,4-bis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-p-tert-**  
59 **butylcalix[4]arene dibromide (4)**  
60

Compound **4** was obtained according to the same procedure as for **3** with **S**<sub>4</sub> (3.050 g, 3.04

mmol) and diphenylmethylphosphine (6.006 g, 30.00 mmol) a white powder in 64 % yield. Mp 148 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 0.79 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.03 (t, 6H, *J* = 6.9, CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 1.31 (s, 18H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.55-1.75 (m, 4H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 1.88-2.04 (m, 4H, CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 2.40-2.58 (m, 4H, CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 3.05 (d, 4H, *J* = 12.5, Ar-CH<sub>2</sub>-Ar), 3.23 (d, 6H, *J* = 14.3, CH<sub>3</sub>-P), 3.62-3.80 (m, 8H, -CH<sub>2</sub>-P and -CH<sub>2</sub>-O), 3.88 (t, 4H, *J* = 5.9, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O), 4.32 (d, 4H, *J* = 12.5, Ar-CH<sub>2</sub>-Ar), 6.43 (s, 4H, Ar-*H*), 7.08 (s, 4H, Ar-*H*), 7.60-7.80 (m, 12H, P-Ar-*H meta, para*), 8.01-8.13 (m, 8H, P-Ar-*H ortho*). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 25.98. *m/z* (MALDI) 1323.7 (M - Br)<sup>+</sup>. Anal. Calcd. for C<sub>84</sub>H<sub>108</sub>O<sub>4</sub>P<sub>2</sub>Br<sub>2</sub>: C, 71.88; H, 7.76. Found: C, 71.99; H, 7.85.

### **Tetrakis-(4-triphenylphosphonium-butoxy)-tetrakis-*p*-tert-butylcalix[4]arene tetrabromide (5)**

Compound **S<sub>5</sub>** (1.184 g, 1.00 mmol) was dissolved in chloroform (30 cm<sup>3</sup>). After a few minutes of stirring triphenylphosphine (5.248 g, 20.00 mmol) and chloroform (20 cm<sup>3</sup>) were added. After 6 days of refluxing the mixture was cooled and solvent was evaporated. The residue was dissolved in dichloromethane. The excess of triphenylphosphine was precipitated from methanol and filtered. The organic layer was evaporated. Chromatography on a silica column with 90:10 dichloromethane: methanol mixture as eluent gave compound **5** (0.67 g, 0.30 mmol) in 30% yield. Mp 132 °C. <sup>1</sup>H NMR (300 MHz; CDCl<sub>3</sub>) δ [ppm]: 1.02 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.56-1.72 (m, 8H, -CH<sub>2</sub>-), 2.24-2.41 (m, 8H, -CH<sub>2</sub>-), 2.91 (d, 4H, *J* = 13.0, Ar-CH<sub>2</sub>-Ar), 3.78-4.01 (m, 16H, -CH<sub>2</sub>-P and Ar-O-CH<sub>2</sub>), 4.23 (d, 4H, *J* = 13.0, Ar-CH<sub>2</sub>-Ar), 6.63 (s, 8H, Ar-*H*), 7.59-7.71- (m, 36H, P-Ar-*H, meta, para*), 7.76-7.88 (m, 24H, P-Ar-*H, ortho*). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 25.76. *m/z* (FAB<sup>+</sup>) 479.5 (M)<sup>4+</sup>; *m/z* (MALDI) 2157.7 (M - Br)<sup>+</sup>. Anal. Calcd. for C<sub>132</sub>H<sub>144</sub>O<sub>4</sub>P<sub>4</sub>Br<sub>4</sub>: C, 70.84; H, 6.49. Found: C, 70.97; H, 6.69.

### **Tetrakis-(4-triphenylphosphonium-butoxy)-*p*-tert-butylcalix[4]arene tetraperchlorate (5a)**

In a 10 cm<sup>3</sup> flask **5** (0.100 g, 0.045 mmol) was dissolved in acetonitrile (1 cm<sup>3</sup>). (0.050 g, 0.241 mmol) of AgClO<sub>4</sub> in acetonitrile (1 cm<sup>3</sup>) were added dropwise to the ligand solution. After 24 hours of stirring at room temperature, the precipitate of AgBr was filtered off. The filtrate was evaporated to give compound **5a** (0.088 g, 0.038 mmol) in 84 % yield. Mp > 120 °C decomposition. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 1.07 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.70-1.85 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.45-2.53 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.95 (d, 4H,

1  
2  
3  $J = 13.0$ , Ar- $CH_2$ -Ar), 3.43-3.60 (m, 8H, - $CH_2$ -P), 4.20-4.30 (m, 8H, - $CH_2$ -O), 4.44 (d,  
4 4H,  $J = 13.0$ , Ar- $CH_2$ -Ar), 6.97 (s, 8H, Ar- $H$ ), 7.56-7.90 (m, 60H, P-Ar- $H$ , *ortho*, *meta*,  
5 *para*),  $^{31}P$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  [ppm]: 24.54.  $m/z$  (MALDI) 2214.2 ( $M - ClO_4$ ) $^+$ .  
6  
7 Anal. Calcd: for  $C_{132}H_{144}O_4P_4(ClO_4)_4$ : C, 68.45; H, 6.27. Found: C, 68.70; H, 6.45.  
8  
9  
10  
11  
12

13  
14 **Tetrakis-(4-triphenylphosphonium-butoxy)-tetrakis-*p*-tert-butylcalix[4]arene tetra-**  
15 **hexafluorophosphate (5b)**  
16

17 Compound **5** (0.100 g, 0.045 mmoles) was dissolved in acetonitrile (1  $cm^3$ ). Then (0.062  
18 g, 0.245 mmol)  $AgPF_6$  were dissolved in acetonitrile and added dropwise to the ligand  
19 solution. After 24h the precipitate of NaBr was removed and the solution was evaporated.  
20 Compound **5b** (0.090 g, 0.036 mmole) was obtained in 80 % yield. Mp 128 °C.  $^1H$  NMR  
21 (500 MHz;  $CDCl_3$ )  $\delta$  [ppm]: 1.12 (s, 36H, - $C(CH_3)_3$ ), 1.65-1.81 (m, 8H, - $CH_2$ -), 2.38-2.49  
22 (m, 8H, - $CH_2$ -), 3.25-3.38 (m, 8H, - $CH_2$ -P), 3.47 (d, 4H,  $J = 13.0$ , Ar- $CH_2$ -Ar), 4.15-4.24  
23 (m, 8H, Ar-O- $CH_2$ ), 4.47 (d, 4H,  $J = 13.0$ , Ar- $CH_2$ -Ar), 7.04 (s, 8H, Ar- $H$ ), 7.66-7.70 (m,  
24 60H, P-Ar- $H$ ).  $^{31}P$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  [ppm]: 24.72 [ $P^+$ ], -143.39 [ $PF_6$ ].  $m/z$   
25 (MALDI) 1447.60 ( $M - PF_6$ ) $^+$ . Anal. Calcd. for  $C_{132}H_{144}O_4P_4(PF_6)_4$ : C, 63.46; H, 5.81.  
26 Found: C, 63.30; H, 5.78.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

37 **Tetrakis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-*p*-tert-butylcalix[4]arene**  
38 **tetrabromide (6)**  
39

40 Into a 100  $cm^3$  flask containing **S5** (1.184 g, 1.00 mmol) dissolved in chloroform (30  
41  $cm^3$ ), (4.004 g, 20.00 mmol) diphenylmethylphosphine in chloroform (20  $cm^3$ ) were  
42 added. After 6 days under reflux the mixture was cooled and the solvent was evaporated.  
43 The product was purified by precipitation from a 1/9 dichloromethane/hexane mixture to  
44 give compound **6** (1.190 g, 0.59 mmol) in 59 % yield. Mp 160 °C.  $^1H$  NMR (300 MHz;  
45  $CDCl_3$ )  $\delta$  [ppm]: 1.03 (s, 36H, - $C(CH_3)_3$ ), 1.55-1.75 (m, 8H, - $CH_2$ - $CH_2$ -P), 2.30-2.48 (m,  
46 8H, - $CH_2$ - $CH_2$ -O), 2.91 (d, 12H,  $J = 13.5$ ,  $CH_3$ -P), 3.03 (d, 4H,  $J = 12.8$ , Ar- $CH_2$ -Ar),  
47 3.70-3.87 (m, 16H, - $CH_2$ -P and  $CH_2$ -O), 4.26 (d, 4H,  $J = 12.8$ , Ar- $CH_2$ -Ar), 6.71 (s, 8H,  
48 Ar- $H$ ), 7.54-7.78 (m, 24H, P-Ar- $H$  *meta*, *para*), 8.02-8.17 (m, 16H, P-Ar- $H$  *ortho*).  $^{31}P$   
49 NMR (400 MHz,  $CDCl_3$ )  $\delta$  [ppm]: 25.98.  $m/z$  (MALDI) 1909.68 ( $M - Br$ ) $^+$ . Anal. Calcd.  
50 for  $C_{112}H_{136}O_4P_4Br_4$ : C, 67.61; H, 6.89. Found: C, 68.59; H, 7.10.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Tetrakis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-*p*-tert-butylcalix[4]arene**



**tetraperchlorate (6a)**

Compound **6a** was obtained according to the same procedure as **5a** with **6** (0.100 g, 0.05 mmol) and AgClO<sub>4</sub> (0.050 g, 0.24 mmol) in 90 % yield. Mp > 120 °C decomposition. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 1.09 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.65-1.80 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.35-2.50 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.50 (d, 12H, *J* = 12.0, CH<sub>3</sub>-P), 3.18-3.28 (m, 8H, -CH<sub>2</sub>-P), 3.38 (d, 4H, *J* = 12.8, Ar-CH<sub>2</sub>-Ar), 4.15-4.28 (m, 8H, -CH<sub>2</sub>-O), 4.45 (d, 4H, *J* = 12.8, Ar-CH<sub>2</sub>-Ar), 6.97 (s, 8H, Ar-*H*), 7.60-7.87 (m, 40H, P-Ar-*H*, *ortho*, *meta*, *para*). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 24.77. *m/z* (MALDI) 1966.24 (M - ClO<sub>4</sub>)<sup>+</sup>. Anal. Calcd. for C<sub>112</sub>H<sub>136</sub>O<sub>4</sub>P<sub>4</sub>(ClO<sub>4</sub>)<sub>4</sub>: C, 65.05; H, 6.63. Found: C, 65.13; H, 6.72.

**Tetrakis-(4-(P,P-diphenyl-P-methylphosphonium)-butoxy)-p-tert-butylcalix[4]arene tetrahexafluorophosphate (6b)**

Compound **6b** was obtained according to the same procedure as for **5b** with **6** (0.100 g, 0.05 mmol) and AgPF<sub>6</sub> (0.069 g, 0.27 mmol) in 80 % yield. Mp 155 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 1.11 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.55-1.70 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-P), 2.24-2.38 (m, 8H, -CH<sub>2</sub>-CH<sub>2</sub>-O), 2.42 (d, 12H, *J* = 12.0, CH<sub>3</sub>-P), 2.95-3.10 (m, 8H, -CH<sub>2</sub>-P), 3.39 (d, 4H, *J* = 12.8, Ar-CH<sub>2</sub>-Ar), 4.05-4.20 (m, 8H, -CH<sub>2</sub>-O), 4.42 (d, 4H, *J* = 12.8, Ar-CH<sub>2</sub>-Ar), 7.00 (s, 8H, Ar-*H*), 7.58-7.80 (m, 40H, P-Ar-*H*, *ortho*, *meta*, *para*). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 24.50 [P<sup>+</sup>], -143.07 [PF<sub>6</sub>], *m/z* (MALDI) 2104.80 (M - PF<sub>6</sub>)<sup>+</sup>. Anal. Calcd. for C<sub>112</sub>H<sub>136</sub>O<sub>4</sub>P<sub>4</sub>(PF<sub>6</sub>)<sub>4</sub>: C, 59.79; H, 6.09. Found: C, 59.87; H, 6.03.

**Protonated tetrakis-(4-(P,P-diphenyl-phosphine)-butoxy)-p-tert-butylcalix[4]arene tetrabromide (7)**

Into a 50 cm<sup>3</sup> flask containing **S<sub>6</sub>** (1.510 g, 0.94 mmol) in dichloromethane (10 cm<sup>3</sup>), 5 cm<sup>3</sup> of HBr (solution of 33 % wt. in glacial acetic acid) in dichloromethane (15 cm<sup>3</sup>) were added. The mixture was left stirred for 24 hours at room temperature. The mixture was then evaporated to give the pure product **7** (0.954 g, 0.49 mmol) in 52 % yield. Mp 175 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ [ppm]: 1.22 (s, 36H, -C(CH<sub>3</sub>)<sub>3</sub>), 1.80-2.15 (m, 16H, O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-P), 2.92-3.10 (m, 4H, Ar-CH<sub>2</sub>-Ar), 3.18-3.30 (m, 8H, CH<sub>2</sub>-O), 3.40-3.55 (m, 8H, -CH<sub>2</sub>-P), 4.05-4.27 (m, 4H, Ar-CH<sub>2</sub>-Ar), 7.05 (s, 8H, Ar-*H*), 7.60-8.05 (m, 40H, P-Ar-*H*, *ortho*, *meta*, *para*), 10.30-10.40 (m, 4H, H-P). <sup>31</sup>P NMR (400 MHz, CDCl<sub>3</sub>) δ [ppm]: 46.32 [P<sup>+</sup>]. *m/z* (MALDI) 1849.04 (M - Br)<sup>+</sup>. Anal. Calcd. for C<sub>108</sub>H<sub>128</sub>O<sub>4</sub>P<sub>4</sub>Br<sub>4</sub>: C, 67.08; H, 6.67. Found: C, 66.84; H, 6.92.

### NMR studies

5 mg of ligand were introduced with 6 equivalents of the solid alkali metal salts ( $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{Cs}^+$ ) or tetraethylammonium perchlorate in a glass vessel and dissolved in a small volume of deuterated chloroform or acetonitrile. After manual shaking for a few minutes, the mixture was left in contact for 24 hours before filtration of the excess of salt if necessary. The solution (or the filtrate) was then sampled in a NMR tube and its spectrum recorded on Bruker SY300 MHz or SY400 MHz spectrometers equipped for  $^1\text{H}$  or  $^{31}\text{P}$  resonances.

### Microcalorimetric studies

Microcalorimetric titrations were performed using a 2277 Thermal Activity Monitor Microcalorimeter (Thermometric). Titrations were carried out at 25 °C on 2.7 cm<sup>3</sup> of 10<sup>-5</sup> to 5×10<sup>-4</sup> M solutions of the ligand in acetonitrile using a glass cell of 4 cm<sup>3</sup>. The heat changes were measured after injection of 15×15 μL of 10<sup>-3</sup> and 10<sup>-2</sup> M  $\text{LiClO}_4$ ,  $\text{LiBr}$ ,  $\text{NaClO}_4$ ,  $\text{NaPF}_6$ , or  $\text{Et}_4\text{NClO}_4$  solutions in the same solvent. Chemical calibration was made by determination of the complexation enthalpy of  $\text{Ba}^{2+}$  with 18C6 in water or of  $\text{Rb}^+$  with 18C6 in methanol, as recommended [37]. Values of the stability constants ( $\beta$ ) and of the enthalpies of complexation ( $\Delta H$ ) were refined simultaneously from these data using the ligand binding analysis program DIGITAM version 4.1 [38] and after correction for the heat of dilution determined in separate experiments by adding the salt solutions to 2.7 cm<sup>3</sup> of pure solvent. The values of the corresponding entropies of complexation ( $\Delta S$ ) were then derived from the expressions  $\Delta G = -RT\ln\beta$  and  $\Delta G = \Delta H - T\Delta S$ .

### Ion selective electrodes

THF was dried and freshly distilled before used for the preparation of the ion selective membranes. PVC (high molecular weight poly(vinyl chloride)), 2-nitrophenyl octyl ether (*o*-NPOE), bis-(2-ethylhexyl)sebacate (BEHS), (2-morpholino)ethanesulfonic acid monohydrate (MES) were from Fluka Selectophore. The  $\text{LiClO}_4$ ,  $\text{CsClO}_4$  and sodium salts:  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$ ,  $\text{ClO}_4^-$ ,  $\text{SCN}^-$ ,  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{CO}_3^{2-}$ ,  $\text{HPO}_4^{2-}$ ,  $\text{PO}_4^{3-}$ ,  $\text{Cr}_2\text{O}_7^{2-}$ , citrate, acetate, benzoate, and oxalate were of p.a. grade. All aqueous salt solutions were prepared with demineralised water (conductivity < 1.0 μS/cm).

The membranes were composed of 4 mg of ionophores **1** – **6** and **8**, 60 mg of PVC and 120 mg of plasticizer. All the components were dissolved in 1.5 cm<sup>3</sup> of dried, freshly distilled

1  
2  
3 THF and the solutions were poured into glass rings of 24 mm in diameter. The solutions were  
4 left for 24 h for slow solvent evaporation giving the mother membranes of thickness about  
5 0.1mm. Several membranes of 7 mm diameter were cut from each mother membrane and  
6  
7 were incorporated into the Ag/AgCl electrodes bodies of IS 561 type (Moeller S.A., Zurich).  
8  
9 The two plasticizers BEHS and *o*-NPOE were used for the preparation of the membranes.  
10  
11 However, electrodes with membranes based on NPOE had the best lifetime and response  
12 characteristics. The EMF measurements were carried out at zero current conditions using a  
13  
14 Lawson Lab 16 EMF station (multi-channel millivoltmeter) or a Metrohm 654 millivoltmeter.  
15  
16 A double-junction reference Radelkis 0P0820P electrode with a 1M CH<sub>3</sub>COOLi solution in  
17  
18 the bridge cell was used. The measurements were carried out using cells of the type:

19  
20 Ag/AgCl | 1M KCl | 1M CH<sub>3</sub>COOLi | sample |  
21  
22 | membrane |  
23  
24 | 0.05 M MES/NaOH, 0.01M NaCl | AgCl/Ag.  
25

26  
27 At least three identical electrodes of the same membrane composition and containing the  
28 same inner electrolyte were prepared [39]. The studies were repeated several times over the  
29 period of one month.  
30

31  
32 To reduce the pH changes during the titrations solutions were prepared with 0.05 M  
33 MES/NaOH buffer of pH = 5.5 ((2-morpholino)ethanesulfonic acid monohydrate (MES)). All  
34 salt solutions contained 10<sup>-2</sup> M NaCl as supporting electrolyte [22].  
35

36  
37 The selectivity coefficients  $K_{A,B}^{pot}$  of the electrodes were determined by the separate  
38 solution method (SSM) and in some cases by the fixed interference method (FIM) [40,41,42].  
39  
40 The calibration curves were obtained by addition of standard solutions of different anions to  
41 50 cm<sup>3</sup> of 0.01 M NaCl in 0.05 M MES/NaOH buffer solution of pH = 5.5. The concentration  
42 of the primary anion [A] was increased from 10<sup>-7</sup> to 10<sup>-2</sup> M. They were also measured by  
43 successive dilution of initial 5×10<sup>-2</sup> M salt solutions until further dilution resulted in no  
44 potential change.  
45  
46  
47  
48  
49  
50  
51  
52

### 53 **Acknowledgements**

54  
55 R.P. gratefully acknowledges the Gdansk University of Technology, Poland, and the  
56 Ministère de la Recherche et des Nouvelles Technologies, France (Cotutelles de thèses),  
57 for financial support.  
58  
59  
60

## References

- [1] Bianchi, A.; Bowman-James, K.; Garcia-Espana, E. Eds., *Supramolecular Chemistry of Anions*; John Wiley-VCH, New York, 1997, ch. 3, pp. 63.
- [2] Kanyo, Z. F.; Christianson, D. W. *J. Biol. Chem.*, **1991**, 266, 4264.
- [3] Pardee, A. P. *J. Biol. Chem.*, **1966**, 241, 5887.
- [4] Puljak L.; Kilic, G. *Biochim. Biophys. Acta*, **2006**, 1762, 404.
- [5] Weiner, M. L.; Salminen, W. F.; Larson, P. R.; Barter, R. A.; Kranetz J. L.; Simon, G. *S. Food and Chemical Toxicology*, **2001**, 39, 759.
- [6] Matzer, W. E. *Environmental Forensic*, **2001**, 2, 301.
- [7] Bondy, C. R.; Gale P. A.; Loeb, S. J. *J. Am. Chem. Soc.*, **2004**, 126, 5030.
- [8] Amendola, V.; Boiocchi, M.; Esteban-Gómez, D.; Fabbri L.; Monzani, E. *Org. Biomol. Chem.*, **2005**, 3, 2632.
- [9] Brooks, S. J.; Edwards, P. R.; Gale P. A.; Light, M. E. *New J. Chem.*, **2006**, 30, 65.
- [10] Turner, D. R.; Paterson M. J.; Steed, J. W. *J. Org. Chem.*, **2006**, 71, 1598.
- [11] Boerrigter, H.; Grave, L.; M. Nissink, J. W.; Chrisstoffels, L. A. J.; van der Maas, J. H.; Verboom, W.; de Jong F.; Reinhoudt, D. N.; *J. Org. Chem.*, **1998**, 63, 4174.
- [12] Lee K. H.; Hong, J. I. *Tetrahedron Lett.*, **2000**, 41, 6083.
- [13] Chmielewski M.; Jurczak, J. *Tetrahedron Lett.*, **2004**, 45, 6007.
- [14] Beer, P. D.; Drew M. G. B.; Gradwell, . K. *J. Chem. Soc., Perkin Trans. 2*, **2000**, 511.
- [15] Dietrich, B. *Pure & Appl. Chem.*, **1993**, 65, 1457.
- [16] Egorov, V. V.; Rakhman'ko, E. M.; Okaev E. B.; Pomelenok E. V.; Nazarov, V. A.; *Talanta*, **2004**, 63, 119.
- [17] Matthews S. E.; Beer, P. D. *Calixarenes 2001*; Asfari, Z.; Böhmer, V.; Harrowfield, J.; Vicens J. Eds., Kluwer Academic Publishers, Dordrecht, 2001, ch. 23, p. 421.
- [18] Miao, R.; Zheng, Q. Y.; Chen C. F.; Huang Z. T. *Tetrahedron Lett.*, **2004**, 45, 4959.

- 1  
2  
3  
4  
5 [19] Jagessar, R. C.; Shang, M.; Scheidt, W. R.; Burns; D. H. *J. Am. Chem. Soc.*, **1998**,  
6 *120*, 11684.  
7  
8 [20] Gale, P. A.; Sessler, J. L.; Kral, V.; Lynch, V. *J. Am. Chem. Soc.*, **1996**, *118*, 5140.  
9  
10 [21] Gale, P. A.; Sessler, J. L.; Allen, W. E.; Tvermoes, N. A.; Lynch, V. *Chem. Commun.*,  
11 **1997**, 665.  
12  
13 [22] Pomecko, R.; Asfari, Z.; Hubscher-Bruder, V.; Bochenska M.; Arnaud-Neu, F.  
14 *Supramol. Chem.*, **2007**, *19*, 459.  
15  
16 [23] Hamdi, A.; Nam, K. C.; Ryu, B.J.; Kim, J. S.; Vicens, J. *Tetrahedron Letters* **2004**, *45*,  
17 4689.  
18  
19 [24] Broder, C. K.; Davidson, M. G.; Forsyth, V. T.; Howard, J. A. K.; Lamb S.; Mason, S.  
20 *A. Crys. Grow. Des.*, **2002**, *3*, 163.  
21  
22 [25] Issleib K.; Muller, D.-W. *Chem. Berichte*, **1959**, *92*, 3175.  
23  
24 [26] Clark, P. W. *Org. Prep. and Proc. Int.*, **1979**, *11*, 103.  
25  
26 [27] Mohr, W.; Horn, C. R.; Stahl J.; Gladysz, J. A. *Synthesis*, **2003**, *8*, 1279.  
27  
28 [28] Baklouti, L.; Cherif J.; Abidi, R.; Arnaud-Neu, F.; Harrowfield J.; Vicens, J. *Org.*  
29 *Biomol. Chem.*, **2004**, *2*, 2786.  
30  
31 [29] Savage, P. B.; Holmgren S. K.; Gellman, S. H. *J. Am. Chem. Soc.*, **1994**, *116*, 4069.  
32  
33 [30] Casabo, J.; Munoz J. A.; Errachid, A. *Sens. Actuators, B*, **1997**, *43*, 206.  
34  
35 [31] Shamsipur, M.; Soleymannpour, A.; Akhond, M.; Sharghi H.; Hasaninejad, A. R. *Sens.*  
36 *Actuators, B*, **2003**, *89*, 9.  
37  
38 [32] Dinten, O.; Spichiger, U. E.; Chaniotakis, N.; Gehrig, P.; Rusterholz, B.; Morf, W. E.;  
39 Simon, W. *Anal. Chem.*, **1991**, *63*, 596.  
40  
41 [33] Lizondo-Sabater, J.; Segui, M. J.; Loris, J. M.; Martinez-Manez, R.; Pardo, T.;  
42 Sancenon, F.; Soto, J. *Sens. Actuators B*, **2004**, *101*, 20.  
43  
44 [34] Segui, M. J.; Lizondo-Sabater, J.; Martinez-Manez, R.; Soto, J. *Analyst*, **2002**, *127*,  
45 387.  
46  
47 [35] Casabo, J.; Escriche, L.; Perez-Jimenez, C.; Munoz, J. A.; Teixidor, F.; Bausells, J.;  
48 Errachid, A. *Anal. Chim. Acta*, **1996**, *320*, 63.  
49  
50 [36] Schaller, U.; Bakker, E.; Spichiger, U. E.; Pretsch, E. *Anal. Chem.*, **1994**, *66*, 391.  
51  
52 [37] Arnaud-Neu, F.; Delgado, R.; Chaves, S. *Pure Appl. Chem.*, **2003**, *75*, 71.  
53  
54 [38] Hallen, D. *Pure Appl. Chem.*, **1993**, *65*, 1527.  
55  
56 [39] Bochenska, M.; Zielenska, A.; Pomecko, R.; Hubscher-Bruder, V.; Arnaud-Neu, F. *J.*  
57 *Incl. Phenom.*, **2005**, *52*, 129.  
58  
59  
60

- 
- 1  
2  
3  
4 [40] Buck, R. P.; Linder, E. *Pure Appl. Chem.*, **1994**, *66*, 2527.  
5  
6 [41] Umezawa, Y.; Buhlmann, P.; Umezawa, K.; Tohda, K.; Amemiya, S. *Pure Appl.*  
7  
8 *Chem.* **2000**, *72*, 1851.  
9  
10 [42] Lindner, E.; Umezawa, Y. *Pure Appl. Chem.* **2008**, *80*, 85.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review Only

**Table 1.** Changes ( $\Delta\delta$  in ppm) in the  $^{31}\text{P}$  NMR spectra of phosphonium ligands in the presence of sodium iodide, thiocyanate and perchlorate in  $\text{CDCl}_3$

	<b>1</b>	<b>2</b>	<b>5</b>	<b>6</b>
Ligand ( $\delta$ )	25.797	25.962	25.756	25.982
Ligand + $\text{I}^-$ ( $\Delta\delta$ )	0.215	0.203	0.210	0.175
Ligand + $\text{SCN}^-$ ( $\Delta\delta$ )	0.470	0.530	0.767	0.965
Ligand + $\text{ClO}_4^-$ ( $\Delta\delta$ )	0.602	0.620	0.992	1.170

**Table 2.** Differences ( $\Delta\delta$ ) in the proton chemical shifts ( $\delta$ )[ppm] in the spectra of ligands **5** and **6**, **5a** and **6a**, **5b** and **6b** in CD<sub>3</sub>CN

	<b>c</b>	<b>e</b>	<b>f</b>	<b>g</b>	<b>j</b>	<b>o</b>
$\delta$ ( <b>5</b> )	6.87	2.93	4.15	3.74	3.81	-
$\delta$ ( <b>5a</b> )	6.83	2.91	4.11	3.76	3.17	-
$\Delta\delta$ ( <b>5a-5</b> )	-0.04	-0.02	-0.04	0.02	-0.64	-
$\delta$ ( <b>5b</b> )	6.80	2.87	4.09	3.77	3.06	-
$\Delta\delta$ ( <b>5b-5</b> )	-0.07	-0.06	-0.06	0.03	-0.75	-
$\delta$ ( <b>6</b> )	6.92	3.02	4.19	3.70	3.58	2.78
$\delta$ ( <b>6a</b> )	6.89	2.98	4.15	3.73	2.85	2.37
$\Delta\delta$ ( <b>6a-6</b> )	-0.03	-0.04	-0.04	0.03	-0.73	-0.41
$\delta$ ( <b>6b</b> )	6.88	2.97	4.12	3.75	2.75	2.32
$\Delta\delta$ ( <b>6b-6</b> )	-0.04	-0.05	-0.07	0.05	-0.83	-0.46



**Table 3.** Characteristics of potentiometric responds for perchlorate of PVC/NPOE electrodes containing ligand **5** and **6** and different amount of KTCIPB. (Inner and conditioning electrolyte MES/NaOH pH=5.5/10<sup>-2</sup>M NaCl)

Ligand	KTCIPB (mol %)	S (mV/decade)	LR (log[A])
<b>5</b>	0	-55.6	-6.0
<b>5</b>	40	-54.6	-6.0
<b>5</b>	120	-55.9	-5.7
<b>6</b>	0	-54.2	-6.0
<b>6</b>	40	-51.8	-6.0
<b>6</b>	120	-49.9	-5.7

**Table 4.** Characteristics of potentiometric responds for perchlorate of PVC/NPOE electrodes containing tetrasubstituted phosphonium ligands with different counterions (Inner and conditioning electrolyte MES/NaOH pH=5.5/10<sup>-2</sup>M NaCl).

Ligand	Counterion	S (mV/decade)	LR (log[A])
<b>5</b>	Br <sup>-</sup>	-56.5	-6.0
<b>5a</b>	ClO <sub>4</sub> <sup>-</sup>	-56.1	-6.0
<b>5b</b>	PF <sub>6</sub> <sup>-</sup>	-38.3	-5.5
<b>6</b>	Br <sup>-</sup>	-54.4	-5.7
<b>6a</b>	ClO <sub>4</sub> <sup>-</sup>	-40.4	-6.0
<b>6b</b>	PF <sub>6</sub> <sup>-</sup>	-36.2	-6.0

**Table 5.** Selectivity coefficients as  $\log K_{\text{ClO}_4^-,X}^{\text{pot}}$  of the PVC/NPOE membrane electrodes based on phosphonium calixarenes **1 - 6** and the monomer **8**

Anion X	$\log K_{\text{ClO}_4^-,X}^{\text{pot}}$						
	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>8</b>
$\text{ClO}_4^-$	0	0	0	0	0	0	0
$\text{SCN}^-$	-1.2	-1.4	-1.1	-0.6	-1.3	-1.4	-1.1
$\text{I}^-$	-1.7	-1.7	-1.3	-0.9	-2.0	-2.0	-1.6
$\text{NO}_3^-$	-2.6	-2.6	-2.4	-1.9	-2.9	-3.0	-2.5
$\text{HCO}_3^-$	-4.6	-4.5	-4.6	-4.4	-4.6	-4.4	nd
$\text{Cr}_2\text{O}_7^{2-}$	-2.2	-2.4	-2.6	-1.7	-2.6	-3.2	-1.9
$\text{HPO}_4^{2-}$	-4.5	-4.5	-4.5	-4.4	-4.5	-4.4	nd
$\text{SO}_4^{2-}$	-4.9	-4.7	-4.7	-4.7	-4.7	-4.5	nd

nd: not determined


**Figure captions**

- 1  
2  
3  
4  
5  
6  
7  
8  
9 **Figure 1.** Chemical structures of the ligands under study
- 10  
11 **Figure 2.**  $^1\text{H}$  NMR spectra of ligand **1** alone and in the presence of  $\text{ClO}_4^-$  in  $\text{CDCl}_3$
- 12  
13  
14 **Figure 3.**  $^1\text{H}$  NMR spectra of ligand **6** alone and in the presence of  $\text{ClO}_4^-$  in  $\text{CDCl}_3$
- 15  
16  
17 **Figure 4.** Potentiometric anion responses of electrodes with PVC/NPOE membrane  
18 containing ligand **1** in MES buffer at pH 5.5.
- 19  
20  
21 **Figure 5.** Potentiometric anion responses of electrodes with PVC/NPOE membrane  
22 containing ligand **6** in MES buffer at pH 5.5.
- 23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Supplementary Online Material

## Anion recognition by phosphonium calix[4]arenes : synthesis and physico-chemical studies

Radoslaw Pomecko,<sup>a,b</sup> Zouhair Asfari,<sup>a</sup> Véronique Hubscher-Bruder,<sup>a</sup> Maria Bochenska<sup>b\*</sup> and Françoise Arnaud-Neu<sup>a\*</sup>



**Table S1.** Changes ( $\Delta\delta$ ) of proton chemical shifts ( $\delta$ ) [ppm] in the spectrum of ligand **1** in the presence of various sodium salts in  $\text{CDCl}_3$ 

	<b>a</b>	<b>b</b>	<b>c</b>	<b>d</b>	<b>e</b>	<b>f</b>	<b>g</b>	<b>h</b>	<b>i</b>	<b>j</b>	<b>k</b>	<b>l</b>	<b>m</b>	<b>n</b>
$\delta(\mathbf{1})$	0.99	1.28	6.79	6.99	3.19	3.90	3.90	2.19	2.19	3.96	7.86	7.59	7.69	7.49
$\delta(\mathbf{1+I}^-)$	0.99	1.28	6.79	6.99	3.19	3.91	3.90	2.19	2.19	3.85	7.84	7.60	7.71	7.49
$\Delta\delta$	0	0	0	0	0	0.01	0	0	0	-0.11	-0.02	0.01	0.02	0
$\delta(\mathbf{1+SCN}^-)$	1.00	1.28	6.79	7.00	3.19	3.92	3.92	2.20	2.20	3.75	7.80	7.65	7.70	7.45
$\Delta\delta$	0.01	0	0	0.01	0	0.02	0.02	0.01	-0.01	-0.21	-0.06	0.06	0.01	-0.04
$\delta(\mathbf{1+ClO}_4^-)$	1.01	1.28	6.82	7.00	3.20	3.92	3.90	2.13	2.05	3.42	7.75	7.60	7.75	7.60
$\Delta\delta$	0.02	0	0.03	0.01	0.01	0.02	0	-0.06	-0.14	-0.54	-0.11	-0.01	0.06	-0.11

**Table S2.** Changes ( $\Delta\delta$ ) of proton chemical shifts ( $\delta$ ) [ppm] in the spectrum of ligand **2** in the presence of various sodium salts in  $\text{CDCl}_3$ 

	<b>a</b>	<b>b</b>	<b>c</b>	<b>d</b>	<b>e</b>	<b>f</b>	<b>g</b>	<b>h</b>	<b>i</b>	<b>j</b>	<b>k</b>	<b>l</b>	<b>m</b>	<b>n</b>	<b>o</b>
$\delta(\mathbf{2})$	0.96	1.29	6.76	7.01	3.23	4.02	3.90	2.18	2.00	3.67	7.98	7.54	7.65	7.31	2.99
$\delta(\mathbf{2+I}^-)$	0.97	1.29	6.77	7.01	3.23	4.01	3.91	2.20	2.05	3.58	7.96	7.54	7.65	7.31	2.96
$\Delta\delta$	0.01	0	0.01	0	0	-0.01	0.01	0.02	0.05	-0.09	-0.02	0	0	0	-0.03
$\delta(\mathbf{2+SCN}^-)$	0.96	1.30	6.76	7.02	3.24	4.01	3.93	2.18	2.05	3.40	7.87	7.57	7.65	7.26	2.82
$\Delta\delta$	0	0.01	0	0.01	0.01	-0.01	0.03	0	0.05	-0.27	-0.11	0.03	0	-0.05	-0.17
$\delta(\mathbf{2+ClO}_4^-)$	0.97	1.29	6.78	7.01	3.23	4.01	3.90	2.15	2.01	3.35	7.87	7.54	7.65	7.32	2.78
$\Delta\delta$	0.01	0	0.02	0	0	-0.01	0	-0.03	0.01	-0.32	-0.11	0	0	0.01	-0.21

**Table S3.** Changes ( $\Delta\delta$ ) of proton chemical shifts ( $\delta$ ) [ppm] in the spectrum of ligand **6** in the presence of various sodium salts in  $\text{CDCl}_3$ 

	<b>a</b>	<b>c</b>	<b>e</b>	<b>f</b>	<b>g</b>	<b>h</b>	<b>i</b>	<b>j</b>	<b>k</b>	<b>l</b>	<b>m</b>	<b>o</b>
$\delta(\mathbf{6})$	1.03	6.71	3.03	4.26	3.79	2.39	1.69	3.79	8.10	7.66	7.66	2.91
$\delta(\mathbf{6+I}^-)$	1.06	6.94	3.33	4.40	4.39	2.42	1.74	3.76	7.99	7.66	7.66	2.80
$\Delta\delta$	0.03	0.23	0.30	0.14	0.60	0.03	0.05	-0.03	-0.11	0	0	-0.11
$\delta(\mathbf{6+SCN}^-)$	1.06	6.94	3.30	4.37	4.28	2.40	1.79	3.50	7.88	7.69	7.69	2.73
$\Delta\delta$	0.03	0.23	0.27	0.11	0.49	0.01	0.10	-0.29	-0.22	0.03	0.03	-0.18
$\delta(\mathbf{6+ClO}_4^-)$	1.07	6.95	3.33	4.38	4.27	2.33	1.66	3.20	7.82	7.63	7.63	2.50
$\Delta\delta$	0.04	0.24	0.30	0.12	0.48	-0.06	-0.03	-0.59	-0.30	-0.03	-0.03	-0.41



**Table S4.** Most important changes ( $\Delta\delta$ ) of protons chemical shifts ( $\delta$ )[ppm] in the spectra of ligands **5** and **5b** in the presence of different perchlorate salts in  $\text{CD}_3\text{CN}$

	<b>c</b>	<b>e</b>	<b>f</b>	<b>g</b>	<b>j</b>
$\delta$ ( <b>5</b> )	6.87	2.93	4.15	3.74	3.81
$\delta$ ( <b>5</b> + <b>LiClO<sub>4</sub></b> )	7.19	3.28	4.21	4.36	3.29
$\Delta\delta$	0.32	0.35	0.06	0.62	-0.52
$\delta$ ( <b>5</b> + <b>NaClO<sub>4</sub></b> )	7.22	3.33	4.21	4.09	3.28
$\Delta\delta$	0.35	0.40	0.06	0.35	-0.53
$\delta$ ( <b>5</b> + <b>CsClO<sub>4</sub></b> )	6.86	2.92	4.14	3.75	3.75
$\Delta\delta$	-0.01	-0.01	-0.01	0.01	-0.06
$\delta$ ( <b>5</b> + <b>NEt<sub>4</sub>ClO<sub>4</sub></b> )	6.93	3.00	4.22	3.83	3.57
$\Delta\delta$	0.06	0.07	0.07	0.09	-0.24
$\delta$ ( <b>5b</b> )	6.80	2.87	4.09	3.77	3.06
$\delta$ ( <b>5b</b> + <b>NaClO<sub>4</sub></b> )	7.19	3.31	4.19	4.06	3.27
$\Delta\delta$	0.39	0.44	0.10	0.29	0.21



**Table S5.** Thermodynamic parameters of the interaction between ligands **5** and **5b** and several sodium and lithium salts.

Ligand	Salt	Complexes <sup>1</sup>	$\log \beta$	$-\Delta H$ [kJ mol <sup>-1</sup> ]	$T\Delta S$ [kJ mol <sup>-1</sup> ]
<b>5</b>	NaClO <sub>4</sub>	ML	4.42	23.8	1.4
		ML <sub>2</sub>	7.95	30.7	14.6
<b>5b</b>	NaClO <sub>4</sub>	ML	3.14	51.7	-33.8
		ML <sub>2</sub>	7.73	40.1	4.0
<b>5</b>	LiClO <sub>4</sub>	ML	3.24	20.3	-1.8
		ML <sub>2</sub>	7.34	35.0	0.8
<b>5b</b>	LiClO <sub>4</sub>	ML	4.29	23.7	6.8
		ML <sub>2</sub>	7.34	35.0	6.8
<b>5</b>	LiBr	ML	3.15	30.1	-12.1
<b>5b</b>	LiBr	ML	4.32	55.4	-30.8
		ML <sub>2</sub>	7.91	72.7	-27.6

**Table S6.** Characteristics of anion response of PVC/NPOE membrane electrodes containing ligands **1** and **2**  
(Inner and conditioning electrolyte: MES/NaOH, pH=5.5/10<sup>-2</sup>M NaCl)

Primary anion	<b>1</b>			<b>2</b>		
	DL (log[A])	LR (log[A])	S (mV/decade)	DL (log[A])	LR (log[A])	S (mV/decade)
ClO <sub>4</sub> <sup>-</sup>	-6.0	-5.5	-56	-6.7	-6.0	-55
SCN <sup>-</sup>	-6.0	-5.5	-56	-5.5	-5.5	-52
I <sup>-</sup>	-5.0	-5.0	-55	-5.5	-5.0	-55
Br <sup>-</sup>	-4.5	-4.2	-31	-4.0	-3.5	-47
NO <sub>3</sub> <sup>-</sup>	-4.4	-4.0	-51	-4.0	-3.6	-39
HPO <sub>4</sub> <sup>2-</sup>	nr	nr	nr	nr	nr	nr
Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup>	-4.8	-4.3	-43	-4.8	-4.2	-43

nr – no response


**Table S7.** Characteristics of perchlorate response of PVC/NPOE membrane electrodes containing ligands **1** – **4**  
(Inner and conditioning electrolyte: MES/NaOH, pH=5.5/10<sup>-2</sup>M NaCl)

Ligand	DL (log[A])	LR (log[A])	S (mV/decade)
<b>1</b>	-6.0	-5.5	-56
<b>3</b>	-5.7	-5.5	-48
<b>2</b>	-6.7	-6.0	-55
<b>4</b>	-5.6	-5.5	-48


**Table S8.** Characteristics of the electrodes containing ligands **5**, **6** and **8**(Inner and conditioning electrolyte: MES/NaOH, pH=5.5/10<sup>-2</sup>M NaCl)

Primary anion	<b>5</b>			<b>6</b>			<b>8</b>		
	DL (log[A])	LR (log[A])	S (mV/ decade)	DL (log[A])	LR (log[A])	S (mV/ decade)	DL (log[A])	LR (log[A])	S (mV/ decade)
ClO <sub>4</sub> <sup>-</sup>	-6.6	-6.4	-56	-6.0	-6.2	-53	-5.6	-5.4	-51
SCN <sup>-</sup>	-5.5	-5.5	-52	-5.6	-5.4	-52	-4.5	-4.8	-48
I <sup>-</sup>	-5.5	-5.2	-55	-5.5	-5.2	-52	-4.5	-4.8	-48
Br <sup>-</sup>	-3.8	-3.8	-44	-3.8	-3.8	-11	-2.6	-3.0	-10
NO <sub>3</sub> <sup>-</sup>	-5.0	-4.5	-51	-4.5	-4.5	-25	-3.4	-4.0	-22
HCO <sub>3</sub> <sup>-</sup>	-3.0	-3.0	-35	-3.0	-3.0	-10	-2.8	-2.8	-10
Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup>	-4.0	-5.5	-31	-5.0	-5.5	-46	-4.0	-4.6	-43





**Figure S1.** Thermograms corresponding to the titrations of ligands **5** and **5b** against  $\text{NaClO}_4$  and  $\text{Et}_4\text{NClO}_4$  in acetonitrile.



**1:** R, R'=Ph  
**2:** R=Ph, R'=Me


**3:** R, R'=Ph  
**4:** R=Ph, R'=Me


**5:** R, R'=Ph, A=Br<sup>-</sup>  
**5a:** R, R'=Ph, A=ClO<sub>4</sub><sup>-</sup>  
**5b:** R, R'=Ph, A=PF<sub>6</sub><sup>-</sup>


**8:** R=Ph

**6:** R=Ph R'=Me, A=Br<sup>-</sup>  
**6a:** R=Ph R'=Me, A= ClO<sub>4</sub><sup>-</sup>  
**6b:** R=Ph R'=Me, A= PF<sub>6</sub><sup>-</sup>

**7:** R=Ph R'=H, A=Br<sup>-</sup>









Review Only






Review Only



Scheme 1

Scheme 1  
169x135mm (600 x 600 DPI)



Scheme 2

Scheme 2  
171x138mm (600 x 600 DPI)