

## **Organomercury compounds in environmental samples: emission sources, toxicity, environmental fate and determination**

**Małgorzata Chmiel<sup>1</sup>, Kinga Dubalska<sup>1</sup>, Gabriela Bajger<sup>1</sup>, Piotr Konieczka<sup>1</sup>, Jacek Namieśnik<sup>1</sup>**

*I-Department of Analytical Chemistry, Chemical Faculty, Gdańsk University of Technology, Gdańsk, Poland, phone: +48 58 347 21 10, fax: +48 58 347-26-94, e-mail: chemanal@pg.gda.pl, malgorzatahelenachmiel@gmail.com*

### *Abstract*

In view of the specific properties of mercury and its capability of forming compounds that can be bioaccumulated and biomagnified at successive levels of the trophic pyramid, it has become necessary to gather detailed information on the sources of emission of this element into the environment and its fate there. Moreover, the increasing awareness of the relationship between the toxicity of mercury and its chemical form has sharpened interest in the identification of its various forms in environmental samples. Investigating the speciation of mercury has therefore become of major importance with respect not only to determining its biogeochemical cycle but also to assessing the scale of this analytical challenge, given the need to design the appropriate analytical methodologies and reference materials that will constitute the tools for obtaining reliable analytical information.

*Key words: methylmercury, ethylmercury, organomercury compounds, bioaccumulation, speciation analysis, speciation of mercury*

### *1. Introduction*

Experts from many different organizations, including the U.S. Environmental Protection Agency (EPA), regard mercury as one of the most toxic elements [1]. The toxicity, biochemical properties and environmental cycle of mercury depend on its concentration and the chemical form in which it is present [2-6].

Every form of mercury can undergo transformation in the environment [7]. The formation of organomercury compounds as a result of naturally occurring processes is

particularly dangerous, since the organic compounds of mercury are usually more toxic than its inorganic ones [8]. The most common organic form of mercury – methylmercury (MeHg) – is a strong neurotoxin that can be bioaccumulated, and its stability in organisms and the environment affects human health and growth as well as nature in general [1,9]. Table 1 lists information on the chemical forms of mercury that most frequently occur in different compartments of the environment.

Mercury can enter the environment naturally and as a result of human actions.

Anthropogenic emissions include:

- the combustion of fossil fuels and the heating of other materials containing mercury,
- gold mining,
- the roasting of sulfide ores,
- the production of paper, drugs and the chloralkali process,
- agriculture [1,12-15].

The most important natural sources of mercury emissions are:

- volcanic eruptions,
- forest fires,
- weathering of rocks,
- tectonic movements and their associated degassing of the Earth's crust [16-19].

In the environment mercury is subject to numerous processes and transformations, such as:

- dry and wet deposition (in this way mercury gets into water bodies and ground waters),
- sorption/desorption,
- re-emission into the atmosphere of elemental mercury and/or its volatile forms,
- biomethylation/demethylation,
- bioaccumulation by fauna and flora,
- biomagnification [20].

Figure 1 illustrates the mercury cycle in diagrammatic form.

The processes that are of the greatest significance as far as the toxicity of mercury and the effect of this element on human and animal health is concerned are its biomethylation and bioaccumulation.

## 2. Methylation

In view of the mercury cycle in nature and the transformations of the element in the environment, it is aquatic ecosystems (oceans, seas, lakes, rivers and bottom sediments) that are the most susceptible to MeHg contamination [21] since practically all forms of mercury can be converted into methylmercury as a result of natural processes [1, 10, 22, 23]. Certain groups of organisms are responsible for this process, e.g. fungi [24], macroalgae and bacteria participating in the ethylation of mercury (sulfate reduction), which mostly form MeHg under anaerobic conditions in the upper layers of bottom sediments (in the top 2 cm of the sediments, where microbiological activity is at its highest [25-34]). But because methylmercury is:

- readily assimilated,
- only slowly eliminated from the body [35],
- and has lipophilic properties,

it makes up from ca 60 to 90% of the total content of mercury in living organisms [36].

The rate and degree of methylation of mercury (II) ions in waters and bottom sediments depends on such factors as:

- the form of mercury,
- the methylation agent,
- the chemical composition of the sediment,
- the quantity of oxygen available in the sediment,
- the pH of the sediment [1].

The major factors capable of retarding the biological synthesis of methylmercury are:

- the difficulty of carrying out methylation reactions in water,
- the possible decomposition of organomercury compounds by solar UV light [1, 37].

Methylmercury can pass from sediment and water to the tissues of aquatic organisms [38], where they may be bioaccumulated and biomagnified [39]. It is for this reason that maximum permissible levels of mercury and methylmercury in food have been defined. Such levels have been established by, *inter alia*, the US Food and Drug Administration (FDA),

which has stipulated a maximum level of methylmercury in fish tissue of 1 mg/kg wet weight [40].

### 3. *Bioaccumulation*

The bioaccumulation of methylmercury takes place up the aquatic food pyramid: sea water – phytoplankton – zooplankton – small herbivorous fish – large predatory fish – marine mammals [41]. Every organism from a given level in the trophic pyramid contains a higher concentration of methylmercury than organisms from lower levels, with the result that very much higher concentrations are reached in animals at the top of the pyramid in comparison with the initial concentration in water, bottom sediment or soil [42-46].

The methylmercury content in soil samples is not normally very high. But there may be from 0.5 to 1.5% methylmercury in the total soil content of mercury [19]. Mercury accumulates mostly in the upper layers of the soil because of its very strong affinity for the organic substances and certain minerals in the soil [47], which means that the bioavailability of mercury in soil is low [48].

In aquatic ecosystems both inorganic and organic forms of mercury tend to accumulate in bottom sediments. Since some sediment organisms participate in the conversion of mercury compounds into methylmercury, levels of this toxic compound in sediments are usually higher than in the soil [49] or in the water itself.

Dangerously high levels of organic mercury have been found in the tissues of fish, and in fish-feeding aquatic and terrestrial birds and mammals. In fish tissue much of this organic mercury is the toxic methylmercury. The most significant factors affecting the degree of MeHg bioaccumulation in fish are the size of the fish and/or its fat content, the protein affinity mechanism [50], and the dissolved oxygen content in the waters that the fish inhabit. In favorable conditions fish can accumulate considerable amounts of methylmercury. Fish are a significant link in the biological circulation of organic mercury compounds in nature because they are not only a basic item in the diet of many aquatic organisms, they are also consumed by humans [51-55].

Figure 2 provides information on levels of total mercury and the percentage of methylmercury in samples of organisms making up a typical food pyramid in an aquatic ecosystem.

Marine mammals have the most diversified diet as they are at the top of the food pyramid. Mercury levels are high, particularly in the liver [56-59]. Studies to determine the



toxic forms of mercury in tissues of marine mammals found that the liver contained far higher levels of mercury than the kidneys [27, 60-62]. This is connected with the storage and transformation of toxic forms of mercury in the liver [45]. The results of various studies indicate that organic forms of mercury, and methylmercury in particular, are converted to less toxic forms, e.g. HgSe, in the livers of marine mammals [45, 63]. The liver is the organ where metals accumulate, where they are detoxified and where metabolic processes involving metals take place.

Environmental contamination by organomercury compounds is not a problem solely of aquatic ecosystems: it can also affect terrestrial ecosystems, especially animals at the upper levels of the trophic pyramid [10, 64, 65].

#### 4. *The harmful action of organomercury compounds*

In the various compartments of the environment mercury is subject to transformation, and contact with any form of mercury has toxic effects. In the case of organomercury compounds, poisoning results mostly from the consumption of toxic methylmercury or other compounds together with food. Particularly dangerous are methylmercury and ethylmercury, which are almost entirely absorbed in the digestive tract [6, 8, 15, 50, 66], readily crossing biological barriers such as the blood-brain barrier, and also the placental barrier, accumulating in the fetus and maternal milk [67-70]. Even though ethylmercury is less toxic than methylmercury, both compounds elicit similar symptoms [71, 72]. Exposure to organic compounds of mercury, methylmercury in particular, in the prenatal period or directly after birth (these compounds are consumed with the mother's milk) affects early childhood development. Figure 3 illustrates the circulation of methylmercury in the mother's body and the fetus.

Following exposure to methylmercury, its metabolites (forming as a result of demethylation in the liver) are excreted in the urine and the stool, but only to a slight extent, because methylmercury is subject to hepato-intestinal recirculation [15].

Because of its strong affinity for sulfur, and hence for sulfhydryl groups, methylmercury reacts with proteins and enzymes causing dysfunction of organs, blockage of enzyme binding sites and protein synthesis, impedes thymidine incorporation into DNA [74] and has an extremely harmful influence on the entire central nervous system of humans and other organisms [23, 75]. That is why methylmercury accumulates to a far greater extent in



tissues rich in the sulfhydryl groups of amino acids than, for example, in fatty tissue [23]. Moreover, short-chain compounds like methylmercury or ethylmercury readily penetrate red blood cells, where they bind to hemoglobin. This is the reason why methylmercury, unlike the inorganic forms of mercury, accumulates in erythrocytes [8].

A less often encountered but more toxic form of mercury – consumption of just 15-20 microliters is lethal – is dimethylmercury (DMM). The physical properties, along with its lipophilicity (stronger than that of MMM), which is due to the presence of the second alkyl group, mean that DMM is rapidly absorbed through the skin, lungs and digestive tract and is accumulated in the fatty tissue, blood serum proteins and the brain.

Once DMM gets into the body, it is first converted into the monomethyl form and is transported to the blood and tissues.

Therefore, the toxic effects of DMM are associated with its dealkylation, and all are exactly the same as those of MMM exposure. Methylmercury reacts with sulfur and the sulfur-containing thiol groups of enzymes, thereby inhibiting their action.

The other organic compounds of mercury are rarely encountered and are rarely investigated by analysts engaged in environmental studies. Table 2 provides information on the toxic effects of some organic forms of mercury.

The toxic effects of inorganic mercury have been known since antiquity [15]. But the first information on mortality that turned out to have been caused by mercury poisoning was published in the 19<sup>th</sup> century. Later reports on the poisoning of humans by organomercury compounds have come from many parts of the world. But most cases of disease or the adverse effects of organomercury compounds on humans have been recorded in Asia [1]. Table 3 lists information on the most important events associated with poisoning by organic compounds of mercury.

##### *5. The content of organomercury compounds in environmental samples.*

The most important factor governing the level of organomercury compounds in the tissues of an animal appears to be the diet [45, 85, 86]. Hence, the type and variety of food consumed is intimately connected with the level of mercury in the tissues and organs of organisms. Table 4 lists literature information on levels of organomercury compounds in samples of water, sediments and the tissues of animals at different levels of the trophic pyramid.



## 6. *Speciation of mercury*

Human activities release into the environment large quantities of mercury, which can take on highly toxic forms. In view of the fact that the toxicity, mobility, bioavailability and bioaccumulation of mercury depends on its chemical form, it is necessary to determine the individual forms of mercury and not the total concentration of the metal in environmental samples. This type of determination is possible with the application of speciation analysis [1, 113].

Speciation as a field of research made its appearance in the 1970s at a time coinciding with the development of numerous analytical procedures and techniques, which enabled the quantitative determination of elements in amounts of the order of  $10^{-6}$  % -  $10^{-7}$  %.

Contemporary speciation analysis focuses primarily on biologically active elements. It is used to establish the metabolism and biological activity of different elements in living organisms. It is also applied in food chemistry, pharmacy, biology, toxicology and environmental studies, and even in studies of historical monuments.

Obtaining reliable measurements of the content of organomercury compounds in environmental samples requires the application of analytical procedures consisting of the following steps:

1. extraction of organic forms of mercury from samples (soil, sediment, living organisms) in such a way as to prevent degradation and chemical changes that could alter the original composition of the compounds in the sample; it should also fulfill the basic requirements of trace analysis with respect to analyte loss and possible sample contamination [1];
2. preconcentration of analytes, as a result of which the concentration ratio or the quantities of the microconstituents (trace constituents) and macroconstituents (matrix) increase in such a manner that it becomes possible to obtain values below the limit of detection of the chosen determination technique [1, 114];
3. isolation of the various forms of mercury in such a way as not to change the concentration of the individual compounds in the sample [1];
4. determination of each of the earlier isolated forms of mercury [1].



## 7. Extraction

The extraction of the various forms of mercury is regarded as one of the most important stages in speciation analysis. Depending on the type and form of the sample, liquid-liquid and liquid-solid extraction are used for preparing samples for speciation.

Extraction should be efficient and effective and, above all, should not alter in any way the forms of mercury present in the sample and should not promote the formation of new compounds in the sample [5, 115]. It does happen, however, that the accuracy of the final results may deteriorate as a result of the formation of methylmercury from inorganic forms of mercury during extraction with a solution of an acid or base [5, 116, 117] or transformations of forms of mercury among themselves [3, 118, 119]. When choosing a suitable extraction technique or extractant, one should be guided by:

- ✓ the chemical properties of the analyte,
- ✓ the chemical form of the analyte,
- ✓ the matrix composition of the sample,
- ✓ the technique for determining the analyte.

To improve the efficacy of extraction some form of assistance is increasingly being applied, for example, with:

- ✓ ultrasound (UAE) [50, 115, 120],
- ✓ elevated pressure (PFE),
- ✓ microwave radiation (MAE) [121-125]

Figure 4 illustrates a scheme for the preliminary preparation of samples prior to the determination of the various forms of mercury [126].

## 8. The conversion of analytes into volatile derivatives

All types of environmental samples are highly complicated research materials because of their complex matrix composition, the diversity of compounds in the sample and the range of analyte concentrations. Despite the application of many different extraction techniques, the determination of some substances in unchanged form in environmental samples is impossible. In such cases, therefore, derivatization has to be resorted to, that is, the analyte is converted into derivatives that have properties enabling them to be determined with a particular analytical technique. Derivatization is very often used to isolate various forms of mercury from a sample or to pre-concentrate an analyte, in that mercury's ability to form vapors when





reacting with a reducing agent is made use of. It is often the case that the conversion of mercury into volatile derivatives is preceded by isolation using gas chromatography.

Mercury compounds are usually derivatized using one of three reactions:

1. Reaction with a reducing agent – sodium borohydride ( $\text{NaBH}_4$ ). This type of reaction preserves the mercury-carbon bond in the target molecule. The reaction with  $\text{NaBH}_4$  is easy to carry out (it takes place in an aqueous medium), but determining mercury compounds after such a reaction may be troublesome because of the instability of the derivatives formed and the possibility of disproportionation [140,141].
2. Alkylation. This requires less time and effort than derivatization with Grignard reagents because there is no need to change the solvents. Moreover, this method is neutral vis-à-vis most reagents, which caused problems during the formation of hydrides, i.e. the presence of metal ions, proteins, fats or humus substances; it can therefore be used for analyzing samples of diverse provenance, including biological ones [G32]. A particular advantage of alkylation is that extraction and derivatization can be performed simultaneously, which considerably shortens the sample preparation time. But there are also drawbacks to this method, the principal one being the small number of commercially available derivatizing reagents. The three main alkylating reagents are:
  - ✓ Sodium tetraethyl borate,
  - ✓ Sodium tetraphenyl borate,
  - ✓ Tetrabutylammonium tetrabutyl borate.

$\text{NaBEt}_4$  is universally used for the speciation of organometallic compounds, but it is of no use for derivatizing inorganic compounds of mercury or ethylmercury, since both forms produce the same compound,  $\text{HgEt}_2$ . These two forms of mercury can be distinguished if sodium phenyl borate is used – this reacts only with mercury derivatives [142-147].

3. Reaction with Grignard reagents [142,148,149] (alkylmagnesium halides, e.g. methyl, ethyl, propyl, butyl, pentyl, hexyl, or also phenylmagnesium chlorides or bromides). A serious problem limiting the use of such reactions for derivatization is the need to perform them in an anhydrous environment, which is time-consuming and labor-intensive. Prior to derivatization, extracts have to be dried and/or transferred to another solvent, and after derivatization the excess Grignard reagent has to be removed.

Since the conversion of an analyte into volatile derivatives may be source of numerous errors, this step is avoided during speciation analysis if at all possible.

### *9. Techniques for isolating and detecting analytes*

Since biological samples have a very complex composition and the organic forms of mercury are present in them at very low levels, selective isolation techniques with sensitive and targeted detection methods are needed. Known as conjugated, hybrid or linked systems, such combinations offer a substantial improvement in sensitivity, and the time of analysis is far shorter [113].

The choice of isolation technique is connected with the physicochemical properties of the target analyte. The most commonly used isolation techniques in speciation analysis are gas chromatography (GC), high-performance liquid chromatography (HPLC) and capillary electrophoresis (CE). Table 5 lists basic information on the techniques used with environmental samples for detecting and determining various speciation forms of mercury in them. Just as the sample preparation technique depends on the type of matrix in which the target analytes are present, so the choice of isolation technique depends on the physicochemical properties of these analytes, i.e. volatility, charge, polarity. This is why GC is used to isolate volatile, thermally stable and neutral compounds (or those that can be converted into volatile and stable derivatives), whereas HPLC is suitable for isolating the remaining compounds.

In view of the wide variety of environmental and biological samples, not to mention the plethora of analytical techniques, procedures should be selected that enable the lowest possible levels of organomercury compounds to be determined. Table 6 lists basic information on the analytical procedures used in speciation studies of mercury in environmental samples.

### *10. Summary*

The hazards arising out of the presence of mercury and its compounds in the environment, especially in aquatic ecosystems, make it necessary to understand the transformation pathways of these compounds and to monitor the contents of organomercury compounds in the tissues of organisms at all levels of the trophic pyramid. The possibility that organic forms of mercury can get into the human body with food is a serious threat because



these compounds are highly toxic. Many cases – some fatal – have been reported of disease among both adults and children following their exposure to organomercury compounds. That is why it is so crucial to monitor levels of organomercury compounds in samples of water, sediments and tissues from all levels of the trophic pyramid using the tool of speciation analysis.

## ACKNOWLEDGEMENT

This work was financially supported by the Foundation for Polish Science (MISTRZ program).

## REFERENCES

- [1] Sánchez Uría J.E., Sanz-Medel A., Inorganic and methylmercury speciation in environmental samples: Review, *Talanta* 47, 509, 1998
- [2] Selin N.E., Global Biogeochemical Cycling of Mercury: A Review, *Annu. Rev. Environ. Resour.* 34, 43, 2009
- [3] Leermakers M., Baeyens W., Quevauviller P., Horvat M., Mercury in environmental samples: speciation, artifacts and validation, *Trends Anal. Chem.* 24, 383, 2005
- [4] Peixoto N.C., Serafim M.A., Flores E.M.M., Bebianno M.J., Pereira M.E., Metallothionein, zinc, and mercury levels in tissues of young rats exposed to zinc and subsequently to mercury, *Life Sci.* 81, 1264, 2007
- [5] López I., Cuello S., Cámara C., Madrid Y., Approach for rapid extraction and speciation of mercury using a microtip ultrasonic probe followed by LC–ICP-MS, *Talanta* 82, 594, 2010
- [6] Duarte F.A., Bizzi C.A., Goldschmidt Antes F., Dressler V.L., de Moraes Flores É.M., Organic, inorganic and total mercury determination in fish by chemical vapor generation with collection on a gold gauze and electrothermal atomic absorption spectrometry, *Spectrochim. Acta Part B* 64, 513, 2009

- [7] Houserová P., Kubáň V., Kráčmar S., Sitko J., Total mercury and mercury species in birds and fish in an aquatic ecosystem in the Czech Republic, *Environ. Pollut.* 145, 185, 2007
- [8] Graeme K.A., Pollack C.V., Heavy metal toxicity, part I: arsenic and mercury, *J. Emerg. Med.* 16, 45, 1998
- [9] Dresslera V.L., Goldschmidt Antes F., Marques Moreira C., Pozebon D., Duarte F.A., As, Hg, I, Sb, Se and Sn speciation in body fluids and biological tissues using hyphenated-ICP-MS techniques: A review, *Int. J. Mass Spectrom.* 307, 149, 2011
- [10] Fu J., Wang Y., Zhou Q., Jiang G., Trophic transfer of mercury and methylmercury in an aquatic ecosystem impacted by municipal sewage effluents in Beijing, China, *J. Environ. Sci.* 22, 1189, 2010
- [11] Vezér T., Papp A., Kurunczi A., Párducz Á., Náray M., Nagymajtényi L., Behavioral and neurotoxic effects seen during and after subchronic exposure of rats to organic mercury, *Environ. Toxicol. Pharmacol.* 19, 785, 2005
- [12] Magos L., Review on the toxicity of ethylmercury, including its presence as a preservative in biological and pharmaceutical products. *J. Appl. Toxicol.* 21, 1, 2001
- [13] Clarkson T.W., The three modern faces of mercury, *Environ. Health Perspect.* 110, 11, 2002
- [14] Morel F.M.M., Kraepiel A.M.L., Amyot M., The chemical cycle and bioaccumulation of mercury, *Annu. Rev. Ecol. Syst.* 29, 543, 1998
- [15] Ceccatelli S., Daré E., Moors M., Methylmercury-induced neurotoxicity and apoptosis, *Chem. Biol. Interact.* 188, 301, 2010
- [16] Schroeder W.H., Munthe J., Atmospheric mercury - an overview, *Atmos. Environ.* 32, 809, 1997
- [17] Pacyna E.G., Pacyna J.M., Steenhuisen F., Wilson S.J., Global anthropogenic mercury emission inventory for 2000, *Atmos. Environ.* 40, 4048, 2006

- [18] Swain E.B., Jakus P., Rice G., Lupi F., Maxson P.A., Pacyna J.M., Penn A., Spiegel S.J., Veiga M.M., Socioeconomic consequences of mercury use and pollution, *Ambio* 36, 46, 2007
- [19] Rieder S.R., Brunner I., Horvat M., Jacobs A., Frey B., Accumulation of mercury and methylmercury by mushrooms and earthworms from forest soils, *Environ. Pollut.* 159, 2861, 2011
- [20] Margetínová J., Houserová – Pelcová P., Kubáň V., Speciation analysis of mercury in sediments, zoobenthos and river water samples by high-performance liquid chromatography hyphenated to atomic fluorescence spectrometry following preconcentration by solid phase extraction, *Anal. Chim. Acta* 615, 115, 2008
- [21] Fitzgerald W.F., Lamborg C.H., Hammerschmidt C.R., Marine biogeochemical cycling of mercury, *Chem. Rev.* 107, 641, 2007
- [22] Yang L., Mester Z., Sturgeon R.E., Determination of methylmercury in fish tissues by isotope dilution SPME-GC-ICP-MS, *J. Anal. At. Spectrom.* 18, 431, 2003
- [23] Vereda Alonso E., Siles Cordero M.T., García de Torres A., Cañada Rudner P., Cano Pavón J.M., Mercury speciation in sea food by flow injection cold vapor atomic absorption spectrometry using selective solid phase extraction, *Talanta* 77, 53, 2008
- [24] Vonk J.W., Sijpesteijn A.K., Studies on the methylation of mercuric chloride by pure cultures of bacteria and fungi, *Antonie van Leeuwenhoek* 39, 505, 1973
- [25] Ullrich S.M., Ilyushchenko M.A., Uskov G.A., Tanton T.W., Mercury distribution and transport in a contaminated river system in Kazakhstan and associated impacts on aquatic biota, *Appl. Geochem.* 22, 2706, 2007
- [26] Benoit J.M., Gilmour C.C., Mason R.P., Riedel G.S., Riedel G.F., The sources and cycling of mercury in the Patuxent estuary, *Biogeochemistry* 40, 249, 1998
- [27] Hajeb P., Jinap S., Ahmad I., Biomagnifications of mercury and methylmercury in tuna and mackerel, *Environ. Monit. Assess.* 171, 205, 2010

- [28] Mason R.P., Lawrence A.L., Concentration, distribution, and bioavailability of mercury and methylmercury in sediments of Baltimore Harbor and Chesapeake Bay, Maryland, USA, *Environ. Toxicol. Chem.* 18, 2438, 1999
- [29] Monson B.A., Brezonik P.L., Seasonal patterns of mercury species in water and plankton from softwater lakes in Northeastern Minnesota, *Biogeochemistry* 40, 147, 1998
- [30] Gilmour C.C., Henry E.A., Mitchell R., Sulfate stimulation of mercury methylation in freshwater sediments, *Environ. Sci. Technol.* 26, 2281, 1992
- [31] Matilainen T., Involvement of bacteria in methylmercury formation in anaerobic lake waters, *Water Air Soil Pollut.* 80, 757, 1995
- [32] St. Louis V.L., Rudd J.W.M., Kelly C.A., Beatly K.G., Flett R.J., Roulet N.T., Production and loss of methylmercury and loss of total mercury from boreal forest catchments containing different types of wetlands, *Environ. Sci. Technol.* 30, 2719, 1996
- [33] Holloway J.M., Goldhaber M.B., Scow K.M., Drenovsky R.E., Spatial and seasonal variations in mercury methylation and microbial community structure in a historic mercury mining area, Yolo County, California, *Chemical Geology* 267, 85, 2009
- [34] Merritt K.A., Amirbahman A., Mercury methylation dynamics in estuarine and coastal marine environments — A critical review, *Earth Sci. Rev.* 96, 54, 2009
- [35] Ebdon L., Foulkers M.E., Le Roux S., Muñoz-Olivas R., Cold vapour atomic fluorescence spectrometry and gas chromatography-pyrolysis-atomic fluorescence spectrometry for routine determination of total and organometallic mercury in food samples, *Analyst* 127, 1108, 2002
- [36] Endo T., Haraguchi K., Sakata M., Mercury and selenium concentrations in the internal organs of toothed whales and dolphins marketed for human consumption in Japan, *Sci. Total Environ.* 300, 15, 2002

[37] Suda I., Sudal M., Hirayama K., Degradation of methyl and ethyl mercury by singlet oxygen generated from sea water exposed to sunlight or ultraviolet light, *Arch. Toxicol.* 67, 365, 1993

[38] Clarkson T.W., Magos L., The toxicology of mercury and its chemical compounds, *Crit. Rev. Toxicol.* 36, 609, 2006

[39] Lawrence A.L., Mc Aloon K.M., Mason R.P., Mayer L.M., Intestinal solubilization of particle associated organic and inorganic mercury as a measure of bioavailability to benthic invertebrates, *Environ. Sci. Technol.* 33, 1871, 1999

[40] Havarinasab S., Hulman P., Organic mercury compounds and autoimmunity, *Autoimmunity Reviews* 4, 270, 2005

[41] Ando T., Yamamoto M., Tomiyasu T., Hashimoto J., Miura T., Nakano A., Akiba S., Bioaccumulation of mercury in a vestimentiferan worm living in Kagoshima Bay, Japan, *Chemosphere* 49, 477, 2002

[42] Dovydaitis T., Fish Consumption During Pregnancy: An Overview of the Risks and Benefits, *Journal of Midwifery & Women's Health* 53, 325, 2008

[43] Kunito T., Nakamura S., Ikemoto T., Anan Y., Kubota R., Tanabe S., Rosas F.C.W., Fillmann G., Readman J.W., Concentration and subcellular distribution of trace elements in liver of small cetaceans incidentally caught along the Brazilian coast, *Mar. Pollut. Bull.* 49, 574, 2004

[44] Feroci G., Badiello R., Fini A., Interactions between different selenium compounds and zinc, cadmium and mercury, *J. Trace Elem. Med. Biol.* 18, 227, 2005

[45] Seixas T.G., Kehrig H.A., Costa M., Fillmann G., Di Benedetto A.P.M., Secchi E.R., Souza C.M.M., Malm O., Moreira I., Total mercury, organic mercury and selenium in liver and kidney of a South American coastal dolphin, *Environ. Pollut.* 154, 98, 2008

[46] Wang Q., Kim D., Dionysiou D.D., Sorial G.A., Timberlake D., Sources and remediation for mercury contamination in aquatic systems - literature review, *Environ. Pollut.* 131, 323, 2004

[47] Ravichandran M., Interactions between mercury and dissolved matter - a review, *Chemosphere* 55, 319, 2004

[48] Tipping E., Lofts S., Hooper H., Frey B., Spurgeon D., Svendsen C., Critical limits for Hg (II) in soils, derived from chronic toxicity data, *Environ. Pollut.* 158, 2465, 2010

[49] Barańkiewicz D., Gramowska H., Specjacja metylortęci i rtęci nieorganicznej w wodzie i osadzie jeziora Swarzędzkiego (in polish), In: Specjacja Chemiczna, Barańkiewicz D., Bulska E., 60, 2009

[50] López I., Cuello S., Cámara C., Madrid Y., Approach for rapid extraction and speciation of mercury using a microtip ultrasonic probe followed by LC-ICP-MS, *Talanta* 82, 594, 2010

[51] Wu Y., Wang W.X., Accumulation, subcellular distribution and toxicity of inorganic mercury and methylmercury in marine phytoplankton, *Environ. Pollut.* 159, 3097, 2011

[52] Guallar E., Sanz-Gallardo M.I., Van't Veer P., Bode P., Aro A., Gomez-Aracena J., Kark J.D., Riemersma A.R., Martín-Moreno J.M., Frans J.K., Mercury, fish oil, and the risk of myocardial infarction, *N. Engl. J. Med.* 347, 1747, 2002

[53] Rissanen T., Voutilainen S., Nyssönen K., Lakka T.A., Salonen J.T., Fish oil-derived fatty acids, docosahexaenoic acid and docosapentanoic acid, and the risk of acute coronary events, *Circulation* 102, 2677, 2000

[54] Lucotte M., Mucci A., Hillaire-Marcel C., Pichet P., Grondin A., Anthropogenic mercury enrichment in remote lakes of northern Québec (Canada), *Water Air Soil Pollut.* 80, 467, 1995

[55] Tremblay A., Lucotte M., Schetagne R., Total mercury and methylmercury accumulation in zooplankton of hydroelectric reservoirs in northern Québec (Canada), *Sci. Total Environ.* 213, 307, 1998





[56] Holsbeek L., Siebert U., Joiris C.R., Heavy metals in dolphins stranded on the French Atlantic coast, *Sci. Total Environ.* 217, 241, 1998

[57] Meador J.P., Ernest D., Hohn A.A., Tilbury K., Gorzelany J., Worthy G., Stein J.E., Comparison of elements in bottlenose dolphins stranded on the beaches of Texas and Florida in the gulf of Mexico over a one-year period, *Arch. Environ. Contam. Toxicol.* 36, 87, 1999

[58] Zhou J.L., Salvador S.M., Liu Y.P., Sequerira M., Heavy metals in the tissue of common dolphins (*Delphinus delphis*) stranded on the Portuguese coast, *Sci. Total Environ.* 273, 61, 2001

[59] Endo T., Haraguchi K., Cipriano F., Simmonds M.P., Hotta Y., Sakata M., Contamination by mercury and cadmium in the cetacean products from Japanese market, *Chemosphere* 54, 1653, 2004

[60] Gerpe M.S., Rodriguez D.H., Moreno V.J., Bastida R.O., Moreno J.A.E., Accumulation of heavy metals in the Franciscana (*Pontoporia blainvillei*) from Buenos Aires Province, Argentina, *Lat. Am. J. Aquat. Mamm.* 1, 95, 2002

[61] Lailson-Brito J., Azeredo M.A.A., Malm O., Ramos R.A., Di Benedetto A.P., Saldanha M.F.C., Trace metals in liver and kidney of the Franciscana (*Pontoporia blainvillei*) from the northern coast of Rio de Janeiro State, Brazil, *Lat. Am. J. Aquat. Mamm.* 1, 107, 2002

[62] Seixas T.G., Kehrig H.A., Fillmann G., Di Benedetto A.P.M., Souza C.M.M., Secchi E.R., Moreira I., Malm O., Ecological and biological determinants of trace elements accumulation in liver and kidney of *Pontoporia blainvillei*, *Sci. Total Environ.* 385, 208, 2007

[63] Ikemoto T., Kunito T., Watanabe I., Yasunaga G., Baba N., Miyazaki N., Petrov E.A., Tanabe S., Comparison of trace element accumulation in Baikal seals (*Pusa sibirica*), Caspian seals (*Pusa caspica*) and northern fur seals (*Callorhinus ursinus*), *Environ. Pollut.* 127, 83, 2004



[64] Cristol D.A., Brasso R.L., Condon A.M., Fovargue R.E., Friedman S.L., Hallinger K.K., Monroe A.P., White A.E., The movement of aquatic mercury through terrestrial food webs, *Science* 320, 335, 2008

[65] Orihel D.M., Paterson M.J., Blanchfield P.J., Bodaly R.A., Hintelmann H., Experimental evidence of a linear relationship between inorganic mercury loading and methylmercury accumulation by aquatic biota, *Environ. Sci. Technol.* 41, 4952, 2007

[66] Evans R.D., Addison E.M., Villeneuve J.Y., MacDonald K.S., Joachim D.G., Distribution of Inorganic and Methylmercury among Tissues in Mink (*Mustela vison*) and Otter (*Lutra canadensis*), *Environ. Res. A* 84, 133, 2000

[67] Amin-Zaki L., Elhassani S., Majeed M.A., Clarkson T.W., Doherty R.A., Greenwood M.R., Intra-uterine methylmercury poisoning in Iraq. *Pediatrics* 54, 587, 1974

[68] Honda S., Hylander L., Sakamoto M., Recent advances in evaluation of health effects on mercury with special reference to methylmercury – a minireview, *Environ. Health Prev. Med.* 11, 171, 2006

[69] Grandjean P., Jorgensen P.J., Weihe P., Human milk as a source of methylmercury exposure in infants, *Environ. Health Perspect.* 102, 74, 1994

[70] Oskarsson A., Schultz A., Skerfving S., Hallen I.P., Ohlin B., Lagerkvist B.J., Total and inorganic mercury in breast milk in relation to fish consumption and amalgam in lactating women, *Arch. Environ. Health* 51, 234, 1996

[71] Eto K., Pathology of Minamata Disease, *Toxicol. Pathol.* 25, 614, 1997

[72] Peralta-Videa J.R., Lopez M.L., Narayan M., Saupe-Torresdey G., The biochemistry of environmental heavy metal uptake by plants: Implications for the food chain, *Int. J. Biochem. Cell Biol.* 41, 1665, 2009

[73] Cernichiari E., Myers G.J., Ballatori N., Zareba G., Vyas J., Clarkson T., The biological monitoring of prenatal exposure to methylmercury, *NeuroToxicology* 28, 1015, 2007



[74] Syversen T., Kaur P., The toxicology of mercury and its compounds, *J. Trace Elem. Med. Biol.* doi:10.1016/j.jtemb.2012.02.004, 2012

[75] Capelo J.L., Maduro C., Mota A.M., Evaluation of focused ultrasound and ozonolysis as sample treatment for direct determination of mercury by FI-CV-AAS. Optimization of parameters by full factorial design, *Ultrason. Sonochem.* 13, 98, 2006

[76] Weihe P., Grandjean P., Debes F., White R., Health implications for Faroe islanders of heavy metals and PCBs from pilot whales, *Sci. Total Environ.* 186, 141, 1996

[77] Grandjean P., Weihe P., White R.F., Debes F., Cognitive performance of children prenatally exposed to “safe” levels of methylmercury, *Environ. Res. A* 77, 165, 1998

[78] HaMai D., Bondy S.C., Dimethylmercury In: Encyclopedia of Toxicology. Wexler P (ed). Second Edition, Academic Press, Bethesda, 2005

[79] Powell P.P., Minamata disease: a story of mercury’s malevolence, *South. Med. J.* 84, 1352, 1991

[80] Hunter D., Russell D.S., Focal cerebral and cerebellar atrophy in a human subject due to organic mercury compounds, *J. Neurol. Neurosurg. Psychiatr.* 17, 235, 1954

[81] O’Shea J.G., “Two minutes with Venus, two years with mercury”: mercury as an antisymphilitic chemotherapeutic agent, *J. R. Soc. Med.* 83, 392, 1990

[82] Bakir F., Damluji S.F., Amin-Zaki L., Murtadha M., Khalidi A., Al-Rawi N.Y., Tikriti S., Dhahir H.I., Clarkson T.W., Smith J.C., Doherty R.A., Methylmercury poisoning in Iraq, *Science* 181, 230, 1973

[83] Bakir F., Rustom H., Tikriti S., Al-Damluji S.F., Shihristani H., Clinical and epidemiological aspects of methylmercury poisoning, *Postgrad. Med. J.* 56, 1, 1980



[84] Harada M., Nakanishi J., Yasoda E., Pinheiro M.C., Oikawa T., de Assis Guimaraes G., da Silva Cardoso B., Kizaki T., Ohno H., Mercury pollution in the Tapajos River basin, Amazon: mercury level of head hair and health effects, *Environ. Int.* 27, 285, 2001

[85] Bustamante P., Caurant F., Fowler S.W., Miramand P., Cephalopods as a vector for the transfer of cadmium to top marine predators in the northeast Atlantic Ocean, *Sci. Total Environ.* 220, 71, 1998

[86] Monaci F., Borre A., Leonzio C., Marsili L., Calzada N., Trace elements in striped dolphins (*Stenella coeruleoalba*) from the western Mediterranean, *Environ. Pollut.* 99, 61, 1998

[87] Molina C.I., Gibon F.M., Duprey J.L., Dominguez E., Guimarães J.R.D., Roulet M., Transfer of mercury and methylmercury along macroinvertebrate food chains in a floodplain lake of the Beni River, Bolivian Amazonia, *Sci. Total Environ.* 408, 3382, 2010

[88] Shia J., Lianga L., Jianga G., Jin X., The speciation and bioavailability of mercury in sediments of Haihe River, China, *Environ. Int.* 31, 357, 2005

[89] Roulet M., Lucotte M., Guimarães J.R.D., Rheault I., Methylmercury in water, seston, and epiphyton of an Amazonian river and its floodplain, Tapajós River, Brazil, *Sci. Total Environ.* 261, 43, 2000

[90] Gómez-Ariza J.R., Lorenzo F., García-Barrera T., Simultaneous determination of mercury and arsenic species in natural freshwater by liquid chromatography with on-line UV irradiation, generation of hydrides and cold vapor and tandem atomic fluorescence detection, *J. Chromatogr. A* 1056, 139, 2004

[91] Kehrig H.A., Seixas T.G., Baêta A.P., Malm O., Moreira I., Inorganic and methylmercury: Do they transfer along a tropical coastal food web?, *Mar. Pollut. Bull.* 60, 2350, 2010

[92] Back R.C., Gorski P.R., Cleckner L.B., Hurley J.P., Mercury content and speciation in the plankton and benthos of Lake Superior, *Sci. Total Environ.* 304, 349, 2003



- [93] Ali I.B, Joiris C.R., Holsbeek L., Total and organic mercury in the starfish *Ctenodiscus crispatus* and the polychaete *Maldanes sarsi* from the Barents Sea, *Sci. Total Environ.* 201, 189, 1997
- [94] Yamashita Y., Omura Y., Okazaki E., Total mercury and methylmercury levels in commercially important fishes in Japan, *Fish. Sci.* 71, 1029, 2005
- [95] Claisse D., Cossa D., Bretaudeau – Sanjuan J., Touchard G., Bombled B., Methylmercury in Molluscs along the French Coast, *Mar. Pollut. Bull.* 42, 329, 2001
- [96] Raimundo J., Vale C., Canário J., Branco V., Moura I., Relations between mercury, methyl-mercury and selenium in tissues of *Octopus vulgaris* from the Portuguese Coast, *Environ. Pollut.* 158, 2094, 2010
- [97] Maggi C., Berducci M.T., Bianchi J., Giani M., Campanella L., Methylmercury determination in marine sediment and organisms by Direct Mercury Analyser, *Anal. Chim. Acta* 641, 32, 2009
- [98] Rahman S.A., Wood A.K., Sarmani S., Majid A.A., Determination of mercury and organic mercury contents in Malaysian seafood, *J. Radioanal. Nucl. Chem.* 217, 53, 1997
- [99] Mikac N., Kwokal Ž., Martinčić D., Branica M., Uptake of mercury species by transplanted mussels *Mytilus galloprovincialis* under estuarine conditions (Krka river estuary), *Sci. Total Environ.* 184, 173, 1996
- [100] Minganti V., Capelli R., De Pellegrini R., Orsi Relini L., Relini G., Total and organic mercury concentrations in offshore crustaceans of the Ligurian Sea and their relations to the trophic levels, *Sci. Total Environ.* 184, 149, 1996
- [101] Storelli M.M., Giacomini-Stuffler R., Marcotrigiano G.O., Total and methylmercury residues in cartilaginous fish from Mediterranean Sea, *Mar. Pollut. Bull.* 44, 1354, 2002
- [102] Ikingura J.R., Akagi H., Total mercury and methylmercury levels in fish from hydroelectric reservoirs in Tanzania, *Sci. Total Environ.* 304, 355, 2003



[103] Jin L., Liang L., Jiang G., Xu Y., Methylmercury, total mercury and total selenium in four common freshwater fish species from Ya-Er Lake, China, *Environ. Geochem. Health* 28, 401, 2006

[104] Kampalath R., Gardner S.C., Méndez-Rodríguez L., Jay J.A., Total and methylmercury in three species of sea turtles of Baja California Sur, *Mar. Pollut. Bull.* 52, 1784, 2006

[105] Ruelas-Inzunza J., Hernández-Osuna J., Páez-Osuna F., Organic and total mercury in muscle tissue of five aquatic birds with different feeding habits from the SE Gulf of California, Mexico, *Chemosphere* 76, 415, 2009

[106] Kim E.Y., Murakami T., Saeki K., Tatsukawa R., Mercury Levels and its Chemical Form in Tissues and Organs of Seabirds, *Arch. Environ. Contam. Toxicol.* 30, 259, 1996

[107] Campbell L.M., Norstrom R.J., Hobson K.A., Muir D.C.G., Backus S., Fisk A.T., Mercury and other trace elements in a pelagic Arctic marine food web (Northwater Polynya, Baffin Bay), *Sci. Total Environ.* 351, 247, 2005

[108] Gardner W.S., Kendall D.R., Odom R.R., Windom H.L., Stephens J.A., The distribution of methylmercury in a contaminated salt marsh ecosystem, *Environ. Pollut.* 15, 243, 1978

[109] Joiris C.R., Holsbeek L., Bolba D., Gascard C., Stanev T., Komakhidze A., Baumgärtner W., Birkun A., Total and organic mercury in the Black Sea Harbour Porpoise *Phocoena phocoena relicta*, *Mar. Pollut. Bull.* 42, 905, 2001

[110] Endo T., Kimura O., Hisamichi Y., Minoshima Y., Haraguchi K., Kakumoto C., Kobayashi M., Distribution of total mercury, methyl mercury and selenium in pod of killer whales (*Orcinus Orca*) stranded in the northern area of Japan: Comparison of mature females with calves, *Environ. Pollut.* 144, 145, 2006

[111] Endo T., Haraguchi K., Hotta Y., Hisamichi Y., Lavery S., Dalebout M.L., Scott Baker C., Total mercury, methyl mercury, and selenium levels in the red meat of small cetaceans sold for human consumption in Japan, *Environ. Sci. Technol.* 39, 5703, 2005

[112] Joiris C.R., Holsbeek L., Bouquegneau J., Bossicart M., Mercury contamination of the harbour porpoise *Phocoena Phocoena* and other cetaceans from the North Sea and the Kattegat, *Water Air Soil Pollut.* 56, 283, 1991

[113] Yan L., Shu-Juan L., Dong-Qing J., Yan J., Xiu-Ping Y., Gas Chromatography-Inductively Coupled Plasma-Mass Spectrometry for Mercury Speciation in Seafood, *Chin. J. Anal. Chem.* 36, 793, 2008

[114] IUPAC Compendium of Chemical Terminology 2nd Edition, 1997

[115] Meng W., Weiyue F., Junwen S., Fang Z., Bing W., Motao Z., Baia L., Yuliang Z., Zhifang C., Development of a mild mercaptoethanol extraction method for determination of mercury species in biological samples by HPLC–ICP-MS, *Talanta* 71, 2034, 2007

[116] Hintelmann H., Falter R., Ilgen G., Evans R.D., Determination of artifactual formation of monomethylmercury ( $\text{CH}_3\text{Hg}^+$ ) in environmental samples using stable  $\text{Hg}^{2+}$  isotopes with ICP-MS detection: Calculation of contents applying species specific isotope addition, *Fresenius J. Anal. Chem.* 358, 363, 1997

[117] Cabañero Ortiz A.I., Madrid Albarrán Y., Cámara Rica C., Evaluation of different sample pre-treatment and extraction procedures for mercury speciation in fish samples, *J. Anal. At. Spectrom.* 17, 1595, 2002

[118] Hirner A.V., Speciation of alkylated metals and metalloids in the environment, *Anal. Bioanal. Chem.* 385, 555, 2006

[119] Maldonado Santoyo M., Landero Figueroa J.A., Wrobel K., Wrobel K., Analytical speciation of mercury in fish tissues by reversed phase liquid chromatography–inductively coupled plasma mass spectrometry with  $\text{Bi}^{3+}$  as internal standard, *Talanta* 79, 706, 2009

[120] de Souza S.S., Rodrigues J.L., de Oliveira Souza V.C., Barbosa Jr F., A fast sample preparation procedure for mercury speciation in hair samples by high-performance liquid chromatography coupled to ICP-MS, *J. Anal. At. Spectrom.* 25, 79, 2010



[121] Reyes L.H., Mizanur Rahman G.M., Skip Kingston H.M., Robust microwave-assisted extraction protocol for determination of total mercury and methylmercury in fish tissues, *Anal. Chim. Acta* 631, 121, 2009

[122] Chang L.F., Jiang S.J., Sahayamb A.C., Speciation analysis of mercury and lead in fish samples using liquid chromatography–inductively coupled plasma mass spectrometry, *J. Chromatogr. A* 1176, 143, 2007

[123] Reyes L.H., Mizanur Rahman G.M., Fahrenholz T., Skip Kingston H.M., Comparison of methods with respect to efficiencies, recoveries, and quantitation of mercury species interconversions in food demonstrated using tuna fish, *Anal. Bioanal. Chem.* 390, 2123, 2008

[124] Cabañero A.I., Madrid Y., Cámara C., Mercury–Selenium species ratio in representative fish samples and their bioaccessibility by an in vitro digestion method, *Biol. Trace. Elem. Res.* 119, 195, 2007

[125] Balarama Krishna M.V., Karunasagar D., Rao S.V., Arunachalam J., Preconcentration and speciation of inorganic and methyl mercury in waters using polyaniline and gold trap-CVAAS, *Talanta* 68, 329, 2005

[126] Issaroa N., Abi-Ghanem C., Bermonda A., Fractionation studies of mercury in soils and sediments: A review of the chemical reagents used for mercury extraction, *Anal. Chim. Acta* 631, 1, 2009

[127] Westöö G., Determination of methylmercury compounds in foodstuffs. I. Methylmercury compounds in fish, identification and determination, *Anal. Scand.* 20, 2131, 1966

[128] Desrosiers M., Planas D., Mucci A., Total mercury and methylmercury accumulation in periphyton of Boreal Shield Lakes: Influence of watershed physiographic characteristics, *Sci. Total Environ.* 355, 247, 2006





[129] Chung S.W., Chan B.T., A reliable method to determine methylmercury and ethylmercury simultaneously in foods by gas chromatography with inductively coupled plasma mass spectrometry after enzymatic and acid digestion, *J. Chromatogr. A* 1218, 1260, 2011

[130] Foy G.P., Pacey G.E., Supercritical fluid extraction of mercury species, *Talanta* 61, 849, 2003

[131] Sun Y.C., Chung Y.T., Mierzwa J., Study of matrix influence on supercritical fluid extraction of polar mercury species from solid samples, *Analyst* 126, 1694, 2001

[132] Blanco R.M., Villanueva M.T., Sánchez Uría J.E., Sanz-Medel A., Field sampling, preconcentration and determination of mercury species in river waters, *Anal. Chim. Acta* 419, 137, 2000

[133] Beauchemin P., Siu K.W.M., Berman S.S., Determination of Organomercury in Biological Reference Materials by Inductively Coupled Plasma Mass Spectrometry using Flow Injection Analysis, *Anal. Chem.* 60, 2587, 1988

[134] Wagemann R., Trebacz E., Boila G., Lockhart W.L., Mercury species in the liver of ringed seals, *Sci. Total Environ.* 261, 21, 2000

[135] Mishra S., Tripathi R.M., Bhalke S., Shukla V.K., Puranik V.D., Determination of methylmercury and mercury (II) in a marine ecosystem using solid-phase microextraction gas chromatography-mass spectrometry, *Anal. Chim. Acta* 551, 192, 2005

[136] Muñoz J., Gallego M., Valcárcel M., Speciation of organometallic compounds in environmental samples by Gas Chromatography after Flow Preconcentration on Fullerenes and Nanotubes, *Anal. Chem.* 77, 5389, 2005

[137] Muñoz J., Gallego M., Valcárcel M., Solid-phase extraction–gas chromatography–mass spectrometry using a fullerene sorbent for the determination of inorganic mercury (II), methylmercury (I) and ethylmercury (I) in surface waters at sub-ng/ml levels, *J. Chromatogr. A* 1055, 185, 2004



[138] Sánchez D.M., Martín R., Morante R., Marín J., Munuera M.L., Preconcentration speciation method for mercury compounds in water samples using solid phase extraction followed by reversed phase high performance liquid chromatography, *Talanta* 52, 671, 2000

[139] Oda C.E., Ingle Jr. J.D., Speciation of Mercury by Cold Vapor Atomic Absorption Spectrometry with Selective Reduction, *Anal. Chem.* 53, 2305, 1981

[140] Rapsomanikis S., Derivatization by ethylation with sodium tetraethylborate for the speciation of metals and organometallics in environmental samples, *Analyst* 119, 1429, 1994

[141] Abalos M., Bayona J.M., Compano R., Granados M., Leal C., Prat M.D., Analytical procedures for the determination of organotin compounds in sediment and biota: a critical review, *J. Chromatogr. A* 788, 1, 1997

[142] Moens L., De Smaele T., Dams R., Sensitive, Simultaneous determination of Organomercury, -lead, and -tin compounds with Headspace Solid Phase Microextraction Capillary Gas Chromatography combined with Inductively Coupled Plasma Mass Spectrometry, *Anal. Chem.* 69, 1604, 1997

[143] Fischer R., Rapsomanikis S., Andreae M.O., Determination of Methylmercury in Fish Samples Using GC/AA and Sodium Tetraethylborate Derivatization, *Anal. Chem.* 65, 763, 1993

[144] Tseng C.M., De Diego A., Martin F.M., Amouroux D., Donard O.F.X., Rapid Determination of Inorganic Mercury and Methylmercury in Biological Reference Materials by Hydride Generation, Cryofocusing, Atomic Absorption Spectrometry After Open Focused Microwave-assisted Alkaline Digestion, *J. Anal. At. Spectrom.* 12, 743, 1997

[145] Cai Y., Monsalud S., Furton K.G., Determination of Methyl- and Ethylmercury compounds using Gas Chromatography Atomic Fluorescence Spectrometry following aqueous derivatization with Sodium Tetraphenylborate, *Chromatographia* 52, 82, 2000



[146] Grinberg P., Campos R.C., Mester Z., Sturgeon R.E., A comparison of alkyl derivatization methods for speciation of mercury based on solid phase microextraction gas chromatography with furnace atomization plasma emission spectrometry detection, *J. Anal. At. Spectrom.* 18, 902, 2003

[147] Gibičar D., Logar M., Horvat N., Marn-Pernat A., Ponikvar R., Horvat M., Simultaneous determination of trace levels of ethylmercury and methylmercury in biological samples and vaccines using sodium tetra(n-propyl)borate as derivatizing agent, *Anal. Bioanal. Chem.* 388, 329, 2007

[148] Ikonomou M.G., Fernandez M.P., He T., Cullon D., Gas chromatography–high-resolution mass spectrometry based method for the simultaneous determination of nine organotin compounds in water, sediment and tissue, *J. Chromatogr. A* 975, 319, 2002

[149] Minganti V., Capelli R., De Pellegrini R., Evaluation of different derivatization methods for the multi-element detection of Hg, Pb and Sn compounds by gas chromatography-microwave induced plasma-atomic emission spectrometry in environmental samples, *Fresenius J. Anal. Chem.* 351, 471, 1995

[150] Li Y., Yan X.P., Dong L.M., Wang S.W., Jiang Y., Jiang D.Q., Development of an ambient temperature post-column oxidation system for high-performance liquid chromatography on-line coupled with cold vapor atomic fluorescence spectrometry for mercury speciation in seafood, *J. Anal. At. Spectrom.* 20, 467, 2005

[151] Hintelmann H., Hempel M., Wilken R.D., Observation of unusual organic Mercury species in soils and sediments of industrially contaminated sites, *Environ. Sci. Technol.* 29, 1845, 1995

[152] Cai Y., Jaffé R., Jones R., Ethylmercury in the soils and sediments of the Florida Everglades, *Environ. Sci. Technol.* 31, 302, 1997



[153] Falter R., Ilgen G., Coupling of the RP C18 preconcentration HPLC-UV-PCO-system with atomic fluorescence detection for the determination of methylmercury in sediment and biological tissue, *Fresenius J. Anal. Chem.* 358, 407, 1997

[154] Falter R., Scholer H.F., A new pyrrolidinedithiocarbamate screening method for the determination of methylmercury and inorganic mercury relation in hair samples by HPLC-UV-PCO-CVAAS, *Fresenius J. Anal. Chem.* 354, 492, 1996

[155] Xia L., Hu B., Wu Y., Hollow fiber-based liquid-liquid-liquid microextraction combined with high-performance liquid chromatography for the speciation of organomercury, *J. Chromatogr. A* 1173, 44, 2007

[156] Hempel M., Hintelman H., Wilken R.D., Determination of organic mercury species in soils by high-performance liquid chromatography with ultraviolet detection, *Analyst* 117, 669, 1992

[157] Palmieri H. E. L., Leonel L.V., Determination of methylmercury in fish tissue by gas chromatography with microwave-induced plasma atomic emission spectrometry after derivatization with sodium tetraphenylborate, *Fresenius J. Anal. Chem.* 366, 466, 2000

[158] Lansens P., Baeyens W., Improvement of the semiautomated headspace analysis method for the determination of methylmercury in biological samples, *Anal. Chim. Acta.* 228, 93, 1990

[159] Lansens P., Meuleman C., Leermakers M., Baeyens W., Determination of methylmercury in natural waters by headspace gas chromatography with microwave – induced plasma detection after preconcentration on a resin containing dithiocarbamate groups, *Anal. Chim. Acta.* 243, 417, 1990

[160] Johansson M., Emteborg H., Glad B., Reinholdsson F., Baxter D.C., Preliminary appraisal of a novel sampling and storage technique for the speciation analysis of lead and mercury in seawater, *Fresenius J. Anal. Chem.* 351, 461, 1995

[161] Berzas Nevado J.J., Rodríguez Martín-Doimeadios R.C., Guzmán Bernardo F.J.,

Jiménez Moreno M., Determination of monomethylmercury in low- and high-polluted sediments by microwave extraction and gas chromatography with atomic fluorescence detection, *Anal. Chim. Acta.* 608, 30, 2008

[162] Cai Y., Jaffé R., Alli A., Jones R.D., Determination of organomercury compounds in aqueous samples by capillary gas chromatography-atomic fluorescence spectrometry following solid-phase extraction, *Anal. Chim. Acta.* 334, 251, 1996

[163] Zoorob G.K. , McKiernan J.W., Caruso J.A., ICP-MS for Elemental Speciation Studies, *Mikrochim. Acta* 128, 145, 1998

[164] Falter R., Schöler H.F., Determination of mercury species in natural waters at picogram level with on-line RP C18 preconcentration and HPLC-UV-PCO-CVAAS, *Fresenius J. Anal. Chem.* 353, 34, 1995

[165] Armstrong L.H.E., Corns W.T., Stockwell P.B., O' Connor G., Ebdon L., Evans E.H., Comparison of AFS and ICP-MS detection coupled with gas chromatography for the determination of methylmercury in marine samples, *Anal. Chim. Acta.* 390, 245, 1999

[166] Yin X.B., Dual-cloud point extraction as a preconcentration and clean-up technique for capillary electrophoresis speciation analysis of mercury, *J. Chromatogr. A* 1154, 437, 2007

[167] Silva da Rocha M., Soldado A.B., Blanco-González E., Sanz-Medel A., Speciation of mercury compounds by capillary electrophoresis coupled on-line with quadrupole and double-focusing inductively coupled plasma mass spectrometry, *J. Anal. At. Spectrom.* 15, 513, 2000

[168] Fan Z., Liu X., Determination of methylmercury and phenylmercury in water samples by liquid-liquid-liquid microextraction coupled with capillary electrophoresis, *J. Chromatogr. A* 1180, 187, 2008

[169] Tu Q., Qvarnström J., Frech W., Determination of mercury species by capillary zone electrophoresis-inductively coupled plasma mass spectrometry: a comparison of two spray chamber-nebulizer combinations, *Analyst* 125, 705, 2000



[170] Lee T.H., Jiang S.J., Determination of mercury compounds by capillary electrophoresis inductively coupled plasma mass spectrometry with microconcentric nebulization, *Anal. Chim. Acta.* 413, 197, 2000

[171] Yan X.P., Yin X.B., Jiang D.Q., He X.W., Speciation of Mercury by hydrostatically modified electroosmotic Flow Capillary Electrophoresis coupled with Volatile Species Generation Atomic Fluorescence Spectrometry, *Anal. Chem.* 75, 1726, 2003

[172] Li Y., Jiang Y., Yan X.P., On-line hyphenation of capillary electrophoresis with flame-heated furnace atomic absorption spectrometry for trace mercury speciation, *Electrophoresis* 26, 661, 2005

[173] Mushak P., Gas-Liquid Chromatography in the analysis of Mercury (II) compounds, *Environ. Health Perspect.* 55, 1973

[174] Bramanti E., Lomonte C., Onor M., Zamboni R., D'Ulivo A., Raspi G., Mercury speciation by liquid chromatography coupled with on-line chemical vapour generation and atomic fluorescence spectrometric detection (LC-CVGAFS), *Talanta* 66, 762, 2005







[175] Liang L.N., Jiang G.B., Liu J.F., Hu J.T., Speciation analysis of mercury in seafood by using high-performance liquid chromatography on-line coupled with cold-vapor atomic fluorescence spectrometry via a post column microwave digestion, *Anal. Chim. Acta* 477, 131, 2003

[176] Vallant B., Kadnar R., Goessler W., Development of a new HPLC method for the determination of inorganic and methylmercury in biological samples with ICP-MS detection, *J. Anal. At. Spectrom.* 22, 322, 2007

[177] Falter R., Ilgen G., Determination of trace amounts of methylmercury in sediment and biological tissue by using water vapor distillation in combination with RP C18 preconcentration and HPLC-HPF/HHPN-ICP-MS, *Fresenius J. Anal. Chem.* 358, 401, 1997



**Table 1:** The main organic forms of mercury.

| NAME                        | EMPIRICAL FORMULA   | STRUCTURAL FORMULA   | APPLICATION  | REFERENCES   |
|-----------------------------|---|--|--|--------------|
| METHYL-MERCURY CATION       | (CH <sub>3</sub> Hg <sup>+</sup> )  | H <sub>3</sub> C—Hg <sup>+</sup> X <sup>-</sup>                                      | -  |              |
| ETHYLMERCURY CATION         | (C <sub>2</sub> H <sub>5</sub> Hg <sup>+</sup> )                                |     | Used in industry, a metabolite of thimerosal.  |              |
| PHENYL-MERCURY CATION       | (C <sub>6</sub> H <sub>5</sub> Hg <sup>+</sup> )                                |     | Used in industry. Phenylmercury acetate is used as a fungicide and in paints.  | [7, 10, 11]  |
| DIMETHYL-MERCURY            | C <sub>2</sub> H <sub>6</sub> Hg  |    | Used in toxicology as a reference toxin and to calibrate NMR instruments during Hg determination                                 |              |
| DIETHYL-MERCURY             | C <sub>4</sub> H <sub>10</sub> Hg   |   | -  |              |
| MERBROMIN, (MERCURO-CHROME) | C <sub>20</sub> H <sub>8</sub> Br <sub>2</sub> HgNa <sub>2</sub> O <sub>6</sub> |  | Used as a disinfectant because of its antibacterial properties and as an analytical reagent.                                     | [12]         |
| THIMEROSAL                  | C <sub>9</sub> H <sub>9</sub> HgNaO <sub>2</sub> S                              |   | Used as a preservative in vaccines, some ointments and other forms of drugs; sometimes as an antifungal drug and in dermatology. | [13, 14, 15] |

**Table 2.** The deleterious effects of exposure to organomercury compounds.

| COMPOUND        | SITE OF ACTION         | TOXIC ACTION   | REFERENCE(S)    |
|-----------------|------------------------|--|-----------------|
| Methylmercury   | Central nervous system | Slurred speech, hypersalivation, shouting, dysphagia, scotoma, neurasthenia, loss of libido, depression, hallucinations, focal cramps, chorea, athetosis, myoclonus, paralysis, stupor, coma, and death.   | [8]             |
| Methylmercury   | Fetus                  | Severe brain damage, profound mental retardation, spasticity, seizures, cerebral palsy, chorea, athetotic movements, ataxia, tremors, cataracts, hearing deficiency, small size, anemia, and renal dysfunction.  | [8, 15, 76, 77] |
| Dimethylmercury | The whole body         | Irritation of the eyes, respiratory tract and skin numbness and tingling of the mouth, hands and legs, joint pains, narrowing of the field of vision, emotional disturbances, lack of coordination, slurred speech, deafness, death.<br>Also crosses biological barriers – toxic action like that of MeHg. | [78]            |

**Table 3.** Information on important events associated with poisoning by mercury and its compounds.

| PERIOD      | DESCRIPTION OF EVENT   | REFERENCE(S)       |
|-------------|--|--------------------|
| 1865        | The deaths of two chemists, who used dimethylmercury during investigations to define the valences of a number of metals.   | [15]               |
| 1940        | The inhalation of methylmercury by four workers at a factory producing fungicides for protecting grain crops. In one of the poisoned men cerebral atrophy with cortical loss was found; since then, this has been known as the Hunter-Russell syndrome.  | [15, 79, 80]       |
| 1920 - 1960 | In a Japanese chemical works mercury was used as a catalyst in the production of acetaldehyde and vinyl chloride. The effluents of this process, containing methylmercury chloride, were discharged into Minamata Bay, on the south-western coast of Kyushu - nearly 150 tons of methylmercury during forty years. Through the consumption of fish and other frutti di mare, harmful methylmercury was ingested by humans, causing damage to the central nervous system, and the | [8, 15, 33, 79-81] |



|                    |   |                  |
|--------------------|---|------------------|
|                    | children of women who had consumed the poisoned fish were afflicted by mental retardation, developmental disruption, hepatic diseases, hypertension, retarded metabolism and even death (chorea, ataxia, tremors and seizures). These symptoms were given the name “Minamata disease” by doctors at the Kumamoto University Hospital.   |                  |
| <b>1965</b>        | At Niigata in Japan industrial effluents containing mercury were discharged into the River Agano. As in Minamata Bay, the water was polluted, and as a consequence, the organisms at successive levels in the aquatic trophic pyramid, and ultimately humans.   | [8, 79]          |
| <b>1959 - 1972</b> | More than 6000 persons in Iraq were hospitalized as a result of methylmercury poisoning. MeHg got into their diet via bread produced from grain that had been sprayed with fungicides containing MeHg. A study of a group of Iraqi children exposed to MeHg before birth, like those at Minamata, exhibited developmental anomalies. Prenatal exposure to MeHg also inhibited neuronal migration. | [15, 50, 82, 83] |
| <b>1997</b>        | The death of Karen Wetterhahn, a world-famous scientist specializing in toxic metal exposure. Death ensued just a few months after a one-off exposure to less than 1 ml of dimethylmercury, which had spilt onto her hand, covered with a latex glove.  | [78]             |
| <b>At present</b>  | In Brazilian Amazonia, the inhabitants of fishing villages (consuming fish poisoned by MeHg) situated near gold mines suffered mild neurological symptoms characteristic of MeHg poisoning, i.e. sensory disorders, tremors and balance disorders.  | [84]             |

**Table 4.** Literature data on the content of organic mercury and methylmercury in environmental samples from different parts of the world.



| Type of sample  | Geographical region                 | Analytical procedure | Kingdom | Phylum | Class | Family | Species | Mean concentration of compound [mg/kg d.w.] * |         | MeHg/THg [%] ** | Reference |
|-----------------|-------------------------------------|----------------------|---------|--------|-------|--------|---------|---|---------|-----------------|-----------|
|                 |                                     |                      |         |        |       |        |         | OHg   | MeHg    |                 |           |
| Bottom sediment | Beni River, Bolivia - Amazonia      | GC - CV- AFS         |         |        |       |        |         | -   | 0.4-1.2 | -               | [87]      |
| River sediment  | Loučka River, Czech Republic        | HPLC - CV-AFS        | -----   | -----  | ----- | -----  | -----   | -   | 0.022   | 18.5            | [20]      |
|                 | Bečva River, Czech Republic         |                      |         |        |       |        |         | -   | 0.031   | 22.3            |           |
|                 | Jihlava River, Czech Republic       |                      |         |        |       |        |         | -   | 0.046   | 32              |           |
|                 | Haihe River, Tianjin, China         | HPLC – CV - AFS      |         |        |       |        |         | -   | 1       | -               | [88]      |
|                 | Dagu Drainage Canal, Tianjin, China |                      |         |        |       |        |         | -   | 21.7    | -               |           |
| Water           | Tapaj'os River, Brazil              | GC-CV - AFS          | -       | -      | -     | -      | -       | 0.02  | 1.8     | [89]            |           |
|                 | Piedras River, SW Spain             | HPLC–CV/HG–mAFS–AFS  | -       | -      | -     | -      | -       | 49.5  | -       | [90]            |           |
|                 | San Pedro River, SW Spain           |                      | -       | 50.4   | -     |        |         |   |         |                 |           |
|                 | Guadalete River, SW Spain           |                      | -       | 51.2   | -     |        |         |   |         |                 |           |
|                 | Carreras River, SW Spain            |                      | -       | 52     | -     |        |         |   |         |                 |           |



| Type of sample  | Geographical region             | Analytical procedure | Kingdom          | Phylum     | Class        | Family        | Species   | Mean concentration of compound [mg/kg d.w.] |               | MeHg/TH g [%] | Reference |
|-----------------|---------------------------------|----------------------|------------------|------------|--------------|---------------|---|---|---------------|---------------|-----------|
|                 |                                 |                      |                  |            |              |               |   | OHg   | MeHg          |               |           |
| Whole organisms | Lake Gaobeidian, Beijing, China | HPLC-UV-AFS          | Plants           | -          | Chlorophytes | Zygnemataceae | <i>Spirogyra</i>  | -   | <0.05         | -             | [10]      |
| Whole organisms | Tapaj'os River, Brazil          | CV-AFS               |                  |            |              |               | -   | -   | Phytoplankton | -             | 0.01      |
| Whole organisms | Guanabara Bay, Brazil           | GC - ECD             | Plants / Animals | -          | -            | -             | Microplankton   | -   | 0.0089        | 33.8          | [91]      |
|                 |                                 |                      |                  |            |              |               | Mesoplankton  | -   | 0.0359        | 75.4          |           |
| Whole organisms | Lake Superior, USA              | GC – CV - AAS        | Animals          | -          | -            | -             | Zooplankton ( <i>Mysis relicta</i> )  | -   | 0.033-0.054   | -             | [92]      |
| Whole organisms | Lake Superior, USA              | GC-CV - AAS          |                  |            |              |               | Zooplankton   | -   | 0.035 - 0.050 | -             |           |
| Whole organisms | Lake Gaobeidian, Beijing, China | HPLC-UV-AFS          |                  |            |              |               | Zooplankton ( <i>Monia rectirostris</i> , <i>Monia micrura</i> , <i>Monia macrocopa</i> ) | -   | <0.05         | -             | [10]      |
| Whole organisms | Tapaj'os River, Brazil          | CV-AFS               |                  |            |              |               | Zooplankton   | -   | 0.073         | 48.25         | [92]      |
| Whole organisms | Bečva River, Czech Republic     | HPLC - CV-AFS        |                  |            |              |               | Zoobenthos  | -   | 0.082         | 33.7          | [20]      |
|                 | Loučka River, Czech Republic    |                      |                  |            |              |               |   | -   | 0.158         | 43.6          |           |
|                 | Jihlava River, Czech Republic   |                      | -                | 0.168      | 66           |               |   |   |               |               |           |
| Whole organisms | Lake Superior, USA              | GC – CV - AAS        | Animals          | Arthropods | Insects      | Chironomidae  | Chironomid larvae   | -   | 0.008         | -             | [92]      |



|                               |   |               |         |             |  |              |   |  |                      |  |      |       |    |      |
|-------------------------------|---|---------------|---------|-------------|--|--------------|---|--|----------------------|--|------|-------|----|------|
| Muscles                       | Kagoshima Bay, Japan                      | CV - AAS      | Animals | Annelids    | Polychaetes                            | Siboglinidae | <i>Lamellibrachia satsuma</i>           | 0.014                                      | -                    | -  | [41] |       |    |      |
| Whole organism                | Coast of Novaya Zemlya, Arctic Ocean      | -             |         | Echinoderms | Starfish                               | -            | -                                       | <i>(Asteroidea)</i>                        | -                    | 0.008  | 2.4  | [93]  |    |      |
|                               |   |               |         |             |  |              |   | <i>Maldanes sarsi</i>                      | -                    | 0.16   | 48.6 |       |    |      |
| Muscles                       | Malaysia                                  | INAA          |         | Mollusks    | Cephalopods                            | Loliginidae  | -                                       | <i>Loligo sp.</i>                          | 0.22                 | -  | 64.4 | [92]  |    |      |
| Muscles                       | USA                                       | -             |         |             |  |              |   | Mussels                                    | Mytilidae /Ostreidae | Pacific oyster ( <i>Crassostrea gigas</i> ), Edible mussel ( <i>Mytilus edulis</i> ) | -    | 0.056 | 66 | [95] |
| Soft tissues                  | Dunkirk and Calais, north coast of France | GC - CV - AFS |         |             |  |              |   |  |                      |  | -    | 0.073 | 37 |      |
|                               | Toulon, southeast coast of France         |               |         |             |  |              |   | -  | 0.094                | 52   |      |       |    |      |
|                               | Basque Region, southwest coast of France  |               |         |             | -                                      | 0.113        | 74                                      |  |                      |  |      |       |    |      |
| Lorient, west coast of France | -   |               |         |             | 0.11-0.75                              | -            | [96]                                    |  |                      |  |      |       |    |      |
| Body integument               | Coast of Portugal                         | GC - AFS      |         |             | Cephalopods                            | Octopoidae   |   | Common octopus ( <i>Octopus vulgaris</i> ) | -                    | 0.18-5.0   | -    |       |    |      |
| Digestive tract               |   |               | -       |             |  |              |   |  |                      |  |      |       |    |      |
| Soft tissues                  | Terra Nova Bay, Antarctica                | AAS           | Mussels | Pectinidae  | Mussels ( <i>Adamussium colbecki</i> ) | -            | 0.295                                   | 49.55                                      | [97]                 |  |      |       |    |      |
| Soft tissues                  | Malaysia                                  | INAA          |         |             |  | Aricidae     | Blood cockle ( <i>Anadara granosa</i> ) | 0.32                                       | -                    | 75.5   | [98] |       |    |      |



|  |                                  |             |         |            |              |                       |   |       |       |       |       |
|--|----------------------------------|-------------|---------|------------|--------------|-----------------------|---|-------|-------|-------|-------|
| Edible part                                    | Mouth of the Krka River, Croatia | CV - AAS    |         |            |              | Mytilidae / Ostreidae | Mediterranean mussel ( <i>Mytilus galloprovincialis</i> ) | -     | 5.925 | 32.5  | [99]  |
| Muscles  | USA                              | -           |         |            |              | Lithodidae            | King crab ( <i>Paralithodes camtschaticus</i> )           | -     | 0.01  | 50    | [94]  |
| Whole organism                                 | Lake Superior, USA               | GC-CV - AAS |         |            |              | -                     | <i>Amphipoda</i>  | -     | 0.032 | -     | [92]  |
| Muscles  | Malaysia                         | INAA        |         |            |              | Penaeidae             | Shrimps ( <i>Penaeus sp.</i> )                            | 0.23  | -     | 62.6  | [98]  |
| Whole organism                                 | Ligurian Sea                     | GF - AAS    | Animals | Arthropods | Malacostraca | Euphausiidae          | Northern krill ( <i>Meganyctiphanes norvegica</i> )       | 0.24  | -     | 71.3  | [100] |
|  |                                  |             |         |            |              | Benthescymidae        | ( <i>Gennadas elegans</i> )                               | 0.252 | -     | 70.9  |       |
|  |                                  |             |         |            |              | Pasiphaeidae          | Shrimp ( <i>Pasiphaea sivado</i> )                        | 0.81  | -     | 84.6  |       |
| Pink shrimps ( <i>Pasiphaea multidentata</i> ) | 2.68                             | -           | 88.22   |            |              |                       |   |       |       |       |       |
| Muscles  |                                  |             |         |            |              |                       | Red shrimps ( <i>Aristeus antennatus</i> )                | 2.933 | -     | 85.28 |       |



|         |   |          |         |             |                |              |  |   |        |      |       |
|---------|---|----------|---------|-------------|----------------|--------------|--|---|--------|------|-------|
| Muscles | Mediterranean Sea                         | GC- ECD  | Animals | Vertebrates | Chondrichthyes | Torpedinidae | Atlantic torpedo( <i>Torpedo nobiliana</i> ) | - | 1.90   | 81   | [101] |
|         |   |          |         |             |                | Chimaeridae  | Rabbit fish ( <i>Chimaera monstrosa</i> )    | - | 2.67   | 83.6 |       |
| Muscles | Nyumba' ya Mungu Nature Reserve, Tanzania | GC - ECD | Animals | Vertebrates | Actinopterygii | Cichlidae    | <i>Tilapia (Tilapia urolepis)</i>            | - | 0.0051 | 83.6 | [102] |
|         |   |          |         |             |                | Clariidae    | <i>(Clarias mossambicus)</i>                 | - | 0.007  | 77.8 |       |
|         |   |          |         |             |                | Mochokidae   | <i>(Synodontis maculipinna)</i>              | - | 0.0078 | 72.9 |       |
| Muscles | Kidatu Nature Reserve, Tanzania           |          |         |             |                | Cichlidae    | <i>Tilapia (Tilapia urolepis)</i>            | - | 0.0086 | 57   |       |

|         |   |          |   |             |                |            |   |           |                                     |                |            |                          |         |             |                |            |
|---------|---|----------|---|-------------|----------------|------------|---|-----------|-------------------------------------|----------------|------------|--------------------------|---------|-------------|----------------|------------|
| Muscles | Hyogo Prefecture, Japan                   | GC       | Animals                                     | Vertebrates | Actinopterygii | Carangidae | Japanese horse mackerel<br>( <i>Trachurus japonicus</i> ) | -         | 0.01                                | 62             | [94]       |                          |         |             |                |            |
| Muscles | Oita Prefecture, Japan                    |          |   |             |                |            |   | -         | 0.02                                | 51             |            |                          |         |             |                |            |
| Muscles | Kagoshima Prefecture, Japan               |          |   |             |                |            |   | -         | 0.02                                | 74             |            |                          |         |             |                |            |
| Muscles | Mie Prefecture, Japan                     |          |   |             |                |            |   | -         | 0.02                                | 50             |            |                          |         |             |                |            |
| Muscles | USA                                       | GC       | Animals                                     | Vertebrates | Actinopterygii | Salmonidae | Sockeye salmon<br>( <i>Oncorhynchus nerka</i> )           | -         | 0.02                                | 67             | [94]       |                          |         |             |                |            |
| Muscles | Kidatu Nature Reserve, Tanzania           | GC - ECD |   |             |                |            |   | Animals   | Vertebrates                         | Actinopterygii | Mochokidae | (Synodontis maculipinna) | -       | 0.0258      | 63.7           | [102]      |
| Muscles | Philippines                               | GLC      |   |             |                |            |   |           |                                     |                |            |                          | Animals | Vertebrates | Actinopterygii | Scombridae |
| Muscles | Kiribati, Pacific Ocean                   |          | -   | 0.03        | 75             |            |   |           |                                     |                |            |                          |         |             |                |            |
| Muscles | Marshall Islands                          |          | Yellowfin tuna ( <i>Thunnus albacares</i> ) | -           | 0.03           | 75         |   |           |                                     |                |            |                          |         |             |                |            |
| Muscles | Hale and Pangani Nature Reserve, Tanzania | GC - ECD | Animals                                     | Vertebrates | Actinopterygii | Clariidae  | (Clarias mossambicus)                                     | -         | 0.0335                              | 100            | [102]      |                          |         |             |                |            |
| Muscles | Mtera Nature Reserve, Tanzania            |          |   |             |                |            |   | Cichlidae | Tilapia ( <i>Tilapia urolepis</i> ) | -              |            | 0.0347                   | 87.45   |             |                |            |



|         |   |          |         |             |                |             |   |   |         |      |       |
|---------|---|----------|---------|-------------|----------------|-------------|---|---|---------|------|-------|
| Muscles | Kidatu Nature Reserve, Tanzania           |          |         |             |                | Claroteidae | <i>(Bagrus orientalis)</i>                  | - | 0.040   | 92.8 |       |
| Muscles | Guanabara Bay, Brazil                     |          |         |             |                | Mugilidae   | <i>(Mugil liza)</i>                         | - | 0.0415  | 59.9 | [91]  |
| Muscles |   |          |         |             |                | Clupeidae   | <i>(Sardinella brasiliensis)</i>            | - | 0.0491  | 59.6 |       |
| Muscles | Lake Ya-Er, China                         | HPLC-AFS |         |             |                | Cyprinidae  | Crucian carp ( <i>Carassius carassius</i> ) | - | 0.05233 | -    | [103] |
| Muscles | Hale and Pangani Nature Reserve, Tanzania | GC – ECD | Animals | Vertebrates | Actinopterygii | Mochokidae  | <i>(Synodontis maculipinna)</i>             | - | 0.0526  | 83.4 | [102] |
| Muscles | Mtera Nature Reserve, Tanzania            |          |         |             |                | Claroteidae | <i>(Bagrus orientalis)</i>                  | - | 0.053   | 90.4 |       |
| Muscles |   |          |         |             |                | Alestidae   | <i>(Brycinus affinis)</i>                   | - | 0.056   | 95.6 |       |





|                   |                                 |          |         |             |                |  |   |            |  |       |         |   |       |
|-------------------|---------------------------------|----------|---------|-------------|----------------|--|---|------------|--|-------|---------|---|-------|
| Muscles           | Mtera Nature Reserve, Tanzania  |          |         |             |                |  |   | -          | 0.08555  | 68.85 |         |   |       |
| Muscles           | Kidatu Nature Reserve, Tanzania |          |         |             |                | Characidae   | Tiger Fish ( <i>Hydrocynus vittatus</i> )         | -          | 0.116  | 97.5  | [102]   |   |       |
|                   | Lake Ya-Er, China               | HPLC-AFS |         |             |                | Channidae  | ( <i>Ophiocephalus argus cantor</i> )             | -          | 0.16438  | -     | [103]   |   |       |
|                   | Perlis, Kedah, Malaysia         | GC       | Animals | Vertebrates | Actinopterygii | Scombridae   | Short mackerel ( <i>Rastrelliger brachysoma</i> ) | -          | 0.179  | 76    | [27]    |   |       |
|                   |                                 |          |         |             |                |  | Longtail tuna ( <i>Thunnus tonggol</i> )          | -          | 0.187  | 81    |         |   |       |
|                   | Indian Ocean                    | GLC      |         |             |                |  | Southern bluefin tuna ( <i>Thunnus maccoyii</i> ) | -          | 0.19   | 71    | [94]    |   |       |
|                   | Atlantic Ocean                  |          |         |             |                |  | Bigeye tuna ( <i>Thunnus obesus</i> )             | -          | 0.19   | 71    |         |   |       |
|                   | Australia                       |          |         |             |                | Southern bluefin tuna (farmed) ( <i>Thunnus maccoyii</i> ) | -   | 0.19       | 64   |       |         |   |       |
| Lake Ya-Er, China | HPLC-AFS                        |          |         |             |                |  |   | Cyprinidae | Silver carp ( <i>Hypophthalmichthys molitrix</i> ) | -     | 0.19515 | - | [103] |



|         |                             |          |         |             |                |            |   |  |        |       |      |      |
|---------|-----------------------------|----------|---------|-------------|----------------|------------|---|--|--------|-------|------|------|
|         | Kuantan, Pahang<br>Malaysia | GC       |         |             |                | Scombridae | Short<br>mackerel( <i>Rastrelliger<br/>brachysoma</i> ) | -  | 0.219  | 70    | [33] |      |
| Liver   | Perlis, Kedah, Malaysia     |          |         |             |                |            | Longtail tuna( <i>Thunnus<br/>tonggol</i> )             | -  | 0.236  | 45    | [27] |      |
| Muscles | Atlantic Ocean              |          |         |             |                |            | Istiophoridae   | Blue marlin ( <i>Makaira<br/>nigricans</i> )         | -      | 0.24  | 43   | [94] |
| Liver   | Perlis, Kedah, Malaysia     |          |         |             |                |            | Scombridae  | Short mackerel<br>( <i>Rastrelliger brachysoma</i> ) | -      | 0.242 | 41   | [27] |
| Muscles | Terra Nova Bay, Antarctica  | AAS      | Animals | Vertebrates | Actinopterygii | Bathypoda  | Ploughfish ( <i>Gymnodraco<br/>acuticeps</i> )          | -  | 0.2543 | 65.45 | [97] |      |
|         | Atlantic Ocean              | GLC      |         |             |                | Scombridae | Atlantic bluefin tuna<br>( <i>Thunnus thynnus</i> )     | -  | 0.29   | 70    | [94] |      |
|         | Guanabara Bay, Brazil       | GC – ECD |         |             |                | Ariidae    | ( <i>Bagre bagre</i> )                                  | -  | 0.2973 | 97.9  | [91] |      |

|         |                            |          |               |   |                |                 |  |      |        |      |      |
|---------|----------------------------|----------|---------------|---|----------------|-----------------|--|------|--------|------|------|
| Muscles | Terra Nova Bay, Antarctica | AAS      | Animals       | Vertebrates                                       | Actinopterygii | Channichthyidae | <i>(Chionodraco hamatus)</i>                             | -    | 0.307  | 81.9 | [97] |
|         | Indonesia                  | GLC      |               |   |                | Scombridae      | Bigeye tuna ( <i>Thunnus obesus</i> )                    | -    | 0.31   | 66   | [94] |
|         | Chile                      | GC       |               |   |                | Nothotenidae    | Patagonian toothfish ( <i>Dissostichus eleginoides</i> ) | -    | 0.31   | 54   |      |
|         | Guanabara Bay, Brazil      | GC – ECD |               |   |                | Haemulidae      | <i>(Orthopristis ruber)</i>                              | -    | 0.3388 | 95.3 | [91] |
|         | Atlantic Ocean             | GC       |               |   |                | Xiphiidae       | Swordfish ( <i>Xiphias gladius</i> )                     | -    | 0.34   | 72   | [94] |
| Liver   | Kuantan, Pahang Malaysia   |          | Scombridae    | Short mackerel ( <i>Rastrelliger brachysoma</i> ) | -              | 0.388           | 40   | [27] |        |      |      |
| Muscles | Atlantic Ocean             |          | Istiophoridae | Striped marlin ( <i>Tetrapturus audax</i> )       | -              | 0.39            | 76   | [94] |        |      |      |





|   |                                       |          |         |             |                |                  |   |            |  |      |      |
|---|---------------------------------------|----------|---------|-------------|----------------|------------------|---|------------|--|------|------|
| Muscles   | Guanabara Bay, Brazil                 | GC - ECD |         |             |                | Trichiuridae     | Largehead hairtail<br>( <i>Trichiurus lepturus</i> )      | -          | 0.678  | 99.2 | [91] |
|   | Pacific Ocean                         | GLC      |         |             |                | Scombridae       | Bigeye tuna ( <i>Thunnus obesus</i> )                     | -          | 0.69   | 71   | [94] |
|   | Chendring, Terengganu, Malaysia       | GC       |         |             |                |                  | Longtail tuna ( <i>Thunnus tonggol</i> )                  | -          | 0.708  | 76   | [27] |
|   | Italy                                 | GLC      |         |             |                |                  | Atlantic bluefin tuna (farmed) ( <i>Thunnus thynnus</i> ) | -          | 1.02   | 70   | [94] |
| Liver   | Shizuoka and Chiba Prefectures, Japan | GC       | Animals | Vertebrates | Actinopterygii | Berycidae        | Splendid alfonsino ( <i>Beryx splendens</i> )             | -          | 1.12   | 32   | [94] |
| Muscles   | Terra Nova Bay, Antarctica            | AAS      |         |             |                | Nototheniidae    | ( <i>Trematomus pennelli</i> )                            | -          | 1.356  | 45.3 | [97] |
|   |                                       |          |         |             |                | Bathypodaconidae | Mawson's dragonfish<br>( <i>Cygnodraco mawsoni</i> )      | -          | 6.702  | 73.9 |      |
|   |                                       |          |         |             |                | Malaysia         | INAA  | Scombridae | Indian mackerel<br>( <i>Rastrelliger kanagurta</i> ) | 0.21 | -    |
| Narrow-barred Spanish mackerel ( <i>Scomberomorus</i> ) | 0.31                                  | -        | 77      |             |                |                  |   |            |  |      |      |



|              |                                     |             |         |             |            |              |  |   |       |   |  |     |     |       |
|--------------|-------------------------------------|-------------|---------|-------------|------------|--------------|--|---|-------|---|--|-----|-----|-------|
|              |                                     |             |         |             |            |              | <i>commersoni</i> )                        |   |       |   |  |     |     |       |
| Fatty tissue | Bay of California                   | CV - AFS    |         |             | Sauropsida | Cheloniidae  | Green sea turtle ( <i>Chelonia mydas</i> ) | - | 0.001 | 17  | [104]  |     |     |       |
| Muscles      |                                     |             |         |             |            |              |  | - | 0.006 | 22  |  |     |     |       |
| Kidneys      |                                     |             |         |             |            |              |  | - | 0.019 | 23  |  |     |     |       |
| Liver        |                                     |             |         |             |            |              |  | - | 0.027 | 19  |  |     |     |       |
| Muscles      |                                     |             |         |             |            |              |  | - | 0.17  | 100   |  |     |     |       |
| Kidneys      |                                     |             |         |             |            |              |  | - | 0.208 | 56  |  |     |     |       |
| Liver        |                                     |             |         |             |            |              |  | - | 0.338 | 41  |  |     |     |       |
| Muscles      | Southeastern Bay of California, USA | CV - AAS    | Animals | Vertebrates | Birds      | Anatidae     | Northern Shoveler ( <i>Anas clypeata</i> ) | - | 0.47  | 100   | [105]  |     |     |       |
| Intestines   | Czech Republic                      | HPLC/CV-AFS |         |             |            | Accipitridae | Common buzzard ( <i>Buteo buteo</i> )      | - | 0.57  | 71.3  | [7]  |     |     |       |
| Muscles      | Chaun River, Siberia, Russia        |             |         |             |            | Laridae      | Herring Gull ( <i>Larus argentatus</i> )   | - | 0.6   | -   | [106]  |     |     |       |
|              | Southeastern Bay of California, USA | CV - AAS    |         |             |            | Anatidae     | Blue-winged Teal ( <i>Anas discors</i> )   | - | 0.77  | 26.1  | [105]  |     |     |       |
| Feathers     | Chaun River, Siberia, Russia        | GC          |         |             |            |              |  |   |       | Long-tailed Duck ( <i>Clangula hyemalis</i> ) | -  | 0.9 | -   | [106] |
| Muscles      | Southern Indian Ocean               |             |         |             |            |              |  |   |       | Procellariidae                                | White-chinned Petrel ( <i>Procellaria aequinoctialis</i> ) | -   | 0.9 |       |



|            |                                     |             |  |         |             |                   |  |  |      |      |       |
|------------|-------------------------------------|-------------|--|---------|-------------|-------------------|--|--|------|------|-------|
| Muscles    | Northern Pacific Ocean              | GC          |  |         |             | Procellariidae    | Northern Fulmar<br>( <i>Fulmarus glacialis</i> )           | -  | 0.9  | -    | [106] |
| Intestines | Czech Republic                      | HPLC/CV-AFS |  |         |             | Phalacrocoracidae | Great Cormorant<br>( <i>Phalacrocorax carbo</i> )          | -  | 0.99 | 82.1 | [7]   |
|            |                                     |             |  |         |             |                   | Accipitridae   | Common buzzard ( <i>Buteo buteo</i> )    | -    | 1.02 |       |
| Muscles    | Southeastern Bay of California, USA | CV - AAS    |  |         |             | Anatidae          | Lesser Scaup ( <i>Aythya affinis</i> )                     | -  | 1.06 | 60.8 | [105] |
|            | Southern Indian Ocean               | GC          |  |         |             | Diomedidae        | Southern Royal Albatross<br>( <i>Diomedea epomophora</i> ) | -  | 1.1  | -    | [106] |
|            | Chaun River, Siberia, Russia        |             |  | Animals | Vertebrates | Birds             | Laridae  | Arctic Tern ( <i>Sterna paradisaea</i> ) | -    | 1.1  |       |
| Feathers   |                                     |             |  |         |             |                   |  | -  | 1.1  | -    |       |



|            |                              |             |  |  |  |                   |  |   |      |      |       |
|------------|------------------------------|-------------|--|--|--|-------------------|--|---|------|------|-------|
| Intestines | Czech Republic               | HPLC/CV-AFS |  |  |  | Podicipedidae     | Great-crested Grebe<br>( <i>Podiceps cristatus</i> ) | - | 1.13 | 75.0 | [7]   |
| Liver      | Chaun River, Siberia, Russia | GC          |  |  |  | Laridae           | Herring Gull( <i>Larus argentatus</i> )              | - | 1.2  | -    | [106] |
|            | Czech Republic               | HPLC/CV-AFS |  |  |  | Accipitridae      | Common buzzard ( <i>Buteo buteo</i> )                | - | 1.24 | 47.8 | [7]   |
| Kidneys    | Chaun River, Siberia, Russia | GC          |  |  |  | Laridae           | Herring Gull( <i>Larus argentatus</i> )              | - | 1.3  | -    | [106] |
| Intestines | Czech Republic               | HPLC/CV-AFS |  |  |  | Phalacrocoracidae | Great Cormorant<br>( <i>Phalacrocorax carbo</i> )    | - | 1.41 | 78.4 | [7]   |
| Kidney     |                              |             |  |  |  | Accipitridae      | Common buzzard ( <i>Buteo buteo</i> )                | - | 1.44 | 71.9 |       |



|            |                                     |             |               |   |   |                   |  |     |      |      |       |
|------------|-------------------------------------|-------------|---------------|---|---|-------------------|--|-----|------|------|-------|
| Muscles    | Southeastern Bay of California, USA | CV - AAS    |               |   |   | Phalacrocoracidae | Neotropic Cormorant ( <i>Phalacrocorax brasilianus</i> ) | -   | 1.49 | 46.9 | [105] |
| Intestines | Czech Republic                      | HPLC/CV-AFS |               |   |   | Podicipedidae     | Great-crested Grebe ( <i>Podiceps cristatus</i> )        | -   | 1.57 | 80.0 | [7]   |
| Muscles    | Chaun River, Siberia, Russia        | GC          |               |   |   | Anatidae          | Long-tailed Duck ( <i>Clangula hyemalis</i> )            | -   | 1.6  | -    | [106] |
|            | Czech Republic                      | HPLC/CV-AFS |               |   |   | Podicipedidae     | Great-crested Grebe ( <i>Podiceps cristatus</i> )        | -   | 1.68 | 83.8 | [7]   |
| Kidneys    | Northern Pacific Ocean              | GC          |               |   |   | Procellariidae    | Northern Fulmar ( <i>Fulmarus glacialis</i> )            | -   | 1.7  | -    | [106] |
|            | Chaun River, Siberia, Russia        |             |               |   |   | Laridae           | Arctic Tern ( <i>Sterna paradisaea</i> )                 | -   | 1.9  | -    |       |
| Muscles    | Czech Republic                      | HPLC/CV-AFS | Podicipedidae | Great-crested Grebe ( <i>Podiceps cristatus</i> ) | - | 1.98              | 85.9   | [7] |      |      |       |

|         |  |             |  |  |  |                   |   |   |      |      |       |
|---------|--|-------------|--|--|--|-------------------|---|---|------|------|-------|
| Muscles | Northern Pacific Ocean                 | GC          |  |  |  | Diomedei<br>dae   | Black-footed Albatross<br>( <i>Phoebastria nigripes</i> ) | - | 2.0  | -    | [106] |
|         | Czech Republic                         | HPLC/CV-AFS |  |  |  | Phalacrocoracidae | Great Cormorant<br>( <i>Phalacrocorax carbo</i> )         | - | 2.11 | 84.3 | [7]   |
| Liver   | Chaun River, Siberia, Russia           | GC          |  |  |  | Anatidae          | Long-tailed Duck<br>( <i>Clangula hyemalis</i> )          | - | 2.2  | -    | [106] |
|         |  |             |  |  |  | Laridae           | Arctic Tern ( <i>Sterna<br/>paradisaea</i> )              | - | 2.3  | -    |       |
| Muscles | Czech Republic                         | HPLC/CV-AFS |  |  |  | Podicipedidae     | Great-crested Grebe<br>( <i>Podiceps cristatus</i> )      | - | 2.44 | 90.3 | [7]   |
|         | Southeastern Bay of California,<br>USA | CV - AAS    |  |  |  | Pelecanidae       | Brown Pelican ( <i>Pelecanus<br/>occidentalis</i> )       | - | 2.85 | 93.9 | [105] |

|         |                                       |             |         |             |       |                   |  |   |      |      |       |
|---------|---------------------------------------|-------------|---------|-------------|-------|-------------------|--|---|------|------|-------|
| Kidney  | Czech Republic                        | HPLC/CV-AFS | Animals | Vertebrates | Birds | Phalacrocoracidae | Great Cormorant<br>( <i>Phalacrocorax carbo</i> )          | - | 2.87 | 70.0 | [7]   |
| Muscles | Southeastern Ryukyu Islands,<br>Japan | GC          |         |             |       | Sulidae           | Brown Booby ( <i>Sula leucogaster</i> )                    | - | 2.9  | -    | [106] |
|         | Czech Republic                        | HPLC/CV-AFS |         |             |       | Phalacrocoracidae | Great Cormorant<br>( <i>Phalacrocorax carbo</i> )          | - | 3.04 | 89.3 | [7]   |
| Liver   | Northern Pacific Ocean                | GC          |         |             |       | Procellariidae    | Northern Fulmar<br>( <i>Fulmarus glacialis</i> )           | - | 3.1  | -    | [106] |
| Kidney  | Czech Republic                        | HPLC/CV-AFS |         |             |       | Podicipedidae     | Great-crested Grebe<br>( <i>Podiceps cristatus</i> )       | - | 3.50 | 74.4 | [7]   |
| Kidneys | Southern Indian Ocean                 | GC          |         |             |       | Diomedidae        | Southern Royal Albatross<br>( <i>Diomedea epomophora</i> ) | - | 3.6  | -    | [106] |



|         |                                       |             |  |  |  |                |  |               |   |      |       |
|---------|---------------------------------------|-------------|--|--|--|----------------|--|---------------|---|------|-------|
| Kidneys | Southeastern Ryukyu Islands,<br>Japan | GC          |  |  |  | Sulidae        | Brown Booby ( <i>Sula leucogaster</i> )                    | -             | 3.6   | -    | [106] |
| Liver   |                                       |             |  |  |  |                |  | -             | 3.7   | -    |       |
| Kidneys | Czech Republic                        | HPLC/CV-AFS |  |  |  | Podicipedidae  | Great-crested Grebe ( <i>Podiceps cristatus</i> )          | -             | 3.71  | 72.7 | [7]   |
|         | Chaun River, Siberia, Russia          | GC          |  |  |  |                |  | -             | 3.8   | -    | [106] |
| Muscles | Baffin Bay, Canada                    | GC - AED    |  |  |  | Laridae        | Glaucous Gull ( <i>Larus hyperboreus</i> )                 | -             | 4.0   | 77.5 | [107] |
| Liver   | Czech Republic                        | HPLC/CV-AFS |  |  |  |                |  | Podicipedidae | Great-crested Grebe ( <i>Podiceps cristatus</i> ) | -    | 4.1   |
| Kidneys | Southern Indian Ocean                 | GC          |  |  |  | Procellariidae | White-chinned Petrel ( <i>Procellaria aequinoctialis</i> ) | -             | 4.3   | -    | [106] |



|         |                        |             |         |             |       |                   |   |   |      |      |       |
|---------|------------------------|-------------|---------|-------------|-------|-------------------|---|---|------|------|-------|
| Liver   | Czech Republic         | HPLC/CV-AFS | Animals | Vertebrates | Birds | Podicipedidae     | Great-crested Grebe<br>( <i>Podiceps cristatus</i> )      | - | 4.34 | 60.3 | [7]   |
| Liver   |                        |             |         |             |       | Phalacrocoracidae | Great Cormorant<br>( <i>Phalacrocorax carbo</i> )         | - | 4.49 | 59.9 |       |
| Kidney  |                        |             |         |             |       | Phalacrocoracidae | Great Cormorant<br>( <i>Phalacrocorax carbo</i> )         | - | 4.56 | 63.3 |       |
| Liver   |                        |             |         |             |       | Podicipedidae     | Great-crested Grebe<br>( <i>Podiceps cristatus</i> )      | - | 4.72 | 55.5 |       |
| Muscles | Georgia, USA           | -           | Animals | Vertebrates | Birds | Rallidae          | Clapper Rail ( <i>Rallus longirostris</i> )               | - | 5.0  | 99   | [108] |
| Kidneys | Northern Pacific Ocean | GC          |         |             |       | Diomedidae        | Black-footed Albatross<br>( <i>Phoebastria nigripes</i> ) | - | 6.2  | -    | [106] |
| Muscles | Georgia, USA           | -           |         |             |       | Ardeidae          | Snowy Egret ( <i>Leucophoyx thula</i> )                   | - | 6.3  | 79   | [108] |
|         |                        |             |         |             |       |                   |   |   |      |      |       |



|                        |                              |             |         |             |  |                   |  |   |        |      |       |
|------------------------|------------------------------|-------------|---------|-------------|--|-------------------|--|---|--------|------|-------|
| Feathers               | Chaun River, Siberia, Russia | GC          |         |             |  | Laridae           | Herring Gull( <i>Larus argentatus</i> )                    | - | 6.5    | -    | [106] |
| Liver                  | Czech Republic               | HPLC/CV-AFS |         |             |  | Phalacrocoracidae | Great Cormorant ( <i>Phalacrocorax carbo</i> )             | - | 6.46   | 15.3 | [7]   |
|                        | Southern Indian Ocean        | GC          |         |             |  | Procellariidae    | White-chinned Petrel ( <i>Procellaria aequinoctialis</i> ) | - | 8.0    | -    | [106] |
|                        |                              |             |         |             |  | Diomedidae        | Southern Royal Albatross ( <i>Diomedea epomophora</i> )    | - | 9.8    | -    |       |
|                        |                              |             |         |             |  |                   | Laysan Albatross ( <i>Phoebastria immutabilis</i> )        | - | 11.2   | -    |       |
| Northern Pacific Ocean |                              |             |         |             | Black-footed Albatross ( <i>Phoebastria nigripes</i> ) | -                 | 20.4   | - |        |      |       |
| Fat                    | Black Sea                    | GLC - ECD   | Animals | Vertebrates | Mammals  | Phocoenidae       | Harbor porpoise ( <i>Phocoena phocoena</i> )               | - | 0.0049 | -    | [109] |
| Muscles                |                              |             |         |             |  |                   |  | - | 0.216  | -    |       |



|         |                 |             |                                      |   |      |             |   |       |      |     |       |
|---------|-----------------|-------------|--------------------------------------|---|------|-------------|---|-------|------|-----|-------|
| Kidney  | Japan           | AFS         |                                      |   |      | Delphinidae | Killer whale ( <i>Orcinus orca</i> )                    | -     | 0.24 | 3.2 | [110] |
| Brain   | Ontario, Canada |             |                                      |   |      | Mustelidae  | North American river otter ( <i>Lontra canadensis</i> ) | -     | 0.25 | -   | [66]  |
|         |                 |             |                                      |   |      |             | American mink ( <i>Mustela vison</i> )                  | -     | 0.26 | -   |       |
| Lungs   | Japan           | Delphinidae | Killer whale ( <i>Orcinus orca</i> ) | -   | 0.31 | -           | [110]   |       |      |     |       |
| Liver   | Black Sea       | GLC - ECD   | Phocoenidae                          | Harbor porpoise ( <i>Phocoena phocoena</i> )            | -    | 0.322       | -   | [109] |      |     |       |
|         | Ontario, Canada | AFS         | Mustelidae                           | North American river otter ( <i>Lontra canadensis</i> ) | -    | 0.87        | -   | [66]  |      |     |       |
| Muscles | Japan           |             | Delphinidae                          | Killer whale ( <i>Orcinus orca</i> )                    | -    | 0.90        | 74  | [110] |      |     |       |
| Kidney  | Ontario, Canada |             | Mustelidae                           | American mink ( <i>Mustela vison</i> )                  | -    | 0.94        | -   | [66]  |      |     |       |



|         |                 |        |             |   |         |             |   |   |      |     |       |
|---------|-----------------|--------|-------------|---|---------|-------------|---|---|------|-----|-------|
| Kidney  | Ontario, Canada | AFS    | Animals     | Vertebrates   | Mammals | Mustelidae  | North American river otter<br>( <i>Lontra canadensis</i> )            | - | 0.94 | -   | [66]  |
| Muscles | Japan           | GC-ECD |             |   |         | Phocoenidae | Dall's porpoise<br>( <i>Phocoenoides dalli</i> )                      | - | 1.02 | 84  | [111] |
| Liver   |                 | AFS    |             |   |         | Delphinidae | Killer whale ( <i>Orcinus orca</i> )                                  | - | 1.11 | 1.9 | [68]  |
| Liver   | Ontario, Canada | AFS    |             |   |         | Mustelidae  | American mink ( <i>Mustela vison</i> )                                | - | 1.21 | -   | [66]  |
| Muscles | Japan           | GC-ECD |             |   |         | Delphinidae | Short-finned pilot whale<br>(N) ( <i>Globicephala macrorhynchus</i> ) | - | 1.25 | 81  | [111] |
|         |                 |        |             |   |         | Ziphiidae   | Baird's beaked whale<br>( <i>Berardius bairdii</i> )                  | - | 1.25 | 78  |       |
|         |                 |        | Delphinidae | Pantropical spotted dolphin ( <i>Stenella attenuata</i> ) | -       | 2.62        | 54  |   |      |     |       |
|         |                 |        |             | Risso's dolphin ( <i>Grampus griseus</i> )                | -       | 3.15        | 74  |   |      |     |       |
|         |                 |        | Delphinidae | Rough-toothed dolphin<br>( <i>Steno bredanensis</i> )     | -       | 3.51        | 74  |   |      |     |       |





|   |                     |        |         |             |         |             |  |             |  |    |       |
|---|---------------------|--------|---------|-------------|---------|-------------|--|-------------|--|----|-------|
| Muscles   | Japan               | GC-ECD | Animals | Vertebrates | Mammals | Delphinidae | Striped dolphin ( <i>Stenella coeruleoalba</i> )                   | -           | 3.74   | 63 | [111] |
|   | North Sea, Denmark  |        |         |             |         | Phocoenidae | Harbor porpoise ( <i>Phocoena phocoena</i> )                       | -           | 4.0  | -  | [112] |
|   | Japan               |        |         |             |         | Delphinidae | Short-finned pilot whale (S) ( <i>Globicephala macrorhynchus</i> ) | -           | 6.45   | 64 | [111] |
| Common bottlenose dolphin ( <i>Tursiops truncatus</i> ) |                     | -      | 6.83    | 54          |         |             |  |             |  |    |       |
| Liver   | North Sea, Denmark  | GC-ECD | Animals | Vertebrates | Mammals | Phocoenidae | Harbor porpoise ( <i>Phocoena phocoena</i> )                       | -           | 6.9  | -  | [112] |
| Muscles   | North Sea - Belgium |        |         |             |         |             |  | -           | 7.3  | -  |       |
| Fur   | Ontario, Canada     | AFS    | Animals | Vertebrates | Mammals | Mustelidae  | North American river otter ( <i>Lontra canadensis</i> )            | -           | 8.24   | -  | [66]  |
| Liver   | North Sea - Belgium | GC-ECD |         |             |         | Phocoenidae | Harbor porpoise ( <i>Phocoena phocoena</i> )                       | -           | 8.6  | -  | [112] |
| Muscles   | Japan               |        |         |             |         |             |  | Delphinidae | False killer whale ( <i>Pseudorca crassidens</i> ) | -  | 11.2  |

|        |                           |          |  |  |  |               |   |           |       |   |      |
|--------|---------------------------|----------|--|--|--|---------------|---|-----------|-------|---|------|
| Fur    | Ontario, Canada           | AFS      |  |  |  | Mustelidae    | American mink ( <i>Mustela vison</i> )                | -         | 11.25 | - | [66] |
| Liver  | South coast of Brazil     | CV - AAS |  |  |  | Pontoporiidae | La Plata dolphin<br>( <i>Pontoporia blainvillei</i> ) | 0.05-4.21 | -     | - | [45] |
|        | Southeast coast of Brazil |          |  |  |  |               |   | 0.12-2.36 | -     | - |      |
| Kidney |                           |          |  |  |  |               |   | 0.16-1.13 | -     | - |      |
| Kidney | South coast of Brazil     |          |  |  |  |               |   | 0.26-1.82 | -     | - |      |

\*,\*\* - the numerical values are given as originally stated in the cited literature

**Table 5.** Basic information on the techniques used in the speciation analysis of mercury.

| ISOLATION TECHNIQUE                           | ADVANTAGES  | DRAWBACKS   | DETECTION TECHNIQUE   | LIMIT OF DETECTION   |         |                          | REFERENCES         |
|---|---|---|-----------------------|--|---------|--------------------------|--------------------|
|   |   |   |                       | MeHg(I)  | EtHg(I) | PhHg(I)                  |                    |
| High-performance liquid chromatography (HPLC) | <ul style="list-style-type: none"> <li>✓ High selectivity</li> <li>✓ A universal, integrated, fully automated method of mercury speciation</li> <li>✓ Quick and simple</li> <li>✓ Does not require conversion of target analytes into volatile derivatives</li> <li>✓ Isolation of mercury compounds possible at room temperature</li> </ul>  | <ul style="list-style-type: none"> <li>✓ Such low LODs as with GC are not obtainable</li> <li>✓ Poorer selectivity in the analysis of complex matrices</li> <li>✓ Poorer sensitivity than with GC</li> <li>✓ Large quantities of organic solvents consumed</li> </ul>   | CV - AFS              | 27 pg<br>0.015 - 0.1 µg  | 26 pg   | -                        | [150]<br>[151-153] |
|   |   |   | CV - AAS              | 0.1-16 µg/dm <sup>3</sup>  | -       | -                        | [1,154]            |
|   |   |   | ICP - MS              | 16- 400 ng/ dm <sup>3</sup>  | -       | -                        | [1]                |
|   |   |   | ICP - AES             | 0.1ng/ cm <sup>3</sup>   | -       | -                        | [1]                |
|   |   |   | (CV) MIP - AES        | 0.35 ng/ cm <sup>3</sup>   | -       | -                        | [1]                |
|   |   |   | UV - PCO - CVAFS      | 0.015 µg/kg  | -       | -                        | [153]              |
|   |   |   | HF – LLLME            | 3.8 ng/ cm <sup>3</sup>  | -       | 0.3 ng/ cm <sup>3</sup>  | [155]              |
|   |   |   | UV - Vis              | 7.0–95.1 µg/dm <sup>3</sup>  |         |                          | [156]              |
| Gas chromatography (GC)                       | <ul style="list-style-type: none"> <li>✓ High precision</li> <li>✓ Far lower LODs obtainable than with HPLC</li> <li>✓ Both organic and inorganic mercury compounds can be determined</li> <li>✓ Usually used to determine MeHg in environmental samples</li> <li>✓ High degree of target analyte isolation</li> <li>✓ Can be hyphenated with many different detection systems</li> </ul> | <ul style="list-style-type: none"> <li>✓ Target analytes in the sample have to be converted into volatile derivatives</li> <li>✓ Specific analytical conditions are required</li> <li>✓ For determining MeHg cleanup is required, to eliminate the interference of organic halides</li> <li>✓ Sample preparation has a significant effect on efficacy and accuracy</li> </ul> | MIP-AES               | 0.01-0.06 µg/g<br>0.04-10 ng/dm <sup>3</sup>   | -       | -                        | [1,157-160]        |
|   |   |   | AFS and GC-(CV) - AFS | 5 pg and 0.01-6 ng/ dm <sup>3</sup> ;<br>0.6-1.3 pg  | 5pg     | -                        | [1,161-163]        |
|   |   |   | (CV) - AAS            | 5-167 pg   | -       | -                        | [1,143,164]        |
|   |   |   | ICP - MS              | 0.9 pg<br>0.5 pg<br>0.12-1 pg  | 1.0 pg  | -                        | [1,163,165]        |
|   |   |   | ICP - AES             | 3 pg. 0.6 ng/dm <sup>3</sup>   | -       | -                        | [1]                |
| Capillary electrophoresis (CE)                | <ul style="list-style-type: none"> <li>✓ Highly effective isolation of analytes</li> <li>✓ Only a small volume of sample is needed</li> <li>✓ Short separation time</li> </ul>  | <ul style="list-style-type: none"> <li>✓ Poor sensitivity</li> <li>✓ Poor stability</li> </ul>  | UV                    | 47.5 ng/ cm <sup>3</sup><br>680 ng/ cm <sup>3</sup>  | -       | 4.1 ng/ cm <sup>3</sup>  | [166,167]          |
|   |   |   | DF – ICP - MS         | 54 ng/ cm <sup>3</sup>   | -       | -                        | [167,168]          |
|   |   |   | ICP - MS              | 2.3 pg<br>128 ng/ cm <sup>3</sup><br>80 ng/ cm <sup>3</sup><br>13.6 ng/ cm <sup>3</sup><br>149 ng/ cm <sup>3</sup> | -       | -                        | [167-170]          |
|   |   |   | VSG - AFS             | 2.5 pg<br>16.5 ng/ cm <sup>3</sup>   | 2.4 pg  | 13.3 ng/ cm <sup>3</sup> | [168,171]          |
|   |   |   | AAS                   | 2.9 pg   | -       | -                        | [172]              |

**Table 6.** Analytical procedures used in the speciation analysis of mercury

| TYPE OF SAMPLE                 | EXTRACTION  | ANALYTE CLEANUP  | EXTRACTION EFFICIENCY AS RECOVERY [%]                              |        |     |     | ANALYTICAL TECHNIQUE | METROLOGICAL PARAMETERS   | REFERENCES |
|--------------------------------|---|--|--|--------|-----|-----|----------------------|---|------------|
|                                |   |  |  | MeHg   | OHg | THg |                      |   |            |
| Water                          | Approximately 30 cm <sup>3</sup> of water was weighed in a Teflon tube, 0.2 ml of 1% ammonium pyrrolidinedithiocarbamat, 0.5 ml 4 mol/dm <sup>3</sup> KBr and 0.5 cm <sup>3</sup> 2 mol/dm <sup>3</sup> H <sub>2</sub> SO <sub>4</sub> .  | Solution distilled at 110° C in the presence of N <sub>2</sub> for 4-5 h. Derivatization of MeHg with NaBEt <sub>4</sub> in a buffer solution at pH 4.5. MeEtHg was trapped on a Tenax column.   | -  | -      | -   | -   | GC – CV - AFS        | 20 pg/dm <sup>3</sup> (for MeHg)  | [89]       |
| Water                          |   | Collection of samples <i>in situ</i> by pumping them through a minicolumn packed with C18 modified with sodium diethyldithiocarbamate. Elution of mercury species with 500 cm <sup>3</sup> 5 % thiourea in 0.5 % HCl from minicolumns. | -  | -      | -   | -   | HPLC – ICP - MS      | 5.6 ng/dm <sup>3</sup> (for MeHg)   | [134]      |
| Sediments, zoobenthos, water   | Microwave assisted extraction 0.5– 4 g of a sample with the extraction agent containing 3 mol /dm <sup>3</sup> HCl + 0.2 mol/dm <sup>3</sup> citric acid + 50 % methanol (10 cm <sup>3</sup> ). Acidity of the filtrated extracts was adjusted to pH 3 by NaOH. Addition of 2-mercaptophenol. | The Speed C18 SPE stationary phase (Applied Separations, Allentown, PA, USA) was used for preparation of SPE microcolumns that also enabled countercurrent-flow elution.   | Method of standard addition (certified reference material CRM 580) | -      | >95 | -   | HPLC – CV - AFS      | 4.3 μg/dm <sup>3</sup> (for MeHg)<br>1.4 μg/dm <sup>3</sup> (for EtHg)<br>0.8 μg/dm <sup>3</sup> (for PhHg) | [20]       |
| Soils, earthworm tissue, fungi | Addition to the samples of 5 % H <sub>2</sub> SO <sub>4</sub> , 18% KBr and 1.0 cm <sup>3</sup> 1M CuSO <sub>4</sub>  | Derivatization of MeHg with 50 cm <sup>3</sup> 1 % NaBEt <sub>4</sub> for 15 min in a buffer   | Method of standard addition (certified reference materials:        | 89-103 | -   | -   | GC – CV - AFS        | 0.05 μg/kg (for MeHg)   | [19]       |





|                |  |  |  |        |   |          |               |                    |       |
|----------------|--|--|--|--------|---|----------|---------------|--------------------|-------|
|                | <p>solution and vigorous shaking for 15 min. Addition of 1 cm<sup>3</sup> CH<sub>2</sub>Cl<sub>2</sub> and the samples shaken again for 15 min. Centrifugation for 5 min at 3200 rpm. Separation of the organic phase (CH<sub>2</sub>Cl<sub>2</sub>) from the aqueous phase. Extraction repeated with an additional 5 cm<sup>3</sup> CH<sub>2</sub>Cl<sub>2</sub>. Addition of 35 cm<sup>3</sup> of Milli-Q water to the combined CH<sub>2</sub>Cl<sub>2</sub>. Evaporation on a water bath at about 90°C. Samples purged with N<sub>2</sub> for 5 min to remove remaining CH<sub>2</sub>Cl<sub>2</sub>.</p> | <p>solution at pH 4.6. Ethylmethyl-Hg was purged onto a Tenax trap for 15 min with Hg-free N<sub>2</sub>.</p>  | DORM-2, TORT-2)                                    |        |   |          |               |                    |       |
| Periphyton     | <p>Replicate MeHg extractions carried out by immersing three freeze-dried filters in 10 cm<sup>3</sup> of 25 % KOH in methanol solution in a screw-cap bottle. The bottles were mounted on a wrist shaker overnight at room temperature.</p>   | <p>0.25 cm<sup>3</sup> of the extract transferred to a glass reaction vessel, and diluted with 50 cm<sup>3</sup> of mains tap water. pH was adjusted to 4.9 with acetate buffer. Derivatization of MeHg with 100 µl of 1 % sodium tetraethylborate solution. The volatile alkylmercury derivatives were trapped on a Tenax column.</p> | Analysis of reference material (DORM-2)            | 98     | - | -        | GC – CV - AFS | 10.8 pg (for MeHg) | [128] |
| Zooplankton    | <p>Digestion of 2-5 mg (d.w.) plankton aliquots in 0.5 ml KOH/MeOH (1g/4 cm<sup>3</sup>) solution for 8 h.</p>   |  |  |        |   |          | GC - AFS      | 9.9 pg (for MeHg)  | [55]  |
| Microorganisms | <p>Alkaline digestion of 2 to 5 mg d.w. of powder in 0.5 ml of KOH/MeOH (1g/4 cm<sup>3</sup>) solution during 8 h at 6°C.</p>  | <p>MeHg converted to MeEtHg with sodium tetraethylborate in a buffer solution at pH 4.5. MeEtHg trapped on a Tenax column.</p>   | Analysis of reference material (TORT-2 and DORM-2) | 88-102 | - | 90.4-110 | GC – CV - AFS | 2 ng/g (for meHg)  | [87]  |

|                       |   |   |  |        |   |    |          |   |          |
|-----------------------|---|---|--|--------|---|----|----------|---|----------|
|                       |   |   |  |        |   |    |          |   |          |
| Starfish, polychaetes | -   | -   | Adding known amounts of mercury to samples -method of standard addition (DORM-1) | 87     | - | 94 | GC - ECD | 0.003 µg/g (in an averaged 1.5 g sample) (for MeHg) | [93]     |
| Mollusks              | Dissolution of the tissue in an alkaline medium.  | Ethylation with sodium tetraethylborate. Volatile ethylated forms of mercury were then subjected to a flow of nitrogen and trapped on a Tenax column. |  |        |   |    | GC - AFS | -   | [95]     |
| Mussels               | Separation of MeHg from the homogenized mussel samples using two techniques: a water vapor distillation technique (1.0 g of homogenized sample in a mixture of H <sub>2</sub> SO <sub>4</sub> /NaCl/H <sub>2</sub> O at 150° C), and an ion exchange method (Dovex 1X8, Cl-form, 100-200 mesh), followed by UV decomposition of methylmercury.  | Ionic mercury reduction by tin (II) chloride (10 % SnCl <sub>2</sub> in 20 % H <sub>2</sub> SO <sub>4</sub> ) to elemental mercury                    | -  | -      | - | -  | CV - AAS | -   | [99]     |
| Cephalopod tissue     | Addition of 2 cm <sup>3</sup> Milli-Q water and 3 cm <sup>3</sup> 6 mol/dm <sup>3</sup> KOH solution to 200 mg of dried sample. The mixture was shaken for 2 h, after which 3 cm <sup>3</sup> 6 mol/dm <sup>3</sup> HCl and 4 cm <sup>3</sup> KBr/CuSO <sub>4</sub> (3:1) solution was added. After 10 minutes of shaking, 5 cm <sup>3</sup> dichloromethane (DCM) was added, the mixture centrifuged and the organic | A weak sulfide solution was used to extract MeHg from the organic phase, then MeHg was back-extracted to DCM.   | Method of standard addition (certified reference materials: DORM-2, TORT-2)      | 92-103 | - | -  | GC - AFS | 0.01 µg/g (for Hg)                                  | [96,173] |



|             |   |   |   |                   |   |      |                 |  |       |
|-------------|---|---|---|-------------------|---|------|-----------------|--|-------|
|             | phase separated.  |   |   |                   |   |      |                 |  |       |
| Fish tissue | Drying, cryogenic grinding and addition of potassium bromide and hydrochloric acid solution (1 mol/dm <sup>3</sup> KBr in 6 mol/ dm <sup>3</sup> HCl) to the samples. Centrifugation.   | Extraction of organomercury compounds from KBr solution using chloroform. Back-extraction with 1 % m/v L-cysteine. Mercury vapor generation from extracts was performed using 1 mol/dm <sup>3</sup> HCl and 2.5 % m/v NaBH <sub>4</sub> solutions and a batch chemical vapor generation system. | -   | -                 | - | -    | CVG - ETAAS     | 5 ng/g (for MeHg)  | [6]   |
| Fish tissue | Acidic digestion with HCl-H <sub>2</sub> O (1+1) and triple extraction with 10 cm <sup>3</sup> of benzene. Extracts were diluted to 25 cm <sup>3</sup> with benzene and mixed with 5g Na <sub>2</sub> SO <sub>4</sub> .           | -   | -   | -                 | - | -    | GC - ECD        | -  | [101] |
| Fish tissue | Alkaline digestion with KOH – ethanol solution at 100° C in a water bath containing ethylene glycol.  | Extraction of MeHg with 0.01 % dithizone–toluene solution, clean up and purification of the Dz-Tol extract.   | -   | -                 | - | -    | GC -ECD         | -  | [102] |
| Fish tissue | Acidic digestion with 2 ml of 14.25 mol/dm <sup>3</sup> H <sub>2</sub> SO <sub>4</sub> (saturated with copper(II) sulfate), 2 ml of 4 mol/dm <sup>3</sup> KBr, and 2.5cm <sup>3</sup> of toluene solution and shaking for 35 min. | Organic extract back-extracted with 1 ml cysteine solution (1.5% w/v). Shaking and centrifugation (2200 rpm) for 10 min.  | Analysis of reference material (CRM 463)                                | 98.6              | - | 96.5 | GC - ECD        | 7 ng/g (for MeHg)  | [27]  |
| Fish tissue | Acidic digestion with 11.3 cm <sup>3</sup> 37 % HCl. Incubation at 100° C for 10 min. Addition of 15 cm <sup>3</sup> toluene. Sonication for 20 min and centrifugation for 25 min.  | Extraction of supernatant with 5 ml 0.1 % cysteine solution. Shaking for 10 min and centrifugation for 25 min   | Analysis of two certificate reference material (DORM-2 and NIES CRM 13) | 97 ± 5-<br>98 ± 6 | - | -    | RPC - CVG - AFS | 18 pg (for MeHg)<br>18 pg (for EtHg)<br>20 pg (for PhHg) | [174] |




|                |  |  |   |        |        |        |                 |   |       |
|----------------|--|--|---|--------|--------|--------|-----------------|---|-------|
| Sea food       | Oxidation of the organomercury species permitted the determination of total mercury. Mercury species separated by the selective retention of inorganic mercury on the chelating resin. The inorganic mercury was removed on-line from the microcolumn with 6 % (m/v) thiourea.   | Mercury cold vapor generation was performed on-line with 0.2 % (m/v) sodium borohydride and 0.05 % (m/v) sodium hydroxide as reducing solution.  | Method of standard addition (certified reference materials: (DOLT-1 , TORT-1) | 98-110 | -      | 94-110 | FI CV-AAS       | 10 ng/dm <sup>3</sup><br>(forMeHg)<br><br>6 ng/dm <sup>3</sup><br>(forMeHg) | [23]  |
| Seafood tissue | Addition of water (1.0 cm <sup>3</sup> ) and KOH (1.5 cm <sup>3</sup> , 6 mol/dm <sup>3</sup> ) solution to each of the samples, shaken for 12 h. Addition of HCl (3 cm <sup>3</sup> , 3mol/dm <sup>3</sup> ) to each vial for neutralization. Once the effervescence and heat had subsided, acidic KBr and CuSO <sub>4</sub> mixture at 3:1 ratio (v/v, 3 cm <sup>3</sup> ) were added. Then, 10 cm <sup>3</sup> of toluene was added, and the mixture was shaken for 3 min and set aside for 10 min. | Centrifugation at 8000 rpm for 2 min. Subsequently, 5 cm <sup>3</sup> of Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> (0.01 mol/dm <sup>3</sup> ) was added to 5 cm <sup>3</sup> of toluene and the mixture was shaken for 3 min and set aside for 10 min. Addition of acidic KBr/CuSO <sub>4</sub> in 3:1 ratio (v/v, 0.4 cm <sup>3</sup> ) to 2 cm <sup>3</sup> of the aquatic layer. After the addition 0.2 cm <sup>3</sup> of toluene, the mixture was shaken for 2 min, then set aside for 10 min. | Method of standard addition (certified reference material: DORM-2)            | -      | 88-106 | -      | GC – ICP - MS   | 0.5 pg (for MeHg)<br>1.0 pg (for EtHg)                                      | [113] |
| Seafood        | Addition of 3 cm <sup>3</sup> 25 % (m/v) KOH (in methanol) to 1.0–2.0 g wet samples with mechanical shaking overnight. Addition of 3 cm <sup>3</sup> 6 mol/dm <sup>3</sup> HCl, 4 cm <sup>3</sup> acidic KBr/CuSO <sub>4</sub> (3:1) and 5 cm <sup>3</sup> CH <sub>2</sub> Cl <sub>2</sub> were added to the tube in sequence, shaking for 2 h to extract organic mercury into the CH <sub>2</sub> Cl <sub>2</sub> phase.  | Centrifugation at 2000 rpm for about 10 min, the CH <sub>2</sub> Cl <sub>2</sub> . Extraction with 1 cm <sup>3</sup> sodium thiosulfate. Shaking for 45 min.   | -   | -      | -      | -      | HPLC – CV - AFS | 0.2 ng(forMeHg)<br>0.17 ng(forEtHg)<br>0.14 ng(forPhHg)                     | [175] |
| Seafood tissue | Acid digestion with H <sub>2</sub> SO <sub>4</sub> + CuSO <sub>4</sub> + KBr.  |  |   |        |        |        | INNA            | 0.02 mg/kg (for MeHg)   | [98]  |






|  |   |   |   |        |    |   |                      |  |       |
|--|---|---|---|--------|----|---|----------------------|--|-------|
|  | Extraction with toluene.  |   |   |        |    |   |                      |  |       |
| Bird tissue<br>(pectoral muscle,<br>intestines, liver<br>and kidney) | Microwave assisted extraction 0.2-1.0 g of sample with 6 mol/dm <sup>3</sup> HCl + 0.1 mol/dm <sup>3</sup> NaCl for 10 min in an Ethos SEL high-pressure microwave digestion unit.  | After filtration and dilution, the supernatant with acetate buffer (pH = 5) was made up to the final volume of 50 cm <sup>3</sup> .   | Analysis of reference material (DORM-2)   | 95-107 |    |   | HPLC – CV - AFS      | 0.2 ppb (for MeHg)<br>0.06 ppb (for PhHg)<br>0.12 ppb (for EtHg) | [7]   |
| Ringed seal tissue   | Homogenization with an aqueous solution of acidic sodium bromide (5 cm <sup>3</sup> of 30 % in 4 N H <sub>2</sub> SO <sub>4</sub> ) and copper(II) sulfate (7.5 cm <sup>3</sup> of 2.5 % in 4 N H <sub>2</sub> SO <sub>4</sub> ). Extraction MeHg and other forms of organic mercury by vortexing the tissue homogenate with a 3:2 v/v mixture of dichloromethane - hexane (5-10 cm <sup>3</sup> ). | A fraction of the organic layer was withdrawn and extracted with (3 - 4 cm <sup>3</sup> ) aqueous thiosulfate (0.005 N) by vortexing for 1 min and centrifuging. Separation by adding an aliquot of thiosulfate (1-2 cm <sup>3</sup> ), KI (0.5 cm <sup>3</sup> , 3 mol/dm <sup>3</sup> ). Back-extraction with toluene (1.5 - 3.0 cm <sup>3</sup> ). Separation with toluene by centrifuging for 2 min at 2500 rpm. The extract was dried over anhydrous sodium sulfate. | Using marine mammal liver tissue with a relatively high organic mercury and MeHg content to test recovery and extraction efficiencies for MeHg. | 92     | 90 | - | GC - ECD             | 10-80 ng/g Hg ww.<br>pg (for MeHg)                               | [134] |
| Biological tissues   | Extraction of mercury species into 10 g/dm <sup>3</sup> EDTA and 0.2 % (v/v) 2-mercaptoethanol solution.  |   | Method of standard addition (certified reference materials: (DORM-2 , DOLT-3)   | 93-99  |    |   | HPLC – ICP - MS      | 0.2–0.3 µg/dm <sup>3</sup> (for MeHg)                            | [122] |
| Biological samples   | Acidic extraction with hydrochloric acid.   | Decantation and the re-extraction. Neutralization with 10 mol/dm <sup>3</sup> sodium hydroxide. Dilution of extracts in the mobile phase and filtration.  | -   | -      | -  | - | HPLC – ICP - MS      | 0.08 µg/dm <sup>3</sup> (for MeHg)                               | [176] |
| Sediment,<br>biological tissue                                       | Water vapor distillation technique for the isolation of methylmercury   |   | Method of standard addition (certified reference materials: IAEA 356, CRM 463, CRM 464,   | >95    | -  | - | HPF/HHPN<br>ICP - MS | 0.025 mg/kg (for MeHg)   | [177] |

|                             |  |  |  |   |                           |   |                                  |                    |       |
|-----------------------------|--|--|--|---|---------------------------|---|----------------------------------|--------------------|-------|
|                             |  |  | DORM – 1)  |   |                           |   |                                  |                    |       |
| Biological tissue, sediment | Hydrolyzed with HBr (47–49 %). Shaking, addition of 20 cm <sup>3</sup> toluene to the samples. Mixed for 20 min. Centrifugation at 2400 rpm for 20 min. Collecting the supernatant, containing organomercuryspecies in falcon tubes. | Back-extraction with 6 cm <sup>3</sup> of 1 % (v/w) l-cysteine aqueous solution to strip methylmercury from toluene. | Analysis of certificate reference materials (CRM-580, IAEA-405, DORM-2, DOLT-3, SRM-2976 and SRM – 2977) | - | > 80 (except DOLT 3 - 74) | - | Direct Mercury Analyzer DMA - 80 | 0.04 ng (for MeHg) | [197] |



**Figure 1.** The mercury cycle in the biosphere.



**Figure 2.** Levels of total mercury in organisms/samples [ $\mu\text{g}/\text{kg}$ ] and the percentage of methylmercury in the total mercury content [7].




Figure 3. The circulation of methylmercury in the maternal and fetal organisms [73].




Figure 4. Scheme for preliminary sample preparation in the speciation analysis of mercury.