

Quantum origins of objectivity

R. Horodecki,^{1,2} J. K. Korbicz,^{1,2} and P. Horodecki^{3,2}

¹*Institute of Theoretical Physics and Astrophysics, University of Gdańsk, 80-952 Gdańsk, Poland*

²*National Quantum Information Centre in Gdańsk, 81-824 Sopot, Poland*

³*Faculty of Applied Physics and Mathematics, Gdańsk University of Technology, 80-233 Gdańsk, Poland*

(Received 10 September 2014; published 30 March 2015)

In spite of all of its successes, quantum mechanics leaves us with a central problem: How does nature create a bridge from fragile quanta to the objective world of everyday experience? Here we find that a basic structure within quantum mechanics that leads to the perceived objectivity is a so-called spectrum broadcast structure. We uncover this based on minimal assumptions, without referring to any dynamical details or a concrete model. More specifically, working formally within the decoherence theory setting with multiple environments (called quantum Darwinism), we show how a crucial for quantum mechanics notion of nondisturbance due to Bohr [N. Bohr, *Phys. Rev.* **48**, 696 (1935)] and a natural definition of objectivity lead to a canonical structure of a quantum system-environment state, reflecting objective information records about the system stored in the environment.

DOI: [10.1103/PhysRevA.91.032122](https://doi.org/10.1103/PhysRevA.91.032122)

PACS number(s): 03.65.Ta, 03.67.—a

I. INTRODUCTION

The emergence of the objective world from quanta has been a longstanding problem, already present from the very beginning of quantum mechanics [1–3]. One of the most promising approaches is the decoherence theory, based on a system-environment paradigm [4,5]: A quantum system is considered interacting with its environment. It recovers, under certain conditions, a classical-like behavior of the system alone in some preferred frame, singled out by the interaction and called a pointer basis [6], and explains it through information leakage from the system into the environment. However, decoherence theory lacks an explanation for the redundancy of information in the classical realm [7]: The same record can exist in a large number of copies and can be independently accessed by many observers and many times. More basically, since quanta cannot be cloned [8] and information redundancy is, from the perspective of the observers' measurements, at the heart of objectivity, then what quantum process lies at the foundation of the objective classical world?

Recently, a crucial step was made in a series of works (see, e.g., [7,9]) introducing the so-called quantum Darwinism idea. Its essence is that in more realistic environments, composed of many independent fractions, decoherence leads to the appearance of multiple copies of the system's state in the environment, accessible to independent observers. Although presenting a convincing physical picture, there is no general model-independent justification of such claims apart from studies under the strict conditions of specific models, e.g., spin-1/2 systems [10] or an illuminated sphere [11]. However, even those studies still do not present totally convincing arguments within the models themselves, as they are based on a scalar information-theoretic condition and so-called partial information plots, which are known to be only a necessary condition for objectivity but with sufficiency still unknown.

Here we take a more fundamental and rigorous position, based solely on what for now provides the most basic description of nature: a quantum state (see also [12]). More precisely, we derive from first principles a universal objectivity-carrying structure of quantum states, using a general approach,

independent of any dynamics (much like, e.g., the S -matrix theory in quantum field theory [13]): Looking at the postinteraction system-environment state, we ask what properties should it have to reflect the objectivity. Surprisingly, the answer comes with the help of Bohr's notion of nondisturbance [14,15], which was originally used to defend the quantum [14,16], whereas here, ironically, it defines the classical. It is obtained through what we call a spectrum broadcast structure, which precisely pinpoints the distributed character of information and makes it essentially classical. We finally illustrate our approach using one of the emblematic examples of decoherence theory: a dielectric sphere illuminated by photons [11,17,18]. It must be mentioned that in the quantum Darwinism literature there appeared similar quantum state structures (so-called branching states). However, they have been at best tacitly postulated [7,9,19], if at all explicitly mentioned. Our results allow us to understand the intimate connection between the perceived objectivity, a specific structure of quantum states, and information broadcasting.

II. GENERAL THEOREM

A. Basic definitions and the main result

We first define the central concepts of our study and state the main result. The basis of our work is the following definition of an objective state [7,20].

Definition 1: Objectivity. A state of the system S exists objectively if many observers can find out the state of S independently and without perturbing it.

As it stands, the above definition is rather informal and has to be made more rigorous. For example, the key concept of perturbation has to be made precise, which we do in the next section.

The second key concept of our study is a spectrum broadcast structure, defined as follows.

Definition 2: Spectrum broadcast structure. A spectrum broadcast structure is the following form of a joint state of the central system S and a collection of subenvironments

E_1, \dots, E_{fN} (denoted by fE):

$$\rho_{S:fE} = \sum_i p_i |i\rangle_S \langle i| \otimes \rho_i^{E_1} \otimes \dots \otimes \rho_i^{E_{fN}}, \quad (1)$$

where $\{|i\rangle\}$ is some basis in the system's space, p_i are probabilities, and all states $\rho_i^{E_k}$ are perfectly distinguishable:

$$\rho_i^{E_k} \rho_{i'}^{E_k} = 0 \quad \forall i \neq i', \quad k = 1, \dots, fN. \quad (2)$$

All the nomenclature will be clarified in the next section. This is a special form of so-called classical-classical state, which has been introduced as a counterpart of separable states in the context of quantification of quantum correlations [21,22]. It originally appeared in the context of quantum channels in [23].

The main result of this work is the establishment of an intimate connection between these two concepts. The two pivotal assumptions we use are a Bohr nondisturbance [14,15] and a strong independence. The first was formulated in [15] (see Sec. II B); by a strong independence we mean that the only correlation between the environments should be the common information about the system. In other words, conditioned by the information about the system, there should be no correlations between the environments. This is in a sense an idealization, which we use since we are interested in the information flow only between the system and each of the environments, but not between the environments themselves. Under these central assumptions (together with some auxiliary ones), we prove in Sec. II C the following theorem.

Theorem 1. Assume that a system undergoes a full decoherence. Then the appearance of a spectrum broadcast structure is a necessary and sufficient condition for objectivity in the sense of Definition 1:

Objective existence \Leftrightarrow Spectrum broadcast structure,

$$\left(\begin{array}{c} \text{Objective} \\ \text{existence} \end{array} \right) + \left(\begin{array}{c} \text{Strong} \\ \text{independence} \end{array} \right) \Rightarrow \left(\begin{array}{c} \text{Spectrum} \\ \text{broadcast} \\ \text{structure} \end{array} \right).$$

B. Formalization of Definition 1

Here we put Definition 1 into a physical frame and make it as precise as possible, which is the hardest work. As the most suitable, we formally choose decoherence theory with multiple environments [7]: The quantum system of interest S interacts with multiple environments E_1, \dots, E_N (denoted collectively as E), also modeled as quantum systems. The environments (or their collection) are assumed to be macroscopic and are monitored by independent observers [9]. The motivation behind such a choice is that in real-life situations there is always present some interaction with the environment (unless very special conditions are met) and we, the observers, usually have access only to a small portion of it, each to a different part. However, as we will see in what follows, no assumptions on the dynamics will be needed. In fact, we may forget about the dynamics altogether and pose a more general question: Which multipartite system-environment states reflect objectively an existing state of the system in the sense of Definition 1?

We only assume that the system-environment interaction is such that it leads to a full decoherence. The standard, or even paradigmatic, case corresponding to the latter is a

physical situation when there exists a time scale τ_D , called the decoherence time, such that asymptotically for interaction times $t \gg \tau_D$ (i) there emerges in the system's Hilbert space a unique, stable in time pointer basis $\{|i\rangle\}$ and (ii) the reduced state of the system ρ_S becomes stable and diagonal in the pointer basis

$$\rho_S \equiv \text{Tr}_E \rho_{S:E} \approx \sum_i p_i |i\rangle \langle i|, \quad (3)$$

where the p_i are probabilities and by \approx we will always denote asymptotic equality in the deep decoherence limit $t/\tau_D \rightarrow \infty$. However, it should be stressed that while we usually will mean the latter situation, our derivation of the structure of objectivity covers also possible situations when the process happens in finite time. We assume here the above explained full decoherence, so that the system decoheres in a basis rather than in higher-dimensional pointer superselection sectors. This is because we want to consider the full objectivization of a given quantum degree of freedom rather than a partial one. Clearly, the environment must be of a large dimension to have a big informational capacity, needed to carry highly redundant records about the decohered system S . Moreover, some loss of information is needed (and of course happens in reality), as otherwise there will be no decoherence, and we assume that some of the environments pass unobserved. The observed environments E_1, \dots, E_{fN} we denote by fE (depending on the context).

Next we specify the observers. Apart from the environmental ones, we also allow for a (possibly hypothetical) direct observer, who can measure the system S directly. Such an observer is needed as a reference to verify that the findings of the environmental observers are the same as if one had a direct access to the system.

By "finding" we mean that the observers are performing von Neumann (as perfectly repeatable, contrary to the generalized) measurements on their subsystems. It should be stressed here that the von Neumann measurement, with its repeatability property, has been chosen since we identify the spectrum broadcast structure as the paradigmatic, ideal structure of the state, responsible for objectivity. Indeed, this is the object to which any real physical state should be compared if we want to know whether the objectivity in a more or less approximate sense (in terms of a state trace distance) takes place. (Note that it can be compared with the ideal singlet as the target state of quantum distillation or the ideal channel, in the case of coding theory, to which the outputs of real protocols or physical situations are compared.)

By the independence requirement of Definition 1, there can be no correlations between them. Consequently, the global von Neumann measurement, resulting from the individual local observer's measurements, must be fully product

$$\Pi_i^{M_S} \otimes \Pi_{j_1}^{M_1} \otimes \dots \otimes \Pi_{j_{fN}}^{M_{fN}}, \quad (4)$$

where M_S, M_1, \dots, M_{fN} denote measurements on S, E_1, \dots, E_{fN} and all Π are mutually orthogonal Hermitian projectors $\Pi_j^{M_k} \Pi_{j' \neq j}^{M_k} = 0$. The observers so determine the probabilities p_i of $|i\rangle$ in (3) (they must know the pointer basis $\{|i\rangle\}$; otherwise they would not know what the information

they get is all about). As explained before Theorem 1, we will actually demand more by assuming the strong independence.

The most crucial clarification needed in Definition 1 is to make precise the word perturbing. We apply here Bohr's notion of nondisturbance [14,15], according to which the given local measurements on the subsystems are nondisturbing if they leave the whole joint state invariant (after forgetting the results). This is a realistic mathematical idealization of a repetitive information extraction, a crucial prerequisite for objectivity.

We recall that Bohr's nondisturbance was formulated in order to save the completeness of quantum theory against the famous Einstein-Podolsky-Rosen (EPR) argument [16]. Bohr argued [14] that the EPR notion of mechanical nondisturbance (which amounts to the no-signaling principle [15]) was too restricted and a broader notion was needed. Hence, accepting the completeness of quantum theory, as we do for the purpose of this work, one is forced to accept Bohr's notion of nondisturbance.

Finally, the independent measurements will typically reveal inconsistent information about the system (see, however, [12]). Indeed, allowing for general correlations may lead to a disagreement: If one of the observers measures first, the ones measuring afterward may find outcomes depending on the result of the first measurement. Thus, we add to Definition 1 an obvious agreement requirement that many observers can find the same state of S independently.

C. Proof of Theorem 1

We are now ready for a proof of Theorem 1 with the additional assumption of strong independence, explained in Sec. II A. We first prove that the spectrum broadcast structure from Definition 2 is a sufficient condition for an objectively existing state of the system, in the sense of Definition 1. Indeed, from (1) projections on $|i\rangle$ and on the disjoint supports of $\varrho_i^{E_k}$ constitute the nondemolition measurements. Performing them independently, all the observers will repeatedly detect the same index i with probabilities $\{p_i\}$, without Bohr disturbing the joint $S:fE$ state, thus making the state $|i\rangle$ exist objectively in the sense of Definition 1 (cf. [24]).

We now prove the reverse. We assume the decoherence has taken place (cf. [12]). Crucial here is the Bohr nondisturbance condition from Sec. II B. Together with the product structure (4), it implies that on each subsystem S, E_1, \dots, E_{fN} there exists a nondemolition von Neumann measurement, leaving the whole asymptotic state $\varrho_{S:fE}(\infty)$ of the system and the observed environment invariant (the symbol ∞ stands here either for the $t/\tau_D \rightarrow \infty$ asymptote or as mentioned before for any time scale, maybe finite, after which the objectivity structure emerges). For S it is defined by the projectors on $|i\rangle$. For the environments we allow for higher-rank projectors $\Pi_j^{M_k}$, $k = 1, \dots, fN$, not necessarily spanning the whole space, as the environments can have inner degrees of freedom not correlating to S .

Consequently, the total joint probability of the results of the Bohr nondisturbing measurements is given by

$$p_{i j_1 \dots j_{fN}} \equiv \text{Tr}[|i\rangle\langle i| \otimes \Pi_{j_1}^{M_1} \otimes \dots \otimes \Pi_{j_{fN}}^{M_{fN}} \varrho_{S:fE}(\infty)]. \quad (5)$$

Now the agreement requirement from Sec. II B leads to a natural conclusion

$$p_{i j_1 \dots j_{fN}} \neq 0 \quad \text{if and only if } i = j_1 = \dots = j_{fN}. \quad (6)$$

Let us more formally show it, considering for simplicity only two observers. If one of them measures first and gets a result i , then the joint conditional state becomes $\varrho_{|i} = (1/p_i)(\Pi_i \otimes \mathbf{1})\varrho(\Pi_i \otimes \mathbf{1})$, $p_i \equiv \text{Tr}(\Pi_i \otimes \mathbf{1}\varrho)$ and the subsequent measurement by the second observer will yield results j with conditional probabilities $p_{j|i} = (1/p_i)\text{Tr}(\Pi_i \otimes \Pi_j \varrho)$. If for some i , $p_{j|i} p_{j'|i} \neq 0$ for $j \neq j'$, then comparing their results after a series of measurements at some later moment, the observers will be confused as to what exactly the state the system S was: With the probability $p_{j|i} p_{j'|i}$ the second observer will obtain different states $j \neq j'$, while the first observer measured the same state i . The observers' findings are not objective unless for every i there exists only one $j(i)$ such that $p_{j(i)|i} \neq 0$ (actually $p_{j(i)|i} = 1$, which follows from the normalization $\sum_i p_{i|j} = 1$, so that the distributions $p_{\cdot|i}$ are all deterministic). Reversing the measurement order and applying the same reasoning, we obtain that for every j there can exist only one $i(j)$ such that $\tilde{p}_{i(j)|j} \neq 0$, where by the Bayes theorem $\tilde{p}_{i(j)|j} = p_{j|i} p_i / \tilde{p}_j$, $\tilde{p}_j \equiv \text{Tr}(\mathbf{1} \otimes \Pi_j \varrho)$. These two conditions imply that the joint probability $p_{ij} = p_i \delta_{ij}$ (after an eventual renumbering). Applying the above argument to all pairs of indices, one obtains (6). This means that the environmental Bohr-nondisturbing measurements must be perfectly correlated with the pointer basis. Hence, after forgetting the results, the asymptotic postmeasurement state $\varrho_{S:fE}^M(\infty)$ reads

$$\begin{aligned} \varrho_{S:fE}^M(\infty) &\equiv \sum_{i, j_1, \dots, j_{fN}} p_{i j_1 \dots j_{fN}} \varrho_{i j_1 \dots j_{fN}}^{S:fE}(\infty) \\ &= \sum_i |i\rangle\langle i| \otimes \Pi_i \varrho_{S:fE}(\infty) |i\rangle\langle i| \otimes \Pi_i, \end{aligned} \quad (7)$$

where $\Pi_i \equiv \Pi_i^{M_1} \otimes \dots \otimes \Pi_i^{M_{fN}}$.

Now we are ready for the key step: We impose the relevant form of the Bohr-nondisturbance condition

$$\sum_i |i\rangle\langle i| \otimes \Pi_i \varrho_{S:fE}(\infty) |i\rangle\langle i| \otimes \Pi_i = \varrho_{S:fE}(\infty), \quad (8)$$

whose only solution [15] are classical-quantum (CQ) states [25]

$$\varrho_{S:fE}(\infty) = \sum_i p_i |i\rangle\langle i| \otimes \mathbf{R}_i^{fE}, \quad (9)$$

where p_i are identified with the probabilities from Eq. (3) and \mathbf{R}_i^{fE} are some residual states in the space of the observed environments with mutually orthogonal supports $\mathbf{R}_i^{fE} \mathbf{R}_{i' \neq i}^{fE} = 0$. Hence, \mathbf{R}_i^{fE} are perfectly distinguishable through the assumed nondisturbing measurements Π_i , projecting on their supports.

Finally, let us look at the residual states \mathbf{R}_i^{fE} in (9). The demand of the independent ability to determine the state of S , already used in (4), completed with the strong independence condition (cf. Sec. II) leads to the following: Once one of the observers finds a particular result i , the resulting conditional

state should be fully product. Since the direct observer is already uncorrelated by (9), this implies that

$$\mathbf{R}_i^{fE} = \varrho_i^{E_1} \otimes \cdots \otimes \varrho_i^{E_{fN}} \quad (10)$$

and $\varrho_i^{E_k}$ must be perfectly distinguishable for each E_k [cf. (2)] since by (8) for any k it holds that $\Pi_i^{M_k} \varrho_i^{E_k} \Pi_i^{M_k} = \varrho_i^{E_k}$ and $\Pi_i^{M_k} \Pi_{i' \neq i}^{M_k} = 0$. This finishes the proof.

Some remarks are in order. First, in the course of the proof we have formulated a broader class of independent environments, in a way paradigmatic in quantum information theory [26]: The environments are independent if and only if the environmental observers may produce the states (10) and (2), exploiting only local operations (equivalent to local trace-preserving maps), i.e., independent environments are those that simulate a strong independence from the perspective of a specific resource (the class of local operations).

Second, the meaning of Theorem 1 is that it provides an ideal reference structure for objectivity, the broadcast structure (1). Any other nonideal situation should be compared to that broadcast structure no matter what figure of merit is taken. On the level of the states, this must be the trace norm, which has the clear probabilistic interpretation, where the degree of objectivity is just the trace norm distance to the broadcast state. The transition of the initial $S:E$ state to the spectrum broadcast structure (1) identifies a basic process, called here state information broadcasting, responsible for the appearance of the perceived objectivity. Formally, it involves broadcasting of a part of information about the system, the spectrum of its state after the decoherence $\text{Sp}_{\varrho_S} \equiv \{p_i\}$ into the environments, and is thus similar to quantum state [27] and spectrum [23] broadcasting. Condition (2) forces the correlations in (1) to be entirely classical and thus the detailed structures of $\varrho_i^{E_k}$ become irrelevant for the correlations. From (1) and (2) it follows that under a suitable convergence

$$I[\varrho_{S:fE}(\infty)] = H_S \quad \text{for every fraction } f, \quad (11)$$

where $I(\varrho_{AB}) \equiv S_{\text{VN}}(\varrho_A) + S_{\text{VN}}(\varrho_B) - S_{\text{VN}}(\varrho_{AB})$ is the quantum mutual information, $S_{\text{VN}}(\varrho) \equiv -\text{Tr}(\varrho \log \varrho)$ stands for the von Neumann entropy, and $H_S \equiv S_{\text{VN}}[\varrho_S(\infty)] = H(\{p_i\})$ is the entropy of the decohered state (3). Condition (11), postulated as a sufficient condition for objectivity in the quantum Darwinism model, has a clear meaning in the classical information theory [28]: Every fraction f carries the same information H_S about the system; the latter is redundantly encoded in the environment. However, in the quantum world its sense remains unclear (see the next section). Here (11) follows automatically from the deeper structure (1).

III. DISCUSSION OF THE ENTROPIC OBJECTIVITY CONDITION AS EVIDENCE OF OBJECTIVITY

Here we show a potential problem with the entropic objectivity condition (11) as a sufficient condition for objectivity (see, e.g., Refs. [7,9] and references therein). Although our example below is not fully conclusive, we argue that at this moment neither is the reasoning of quantum Darwinism studies.

Condition (11) has been shown to hold in several models, including the illuminated sphere [11,18] and spin baths [10].

For finite times t , the equality (11) is not strict and holds within some error $\delta(t)$, which defines the redundancy $R_\delta(t)$ as the inverse of the smallest fraction of the environment $f_{\delta(t)}$ for which $I[\varrho_{S:f_{\delta(t)}E}(t)] = [1 - \delta(t)]H_S$. When satisfied, (11) implies that the mutual information between the system and the environment fraction is a constant function of the fraction size f (up to an error δ for finite times) and the plot of I against f exhibits a characteristic plateau, called the classical plateau (see, e.g., Ref. [7]). The appearance of this plateau has been heuristically explained in the quantum Darwinism literature as a consequence of the redundancy: Classical information about the system exists in many copies in the environment fractions and can be accessed independently and without perturbing the system by many observers, thus leading to the objective existence of a state of S [7]. That would certainly be the case in the classical information setting: The condition (11) there is equivalent to a perfect correlation of both systems [28], i.e., for every f the environment fraction has full information about the system and indeed this information thus exists objectively in the sense of our definition.

However, in the quantum world the situation may be different and the condition (11) alone may not be sufficient to guarantee objectivity due to the holistic nature of quantum correlations [29]. It is clear that the spectrum broadcast states (1) satisfy (11), but there may also be entangled states satisfying it, thus violating the spectrum broadcast form, derived as a necessary condition for objectivity. As a simple example in favor of such a statement consider the following state of two qubits, where one is the system S and the second the environment E :

$$\varrho_{S:E} \equiv pP_{(a|00)+b|11)} + (1-p)P_{(a|01)+b|10)}, \quad (12)$$

where $P_\psi \equiv |\psi\rangle\langle\psi|$, $p \neq 1/2$, $a = \sqrt{p}$, and $b = \sqrt{1-p}$. Then the partial state $\varrho_S = \tilde{p}|0\rangle\langle 0| + (1-\tilde{p})|1\rangle\langle 1|$, $\tilde{p} \equiv pa^2 + (1-p)b^2$ is diagonal in the basis $|0\rangle, |1\rangle$ and moreover $S_{\text{VN}}(\varrho_S) = S_{\text{VN}}(\varrho_{S:E}) \equiv h(\tilde{p})$ (the binary Shannon entropy [28]), so a form of the entropic condition holds, $I(\varrho_{S:E}) = S_{\text{VN}}(\varrho_S) = H_S$, $H_S = h(\tilde{p})$, but the systems are nevertheless entangled, which one verifies directly through the positive partial transpose criterion [30].

The above example is of course not conclusive, as there is only one environment, but it suggests that the functional condition (11) in principle might indeed be insufficient to show objectivity, as defined in previously. We leave this, in general difficult, question open for further research. In the above context, the paradigmatic shift we propose here, with respect to the earlier works on decoherence and quantum Darwinism models, can already be seen: It is the pivotal observation governing our approach that the core object of the analysis should be a derived structure of the full quantum state of the system S and the observed environment fE rather than the partial state of the system only (decoherence theory) or information-theoretic functions (quantum Darwinism).

IV. SPECTRUM BROADCAST STRUCTURE IN THE ILLUMINATED SPHERE MODEL

We exemplify the general findings from Sec. II on one of the central models of decoherence (see, e.g., [11,17,18]): a dielectric sphere illuminated by photons (for details see the

Appendix). We show that in the course of the evolution, a broadcast state (1) is asymptotically formed in this model, assuming for simplicity pure environments (see [18] for a more general analysis). The sphere is initially in a state without a well-defined position [e.g., in $|\psi_0^S\rangle = (|\vec{x}_1\rangle + |\vec{x}_2\rangle)/\sqrt{2}$]. Photons scatter elastically and slightly differently depending on where the sphere is, but this difference is vanishingly small for each individual scattering: If the observed fraction is too small, the postscattering states $|\Psi_i^{\text{mic}}\rangle \equiv \mathbf{S}_i|\vec{k}_0\rangle$ (\mathbf{S}_i are the scattering matrices) become identical in the thermodynamic limit $\langle\Psi_2^{\text{mic}}|\Psi_1^{\text{mic}}\rangle \equiv \langle\vec{k}_0|\mathbf{S}_2^\dagger\mathbf{S}_1|\vec{k}_0\rangle \xrightarrow{\text{therm}} 1$ and the joint postscattering state approaches effectively a product $(\sum_{i=1,2} p_i |\vec{x}_i\rangle\langle\vec{x}_i|) \otimes |\Psi^{\text{mic}}\rangle\langle\Psi^{\text{mic}}|^{\otimes\mu}$, where probabilities $p_i \equiv |\langle\psi_0^S|\vec{x}_i\rangle|^2$ form the spectrum of the decohered state [cf. (3)]. The photons thus force the sphere to be in a definite position \vec{x}_i with probability p_i , but the observed fraction carries no information about it (a product phase; see the Appendix, Sec. 3).

However, when grouped into macroscopic fractions, the photons become almost perfectly resolving. Imagine that we divide all the photons scattered up to time t , N_t , into M macrofractions of mN_t , $0 < m < 1$, photons (Fig. 1). Then the macroscopic postscattering states $|\Psi_i^{\text{mac}}(t)\rangle \equiv (\mathbf{S}_i|\vec{k}_0\rangle)^{\otimes mN_t}$ become asymptotically perfectly distinguishable

$$|\langle\Psi_2^{\text{mac}}(t)|\Psi_1^{\text{mac}}(t)\rangle| \xrightarrow{\text{therm}} e^{-m(t/\tau_D)}, \quad (13)$$

where τ_D is the decoherence time [11,17]. If we observe fM , $0 < f < 1$, macrofractions out of M , then the joint postscattering state has asymptotically the spectrum broadcast

structure (1),

$$\begin{aligned} \rho_{S:fE}(0) &= \rho_0^S \otimes \underbrace{\rho_0^{\text{mac}} \otimes \dots \otimes \rho_0^{\text{mac}}}_{fM} \xrightarrow[t_{\text{therm}}]{t \gg \tau_D} \rho_{S:fE}(\infty) \\ &= \sum_{i=1,2} p_i |\vec{x}_i\rangle\langle\vec{x}_i| \otimes \underbrace{|i^{\text{mac}}\rangle\langle i^{\text{mac}}| \otimes \dots \otimes |i^{\text{mac}}\rangle\langle i^{\text{mac}}|}_{fM}, \end{aligned} \quad (14)$$

where $|i^{\text{mac}}\rangle \equiv |\Psi_i^{\text{mac}}(\infty)\rangle$ emerges, due to (13), as the nondisturbing environmental basis in the space of each macrofraction. Equation (14) identifies the state information broadcasting process: The information about the sphere's localization $\{p_i\}$ is redundantly transferred into the environment and becomes available in multiple copies through the measurements in $\{|i^{\text{mac}}\rangle\}$. The process consists of (i) decoherence [17] and (ii) orthogonalization (13) and defines a broadcasting phase (see the Appendix, Sec. 3) corresponding to the classical plateau of [11]. From Fannes-Audenaert [31] and Alicki-Fannes [32] inequalities, the entropic condition (11) follows as a consequence of (14) (see the Appendix, Sec. 4). Finally, if all the photons are observed, the postscattering state maintains the full quantum correlation with the system and $I[\rho_{S:fE}(\infty)] = I_{\text{max}}$ (a full information phase).

V. DISCUSSION

In conclusion, based on a universal approach, independent of any dynamics or a concrete model, we have identified the primitive state information broadcasting process responsible for the emergence of the perceived objectivity (for a possible loosening of some of our assumptions see [12]). Our main result (Theorem 1) suggests that the states of the form (14) are notoriously formed in nature. In a laboratory, this can in principle be directly verified via, e.g., quantum state tomography [33]. Moreover, it naturally leads to the view that in fact there may be no quantum-to-classical transition; what we perceive as classical, e.g., objective information, may be merely a reflection of some specific properties of the underlying quantum states, like the spectrum broadcast structure, a view further strengthened by [34].

There appears to be a deep connection between the nonsignaling principle and objective existence in the sense of Definition 1: The core fact that it is at all possible for observers to determine independently the classical state of the system is guaranteed by the nonsignaling principle $\text{Tr}(\mathbf{1}_S \otimes \Pi_{EQS:E}) = \text{Tr}_E(\Pi_{EQE})$. There is no contradiction with the Bohr nondisturbance, as the latter is a strictly stronger condition than the nonsignaling principle [15] (this is the core of Bohr's reply [14] to Einstein, Podolsky, and Rosen). In fact, the above connection reaches deeper than quantum mechanics. In a general theory, where it is possible to speak of probabilities $p(ij|MN)$ of obtaining results i, j when performing measurements M, N (however defined), whatever the definition of objective existence may be, the requirement of the independent ability to locally determine probabilities by each party seems indispensable. This is guaranteed in the nonsignaling theories, where all $p(ij|MN)$ have well-defined marginals. In this sense nonsignaling seems a prerequisite of cognition. In this context, we also believe that our approach




FIG. 1. (Color online) Coarse graining of the photonic environment. The photons (green) scattered in time t are grouped into M equal macroscopic fractions mN_t . Only one fraction (bounded by the red cage) is shown; L is the edge of an artificial box used for quantization (and removed later; see the Appendix). The macrofractions may be thought of as representing a sensitivity of the photon detectors (e.g., an eye) and their exact size mN_t is irrelevant; scaling with the total photon number N_t suffices.

to objectivity will present a different perspective on the celebrated Bell theorem [35].

The emergence of redundantly encoded information in the structure of quantum states may also shed light on the life phenomenon. Since self-replication of the DNA information is indispensable for the existence of life, it cannot be excluded that the state information broadcasting may indeed open a classical window for life processes within quantum mechanics [36].

ACKNOWLEDGMENTS

We thank W. H. Zurek and C. J. Riedel for discussions and comments and M. Piani for discussions about strong independence. P.H. and R.H. acknowledge discussions with K. Horodecki, M. Horodecki, and K. Życzkowski. This research was supported by ERC Advanced Grant QO-LAPS and National Science Centre project Maestro DEC-2011/02/A/ST2/00305.

APPENDIX: TECHNICAL DETAILS OF THE ILLUMINATED SPHERE MODEL FOR PURE ENVIRONMENTS

1. Description of the model

Here we present a detailed derivation of the spectrum broadcast structure (1) in the illuminated sphere model for pure environments (see [18] for a more general situation). We first recall the basics of the model, following the usual treatment (see, e.g., Refs. [11,17,37,38]). The system S is a sphere of radius a and relative permittivity ϵ , bombarded by a constant flux of photons, which constitute the multiple environments and decohere the sphere. The sphere can be located only at two positions \vec{x}_1 or \vec{x}_2 , so effectively its state space is that of a qubit $\mathcal{H}_S \equiv \mathbb{C}^2$ with a preferred orthonormal (due to the mutual exclusiveness) basis $|\vec{x}_1\rangle, |\vec{x}_2\rangle$, which will become the pointer basis. This greatly simplifies the analysis, yet allows the essence of the effect to be observed. The sphere is sufficiently massive compared to the energy of the radiation, so the recoil due to the scattering can be totally neglected and the photons' energy is conserved, i.e., the scattering is elastic.

The environmental photons are assumed to be not energetic enough to individually resolve the sphere's displacement $\Delta x \equiv |\vec{x}_2 - \vec{x}_1|$:

$$k\Delta x \ll 1, \quad (\text{A1})$$

where $\hbar k$ is the characteristic photon momentum. Otherwise, each individual photon would be able to resolve the position of the sphere and studying multiple environments would not bring anything new. On the technical side, following the traditional approach [11,17,37,38], we describe the photons in a simplified way using box normalization: We assume that the sphere and the photons are enclosed in a large box of edge L and volume $V = L^3$ and photon momentum eigenstates $|\vec{k}\rangle$ obey periodic boundary conditions. Although a more rigorous treatment was developed in Ref. [39] with well-localized photon states, we choose this traditional heuristic approach as, at the expense of mathematical rigor, it allows us to expose the physical situation more clearly, without unnecessary mathematical details (we remark that the findings of Ref. [39]

agree with the previous works using box normalization [40]). After dealing with formally divergent terms, we remove the box through the thermodynamic limit (signified by \cong) [11,38]

$$V \rightarrow \infty, \quad N \rightarrow \infty, \quad \frac{N}{V} = \text{const}, \quad (\text{A2})$$

that is, we expand the box and add more photons, keeping the photon density constant, as the relevant physical quantity is the radiative power, proportional to N/V . The thermodynamic limit is crucial in the sense that it defines microscopic and macroscopic regimes, which will in turn be qualitatively very distinct.

The detailed dynamics of each individual scattering is irrelevant; the individual scatterings are treated asymptotically in time. The interaction time t enters the model differently, through the number of scattered photons. It may be called macroscopic time. Assuming photons come from the area of L^2 at a constant rate of N photons per volume V per unit time, the amount of scattered photons from $t = 0$ to t is

$$N_t \equiv L^2 \frac{N}{V} ct, \quad (\text{A3})$$

where c is the speed of light. Throughout the calculations we work with a fixed time t and pass to the asymptotic limit $t/\tau_D \rightarrow \infty$ (signified by \approx or ∞) at the very end.

Since multiphoton scatterings can be neglected and all the photons are treated equally (symmetric environments), the effective sphere-photon interaction up to time t is of a controlled-unitary form

$$U_{S:E}(t) \equiv \sum_{i=1,2} |\vec{x}_i\rangle \langle \vec{x}_i| \otimes \underbrace{\mathbf{S}_i \otimes \cdots \otimes \mathbf{S}_i}_{N_t}, \quad (\text{A4})$$

where (assuming translational invariance of the photon scattering) $\mathbf{S}_i \equiv \mathbf{S}_{\vec{x}_i} = e^{-i\vec{x}_i \cdot \hat{\mathbf{k}}} \mathbf{S}_0 e^{i\vec{x}_i \cdot \hat{\mathbf{k}}}$ is the scattering matrix when the sphere is at \vec{x}_i , \mathbf{S}_0 is the scattering matrix when the sphere is at the origin, and $\hat{\mathbf{k}}$ is the photon momentum operator. Due to the elastic scattering, the \mathbf{S}_i have nonzero matrix elements only between the states $|\vec{k}\rangle$ of the same energy $\hbar c|\vec{k}|$. In the sector (A1) the interaction (A4) is vanishingly small at the level of each individual photon [38]: In the thermodynamic limit $\mathbf{S}_1 \cong \mathbf{S}_2$ (in a suitable sense we clarify later) and hence $\sum_i |\vec{x}_i\rangle \langle \vec{x}_i| \otimes \mathbf{S}_i \cong \mathbf{1} \otimes \mathbf{S}$. Surprisingly, this will not be true for macroscopic groups of photons. We also note that unlike in the previous treatments [11,17,37–39], already at this moment we explicitly include in the description *all* the photons scattered up to the fixed time t . Finally, the preferred role of the basis $|\vec{x}_i\rangle$ is already singled out now by the form of the interaction (A4) [7].

The initial prescattering in state is as usually assumed to be a full product

$$\varrho_{S:E}(0) \equiv \varrho_0^S \otimes (\varrho_0^{\text{ph}})^{\otimes N_t}, \quad (\text{A5})$$

with ϱ_0^S having coherences in the preferred basis $|\vec{x}_i\rangle$ and ϱ_0^{ph} some initial states of the photons (the environments are by assumption symmetric). Next we introduce a crucial environment coarse graining [7]: The full environment (i.e., all the N_t photons) is divided into a number of macroscopic fractions, each containing mN_t photons, $0 \leq m \leq 1$. By

macroscopic we will always understand scaling with the total number of photons N_t . By definition, these are the environment fractions accessible to the independent observers. Such a division may seem artificial and arbitrary as, e.g., the choice of m is unspecified. However, observe that in typical situations detectors used to monitor fractions of the environment, e.g., eyes, have some minimum detection thresholds; some minimum amount of radiative energy delivered in a given time interval is needed to trigger the detection. Each macroscopic fraction mN_t is meant to reflect that detection threshold. Its concrete value (the fraction size m) is for our analysis irrelevant; it is enough that it scales with N_t . This coarse-graining procedure is analogous to the one used, e.g., in the description of liquids: Each point of a liquid (a macrofraction m here) is in reality composed of a suitable large number of microparticles (individual photons). It is also employed in the mathematical approach to von Neumann measurements using so-called macroscopic observables (see, e.g., Ref. [41] and references therein). Thus, we divide the detailed initial state of the environment $(\varrho_0^{\text{ph}})^{\otimes N_t}$ into $M \equiv 1/m$ macroscopic fractions

$$\underbrace{\varrho_0^{\text{ph}} \otimes \dots \otimes \varrho_0^{\text{ph}}}_{N_t} = \underbrace{\varrho_0^{\text{ph}} \otimes \dots \otimes \varrho_0^{\text{ph}}}_{mN_t} \otimes \dots \otimes \underbrace{\varrho_0^{\text{ph}} \otimes \dots \otimes \varrho_0^{\text{ph}}}_{mN_t} \equiv \underbrace{\varrho_0^{\text{mac}} \otimes \dots \otimes \varrho_0^{\text{mac}}}_M, \quad (\text{A6})$$

where $\varrho_0^{\text{mac}} \equiv (\varrho_0^{\text{ph}})^{\otimes mN_t}$ is the initial state of each macroscopic fraction (macrostate for brevity).

2. Dynamical formation of broadcast structure

After all N_t photons have scattered, the asymptotic (in the sense of the scattering theory) out state $\varrho_{S:E}(t) \equiv U_{S:E}(t)\varrho_{S:E}(0)U_{S:E}(t)^\dagger$ is given, from Eqs. (A4)–(A6), by

$$\varrho_{S:E}(t) = \sum_{i=1,2} \langle \vec{x}_i | \varrho_0^S \vec{x}_i \rangle | \vec{x}_i \rangle \langle \vec{x}_i | \otimes \underbrace{\varrho_i^{\text{mac}}(t) \otimes \dots \otimes \varrho_i^{\text{mac}}(t)}_M \quad (\text{A7})$$

$$+ \sum_{i \neq j} \langle \vec{x}_i | \varrho_0^S \vec{x}_j \rangle | \vec{x}_i \rangle \langle \vec{x}_j | \otimes \underbrace{(\mathbf{S}_i \varrho_0^{\text{ph}} \mathbf{S}_j^\dagger)^{\otimes mN_t} \otimes \dots}_M, \quad (\text{A8})$$

where

$$\varrho_i^{\text{mac}}(t) \equiv (\mathbf{S}_i \varrho_0^{\text{ph}} \mathbf{S}_i^\dagger)^{\otimes mN_t}, \quad i = 1, 2. \quad (\text{A9})$$

In order for the decoherence to take place, some of the environment must be traced out. In the current model it is important that the forgotten fraction must be macroscopic: We assume that fM , $0 \leq f \leq 1$, out of all M macrofractions of Eq. (A6) are observed, while the rest $(1-f)M$ are traced out. The resulting partial state reads [cf. Eqs. (A7) and (A8)]

$$\varrho_{S:fE}(t) = \sum_{i=1,2} \langle \vec{x}_i | \varrho_0^S \vec{x}_i \rangle | \vec{x}_i \rangle \langle \vec{x}_i | \otimes [\varrho_i^{\text{mac}}(t)]^{\otimes fM} \quad (\text{A10})$$

$$+ \sum_{i \neq j} \langle \vec{x}_i | \varrho_0^S \vec{x}_j \rangle (\text{Tr} \mathbf{S}_i \varrho_0^{\text{ph}} \mathbf{S}_j^\dagger)^{(1-f)N_t} | \vec{x}_i \rangle \langle \vec{x}_j | \otimes (\mathbf{S}_i \varrho_0^{\text{ph}} \mathbf{S}_j^\dagger)^{\otimes fN_t}. \quad (\text{A11})$$

We finally demonstrate that in the soft scattering sector (A1), the above state is asymptotically of the broadcast form (1) by showing that in the deep decoherence regime $t \gg \tau_D$ two effects take place: (i) The coherent part $\varrho_{S:fE}^{i \neq j}(t)$ given by Eq. (A11) vanishes in the trace norm

$$\|\varrho_{S:fE}^{i \neq j}(t)\|_{\text{tr}} \equiv \text{Tr} \sqrt{[\varrho_{S:fE}^{i \neq j}(t)]^\dagger \varrho_{S:fE}^{i \neq j}(t)} \approx 0 \quad (\text{A12})$$

and (ii) the postscattering macroscopic states $\varrho_i^{\text{mac}}(t)$ [cf. Eq. (A9)] become perfectly distinguishable

$$\varrho_1^{\text{mac}}(t) \varrho_2^{\text{mac}}(t) \approx 0 \quad (\text{A13})$$

or, equivalently, using the generalized overlap [42]

$$B[\varrho_1^{\text{mac}}(t), \varrho_2^{\text{mac}}(t)] \equiv \text{Tr} \sqrt{\sqrt{\varrho_1^{\text{mac}}(t)} \varrho_2^{\text{mac}}(t) \sqrt{\varrho_1^{\text{mac}}(t)}} \approx 0, \quad (\text{A14})$$

despite the individual (microscopic) states becoming equal in the thermodynamic limit.

The first mechanism above is the usual decoherence of S by fE , the suppression of coherences in the preferred basis $|\vec{x}_i\rangle$. Some form of quantum correlations may still survive it since the resulting state (A10) is generally of a CQ form [25]. Those relict forms of quantum correlations are damped by the second mechanism: the asymptotic perfect distinguishability (A13) of the postscattering macrostates $\varrho_i^{\text{mac}}(t)$. Thus, the state $\varrho_{S:fE}(\infty)$ becomes of the spectrum broadcast form (1) for the distribution

$$p_i = \langle \vec{x}_i | \varrho_0^S \vec{x}_i \rangle. \quad (\text{A15})$$

We demonstrate the mechanisms (A12) and (A13) and hence the formation of the broadcast state (1) for pure initial environments

$$\varrho_{ph}^0 \equiv |\vec{k}_0\rangle \langle \vec{k}_0|, \quad k_0 \Delta x \ll 1, \quad (\text{A16})$$

i.e., all the photons come from the same direction and have the same momenta $\hbar k_0$, $k_0 \equiv |\vec{k}_0|$, satisfying (A1). To show (A12), observe that $\varrho_{S:fE}^{i \neq j}(t)$, defined by Eq. (A11), is of a simple form in the basis $|\vec{x}_i\rangle$,

$$\varrho_{S:fE}^{i \neq j}(t) = \begin{bmatrix} 0 & \gamma C \\ \gamma^* C^\dagger & 0 \end{bmatrix}, \quad (\text{A17})$$

where $\gamma \equiv \langle \vec{x}_1 | \varrho_0^S \vec{x}_2 \rangle (\text{Tr} \mathbf{S}_1 \varrho_0^{\text{ph}} \mathbf{S}_2^\dagger)^{(1-f)N_t}$ and $C \equiv (\mathbf{S}_1 \varrho_0^{\text{ph}} \mathbf{S}_2^\dagger)^{\otimes fN_t}$. Since the \mathbf{S}_i are unitary and $\varrho_0^{\text{ph}} \geq 0$, $\text{Tr} \varrho_0^{\text{ph}} = 1$, we obtain

$$\|\varrho_{S:fE}^{i \neq j}(t)\|_{\text{tr}} = |\gamma| \text{Tr} (\mathbf{S}_1 \varrho_0^{\text{ph}} \mathbf{S}_1^\dagger)^{\otimes fN_t} + |\gamma| \text{Tr} (\mathbf{S}_2 \varrho_0^{\text{ph}} \mathbf{S}_2^\dagger)^{\otimes fN_t} \quad (\text{A18})$$

$$= 2 |\langle \vec{x}_1 | \varrho_0^S \vec{x}_2 \rangle| |\text{Tr} \mathbf{S}_1 \varrho_0^{\text{ph}} \mathbf{S}_2^\dagger|^{(1-f)N_t}. \quad (\text{A19})$$

The decoherence factor $|\text{Tr} \mathbf{S}_1 \varrho_0^{\text{ph}} \mathbf{S}_2^\dagger|^{(1-f)N_t}$ for the pure case (A16) has been extensively studied before (see, e.g., Refs. [11,17,37–39]). Let us briefly recall the main results. Under the condition (A1) and using the classical cross section of a dielectric sphere in the dipole approximation $k_0 a \ll 1$,

one obtains in the box normalization

$$\begin{aligned} \langle \vec{k}_0 | \mathbf{S}_2^\dagger \mathbf{S}_1 | \vec{k}_0 \rangle &= 1 + i \frac{8\pi \Delta x k_0^5 \tilde{a}^6}{3L^2} \cos \Theta - \frac{2\pi \Delta x^2 k_0^6 \tilde{a}^6}{15L^2} \\ &\times (3 + 11 \cos^2 \Theta) + O\left(\frac{(k_0 \Delta x)^3}{L^2}\right), \end{aligned} \quad (\text{A20})$$

where Θ is the angle between the incoming direction \vec{k}_0 and the displacement vector $\Delta x \equiv \vec{x}_2 - \vec{x}_1$ and $\tilde{a} \equiv a[(\epsilon - 1)/(\epsilon + 2)]^{1/3}$. This implies that

$$\begin{aligned} |\text{Tr} \mathbf{S}_1 \varrho_0^{\text{ph}} \mathbf{S}_2^\dagger|^{(1-f)N_i} &= |\langle \vec{k}_0 | \mathbf{S}_2^\dagger \mathbf{S}_1 | \vec{k}_0 \rangle|^{(1-f)N_i} \\ &\cong \left[1 - \frac{2\pi \Delta x^2 k_0^6 \tilde{a}^6}{15L^2} (3 + 11 \cos^2 \Theta) \right]^{L^2(1-f)(N/V)ct} \end{aligned} \quad (\text{A21})$$

$$\xrightarrow{\text{therm}} e^{-(1-f)(t/\tau_D)}. \quad (\text{A22})$$

In the second line above we used Eq. (A20) up to the leading order in $1/L$; in the last line we removed the box normalization through the thermodynamical limit (A2) and thus obtained the decoherence time [11,38]

$$\tau_D^{-1} \equiv \frac{2\pi}{15} \frac{N}{V} \Delta x^2 c k_0^6 \tilde{a}^6 (3 + 11 \cos^2 \Theta). \quad (\text{A23})$$

Equations (A19) and (A22) imply that $\|\varrho_{S:fE}^{i \neq j}(t)\|_{\text{tr}} \leq 2e^{-(1-f)t/\tau_D} |\langle \vec{x}_1 | \varrho_0^S | \vec{x}_2 \rangle|$ since the sequence $(1 + x/N)^N$ is monotonically increasing. As a result, whenever we forget a macroscopic fraction of the environment ($f < 1$), the resulting coherent part $\varrho_{S:fE}^{i \neq j}(t)$ decays in the trace norm exponentially, with the characteristic time $\tau_D/(1-f)$. This completes the first step (A12).

The asymptotic orthogonalization (A13) is also straightforward to show in the case of pure environments. The postscattering states of the environment macrofractions [Eq. (A9)] are all pure:

$$\varrho_i^{\text{mac}}(t) = (\mathbf{S}_i | \vec{k}_0 \rangle \langle \vec{k}_0 | \mathbf{S}_i^\dagger)^{\otimes m N_i} \equiv |\Psi_i^{\text{mac}}(t)\rangle \langle \Psi_i^{\text{mac}}(t)|, \quad (\text{A24})$$

so it is enough to consider their overlap (see Fig. 2)

$$\langle \Psi_2^{\text{mac}}(t) | \Psi_1^{\text{mac}}(t) \rangle = |\langle \vec{k}_0 | \mathbf{S}_2^\dagger \mathbf{S}_1 | \vec{k}_0 \rangle|^{L^2 m(N/V)ct} \quad (\text{A25})$$

$$\xrightarrow{\text{therm}} e^{-m(t/\tau_D)}. \quad (\text{A26})$$

Thus, for $t \gg \tau_D$ the states of the macrofractions $\Psi_i^{\text{mac}}(t)$ asymptotically orthogonalize and moreover on the same time scale τ_D as the decay of the coherent part described by Eq. (A26) [note that $0 < m$ and $f \leq 1$ so the time scales from Eqs. (A22) and (A26) do not differ considerably]. This shows the asymptotic formation of the broadcast state (1) with pure encoding states $\varrho_i^{E_k}$:

$$\begin{aligned} \varrho_{S:fE}(0) &= \varrho_0^S \otimes \underbrace{\varrho_0^{\text{mac}} \otimes \dots \otimes \varrho_0^{\text{mac}}}_{fM} \xrightarrow[t \gg \tau_D]{\text{therm}} \varrho_{S:fE}(\infty) \\ &= \sum_{i=1,2} p_i |\vec{x}_i\rangle \langle \vec{x}_i| \otimes \underbrace{|\dot{i}^{\text{mac}}\rangle \langle \dot{i}^{\text{mac}}| \otimes \dots \otimes |\dot{i}^{\text{mac}}\rangle \langle \dot{i}^{\text{mac}}|}_{fM}, \end{aligned} \quad (\text{A27})$$

Microscopic

Macroscopic
($t \gg \tau_D$)




FIG. 2. (Color online) Orthogonalization of macroscopic states. At the microscopic level, the individual postscattering states $|\Psi_i^{\text{mic}}\rangle \equiv \mathbf{S}_i |\vec{k}_0\rangle$, corresponding to the sphere being at \vec{x}_i (represented by the small solid slabs on the left) become identical in the thermodynamic limit [see Eq. (A30)] and hence completely indistinguishable. They carry a vanishingly small amount of information about the sphere's localization, which is due to the assumed weak coupling between the sphere and each individual environmental photon (A1). On the other hand, the collective states of macroscopic fractions $|\Psi_i^{\text{mac}}(t)\rangle \equiv (\mathbf{S}_i |\vec{k}_0\rangle)^{\otimes m N_i}$ (represented by the big solid slabs on the right) become by Eq. (A26) more and more distinguishable in the thermodynamic (A2) and the deep decoherence $t \gg \tau_D$ limits. Together with the decoherence mechanism (A12), this leads to the formation of the spectrum broadcast state (1) with pure environmental states and hence to the objective existence of the (classical) state of the sphere.

where p_i is given by Eq. (A15) and $|\dot{i}^{\text{mac}}\rangle \equiv |\Psi_i^{\text{mac}}(\infty)\rangle$ emerges as the nondisturbing environmental basis in the space of each macrofraction, spanning a two-dimensional subspace, which carries the correlation between the macrofraction and the sphere (this basis depends on the initial state $|\vec{k}_0\rangle$). Thus, the correlations become effectively among the qubits. The full process (A27) is a combination of the measurement of the system in the pointer basis $|\vec{x}_i\rangle$ and spectrum broadcasting of the result, described by a CC-type channel [23]

$$\Lambda_\infty^{S \rightarrow fE}(\varrho_0^S) \equiv \sum_i |\langle \vec{x}_i | \varrho_0^S | \vec{x}_i \rangle| |\dot{i}^{\text{mac}}\rangle \langle \dot{i}^{\text{mac}}| \otimes fM. \quad (\text{A28})$$

The entropic objectivity condition and the classical plateau follow now from Eq. (A27),

$$I[\varrho_{S:fE}(t)] \approx H_S, \quad (\text{A29})$$

because of the conditions (A12) and (A14) (see the next section for details). Thus the mutual information becomes asymptotically independent of the fraction f (as long as it is macroscopic).

In quantum Darwinism simulations for finite fixed times t (see, e.g., Refs. [11,38]), one can observe that the formation of the plateau is more strongly driven by increasing the time rather than the macrofraction f (keeping all other parameters equal). This can be straightforwardly explained by looking at Eqs. (A22) and (A26): The fractions f and m are by definition at most 1 and hence have little effect on the decay of the exponential factors, while t can be arbitrarily greater than τ_D , thus accelerating the formation of the broadcast state (A27).




FIG. 3. (Color online) Information-theoretic phases of the sphere model (see [11]). Schematic phase diagram, showing three different phases of the illuminated sphere model, appearing in the thermodynamic and the deep decoherence $t \gg \tau_D$ limits. The horizontal axis is the observed fraction f of the total photon number, understood modulo a microfraction. The vertical axis is the asymptotic mutual information between the sphere S and the fraction fE , $I[\rho_{S:fE}(\infty)]$. This is the limiting diagram for those from [11] obtained for finite t . There are two phase transitions: at $f = 0$ from the singular product phase (represented by the black point at zero) to the broadcasting phase (the black line at H_S) and at $f = 1$ from the broadcasting phase to the singular full information phase (black dot at I_{\max}).

3. Information-theoretic phases

There is a very distinct difference in the macroscopic and microscopic behaviors of the environment, already alluded to in Refs. [11,38] and summarized in Fig. 3. From Eq. (A20) it follows that within the sector (A1) the postscattering states of individual photons (microstates) $|\Psi_i^{\text{mic}}\rangle \equiv \mathbf{S}_i|\vec{k}_0\rangle$ become identical in the thermodynamic limit and hence encode no information about the sphere’s localization:

$$\langle \Psi_2^{\text{mic}} | \Psi_1^{\text{mic}} \rangle \equiv \langle \vec{k}_0 | \mathbf{S}_2^\dagger \mathbf{S}_1 | \vec{k}_0 \rangle \xrightarrow{\text{therm}} 1. \quad (\text{A30})$$

This is not surprising due to the condition (A1). On the other hand, despite (A1), by Eq. (A26) macroscopic groups of photons are able to resolve the sphere’s position and in the asymptotic limit resolve it perfectly. This leads to the appearance of the different information-theoretic phases in the model, which we now describe. We stress that the macrofraction m can be arbitrarily small [which only prolongs the orthogonalization time; cf. Eq. (A26)], but must scale with the total number of photons N_t . Indeed, for a microscopic, i.e., not scaling with N_t , fraction μ the limit (A30) still holds: $[\langle \vec{k}_0 | \mathbf{S}_2^\dagger \mathbf{S}_1 | \vec{k}_0 \rangle]^\mu \xrightarrow{\text{therm}} 1$. Thus, if the observed portion of the environment is microscopic, the asymptotic postscattering state is in fact a product one:

$$\begin{aligned} \rho_{S:\mu E}(0) &= \rho_0^S \otimes (\rho_0^{\text{mac}})^{\otimes \mu} \xrightarrow[t \gg \tau_D]{\text{therm}} \rho_{S:\mu E}(\infty) \\ &= \sum_{i=1,2} p_i |\vec{x}_i\rangle \langle \vec{x}_i| \otimes (\mathbf{S}_i |\vec{k}_0\rangle \langle \vec{k}_0| \mathbf{S}_i^\dagger)^{\otimes \mu} \end{aligned} \quad (\text{A31})$$

$$= \left(\sum_{i=1,2} p_i |\vec{x}_i\rangle \langle \vec{x}_i| \right) \otimes |\Psi^{\text{mic}}\rangle \langle \Psi^{\text{mic}}|^{\otimes \mu}, \quad (\text{A32})$$

where $|\Psi^{\text{mic}}\rangle \equiv \mathbf{S}_1|\vec{k}_0\rangle \cong \mathbf{S}_2|\vec{k}_0\rangle$ because of Eq. (A30) [\cong denotes equality in the thermodynamic limit (A2)]. This is the product phase, in which $I[\rho_{S:\mu E}(\infty)] = 0$.

Conversely, if we have access to the full environment, ignoring perhaps only a microscopic fraction μ , the arguments leading to Eqs. (A22) and (A26) do not work anymore, since from Eq. (A30)

$$|\text{Tr} \mathbf{S}_1 \rho_0^{\text{ph}} \mathbf{S}_2^\dagger|^\mu \xrightarrow{\text{therm}} 1 \quad (\text{A33})$$

and thus there is no decoherence or orthogonalization. The postscattering state contains then the full quantum information about the system due to the unsuppressed system-environment entanglement produced by the controlled-unitary interaction (A4). As a result, the mutual information attains in the thermodynamical limit its maximum value $I_{\max} = 2H_S$ [for a pure ρ_0^S , since the interaction is of a controlled-unitary form (A4)] and this defines the full information phase. We note that the rise of $I_{S:fE}$ above H_S certifies the presence of entanglement [43]. The intermediate phase described by Eq. (A27) is the broadcasting phase (see Fig. 3).

The quantity experiencing discontinuous jumps is the mutual information between the system S and the observed environment fE and the parameter that drives the phase transitions is the fraction size f . As discussed above, each value of f has to be understood modulo a microfraction. The appearance of the phase diagram is a reflection of both the thermodynamic and the deep decoherence limits and its form is in agreement with the previously obtained results (see, e.g., Refs. [11,38]).

4. Derivation of the entropic objectivity condition in the illuminated sphere model

Here we present an independent derivation of the entropic objectivity condition

$$I[\rho_{S:fE}(t)] \approx H_S \quad (\text{A34})$$

for the illuminated sphere model. Although illustrated on a concrete model, our derivation is indeed more general: Instead of a direct asymptotic calculation of the mutual information $I[\rho_{S:fE}(t)]$ in the model (cf. Refs. [9,11,38]), we will show that Eq. (A34) follows from the mechanisms of (i) decoherence [Eq. (A12)] and (ii) distinguishability [Eq. (A14)], once they are proven. In light of our findings, this gives a clear physical meaning to Eq. (A34): It is a consequence of the state information broadcasting. Most of the proof is for general mixed states.

Let the postinteraction $S:fE$ state for a fixed finite box L and time t be $\rho_{S:fE}(L,t)$. It is given by Eqs. (A10) and (A11) and now we explicitly indicate the dependence on L in the notation. Then

$$|H_S - I[\rho_{S:fE}(L,t)]| \leq |I[\rho_{S:fE}(L,t)] - I[\rho_{S:fE}^{i=j}(L,t)]| \quad (\text{A35})$$

$$+ |H_S - I[\rho_{S:fE}^{i=j}(L,t)]|, \quad (\text{A36})$$

where $\rho_{S:fE}^{i=j}(L,t)$ is the decohered part of $\rho_{S:fE}(L,t)$, given by Eq. (A10). We first bound the difference (A35), decomposing the mutual information using conditional information

$$S_{\text{VN}}(\varrho_{S:fE}|\varrho_{fE}) \equiv S_{\text{VN}}(\varrho_{S:fE}) - S_{\text{VN}}(\varrho_{fE}):$$

$$I(\varrho_{S:fE}) = S_{\text{VN}}(\varrho_S) - S_{\text{VN}}(\varrho_{S:fE}|\varrho_{fE}), \quad (\text{A37})$$

so that

$$|I[\varrho_{S:fE}(L,t)] - I[\varrho_{S:fE}^{i=j}(L,t)]| \leq |S_{\text{VN}}[\varrho_S(L,t)] - S_{\text{VN}}[\varrho_S^{i=j}(L,t)]| \quad (\text{A38})$$

$$+ |S_{\text{VN}}[\varrho_{S:fE}(L,t)|\varrho_{fE}(L,t)] - S_{\text{VN}}[\varrho_{S:fE}^{i=j}(L,t)|\varrho_{fE}^{i=j}(L,t)]|. \quad (\text{A39})$$

From Eq. (A1), the total $S:fE$ Hilbert space is finite dimensional for a finite L,t : There are $fN_i = fL^2(N/V)ct$ photons [cf. Eq. (A3)] and the number of modes of each photon is approximately $(4\pi/3)(L/2\pi\Delta x)^3$. Hence, the total dimension is $2 \times L^2 f(N/V)ct \times (1/6\pi^2)(L/\Delta x)^3 < \infty$ and we can use the Fannes-Audenaert [31] and the Alicki-Fannes [32] inequalities to bound (A38) and (A39), respectively (cf. Ref. [9]). For (A38) we obtain

$$|S_{\text{VN}}[\varrho_S(L,t)] - S_{\text{VN}}[\varrho_S^{i=j}(L,t)]| \leq \frac{1}{2}\epsilon_E(L,t) \log(d_S - 1) + h\left[\frac{\epsilon_E(L,t)}{2}\right], \quad (\text{A40})$$

where $h(\epsilon) \equiv -\epsilon \log \epsilon - (1 - \epsilon) \log(1 - \epsilon)$ is the binary Shannon entropy and

$$\epsilon_E(L,t) \equiv \|\varrho_S(L,t) - \varrho_S^{i=j}(L,t)\|_{tr} \quad (\text{A41})$$

$$= \|\varrho_S^{i \neq j}(L,t)\|_{tr} \cong 2|c_{12}| \times \left[1 - \frac{1}{c\tau_D L^2} \left(\frac{N}{V}\right)^{-1}\right]^{L^2(N/V)ct} \quad (\text{A42})$$

with $c_{12} \equiv \langle \vec{x}_1 | \varrho_0^S | \vec{x}_2 \rangle$, where we have used the reasoning (A17)–(A22), but with $f = 0$. For (A39) the same reasoning and the Alicki-Fannes inequality give

$$|S_{\text{VN}}[\varrho_{S:fE}(L,t)|\varrho_{fE}(L,t)] - S_{\text{VN}}[\varrho_{S:fE}^{i=j}(L,t)|\varrho_{fE}^{i=j}(L,t)]| \leq 4\epsilon_{fE}(L,t) \log d_S + 2h[\epsilon_{fE}(L,t)], \quad (\text{A43})$$

with

$$\epsilon_{fE}(L,t) \equiv \|\varrho_{S:fE}(L,t) - \varrho_{S:fE}^{i=j}(L,t)\|_{tr} \quad (\text{A44})$$

$$= \|\varrho_{S:fE}^{i \neq j}(L,t)\|_{tr} \quad (\text{A45})$$

$$\cong 2|c_{12}| \left[1 - \frac{1}{c\tau_D L^2} \left(\frac{N}{V}\right)^{-1}\right]^{L^2(1-f)(N/V)ct}. \quad (\text{A46})$$

Above L and t are big enough so that $\epsilon_E(L,t), \epsilon_{fE}(L,t) < 1$. Equations (A38)–(A46) give an upper bound on the difference (A35) in terms of the decoherence speed (A12).

To bound the orthogonalization part (A36) (see Ref. [9] for a related analysis), we note that since $\varrho_{S:fE}^{i=j}(L,t)$ is a CQ state [cf. Eq. (A10)], its mutual information is given by the Holevo quantity [44]

$$I[\varrho_{S:fE}^{i=j}(L,t)] = \chi\{p_i, \varrho_i^{\text{mac}}(t)^{\otimes fM}\}, \quad (\text{A47})$$

where p_i is given by Eq. (A15). From the Holevo theorem it is bounded by [44]

$$I_{\text{max}}(t) \leq \chi\{p_i, \varrho_i^{\text{mac}}(t)^{\otimes fM}\} \leq H(\{p_i\}) \equiv H_S, \quad (\text{A48})$$

where $I_{\text{max}}(t) \equiv \max_{\mathcal{E}} I[p_i \pi_{j|i}^{\mathcal{E}}(t)]$ is the fixed-time maximal mutual information, extractable through generalized measurements $\{\mathcal{E}_j\}$ on the ensemble $\{p_i, \varrho_i^{\text{mac}}(t)^{\otimes fM}\}$, and the conditional probabilities read

$$\pi_{j|i}^{\mathcal{E}}(t) \equiv \text{Tr}[\mathcal{E}_j \varrho_i^{\text{mac}}(t)^{\otimes fM}] \quad (\text{A49})$$

(here and below i labels the states and j the measurement outcomes). We now relate $I_{\text{max}}(t)$ to the generalized overlap $B[\varrho_1^{\text{mac}}(t)^{\otimes fM}, \varrho_2^{\text{mac}}(t)^{\otimes fM}]$ [cf. Eq. (A14)], which we have calculated for pure states in Eqs. (A25) and (A26). Using the method of Ref. [42], slightly modified to unequal *a priori* probabilities p_i , we obtain for an arbitrary measurement \mathcal{E}

$$I(\pi_{j|i}^{\mathcal{E}} p_i) = I(\pi_{i|j}^{\mathcal{E}} \pi_j^{\mathcal{E}}) = H(\{p_i\}) - \sum_{j=1,2} \pi_j^{\mathcal{E}} h(\pi_{i|j}^{\mathcal{E}}) \quad (\text{A50})$$

$$\geq H(\{p_i\}) - 2 \sum_{j=1,2} \pi_j^{\mathcal{E}} \sqrt{\pi_{i|j}^{\mathcal{E}} (1 - \pi_{i|j}^{\mathcal{E}})} \quad (\text{A51})$$

$$= H(\{p_i\}) - 2\sqrt{p_1 p_2} \sum_{j=1,2} \sqrt{\pi_{j|1}^{\mathcal{E}} \pi_{j|2}^{\mathcal{E}}}, \quad (\text{A52})$$

where we used first the Bayes theorem $\pi_{i|j}^{\mathcal{E}} = (p_i/\pi_j^{\mathcal{E}})\pi_{j|i}^{\mathcal{E}}$, $\pi_j^{\mathcal{E}} \equiv \sum_i \pi_{j|i}^{\mathcal{E}} p_i = \text{Tr}(\mathcal{E}_j \sum_i \varrho_i)$, then the fact that we have only two states $\pi_{2|j}^{\mathcal{E}} = 1 - \pi_{1|j}^{\mathcal{E}}$, so that $H(\pi_{i|j}^{\mathcal{E}}) = h(\pi_{i|j}^{\mathcal{E}})$, and finally $h(p) \leq 2\sqrt{p(1-p)}$. On the other hand, $B(\varrho_1, \varrho_2) = \min_{\mathcal{E}} \sum_j \sqrt{\pi_{j|1}^{\mathcal{E}} \pi_{j|2}^{\mathcal{E}}}$ [42]. Denoting the optimal measurement by $\mathcal{E}_*^B(t)$ and recognizing that $H(\{p_i\}) = H_S$, we obtain

$$I_{\text{max}}(t) \geq I[p_i \pi_{j|i}^{\mathcal{E}_*^B(t)}(t)] \quad (\text{A53})$$

$$\geq H_S - 2\sqrt{p_1 p_2} B[\varrho_1^{\text{mac}}(t)^{\otimes fM}, \varrho_2^{\text{mac}}(t)^{\otimes fM}] \quad (\text{A54})$$

$$= H_S - 2\sqrt{p_1 p_2} B[\varrho_1^{\text{mac}}(t), \varrho_2^{\text{mac}}(t)]^{fM}. \quad (\text{A55})$$

Inserting the above into the bounds (A48) gives the desired upper bound on the difference (A36):

$$|H_S - I[\varrho_{S:fE}^{i=j}(L,t)]| \leq 2\sqrt{p_1 p_2} B[\varrho_1^{\text{mac}}(t), \varrho_2^{\text{mac}}(t)]^{fM}, \quad (\text{A56})$$

where the generalized overlap is given by Eqs. (A25) and (A26):

$$B[\varrho_1^{\text{mac}}(t), \varrho_2^{\text{mac}}(t)] = |\langle \Psi_2^{\text{mac}}(t) | \Psi_1^{\text{mac}}(t) \rangle| \cong \left[1 - \frac{1}{c\tau_D L^2} \left(\frac{N}{V}\right)^{-1}\right]^{L^2 m(N/V)ct}. \quad (\text{A57})$$

Gathering all the above facts together finally leads to a bound on $|H_S - I[\varrho_{S:fE}(L,t)]|$ in terms of the speed of (i)

decoherence (A12) and (ii) distinguishability (A14):

$$|H_S - I[\varrho_{S:fE}(L,t)]| \leq h \left[\frac{\epsilon_E(L,t)}{2} \right] + 2h[\epsilon_{fE}(L,t)] \quad (A58)$$

$$+ 4\epsilon_{fE}(L,t) \log 2 + 2\sqrt{p_1 p_2} B[\varrho_1^{\text{mac}}(t), \varrho_2^{\text{mac}}(t)]^{fM}, \quad (A59)$$

where $\epsilon_E(L,t)$, $\epsilon_{fE}(L,t)$, and $B[\varrho_1^{\text{mac}}(t), \varrho_2^{\text{mac}}(t)]$ are given by Eqs. (A42), (A46), and (A57), respectively. Choosing L and t big enough so that $\epsilon_E(L,t), \epsilon_{fE}(L,t) \leq 1/2$ [when the binary entropy $h(\cdot)$ is monotonically increasing], we remove the unphysical box and obtain an estimate on the speed of convergence of $I[\varrho_{S:fE}(L,t)]$ to H_S :

$$\lim_{L \rightarrow \infty} |H_S - I[\varrho_{S:fE}(L,t)]| \leq h(|c_{12}|e^{-(t/\tau_D)}) \quad (A60)$$

$$+ 2h(2|c_{12}|e^{-(1-f)(t/\tau_D)} + 8|c_{12}|e^{-(1-f)(t/\tau_D)}) \log 2 \quad (A61)$$

$$+ 2\sqrt{p_1 p_2} e^{-f(t/\tau_D)}. \quad (A62)$$

This finishes the derivation of the condition (A34).

We note that the result (A58) and (A59) is in fact a general statement, valid in any model where (i) the system S is

effectively a qubit and (ii) the system-environment interaction is of a environment-symmetric controlled-unitary type.

Lemma. Let a two-dimensional quantum system S interact with N identical environments, each described by a d -dimensional Hilbert space, through a controlled-unitary interaction

$$U(t) \equiv \sum_{i=1,2} |i\rangle\langle i| \otimes U_i(t)^{\otimes N}. \quad (A63)$$

Let the initial state be $\varrho_{S:E}(0) = \varrho_0^S \otimes (\varrho_0^E)^{\otimes N}$ and $\varrho_{S:E}(t) \equiv U(t)\varrho_{S:E}(0)U(t)^\dagger$. Then for any $0 < f < 1$ and t big enough

$$|H(\{p_i\}) - I[\varrho_{S:fE}(t)]| \leq h \left[\frac{\epsilon_E(t)}{2} \right] + 2h[\epsilon_{fE}(t)] \quad (A64)$$

$$+ 4\epsilon_{fE}(t) \log 2 + 2\sqrt{p_1 p_2} B[\varrho_1(t), \varrho_2(t)]^{fN}, \quad (A65)$$

where

$$p_i \equiv \langle i|\varrho_0^S|i\rangle, \quad \varrho_i(t) \equiv U_i(t)\varrho_0^E U_i(t)^\dagger, \quad (A66)$$

$$\epsilon_E(t) \equiv \|\varrho_S(t) - \varrho_S^{i=j}\|_{\text{tr}}, \quad (A67)$$

$$\epsilon_{fE}(t) \equiv \|\varrho_{S:fE}(t) - \varrho_{S:fE}^{i=j}(t)\|_{\text{tr}}. \quad (A68)$$

[1] N. Bohr, in *Albert Einstein: Philosopher-Scientist*, edited by P. A. Schilpp, The Library of Living Philosophers Vol. VII (Open Court, La Salle, 1949).

[2] W. Heisenberg, *Philosophic Problems in Nuclear Science*, translated by F. C. Hayes (Faber and Faber, London, 1952).

[3] E. Joos *et al.*, *Decoherence and the Appearance of a Classical World in Quantum Theory* (Springer, Berlin, 2003).

[4] H. D. Zeh, in *Quantum Decoherence*, edited by B. Duplantier, J.-M. Raimond, and V. Rivasseau (Birkhäuser, Basel, 2006).

[5] M. Schlosshauer, *Decoherence and the Quantum-to-Classical Transition* (Springer, Berlin, 2007).

[6] W. H. Zurek, *Phys. Rev. D* **24**, 1516 (1981).

[7] W. H. Zurek, *Nat. Phys.* **5**, 181 (2009).

[8] W. Wootters and W. H. Zurek, *Nature (London)* **299**, 802 (1982).

[9] M. Zwolak and W. H. Zurek, *Sci. Rep.* **3**, 1729 (2013).

[10] M. Zwolak, H. T. Quan, and W. H. Zurek, *Phys. Rev. A* **81**, 062110 (2010).

[11] C. J. Riedel and W. H. Zurek, *Phys. Rev. Lett.* **105**, 020404 (2010).

[12] F. G. S. L. Brandao, M. Piani, and P. Horodecki, [arXiv:1310.8640](https://arxiv.org/abs/1310.8640).

[13] T. Y. Cao, *Conceptual Developments of 20th Century Field Theories* (Cambridge University Press, Cambridge, 1997).

[14] N. Bohr, *Phys. Rev.* **48**, 696 (1935).

[15] H. M. Wiseman, *Ann. Phys. (NY)* **338**, 361 (2013).

[16] A. Einstein, B. Podolsky, and N. Rosen, *Phys. Rev.* **47**, 777 (1935).

[17] E. Joos and H. D. Zeh, *Z. Phys. B* **59**, 223 (1985).

[18] J. K. Korbicz, P. Horodecki, and R. Horodecki, *Phys. Rev. Lett.* **112**, 120402 (2014).

[19] R. Blume-Kohout and W. H. Zurek, *Phys. Rev. A* **73**, 062310 (2006).

[20] H. Ollivier, D. Poulin, and W. H. Zurek, *Phys. Rev. Lett.* **93**, 220401 (2004).

[21] J. Oppenheim, M. Horodecki, P. Horodecki, and R. Horodecki, *Phys. Rev. Lett.* **89**, 180402 (2002).

[22] M. Piani, P. Horodecki, and R. Horodecki, *Phys. Rev. Lett.* **100**, 090502 (2008).

[23] J. K. Korbicz, P. Horodecki, and R. Horodecki, *Phys. Rev. A* **86**, 042319 (2012).

[24] W. H. Zurek, *Phys. Rev. A* **87**, 052111 (2013).

[25] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, *Rev. Mod. Phys.* **84**, 1655 (2012).

[26] P. Horodecki and R. Horodecki, *Quantum Inform. Comput.* **1**, 45 (2001).

[27] H. Barnum, C. M. Caves, C. A. Fuchs, R. Jozsa, and B. Schumacher, *Phys. Rev. Lett.* **76**, 2818 (1996).

[28] T. M. Cover and J. A. Thomas, *Elements of Information Theory* (Wiley, New York, 1991).

[29] M. Horodecki, J. Oppenheim, and A. Winter, *Nature (London)* **436**, 673 (2005).

[30] A. Peres, *Phys. Rev. Lett.* **77**, 1413 (1996); M. Horodecki, P. Horodecki, and R. Horodecki, *Phys. Lett. A* **223**, 1 (1996).

[31] K. M. R. Audenaert, *J. Phys. A: Math. Theor.* **40**, 8127 (2007).

[32] R. Alicki and M. Fannes, *J. Phys. A: Math. Gen.* **37**, L55 (2004).

[33] *Quantum State Estimation*, edited by M. Paris and J. Řeháček, Lecture Notes in Physics Vol. 649 (Springer, Berlin, 2004).

[34] M. F. Pusey, J. Barrett, and T. Rudolph, *Nat. Phys.* **8**, 476 (2012).

[35] J. Bell, *Physics* **1**, 195 (1964).

- [36] E. P. Wigner, in *The Logic of Personal Knowledge: Essays Presented to Michael Polanyi on his Seventieth Birthday* (Routledge & Kegan Paul, London, 1961).
- [37] M. R. Gallis and G. N. Fleming, *Phys. Rev. A* **42**, 38 (1990).
- [38] C. J. Riedel and W. H. Zurek, *New J. Phys.* **13**, 073038 (2011).
- [39] K. Hornberger and J. E. Sipe, *Phys. Rev. A* **68**, 012105 (2003).
- [40] S. L. Adler, *J. Phys. A* **39**, 14067 (2006).
- [41] G. Sewell, *Rep. Math. Phys.* **56**, 271 (2005).
- [42] C. A. Fuchs and J. van de Graaf, *IEEE Trans. Inf. Theor.* **45**, 1216 (1999).
- [43] R. Horodecki and P. Horodecki, *Phys. Lett. A* **194**, 147 (1994).
- [44] A. S. Holevo, *Probl. Inform. Transm.* **9**, 177 (1973).