


Fusion of DNA-binding domain of *Pyrococcus furiosus* ligase with *Taq*Stoffel DNA polymerase as a useful tool in PCR with difficult targets

Marta Śpibida¹ · Beata Krawczyk¹ · Beata Zalewska-Piątek¹ · Rafał Piątek¹ · Magdalena Wysocka¹ · Marcin Olszewski¹ 

Received: 17 May 2017 / Revised: 28 August 2017 / Accepted: 27 September 2017
© The Author(s) 2017. This article is an open access publication

Abstract The DNA coding sequence of *Taq*Stoffel polymerase was fused with the DNA-binding domain of *Pyrococcus furiosus* ligase. The resulting novel recombinant gene was cloned and expressed in *E. coli*. The recombinant enzyme was purified and its enzymatic features were studied. The fusion protein (*Pfu*DBDlig-*Taq*S) was found to have enhanced processivity as a result of the conversion of the *Taq*DNA polymerase from a relatively low processive to a highly processive enzyme. The abovementioned processivity enhancement was about threefold as compared to the recombinant *Taq*Stoffel DNA polymerase (*Taq*S), and the recombinant fusion protein was more thermostable. It had a half-life of 23 min at 99 °C as compared to 10 min for *Taq*S. The fusion protein also showed a significantly higher resistance to PCR inhibitors such as heparin or lactoferrin and the fusion polymerase-amplified GC-rich templates much more efficiently and was efficient even with 78% GC pairs.

Keywords PCR · DNA polymerase · Fusion DNA polymerases · Archaeon

Introduction

At the core of PCR-based techniques is the use of a DNA polymerase which plays the major role in molecular biological applications including DNA amplification or sequencing

(Terpe 2013). PCR requires a high temperature, and therefore, it is necessary to use a thermostable enzyme for DNA amplification; the most commonly used enzyme is a DNA polymerase derived from the thermophilic *Thermus aquaticus*. Its native form is produced in *Thermus aquaticus* and its recombinant form by *Escherichia coli*. The *Taq*'s half-life is short compared to that of the other polymerases, e.g., these isolated from Archaea: only 40 min at 95 °C or 9 min at 97.5 °C (Lawyer et al. 1993). Furthermore, the *Taq* DNA polymerase is not able to synthesize products which are longer than 4 kb (Hamilton et al. 2001). In general, *Taq* amplification efficiencies for targets shorter than 1 kb, with 45 to 56% CG content, lie at about 80% (Arezi et al. 2003). Usually, product yields decrease when the amplicon size becomes greater than 1 kb because of a relatively low processivity and thermostability of a wild-type enzyme. A novel strategy used to enhance the processivity and performance of the *Taq* is to fuse it with a thermostable DNA-binding protein such as a Sso7d DNA-binding protein derived from *Sulfolobus solfataricus* (Wang et al. 2004).

We have tried to improve the properties of the *Taq* polymerase by creating a fusion protein which contained a highly thermostable DNA-binding domain of the *Pyrococcus furiosus* ligase, a 6-amino acid linker and a sequence of the *Taq*Stoffel polymerase.

Materials and methods

Construction of recombinant plasmids

The *T. aquaticus* strain (ATCC25104) was used to isolate a genomic DNA which was used as a template for the amplification of a *taq*Stoffel fragment gene with the use of a standard PCR amplification protocol with the Marathon DNA

✉ Marcin Olszewski
molszewski@pg.gda.pl

¹ Department of Molecular Biotechnology and Microbiology, Gdańsk University of Technology, ul. G. Narutowicza 11/12, 80-233 Gdańsk, Poland

polymerase (A&A Biotechnology, Gdansk, Poland). A DNA fragment of *taq*Stoffel corresponding to nucleotides 997 to 2626 was obtained in PCR with the use of primers: 5' AATT TTGTTAACTTTAAGAAGGAGATATACATATGGCCC TGGAGGAGGCC

(forward) and 5' *GCAAGCTTGTGCGACGGAGCT CGAATTCGGATCCTTAatggtggtggtggtggtgCTCC TTGGCGGAGAGCCAG* (reverse). The primers contained sequences complementary to the *taq*Stoffel gene (underlined), a sequence complementary to the pET-30 Ek/LIC vector (italics), and a sequence for the oligohistidine tag (lowercase). The stop codon (TTA) was added to the reverse primer immediately after the oligohistidine sequence.

The DNA-binding domain of *Pyrococcus furiosus* ligase gene was fused with a DNA fragment of *taq*Stoffel (corresponding to nucleotides 997 to 2626 of *taq* gene) in PCR using primers: 5' TATTGGCTTTCGGAAGCGGAGGGGT CGACGCCCTGGAGGAGGCC (forward) and 5' *GCAAGCTTGTGCGACGGAGCTCGAATTCGGATCCTTAatggtggtggtggtggtgCTCCTTGGCGGAGAGCCAG* (reverse). The primers contained sequences complementary to the *taq*Stoffel gene (underlined), a sequence complementary to the pET-30 Ek/LIC vector (italics), a sequence for a 6-amino acid residue linker (bolded), and a sequence for the oligohistidine tag (lowercase). The stop codon (TTA) was added to the reverse primer immediately after the oligohistidine sequence.

The genomic DNA of *Pyrococcus furiosus* (DSM 3638) was used as a template for the amplification of the DNA-binding domain of the ligase gene (accession number: NC_003413) using a standard PCR amplification protocol. The forward primer was 5' ATTTTGTAACTTTAAGAA GGAGATATACATATGCCATTGTGGGCTTTATCTTCAACCA (containing a sequence complementary to the DBD fragment of the ligase gene (underlined), and a sequence complementary to the pET-30 Ek/LIC vector (italics) and the reverse primer was 5'.

TCCCTCCAGGGCGTTCGACCCCTCCGCTTCCGAAAGCCAATAAAGCCAATGCTTGC (containing a sequence complementary to the DBD fragment of the ligase gene (underlined) and a sequence for a 6-amino acid residue linker (bolded). As a result of the PCR amplification, the following two products were obtained: the *taq*Stoffel gene (1703 bp) and the DBD fragment of the ligase gene (685 bp). The PCR products were then mixed together with the DNA of the pET-30 Ek/LIC vector (Novagen, Madison, WI, USA) digested with the use of *Bam*HI and *Nde*I enzymes (NEB, UK), and the mixture was used for cloning with the use of the OverLap Assembly kit (A&A Biotechnology, Poland). The resulting pET30/PfuDBDlig-TaqS plasmid had a reading frame which contained the DBD sequence of *P. furiosus* ligase (amino acid residues 218 to 424), a 6-amino acid linker GSGGVD), a sequence of the *Taq*Stoffel polymerase (amino

acid residues 317 to 832), and a His tag domain which was necessary for the purification of the recombinant protein by metal affinity chromatography.

The nucleotide sequences of the resulting recombinant plasmids, pET30/TaqS and pET30/PfuDBDlig-TaqS, were confirmed by DNA sequencing (Genomed, Poland).

Expression and purification of *Taq*S and *Pfu*DBDlig-*Taq*S polymerases

The pET30/TaqS and pET30/PfuDBDlig-TaqS plasmids were transformed into *E. coli* BL21 (DE3) RIL (Novagen, USA). The cells which carried the required plasmids were grown at 37 °C in the Luria-Bertani medium, supplemented with 50 µg/ml of kanamycin and 50 µg/ml of chloramphenicol to an OD 600 of 0.4 and were induced by incubation in the presence of IPTG, at a final concentration of 1 mM, for 24 h. The cells were then harvested by centrifugation and the pellets were resuspended in 20 ml of buffer A (50 mM Tris-HCl pH 9, 0.5 M NaCl and 5 mM imidazole). The samples were sonicated three times for 30 s at 4 °C and centrifuged for 15 min at 10000×g. The supernatant was heat-treated for 30 min at 70 °C, and the denatured host proteins were removed by centrifugation. The protein was next purified in a one-step procedure using Ni²⁺-affinity chromatography. The supernatant containing the protein was loaded directly into a His•Bind Column (Novagen, USA) prepared and equilibrated using the A buffer. The recombinant proteins were washed twice with the washing buffer B (50 mM Tris-HCl pH 9, 0.5 M NaCl and 40 mM imidazole) and then eluted with the elution buffer E (50 mM Tris-HCl pH 9, 0.5 M NaCl and 300 mM imidazole). The eluted fractions were dialyzed three times against the buffer D (100 mM Tris-HCl pH 8, 100 mM KCl, 0.2 mM EDTA). To remove the trace amounts of the genomic bacterial DNA, 25 U of Benzonase® endonuclease (Merck, Darmstadt, Germany) and MgCl₂ were added at the final concentration of 5 mM and the preparation was incubated for 1 h at 37 °C. Next, Benzonase was inactivated by incubation for 15 min at 70 °C, and following this, the denatured proteins were removed by centrifugation. Finally, the preparations were dialyzed once against the storage buffer (50 mM Tris-HCl pH 8, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 1% Tween 20, 1% Nonidet P-40 and 50% glycerol).

DNA polymerase activity assay

The activity of the DNA polymerase was measured in an isothermal reaction at 72 °C using a real-time PCR apparatus (IT-IS International Ltd., UK) and EvaEZFluorometric Polymerase Activity Assay Kit (Biotium, Hayward, USA) in accordance with the definition of one unit of enzyme

activity by Tveit and Kristensen (“One unit of DNA polymerase activity is conventionally defined as the amount of enzyme that will incorporate 10 nmol of nucleotides during a 30-min incubation”). When the DNA polymerase is active, the primer is extended to form a double-stranded product which binds the EvaGreen dye, with the resulting increase in fluorescence. The rate of increase shows a positive correlation with the polymerase activity (Tveit and Kristensen 2001; Driscoll et al. 2014; Ma et al. 2006). The activity was determined using the commercial *Taq* polymerase (Thermo Scientific, USA) with an activity of 1 U/ μ l as a reference.

Optimization of PCR amplification

To optimize the amplification process, the polymerase activity was measured using various concentrations of $MgCl_2$, KCl and $(NH_4)_2SO_4$ in the buffer and various pHs. All reactions were performed using 1 mM of each dNTP, 0.4 mM of each primer and, as a template for PCR, the pET 30 plasmid DNA containing a known target sequence (PCR product of 300 bp). The PCR experiment was performed using 1 U of purified *Pfu*DBDlig-*TaqS* DNA or *TaqS* DNA polymerase in 20 μ l of the reaction mixture containing 50 ng of the DNA template. The PCR experiment was conducted as follows: 1 min at 94 °C; 25 cycles of 15 s at 94, 55, and 72 °C. To determine the optimum $MgCl_2$ concentration, the PCR was performed at increasing concentrations of $MgCl_2$ (0–9 mM) with the use of the Tris–HCl buffer. Furthermore, the PCR experiment was carried out using various concentrations of KCl (10–90 mM) and $(NH_4)_2SO_4$ (10–90 mM), at various pHs (7.0–9.0), with the use of the Tris–HCl buffer.

Temperature stability was assayed as described by Dabrowski and Kur (1998). One unit of purified *Pfu*DBDlig-*TaqS* and *TaqS* DNA polymerases were heated at 95 and 99 °C for 1, 5, 10, 20, 40, and 60 min. The same amount of enzyme was used to amplify a 300-bp target fragment under the optimal reaction conditions (in the same PCR experiment and under the conditions as determined in the optimization process): 20 mM Tris–HCl (pH 8.0), 4 mM $MgCl_2$, 10 mM $(NH_4)_2SO_4$, and 10 mM KCl.

Measurements of PCR amplification rates

The PCR amplification rates were measured using the method described by Lee et al. 2010, after some modifications. *Pfu*DBDlig-*TaqS* and *TaqS* DNA polymerases were used to amplify the PCR products under the same conditions as those determined in the optimization process and with the use, as a template for PCR, of the pET 30 plasmid DNA containing a known target sequence (PCR product of 300, 500, and 1000 bp). Each PCR cycle started with an initial denaturation

at 94 °C for 2 min and included 25 cycles of 15 s at 94 and 55 °C and 25 cycles of 5 to 60 s at 72 °C.

Processivity

The processivity test was carried out as described by Elshawadfy et al. 2014, with some modifications. Eighty-five microliters of reaction buffer (20 mM Tris–HCl pH 8.3, 10 mM KCl, 10 mM $(NH_4)_2SO_4$, 0.1% Triton X-100), 290 μ M of each of the four dNTPs, 40 nM primer template (5'-GGGGATCCTCTAGAGTCGACCTGC and 5' TATCGGTCCATGAGACAAGCTTGCTTGCCAGCAGGTCGACTCTAGAGGATCCCC), 3 μ l of EvaGreen Fluorescent DNA stain (Jena Bioscience, Jena, Germany), and 1 U of tested polymerase were pre-incubated at 50 °C for 5 min. Reactions were initiated by the simultaneous addition of 7.5 μ l of 50 mM $MgCl_2$ and 7.5 μ l of 0.6 mg/ μ l of a heparin trap, and the polymerization process was allowed to proceed at 72 °C. Aliquots of 10 μ l were withdrawn to cold tubes (4 °C) at 0 (before polymerization process was allowed), 1, 2, 5, and 10 min after polymerization process was allowed and the extension was determined by melting the obtained products with the use of a real-time PCR MyGo instrument (IT-IS International Ltd., GB). The reaction included a pre-melt hold at 95 °C for 10 s (a ramp rate of 5 °C/s), an initial stage at 60 °C for 60 s (a ramp rate of 4 °C/s) and the final stage at 97 °C for 1 s (a ramp rate 0.201 °C/s). The processivity was determined by comparing the melting temperature profiles of the obtained products with the profiles of products with a known length, serving as markers in the reaction. Marker products were obtained in reactions which used the same primer and synthetic templates, which allowed the formation of products which were longer than the primer by 1, 2, 3...22 nt.

Sensitivity

Polymerase sensitivity (affinity for the template) was measured using a PCR in accordance with the protocol described by Halley and Prezioso (2003), after some modifications. The PCR was conducted under conditions optimized for the fusion polymerases *Pfu*DBDlig-*TaqS* and *TaqS*. Plasmid DNA pET28a(+) (Novagen, Madison, USA) was used as a template along with the primers GATGCTGCTGGCTACCCTG (forward) and TCAAGAACTCTGTAGCACCGC (reverse), and the product of the reaction was 1264 bp. The reaction was conducted at decreasing template concentrations (serial two-fold dilutions of template) and the reaction included an initial denaturation at 94 °C for 2 min followed by 25 cycles of 15 s at 94 and 55 °C and of 60 s at 72 °C. The amplified products were analyzed in a 1.2 % agarose gel stained with ethidium bromide.

Resistance to inhibitors

The effect of PCR inhibitors such as lactoferrin (Sigma-Aldrich, St. Louis, USA) in a range from 54 to 0.84 μg , heparin (Sigma-Aldrich, St. Louis, USA) in a range from 600 to 4.7 ng, and human blood (from a healthy individual) in a range from 10 to 0.15%, on the catalytic activity of the *Pfu*DBDlig-*TaqS* and *TaqS* DNA polymerases, was assessed by a PCR reaction using the genomic DNA of *Staphylococcus aureus* as a template and primers for the specific *nuc* gene detection as described by Barski et al. 1996.

Efficiency of a long-range PCR

The efficiency of a long-range PCR was measured as described by Kwona et al. 2016, after some modifications. The PCR experiment was performed with the use of 1 U of purified *Pfu*DBDlig-*TaqS* DNA polymerase and also with the use of *TaqS* DNA polymerase which were placed in a 20 μl reaction mixture containing 50 ng of the *E. coli* genomic DNA as a template. Primers were used to amplify the following four DNA fragments: 5 kbp (5' GCACCATC AACATAAAGGCGC and 5' TTCCGCTAATGCCA TGGTGATAG), 8 kbp (5' GCACCATCAACAAT AAAGGCGC and 5' AACGATGCGATATAGCCGAC AC), and 10 kbp (5' GCACCATCAACAATAAAGGCGC and 5' AACGATGCGATATAGCCGACAC). The PCR experiment was conducted as follows: 2 min at 94 $^{\circ}\text{C}$; 30 cycles of 30 s at 94 and 56 $^{\circ}\text{C}$, and 60 s/kb at 72 $^{\circ}\text{C}$. The amplified products were analyzed in a 1% agarose gel stained with ethidium bromide.

GC-rich templates

The efficiency of the amplification of the products which are rich in GC pairs was evaluated using a GC-rich template of *Mycobacterium tuberculosis*. The use of the forward primer CCGCCGTTACCACCCTTACCACCGTT and the reverse primer GCACCGCACCCACCAGCGGC enabled the production of a target with a length of 301 bp and with a GC content of 78% (Kotłowski 2015). The reaction took place under the conditions which were optimized for the fusion polymerase *Pfu*DBDlig-*TaqS* and reference *TaqS* at an increasing polymerase activity. It started with 3 min at 95 $^{\circ}\text{C}$ followed by 35 cycles of 15 s at 94, 64 and 72 $^{\circ}\text{C}$. The amplified products were analyzed in a 2% agarose gel stained with ethidium bromide.

Results

Expression and purification of *TaqS* and *Pfu*DBDlig-*TaqS* polymerases

The gene encoding a Stoffel fragment of the *Taq* polymerase was cloned into a pET-30 Ek/LIC vector to generate a pET30/*TaqS* plasmid, leading to the expression of the enzyme as a fusion protein with a C-terminal polyhistidine tag. To achieve fusion with the DNA-binding domain of *Pyrococcus furiosus* ligase gene, two PCR products were mixed together with the DNA of the pET-30 Ek/LIC vector DNA to generate the pET30/*Pfu*DBDlig-*TaqS* plasmid coding the fusion enzyme with a C-terminal polyhistidine tag. *E. coli* BL21 (DE3) RIL cultures harboring recombinant plasmids were generated, and cells were harvested and sonicated. The recombinant DNA polymerases were purified by passing the heat-denatured supernatant through a His•BindNi²⁺ affinity column. The specific activities of the purified *TaqS* and *Pfu*DBDlig-*TaqS* polymerases were found to be 1800 and 1600 U/mg, respectively. These results indicated that the *Pfu*DBDlig fusion did not have a negative effect on the catalytic activity of the *TaqS* DNA polymerase. After each purification step, the purity of the DNA polymerase was monitored by SDS–polyacrylamide gel electrophoresis (Fig. 1) which separated the following major protein bands: 62 or 86 kDa for *TaqS* and *Pfu*DBDlig-*TaqS*, respectively; this result was in agreement with the molecular masses of 61.8 or 86.1 kDa which were calculated based on the amino acid sequences. The *E. coli* overexpression system used in this study enabled the production of 30 mg of *TaqS* polymerase and 20 mg of *Pfu*DBDlig-*TaqS* fusion protein per 1 l of induced culture. The production efficiency of the *TaqS* and *Pfu*DBDlig-*TaqS* polymerase in this study was quite satisfactory.

Characterization of the fusion *Pfu*DBDlig-*TaqS* DNA polymerase

In our experiments, the activity of 1 U/ μl was determined for *TaqS* and *Pfu*DBDlig-*TaqS* DNA polymerases and was compared to the commercial *Taq* polymerase with an activity of 1 U/ μl , using the EvaEZFluorometric Polymerase Activity Assay Kit (Biotium, Hayward, USA) in an isothermal reaction at 72 $^{\circ}\text{C}$ on a real-time PCR apparatus (IT-IS International Ltd., UK).

For characterization purposes, the polymerase activity was measured by PCR using various buffer compositions and concentrations of MgCl_2 , KCl, or $(\text{NH}_4)_2\text{SO}_4$ and various pHs (Fig. 2).

MgCl_2 concentration had a strong effect on the activity of *TaqS* and *Pfu*DBDlig-*TaqS*, which was optimal for MgCl_2 equal 2 to 5 mM and 1 to 5 mM, respectively (Fig. 2a).

The DNA polymerase activity was completely inhibited when KCl concentrations exceeded 60 mM for *TaqS* and 30 mM for *Pfu*DBDlig-*TaqS* (Fig. 2b). *TaqS* and




Fig. 1 Expression and purification of the TaqS (a) and the PfuDBDlig-TaqS (b) polymerases. The proteins were analyzed on a 10% polyacrylamide gel (SDS-PAGE). Lane M, unstained protein weight marker 26.610 (14.4–116) (ThermoScientific, USA), with the molecular

mass of proteins marked. Lane 1, sonicated extract of induced cells; lane 2, heat treatment; lane 3, fraction of unbound proteins; lane 4, fraction after second washing with buffer B; lane 5, purified protein after elution with buffer E

PfuDBDlig-TaqS DNA polymerase activities were also strongly affected by $(\text{NH}_4)_2\text{SO}_4$ and were completely inhibited at concentrations of over 30 and 20 mM, respectively (Fig. 2c). The fusion of *PfuDBDlig* to the *TaqS* DNA polymerase resulted in a low tolerance to salt inhibition in PCR.

The effect of pH on the activity of *TaqS* and *PfuDBDlig-TaqS* DNA polymerases was evaluated using buffers with a various pH ranging from 7 to 9. Both polymerases had the highest enzyme activity at pH 8.0 (Fig. 2d).

Based on the results presented in this study, we found that the optimal PCR buffer for the *PfuDBDlig-TaqS* DNA polymerase consisted of 20 mM Tris-HCl (pH 8.0), 4 mM MgCl_2 , 10 mM $(\text{NH}_4)_2\text{SO}_4$, and 10 mM KCl.

Thermostability of *TaqS* and *PfuDBDlig-TaqS* DNA polymerases was tested by measuring the decrease in their activity after preincubation at 95 or 99 °C and was found to be remarkably higher for *PfuDBDlig-TaqS*. *TaqS* and *PfuDBDlig-TaqS* DNA were also found to have a half-life of 10 and 23 min at 99 °C, respectively (Fig. 2e).

PCR amplification rate and processivity

The fusion *PfuDBDlig-TaqS* DNA polymerase was found to replicate template strands at a faster rate than did the *TaqS* DNA polymerase (Fig. 3), namely, it replicated a 300-bp template within 15 s, a 500-bp template within





Fig. 2 Characterization of the fusion *PfuDBDlig-TaqS* DNA polymerase in comparison to the *TaqS* DNA polymerase. The effect of **a** MgCl_2 , **b** KCl, **c** $(\text{NH}_4)_2\text{SO}_4$, **d** pH, and **e** temperature on polymerase activity: gray

95 °C; black 99 °C. Error bars for *PfuDBDlig-TaqS* DNA polymerase have the end bar, for *TaqS* DNA polymerase, do not have the end bar

Fig. 3 Comparison of PCR amplification rates of the fusion *Pfu*DBDlig-*Taq*S DNA polymerase (**a**) and the *Taq*S DNA polymerase (**b**). The elongation times used for PCR amplification are shown at the top. Lane M, DNA molecular size marker (100–1000 bp) (Blirt, Gdańsk, Poland). The amplified products were analyzed in a 1.5% agarose gel stained with ethidium bromide



20 s, and a 1000-bp template within 30 s, while the *Taq*S DNA polymerase required 20 s for a 300-bp template, 35 s for a 500-bp template, and 1 min for a 1000-bp template. This showed that the fusion of the *Pfu*DBDlig protein with the *Taq*S DNA polymerase was two times more efficient and the amplification of DNA required less time.

To determine processivity, the primer template and the polymerase were pre-incubated in the absence of Mg^{2+} and the reaction was initiated by the simultaneous addition of a metal and heparin trap. Next, processivity was determined by comparing the melting temperature profiles of the targets with the profiles of the products of a known length which served as markers in the reaction. In the case of the *Pfu*DBDlig-*Taq*S DNA polymerase, a prominent melting peak is seen resulting from the incorporation of 22 nucleotides and representing processivity (Fig. 4). The control *Taq*S DNA polymerase had a prominent melting peak which indicated lower processivity in the order of 8 nucleotides. This shows that a fusion of the *Taq*S DNA polymerase with the *Pfu*DBDlig strongly improves processivity.

Sensitivity

Sensitivity was measured in a PCR with the use of *Pfu*DBDlig-*Taq*S and *Taq*S polymerases with two-fold serial dilutions of template, and the size of the PCR product was 1264 bp. The experiment showed that the fusion polymerase was more sensitive than the reference polymerase. In the case of the fusion polymerase, it was sufficient to use 0.47 ng of plasmid DNA, while the *Taq*S polymerase required at least 7.5 ng of the plasmid. The results are presented in Fig. 5.

Tolerance of the fusion *Pfu*DBDlig DNA polymerase to PCR inhibitors

The fusion *Pfu*DBDlig-*Taq*S enzyme and the *Taq*S DNA polymerase were PCR-tested for their resistance to serial dilutions of whole human blood, lactoferrin, and heparin, which have been reported to be PCR inhibitors in many publications (Al-Soud and Rådström 1998, 2000, 2001; Yokota et al. 1999; Kermekchiev et al. 2008). It was shown that the fusion *Pfu*DBDlig-*Taq*S DNA polymerase was significantly more resistant to lactoferrin and heparin than the *Taq*Stoffel enzyme and had no resistance to blood. The *Pfu*DBDlig-*Taq*S polymerase remained functional in the presence of 0.84–27.0 μ g of lactoferrin vs. 1.68 μ g for the *Taq*S enzyme (Fig. 6a) and 4.7–150 ng of heparin vs. 9.4 ng for the *Taq*S enzyme (Fig. 6b).




Fig. 4 Determination of processivity by measuring the melting temperature of DNA products in the presence of a heparin trap




Fig. 5 Electrophoretic separation showing the products of plasmid DNA amplification as a function of a template concentration with the use of the fusion polymerase PfuDBDlig-Taqs polymerase (a) and the Taqs

polymerase (b). Lane M, marker HyperLadder II (Bioline, UK). The amplified products were analyzed in a 1.2% agarose gel stained with ethidium bromide

Long-range PCR efficiency

The long-term PCR amplification efficiency of *PfuDBDlig-Taqs* was tested using DNA fragments of various sizes, including 5-, 8-, and 10-kbp DNA targets, various primer combinations and the abovementioned optimal PCR buffer. The results were compared with what was achieved with the use of the *TaqS* DNA polymerase. The PCR amplification efficiency of *PfuDBDlig-Taqs* DNA polymerase using 5 and 8 kbp targets was similar to the efficiency of the *TaqS* DNA polymerase (Fig. 7). However, the fusion *PfuDBDlig-Taqs* DNA polymerase amplified longer DNA fragments more efficiently than *TaqStoffel*: 10 and 8 kbp, respectively.

GC-rich templates


The potential of the fusion polymerase to amplify products with a high GC content was evaluated with the use of *PfuDBDlig-Taqs* and *TaqS* under the optimal conditions. A genomic DNA fragment of *Mycobacterium tuberculosis*

with a GC content of 78% was amplified with the use of various polymerase concentrations: 0.5–8 U. Figure 8 presents the results of the experiment. It was shown that the fusion polymerase *PfuDBDlig-Taqs* had the ability to efficiently amplify a product with a high GC content with the use of only 2 U of polymerase, while the reference *TaqS* polymerase was shown not to work even where 8 U of polymerase were used.

Discussion

Currently, the PCR method is widely used in diagnostics, molecular biology, and genetic engineering. PCR effectiveness, or in other words amplification yield and accuracy, depends on the type of polymerase and PCR conditions. Many problems are encountered during the PCR amplification of so-called difficult templates. These primarily include large products, products with a high content of high-melting GC pairs or templates of clinical or environmental samples containing

Fig. 6 The effect of lactoferrin (a), heparin (b), and blood (c) inhibitors on DNA amplification using the genomic DNA of *S. aureus* as a template and primers for *S. aureus*-specific *nuc* gene detection. Control reactions were carried out without any inhibitors. Lane M, DNA standards ladder (100–1000 bp) (Blirt, Gdańsk, Poland). The amplified products were analyzed in a 2% agarose gel stained with ethidium bromide






Fig. 7 Comparison of the PCR efficiency of the *PfuDBDlig-TaqS* DNA polymerase (a) and the *TaqS* DNA polymerase (b). DNA polymerases were used to amplify 5-, 8-, and 10-kbp targets from *E. coli* genomic DNA with the use of various extension times—72 °C, 1 min/kb. Lane M, GeneRuler DNA ladder SM 1331 (75–20,000 bp) (ThermoScientific, USA) The amplified products were analyzed in a 1% agarose gel stained with ethidium bromide

numerous DNA polymerase inhibitors. One of the solutions could be the use of a fusion polymerase with an additional DNA-binding domain. Research has indicated (Oscorbin et al. 2015) that the addition of a DNA-binding domain of DNA ligase from *Pyrococcus abyssi* to a PCR mixture has a positive effect on the activity of the *Taq* DNA polymerase in the presence of difficult templates. The literature indicates primarily an improvement in the PCR efficiency for long products (4 and 8 kbp) and a considerable reduction in the inhibitory effect of heparin in the reaction mixture.

Our studies have confirmed the results obtained by Oscorbin et al. It was shown that the fusion of a DBD of ligase (from *Pyrococcus furiosus*) to DNA polymerases improves basic enzyme properties such as thermostability, processivity, or reaction sensitivity. Polymerases containing a DBD of ligase are an ideal solution for PCR amplification of difficult templates.

The half-life of the fusion *PfuDBDlig-TaqS* polymerase doubled as compared to the reference *TaqS*, from 10 to

23 min at 99 °C, which predicts that the fusion polymerase will have a greater ability to amplify products with a high GC content. The *PfuDBDlig-TaqS* polymerase is able to efficiently amplify products with a GC content of up to 78%, while *TaqS*, which does not contain an additional domain, fails with targets with such a high GC content.

A two-fold increase in the amplification speed (from 16 to 33 nt/s) and an almost three-fold improvement of the processivity (from 8 to 22 nt) were found to increase the fusion polymerase's ability to amplify large products by 25% (from 8 to 10 kb). Based on the results of the studies, it appears that the *TaqStoffel* polymerase is able to amplify targets with a size of 8 kb, while the fusion *PfuDBDlig-TaqS* polymerase, under the same conditions, of up to 10 kb.

Another example of problematic amplification is a PCR carried out in the presence of inhibitors. The fusion polymerase appeared again to have better properties than the reference *TaqS*. The *PfuDBDlig-TaqS* polymerase tolerates, in a reaction mixture, an almost 16-fold higher concentration of lactoferrin and an eightfold higher concentration of heparin which are both primary inhibitors found in clinical and environmental samples (Al-Soud and Rådström 1998, 2001; Yokota et al. 1999). Such an improvement may be due to a 16-fold increase in sensitivity of the fusion polymerase which therefore has a higher affinity for the template. It seems that the template is picked up more selectively from the mixture in the presence of a binding domain and the PCR is more efficient even when inhibitors are present.

The fusion *PfuDBDlig-TaqS* DNA polymerase consisting of a *TaqStoffel* polymerase and the DNA-binding domain of a ligase obtained from *Pyrococcus furiosus* exhibits a higher extension rate, higher processivity, thermostability, sensitivity, and a higher tolerance to some PCR inhibitors and is able to amplify templates with a higher GC content and longer PCR products than the *TaqStoffel* DNA polymerase. For this reason, it may be a better solution to use the *PfuDBDlig-TaqS* DNA polymerase instead of the *Taq* DNA polymerase in many PCR applications with difficult targets.




Fig. 8 Comparison of *PfuDBDlig-TaqS* DNA polymerase (a) and the *TaqS* DNA polymerase (b) PCR efficiency for templates with high content of GC. DNA polymerases were used to amplify 78% GC-rich target from *M. tuberculosis* genomic DNA for various polymerase

concentrations (0.5–8 U). Lane M, DNA ladder (100–500 bp) (Blirt, Gdańsk, Poland). The amplified products were analyzed in a 2% agarose gel stained with ethidium bromide

Authors' contributions MŚ carried out the cloning, protein expression, and purification experiments, characterized the DBD-*TaqS* DNA polymerase, and analyzed the data. BK coordinated the study and analyzed the data. BZP took part in the protein expression and purification experiments. MW participated in the preparation of the characterization of polymerases. RP took part in the characterization of polymerases. MO participated in the protein expression and characterization, coordinated the study, drafted the manuscript, and revised the manuscript. All the authors have read and approved the final version of the manuscript.

Compliance with ethical standards

Competing interests The authors declare that they have no competing interests.

Ethics approval This article does not contain any studies with human participants or animals performed by any of the authors.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Al-Soud WA, Rådström P (1998) Capacity of nine thermostable DNAPolymerases to mediate DNA amplification in the presence of PCR inhibiting samples. *Appl Environ Microbiol* 64:3748–3753
- Al-Soud WA, Rådström P (2000) Effect of amplification facilitators on diagnostic PCR in the presence of blood, feces and meat. *J Clin Microbiol* 38:4463–4470
- Al-Soud WA, Rådström P (2001) Purification and characterization of PCR-inhibitory components in blood cells. *J Clin Microbiol* 39: 485–493
- Arezi B, Xing W, Sorge JA, Hogrefe HH (2003) Amplification efficiency of thermostable DNA polymerases. *Anal Biochem* 321:226–235
- Barski P, Piechowicz L, Galiński J, Kur J (1996) Rapid assay for detection of methicillin-resistant *Staphylococcus aureus* using multiplex PCR. *Mol Cell Probes* 10:471–475
- Dabrowski S, Kur J (1998) Recombinant His-tagged DNA polymerase II. Cloning and purification of *Thermus aquaticus* recombinant DNA polymerase (Stoffel fragment). *Acta Biochim Pol* 45:661–667
- Driscoll MD, Rentergent J, Hay SA (2014) Quantitative fluorescence-based steady-state assay of DNA polymerase. *FEBS J* 281(8): 2042–2050
- Elshawadfy AM, Keith BJ, EeOoi H, Kinsman T, Heslop P, Connolly BA (2014) DNA polymerase hybrids derived from the family-B enzymes of *Pyrococcus furiosus* and *Thermococcus kodakarensis*: improving performance in the polymerase chain reaction. *Front Microbiol* 5:224
- Halley G, Prezioso V (2003) Eppendorf HotMaster—an innovative hot start/cold stop technology for better PCR* results. *Tech Notes (Eppendorff-HotMaster)* 76–79
- Hamilton SC, Farchaus JW, Davis MC (2001) DNA polymerases as engines for biotechnology. *BioTechniques* 31:370–383
- Kermekchiev MB, Kirilova LI, Vail EE, Barnes WM (2008) Mutants of *Taq* DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. *Nucleic Acids Res* 37:e40
- Koźłowski R (2015) A novel method of *Mycobacterium tuberculosis* complex strain differentiation using polymorphic GC-rich gene sequences. *Acta Biochim Pol* 62:317–322
- Kwona KM, Kang SG, Sokolovac TG, Chod SS, Kimb YJ, Kima CH (2016) Characterization of a family B DNA polymerase from *Thermococcus barophilus* CH5 and its application for long and accurate PCR. *Enzyme Microb Technol* 86:117–126
- Lawyer FC, Stoffel S, Saiki RK, Chang SY, Landre PA, Abramson RD, Gelfand DH (1993) High-level expression, purification, and enzymatic characterization of full-length *Thermus aquaticus* DNA polymerase and a truncated form deficient in 5' to 3' exonuclease activity. *PCR Methods Appl* 2:275–287
- Lee J II, Cho SS, Kil E-J, Kwon S-T (2010) Characterization and PCR application of a thermostable DNA polymerase from *Thermococcus pacificus*. *Enzyme Microb Technol* 47:147–152
- Ma C, Tang Z, Wang K, Tan W, Li J, Li W, Li Z, Yang X, Li H, Liu L (2006) Real-time monitoring of DNA polymerase activity using molecular beacon. *Anal Biochem* 353(1):141–143
- Oscorbin IP, Boyarskikh UA, Zakabunin AI, Khrapov EA, Filipenko ML (2015) DNA-binding domain of DNA ligase from the Thermophilic Archaeon *Pyrococcus abyssi*: improving long-range PCR and neutralization of Heparin's inhibitory effect. *Appl Biochem Biotechnol* 176(7):1859–1869
- Terpe K (2013) Overview of thermostable DNA polymerases for classical PCR applications: from molecular and biochemical fundamentals to commercial systems. *Appl Microbiol Biotechnol* 97: 10243–10254
- Tveit H, Kristensen T (2001) Fluorescence-based DNA polymerase assay. *Anal Biochem* 289(1):96–98
- Wang Y, Prosen DE, Mei L, Sullivan JC, Finney M, Vander Horn PB (2004) A novel strategy to engineer DNA polymerases for enhanced processivity and improved performance in vitro. *Nucleic Acids Res* 32:1197–1207
- Yokota M, Tatsumi N, Nathalang O, Yamada T, Tsuda I (1999) Effects of heparin on polymerase chain reaction for blood white cells. *J Clin Lab Anal* 13:133–140