

Literature update of analytical methods for biogenic amines determination in food and beverages

Myrsini Papageorgiou^a, Dimitra Lambropoulou^a, Calum Morrison^b, Ewa Kłodzińska^c, Jacek Namieśnik^d, Justyna Płotka-Wasyłka^{d,*}

^a Department of Chemistry, Environmental Pollution Control Laboratory, Aristotle University of Thessaloniki, 54 124 Thessaloniki, Greece

^b Forensic Medicine and Science, School of Medicine, Dentistry and Nursing, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow G12 8QQ, United Kingdom

^c Institute of Sport e National Research Institute, Department of Anti-Doping Research, Trylogii 2/16 Street, 01-982 Warsaw, Poland

^d Department of Analytical Chemistry, Faculty of Chemistry, Gdańsk University of Technology, 11/12 Narutowicza Street, 80-233 Gdańsk, Poland

a b s t r a c t

Biogenic amines (BAs) have been reported in a variety of foods, such as fish, meat, cheese, and wines. The formation of BAs in food by the microbial decarboxylation of amino acids can result in human allergic reactions, characterized by difficulty in breathing, rash, vomiting, and hypertension. Control measures to prevent biogenic amine formation in foods and/or reduce their levels should be considered. Therefore, monitoring of BAs in food samples with the application of analytical techniques is of high importance.

This review is based on literature data from 2010 until today and refers to food samples and alcoholic beverages. The rationale of this study is to provide data for the occurrence of BAs in food and beverages samples and a comparison of the analytical techniques and challenges in liquid and solid matrices. Importantly, BAs can be used as future markers for quality and freshness of the food products and alcoholic beverages.

Keywords: Biogenic amines, Food samples, Chromatographic techniques, Capillary electrophoresis, Green assessment

1. Introduction

Nowadays, food products are manufactured and distributed worldwide, while deficiencies in regulation increases the concern for consumer's health. Heavy metals, pesticides, pharmaceuticals, personal care products and biogenic amines have been detected in different food matrices worldwide. A so called -chemical contamination- include multi-residues, may enter the food chain from different routes. Some sources linked with chemical contamination potentially include the application of pesticides in the field, veterinary medicines in animals, natural toxins and residues formed during the food processing or intentional contamination, such as food adulteration or contamination from sources which have not yet been identified.

Amino acids are the main compounds which offer nutrition value, aroma and flavor in products such as cheese, wine, honey and

other fermented foodstuffs after microbial or enzymatic conversion [1]. Therefore, the bacterial decarboxylation of amino acids or transamination of aldehydes and ketones leads to formation of active biogenic amines (BAs) [2,3]. BAs are nitrogenous organic bases of low molecular weight, polar or semi-polar compounds with aliphatic (putrescine, cadaverine, spermine and spermidine), aromatic (tyramine, phenylethylamine) or heterocyclic (histamine, pyrrolidine) structure.

The demand for safer foods has promoted more research into biogenic amines over the past few years, and thus many countries are involved in the analysis of food samples in terms of biogenic amines occurrence (Fig. 1). However, there are still some questions that remain unanswered. Despite the fact that biogenic amines are present in foods and beverages and can cause toxic effect to the body, a shared regulation limiting the amounts of biogenic amines in food and beverages is still lacking. Knowledge on their presence in wine is also important for the food trade sector.

Referring to BAs, their occurrence has been detected in a broad range of protein-rich food matrices (from plant or animals origin) that are regularly consumed by humans and placed within a daily dietary program, such as meat [4–7], fish [4,5,8–14], dairy products

* Corresponding author. Department of Analytical Chemistry, Chemical Faculty, Gdańsk University of Technology, 11/12 Narutowicza Street, 80-233, Gdańsk, Poland.

E-mail addresses: plotkajustyna@gmail.com, justyna.wasyłka@pg.edu.pl (J. Płotka-Wasyłka).




Fig. 1. Percentage breakdown of biogenic amine researches related to food products published since 2010 categorized with the geographical region of the analyzed samples.

[15] and alcoholic beverages. This could place them as markers for food and beverages quality and freshness being connected to the degree of food storage, processing or degradation [16]. Therefore, any food stuff produced by fermentation or exposed to microbial contamination during processing or storage may contain BAs [17].

With respect to health risks, little legislation exists; in low concentration BAs are essential for many physiological functions in humans, acting as hormones or neurotransmitters. The key roles of BAs are important for growth, renewal, and metabolism in organs, for high metabolic activity of the normal functioning and immunological system of the gut and for temperature regulation [18–20]. While in high concentrations, effects such as headache, nausea, hypo- or hypertension, migraine, skin allergy and digestive problems of food poisoning have been observed by biogenic amines [21,22]. Also, Biogenic Amines (BAs) are precursors of nitrosamines, which have been linked to carcinogenic and mutagenic activity. Generally, BAs are not destroyed during cooking, treatment with high temperature, storage and are stable and difficult to degrade [23].

The determination of biogenic amines is laborious and challenging due in part to the physicochemical properties of these compounds. Numerous analytical methods have been developed while few of them can overcome the step of the derivatization process easily. A pre-concentration step is generally needed, when the analysts deal with complex matrices. During the last decade, new articles have developed extractions which can be more environmental friendly, with the use of less toxic, organic solvents combined with sensitive and selective analytical instrumentation using mass spectrometer and tandem mass spectrometer detectors [5,24,25].

The aim of this review is to summarize the concentration of biogenic amines in different food and beverage products from 2010 until today, to highlight which matrices have not been analyzed and also, which matrices are the most popular and should continue to be monitored for their toxicity. Regarding the physicochemical characteristics of BAs, an overview of the analytical methodologies, cleanup and pre-concentration techniques, a comparison between the derivatization agents and environmental assessment of the analytical methodologies have been made, so we can underline the development of more green, environmental friendly extraction procedures. Finally, the regulation policy and the toxicity of BA's have been discussed. The study concludes with a focus on the main issues that should be further investigated.

2. Function and occurrence of biogenic amines

Generally, BAs are sources of nitrogen and precursors for the synthesis of many specific compounds in organisms such as alkaloids, hormones, nucleic acids, and proteins. Moreover, they are also responsible for several processes in the organism such as the regulation of body temperature, intake of nutrition, increase/decrease of blood pressure [26]. In plants, putrescine, spermidine and spermine are implicated in a numerous of physiological processes, such as cell division, fruit development, flowering, response to stress and senescence. Generally, polyamines are important for processes of every organ in the body including growth, renovation, metabolism and are essential for maintaining the high metabolic activity of the normal functioning and immunological system of gut [26].

It has been shown that BAs are potential precursors for the formation of carcinogenic N-nitroso compounds. The reaction of primary amines and nitrosating-agents produces short-lived alkylating species, which can react with other components present in the food matrix to generate products for example alcohols, devoid of toxic activity in the relevant contents [26]. The secondary amines such as agmatine, spermine, spermidine and others can react with nitrile and, in sum, generate the nitrosamines, while tertiary amines produce a range of labile N-nitroso products [27]. In foods contain high level of fat (e.g. bacon), in the presence of water and at high temperature, the carcinogen N-nitrosopyrrolidine can be formed from putrescine or spermidine. Moreover, several of BA including putrescine, spermidine and cadaverine, can act as free radical scavengers, while tyramine has a remarkable antioxidative activity, which increases with its content. Thus, inhibiting effect depends on amino and hydroxy groups. Another function of BAs is impact on flavor and taste of food [26,28].

Theoretically, all foods that contain free amino acids or protein and are exposed to conditions enabling microbial or biochemical activity, BA presence can be expected [28]. BAs generally occur in fermented products such as cheese (5–4500 mg/kg), wine (5–50 mg/dm³), beer (2.8–13 mg/dm³), sauerkraut (110–300 mg/kg), but they can be present in food that is not properly stored, for example fish (2400–5000 mg/kg), beef liver (340 mg/kg), prepared meats (10–700 mg/kg) [4,29]. In fact, BAs are found as indicators and markers of food decomposition of non-fermented foods [1,26]. Spoiled foods are also rich in BA and usually contain high levels of putrescine and cadaverine [1,26].

It is worth to notice, that amine content and profiles may vary depending on several extrinsic and intrinsic factors during the manufacturing process, such as the ripening conditions, formulation, physico-chemical and proteolytic parameters, as well as microflora development and its decarboxylase activity. In fact, every country has its own traditions of food, which differ in both the microbial characteristics, the way of production and processing methods. In addition, each country is characterized by a specific climate, which also affects the presence of biogenic amine in the substrate, for example, beverages. These differences, can be observed in the following section of this review [30,31].

2.1. Biogenic amines in seafood and its products

Since the maximum levels for histamine are legally established for seafood and its products, most research is related to these products. Seafood is a specific type of food which may harbor several biological, chemical, and physical hazards, the most common of which are biotoxins, pathogenic bacteria and viruses, metal inclusion and biogenic amines [32].

It is proven that both specific as well as non-specific microbial contaminants originating from the natural environment or being acquired during processing impact on the microbiological complexity of seafood [32]. The wide range of environmental habitats and the variety of processing practices (iced seafood products to canned products) are all important parameters in determining the initial contamination of fish and their products. The microorganisms of seafood intended for human consumption depend on the environmental conditions of its natural habitat, but here can be mentioned, Gram-negative bacteria belonging to the genera *Pseudomonas*, *Moraxella*, *Acinetobacter*, *Shewanella*, *Flavobacterium*, *Vibrionaceae* (*Vibrio* and *Photobacterium*) and the *Aeromonadaceae*. Although Gram-negative bacteria are the predominant microorganisms, Gram-positive bacteria such as *Bacillus*, *Micrococcus*, *Clostridium*, *Lactobacillus*, and *Coryneforms* can also be found at various levels. In polluted waters, high numbers of *Enterobacteriaceae* may be found. In clean temperate waters, these organisms disappear rapidly, but it has been shown that *Escherichia coli* and *Salmonella* can survive for long periods in tropical waters, and once introduced, may become indigenous to the environment [32].

The composition of seafood microbiota changes quite dramatically during spoilage. For example, *Shewanella putrefaciens* and *Pseudomonas aeruginosa* have been identified as the prominent spoilage bacteria of fresh fish, while at ambient temperature (25°C), the microbiota at the point of spoilage is dominated by mesophilic *Vibrionaceae* and, particularly if the fish is caught in polluted waters, *Enterobacteriaceae* [33].

The important biogenic amines in seafood are tyramine, histamine, putrescine, tryptamine, and cadaverine. These are formed from their respective free amino acids tyrosine, histidine, ornithine, tryptophan, and lysine. Spermidine and spermine are produced from putrescine [34]. Thus, the concentration of free amino acids especially histidine is important since histidine is the precursor in the biosynthesis of histamine. Glutamic acid together with lysine, arginine, leucine, aspartic acid, isoleucine, glycine, alanine, threonine and valine represented 77.6% of total amino acids in naturally fermented fish [34].

Histamine levels in freshly caught fish are generally low, usually below 0.1 mg/100 g [32]. However, exposure of certain fish to elevated temperatures after the catch and before consumption can cause formation of histamine from histidine by bacterial histidine decarboxylases and thus, the level of histamine can increase.

Some technological processes such as salting, ripening, fermentation or marination can increase the possibility of BA

formation. BAs can also be produced throughout the manufacturing process, as well as during storage of the end product if improper holding temperatures are employed [32].

In studies conducted in fish and fish products, the presence of BAs has been also reported. Among the analyzed samples were smoked mackerel and salmon, tuna, anchovies, tinned tuna, canned fish, mackerel, sardines, tuna, marinated anchovies and shrimp sauce. Bilgin et al., 2015 [13] detected the mean concentration of cadaverine (up to 122.18 ± 68 mg/kg) in canned tuna and sardines from Turkey, tryptamine (up to 190.61 ± 7.67 mg/kg) in marinated anchovies, histamine in maximum concentration (up to 110.33 ± 9.87 mg/kg) and putrescine (up to 116.53 ± 2.90 mg/kg) in canned tuna. Bilgin et al., 2015 connected the high values of BAs (upper the safe level) with the bad quality of the raw fish and the processes needed for the formation of these products. Also, Palermo et al., 2013 [9] detected trimethylamine in fresh anchovy samples from the Italian market in concentrations up to 120 ± 1 mg/kg indicating poor storage conditions. Lower concentrations were observed for the analyzed BAs in all the other fish samples from other regions. Only one survey was conducted in shrimp sauce from Malaysia detecting tryptamine, putrescine and tyramine up to 57.9 mg/L, followed by 2-phenylethylamine, histamine and spermidine (up to 5.4 mg/L) [35].

2.2. Biogenic amines in meat

Meat and meat products are an important component of the diet in developed countries. This type of food has repeatedly been reported to contain biogenic amines [16]. The most prevalent BAs in these products are putrescine, tyramine, cadaverine, and also histamine. The only amines present at significant levels in fresh meat are spermidine and spermine. Some amines such as tyramine, putrescine and cadaverine can be formed during storage of meat [36]. It is reported that tyramine concentrations in stored beef was highest on the meat surface, however, it can be reduced effectively by washing [37]. Generally, fermented meat products are an example of products which considerable amounts of BAs can be found as a consequence of the use of poor quality raw materials, contamination and inappropriate conditions during processing and storage [16]. Moreover, the microorganisms responsible for the fermentation process may contribute to BAs accumulation. In fact, the non-protein nitrogen fraction which increases during fermentation includes the presence of free amino acids, precursors of BAs. The major protease activity is derived from endogenous meat enzymes. Proteolysis is favored by the denaturation of proteins as a consequence of acidity increase, dehydration and action of sodium chloride [16]. Fermented products with comparable microbiological profiles may differ in their BAs content, indicating that the production of such compounds depends on a complex interaction of factors [16]. Raw meat material is the natural source of the substrate from which BAs are produced. It also is the largest component of the matrix in which the decarboxylation reactions take place and any conditions that alter its nature and characteristics will influence the formation of biogenic amines.

The highest concentration of BAs in meat and meat products occurred in meat sausages from Greece. Papavergou et al., 2012 [7] reported the occurrence of cadaverine, putrescine and histamine in concentrations up to 689.89 mg/kg in fermented sausages, in contrast to dry meat ripened products with spermine to be the most frequent detected BA with concentration range from 11.69 ± 0.77 to 46.32 ± 0.39 mg/kg. Lower concentrations were detected in sausages from China and Poland, with the most dominant BAs being spermine, histamine and tryptamine ranging up to 104.28 mg/kg (China) [38] and 2-phenylethylamine, tyramine and putrescine ranging up to 17.1 mg/kg (Poland) [4]. Moreover, Lazaro

et al., 2015 [39] detected tyramine up to 356.8 mg/kg, followed by putrescine, cadaverine, spermine and spermidine with concentrations up to 54.8 mg/kg in chicken and quail samples from Brazil. Furthermore, Lazaro et al., 2015 [39] supported that the correlation between the BAs and bacterial growth does not present a specific profile for the BAs but depends on the meat sample. In salami from Germany, putrescine and tyramine were the most detected BAs (up to 77.14 mg/kg) [5] in contrast to salami from Italy, agmatine and spermine appeared in lower concentrations up to 8.3 mg/kg [9]. Slightly higher concentrations were presented in 4 different meat samples (fresh beef, chicken, lamb, rabbit) from Italy, with the presence of spermine not exceeding 22.22 mg/kg [6].

2.3. Biogenic amines in dairy products

Dairy products are important components in the human diet. Their current consumption is relatively high and is expected to increase steadily during the next few decades [40]. Thus, the provision of wholesome and safe dairy products to consumers is expected to be more challenging with the anticipated increased consumption, as the risk increases with the exposure to hazards, such as biogenic amines (BAs), potentially present in the product [40].

Together with wine, dairy products can accumulate high levels of BAs. Biogenic amines continue to raise concern due to their frequent detection at high levels in various types of dairy products and to increased awareness of their actual or potential adverse health effects [40]. Also, the fact that BAs are produced not only by microbial dairy contaminants of different origins but also by the technological microbiota used in the fermentation and/or ripening of dairy products, including lactic acid bacteria (LAB), yeasts, and molds, complicates their control by conventional means.

In the raw material (milk) polyamines are the most abundant; however, also in the final product such amines as cadaverine, histamine, putrescine, tyramine, β -phenylethylamine and tryptamine are all detected. The content of BAs in different types of cheese is different. Moreover, these differences can also appear within the same type of cheese and even between different sections of the same cheese.

As was previously mentioned, the occurrence and amount of BAs in food depends on many factors among which the availability of the precursor amino acid(s) is a limiting factor. It is noted that the precursor amino acids may be naturally present in milk in a free state or be released from milk proteins by hydrolysis. Proteolytic activities impact on the formation of precursor amino acids in dairy products may result from different sources acting independently or in combination, including proteolytic strains of micro-organisms present in dairy products, proteases used for coagulating milk in cheese-making, the milk-native heat-stable protease plasmin, and other proteases liberated from somatic cells. Subsequently, BA-producing microorganisms will continue the formation process of BAs, which are then released into the matrix of dairy products [41].

It is reported that Gram-positive bacteria are the main BA producers in cheese. Several strains of such genera as *Enterococcus*, *Streptococcus*, *Leuconostoc* and *Lactobacillus* are indicated as BA producers, and can be present in microbiota of milk or introduced through contamination before, during or after the dairy products processing. In fact, BA⁺-LAB may even form part of the starters or adjunct cultures, thus, including the inability to produce BAs as an indispensable condition of strains intended to be used as starters is of high importance. This information was the basis to introduce a system for a pre-market safety assessment of selected taxonomic groups of microorganisms leading to a 'Qualified Presumption of Safety' (QPS) by the European Food Safety Agency (EFSA) [42]. And thus, several bacteria associated with food e.g. *Lactobacillus*,

including *L. buchneri*, *L. brevis*, *L. hilgardii*, have obtained a QPS status [42], although some strains of these species have been indicated as BAs producers.

The presence of BAs in different types of cheese has been described in a lot of surveys. More than 15 BAs have been analyzed [5,9,37,43–47]. Among different types of cheese; cadaverine was found to be more dominant than other BAs. In 7 different types of cheese in Austria, cadaverine ranged from 2 to 748.2 mg/kg [5]. The occurrence of cadaverine in Formaggio di Fossa ranged up to 1303 \pm 5.02 mg/kg [45] and in Cabrales 774.51 mg/kg from Italy [44]. The highest concentration presented in Otlu peynior cheese from Turkey (1844.5 mg/kg) [43]. Moreover, histamine presented up to 1159.7 mg/kg [3], tyramine up to 1125.2 mg/kg [43] in Turkey and up to 2519.98 mg/kg in cabrales cheese from Italy [44], while slightly lower concentration were observed for other cheese types and regions with BAs concentration not exceeding 847 mg/kg [43].

Regarding BAs, in all the other dairy products [15] yogurt [5,38], milk (cow and goat), kefir, fermented cream and buttermilk [46], only putrescine, histamine and agmatine has been detected in dairy products from USA with maximum concentration up to 3.2 mg/L, while tyramine has been detected in cow and goat milk from Brazil at concentrations up to 337.11 mg/kg [47]. Lower concentrations observed for all the other regions with BAs maximum concentrations of BAs up to 26.1 mg/kg in yogurt, 5 mg/kg in buttermilk, 15.4 mg/kg in fermented cream, 14.3 mg/kg in kefir from Czech Republic [46].

2.4. Biogenic amines in chocolate and coffee

Chocolate contains many pharmacological agents that separately or as a group, could evoke physiological sensations and may be the driving force behind chocolate cravings. In the past few years, the psychopharmacological effects of chocolate have been a topic of increasing interest among nutrition neuroscientists as evidence continues to build for the localization of chocolate's actions and the precise biomolecules involved.

One of the groups of compounds that occur in chocolate and may impact negatively on human body is biogenic amines. Several endogenous biogenic amines are found in chocolate, most notably, tyramine and phenylethylamine, tryptamine, clovamide, and serotonin. It is reported that chocolate has the highest level of phenylethylamine among tested foods. This compound is produced by brain tissue and is rapidly metabolized by monoamine oxidase- β and aldehyde dehydrogenase to phenylacetic acid, the major metabolite of this compound in the brain. When consumed at high levels, effect such as blushing, headaches and increase blood pressure can occur. Consequently, Mayr et al., 2012 [5] reported that the dominant BAs were spermidine, ethanolamine, tyramine, and 2-phenylethylamine with maximum concentration up to 7.40 \pm 3.4 mg/kg in chocolate from Germany. Sparse information exists for all the other regions.

Coffee is one of the most common beverages and composed of carbohydrates, fiber, proteins, free amino acids, lipids, minerals, organic acids, chlorogenic acid, trigonelline and caffeine. Three different extraction techniques, decoction, infusion and pressure methods were used for coffee brewing. Many factors influence the process of brewing, such as ground roast coffee composition, grid, brewing method, coffee/water ratio, water hardness and temperature, duration of the coffee–water contact, and filter material [48]. Putrescine appeared to be the most dominant BAs in coffee beans and in coffee beverage followed by spermidine, spermine, and serotonin, while cadaverine and tyramine are generally present in smaller amounts. Ground coffee samples from Turkey and Brazil were analyzed. Higher concentrations occurred for serotonin (170.8 mg/kg), tyramine (83.94 mg/kg), cadaverine (75.08 mg/kg)



and tryptamine (37.85 mg/kg) in Turkey ground coffee [49], while different pattern observed in Brazil [50]. The highest concentration observed for putrescine (61.5 mg/kg), followed by spermidine (8.2 mg/kg) and spermine (7.2 mg/kg). Moreover, in brewed coffee from Turkey [49] were detected higher concentrations of tyramine (19.7 mg/L) and serotonin (13.55 mg/L), followed by tryptamine (9.185 mg/L), cadaverine (9.059 mg/L) and 2-phenylethylamine (4.944 mg/L) compared with the brewed coffee from Italy (up to 1.95 mg/L) [48].

2.5. Biogenic amines in fruits, vegetables and its products

Fresh fruits, vegetables and their products like juices and sauces are a group, common added in the daily dietary habits, expresses the culture, history and lifestyle of every country. Harvesting processes, storage conditions and further management of the fruits and vegetables (mashing, mixing, fermentation) varies greatly between countries. Fresh food and vegetables contain BAs as endogenous components and due to uncontrolled microbial enzymatic activity they can be accumulated [4]. Until now, in fresh fruit and vegetables, in their fermented products and juices have been detected 22 BAs. The most detected BAs for this group are tyramine, putrescine, cadaverine, histamine, spermine and spermidine, while serotonin and agmatine has not detected in any of the analyzed samples. Also, the highest concentration of BAs found in fresh fruits instead of juices. Among this group apple, banana, cabbage (raw and fermented), olives, pineapples, pomelo, red grapes, white grapes and grape, apple and plum, apricot, black and red currant, cherry grape, grapefruit, litchi, mango, orange, peach, pear, pineapple juices have been analyzed. In fresh fruits and vegetables, higher concentrations were observed for cadaverine (36.2 mg/kg) and tyramine (23.5 mg/kg) in bananas from Poland [4], followed by spermidine (22.3 mg/kg) in raw cabbage from Spain and putrescine in olives (14.63 mg/kg) from Italy [9]. Regarding the fermented cabbage, higher concentrations were noted with putrescine (108.9 mg/kg), tyramine (60.66 mg/kg) and histamine (37.01 mg/kg) from Germany [5]. The only high levels in juices was the concentration of cadaverine in apricot, peach and pear (up to 17.22 mg/kg) from Italy [51] and putrescine in peach (7.22 mg/kg) and pineapple (5.02 mg/kg) from Italy and Poland [4], respectively.

2.6. Biogenic amines in soybean products

Soybean products are a diverse group of food products including tofu, soya sauce, sufu, doubanjiang, natto, soymilk, soy sprouts, tempeh, miso, tamari, soybean paste, soya bean, bean curd used in Chinese cuisine in Europe, America and Asia.

Different profile of BAs presented in different products. The differences in BAs profile between the soybean products may be attributed to different processes or hygienic conditions in the manufacture [52] and distribution. However, other factors in the variation of BA content could be the quantity of raw soybean product used, the fermentation processes or other technological processes, the proteolytic activity from bacteria [53] or other microorganism contaminants with potential activity [54].

In sufu the most dominant BAs were putrescine (316.9 mg/kg) and tyramine (446.6 mg/kg) [55], while another profile of BAs observed in sufu products from China, with cadaverine (up to 883.7 mg/kg), putrescine (177.6 mg/kg) and spermidine (154.8 mg/kg), higher concentrations were detected in Spain up to 1730 mg/kg for tyramine and histamine [54]. Also, Yang et al., 2014 [56] reported the presence of six out of eight BAs in white sufu from China and the detection of unsafe level in 8 samples, although the authors conclude that further investigation is needed. Additionally, Kim et al., 2012 [57] detected 2-phenylethylamine, spermidine and

tyramine in higher concentrations from safe level of human health in some natto products from Korea, while lower concentrations of biogenic amines were detected in natto products from Spain (spermidine up to 75.21 mg/kg). Among the other soybean products only soybean paste from Korea was presented remarkable concentrations in all of the nine analyzed BAs [58]. Lower concentrations were observed in soybean paste from Spain [54].

2.7. Biogenic amines in beer and other alcoholic beverages

Alcoholic beverages are a major category, including beers (all types), liqueurs, gin, rum, brandy, whiskey and ciders. Beer is produced mainly by mixing malted barley and hops and selected yeast strains, such as *Saccharomyces cerevisiae* (top fermenting), *Saccharomyces carlsbergensis* with wild yeast and lactic acid bacteria. It was observed that the high concentration of histamine and tyramine linked with microbial contamination during brewing. On the other hand, putrescine, agmatine, spermidine, spermine and 2-phenylethylamine could be natural constituents either from malt or hop.

Ciders also produced from apple juice with indigenous yeast and LAB. A lot of strains have been isolated and identified as BA producers. The other alcohol beverage, liqueurs is a mixture of ethyl alcohol (e.g. cognac, rum, or whiskey) or a neutral distillate of agricultural origin flavored with fruit, cream, herbs, spices, flowers or nuts and bottled with added sugar, honey, or high-fructose corn syrup, produced either by heat, infusion, cold or maceration process or distillation [59].

Besides the extended consumption worldwide of beer and other alcoholic beverages, scarce data exists concerning the detected BAs. Almeida et al., 2012 reported the detection and analysis of eighteen BAs in beer samples from Portugal [59]. The most dominant ones were putrescine, tyramine and dimethylamine with concentrations up to 12.777 mg/L, which are in agreement with published data from Iran and Brazil [17]. Aflaki et al., 2014 [17] observed very low concentrations for non-alcohol beers from Iran with concentrations below 1.17 mg/L for the analyzed BAs. For all the other alcoholic beverages have been conducted only one survey, so further investigation is needed. Among, rum, gin, whiskey and brandy were highlighted only the BAs in rum. On the other hand, in liqueurs were detected very low concentrations in contrast to the other beverages.

2.7.1. Biogenic amines in wine

Literature data reports that more than 15 amines have been identified in wine and their total concentration has been reported to range from a few ng/L to about 50 mg/L, depending on the quality of the wine. Several factors impact on the type of BA that could be found in wine including differences in the wine-making process, time and storage conditions, raw material quality and possible microbial contamination during winery operations. Generally, they are two different sources of BAs in wine: raw materials and fermentation processes. It has been shown that some amines occur in grapes, namely, histamine and tyramine, as well as several polyamines and volatile amines. Although, histamine, putrescine and tyramine are present in higher concentration level in wine, other compounds including phenylethylamine, cadaverine and isoamylamine are also present in smaller amounts. Putrescine and cadaverine are normally associated with poor sanitary conditions of grapes.

Generally, lactic acid bacteria [10] can produce metabolic energy and/or increase their acid resistance by using catabolic pathways that convert amino acids into amine-containing compounds including BAs. Therefore, many researchers are focused on studies of correlation of BAs production in wine with species of LAB involved in the wine-making process. It is widely known that several bacteria are implicated in BAs production in wine including

Lactobacillus, *Pediococcus*, *Oenococcus* spp. and *Leuconostoc* [17]. It is reported that *Leuconostoc mesenteroides* has a high potential to produce tyramine or histamine in wine [17], while *Oenococcus oeni* is able to significantly contribute to the overall BA content of wines, mainly producing histamine, however, the ability of each bacteria type to produce BAs varies among strains [17]. Moreover, LAB strains have the ability to simultaneously produce different amines, suggesting that some strains might possess more than one amino acid decarboxylase activity under specific culture conditions.

According to the literature, in white wines the range of BAs ranged from n.d. to 12.8 mg/L. High concentrations were found in Italy for histamine, putrescine, tyramine and ethylamine while in Portugal only histamine presented in 8.94 mg/L 15 out of 16 BAs occurred in red wines worldwide. Higher concentrations were detected for histamine 23.1 ± 2.2 mg/L in Portugal [60], in Italy 10.8 mg/L [61], in France 14.05 mg/L [62] and in different EU countries (Italy, France, Germany, Switzerland, Austria and Spain) 16.2 mg/L [63]. Also, putrescine presented the highest concentration in three different surveys in Italy with 31.8, 11.13 and 16.2 mg/L respectively [51,61,63] and in France 48.72 mg/L [62]. Lower concentrations were detected for the above BAs in Brazil and China. Slightly lower concentration of tyramine were present and ranged up to 18.8 mg/L [61]. Ethylamine and methylamine presented the highest concentrations in France up to 10.46 and 36.64 mg/L, respectively [62]. Ethanolamine was analyzed and detected in concentrations up to 17.44 μ g/L in China [64] and serotonin was absent from the analyzed samples in Italy and in China [51,64]. Moreover, the most persistent BAs in rose wines were histamine in Portugal with concentrations ranged up to 15.1 mg/L [60] and tyramine (10.8 mg/L) in Italy [61], while histamine and spermine detected in China in concentrations up to 18.35 mg/L in rice wine [38].

However, studies have shown that higher concentrations of BAs occurred in red wine samples instead of white, rose, rice and Porto wines. Comuzzo et al., 2013 [63] supported that climatic conditions could be connected with these variations but also winemaking technologies could play a key role in the production of BAs.

2.7.2. Biogenic amines in vinegar

Vinegar constituted after the bacteria conversion of ethanol to acetic acid. Every alcoholic mixture, with ratio alcohol-water to wine can produce vinegar [65]. Although these two food products are of great importance, data concerning the occurrence of BAs is limited. However, spermine and tyramine were detected in both black and green tea from Asia with concentrations up to 0.336 mg/L. While higher concentrations of 3-aminopropionic acid, 4-aminobutanoic acid and agmatine were detected in herbal tea from USA [15], so further investigation should be consider. Furthermore putrescine was detected in vinegars from USA (up to 3.2 mg/L) and Spain (0.525), while histamine presented slightly lower concentration in Spanish vinegars (0.309 mg/L) [65].

2.8. Biogenic amines in tea

Tea is a widespread product worldwide and is available and consumed regularly in large amounts. The leaves and buds of *Camellia sinensis* mixed with water constitute the tea extract. Different varieties, quantities, qualities of the plant and processes applied in every country. Two different categories are well known, green tea after heat or steam treatment and fast drying of young leaves and buds of the plant and black tea after procedures such as weathering, destruction of plant tissues by various rolling, crushing and/or tearing and finally drying. After these processes, the enzymes which are responsible for oxidation and degradation released and form polyphenols and the color compounds.

3. Physiological and toxicological aspects of biogenic amines

Many BA's are important for normal function of biological systems. For example in eukaryote cells they are important precursors for a variety of precursors, some play an important roles as neurotransmitters and others such as putrescine and spermidine are involved in critical biological functions. However, in larger concentrations, usually from an accumulation in a food source, these compounds can be toxic.

Biogenic amines may be a constituent of many foods and consumption can be problematic for human health. High BA concentrations can cause flushes, headache, nausea, palpitations of the heart and changes in blood pressure and many other physiological problems. Important BA's found in food include histamine, tyramine, putrescine, cadaverine and phenylethylamine. Polyamines, such as putrescine, cadaverine, agmatine, spermine, and spermidine, are naturally present in food and are involved in growth and cell proliferation [66].

The toxicological effects of many of these compounds has been reported for many reasons including the effect they can have on human health.

Many of these amines in the presence of nitrites can be potential carcinogens when converted to nitrosamines [66], but nitrosamines formed from the polyamines only become a health risk as toxicity is reached after consumption of large amounts, and more than expected in a daily meal.

In general, there is little research on the toxicology of BA's, but that which exists is predominately on histamine. Consuming large amounts of histamine (greater than 500 ppm) can lead to poisoning which may manifest as an allergen reaction with symptoms including breathing, itching, rash, vomiting, fever, and hypertension. An intake of 5–10 mg of histamine may be considered as having some effect to some sensitive people while 10 mg is generally considered as tolerable limit, 100 mg can induce a medium toxicity and 1000 mg is highly toxic [25]. In fact histamine and tyramine are considered most toxic and particularly relevant for food safety, with putrescine and cadaverine potentiating the effects by inhibition of histamine detoxifying enzymes [25].

BA's ingested with food are detoxified by amine oxidases in the GI tract, and particularly the intestinal mucosa. Under normal conditions, amines ingested are metabolized by conjugation or oxidation reactions with amine oxidase enzymes such as monoamine oxidase (MAO), di-amine oxidase (DAO), polyamine oxidases (PAO) and N-methyl transferase [67].

Therefore, it can be seen that MOA or DOA inhibitor medication (antihistamines, antimalarial agents, anti-depressant drugs), alcohol or genetics (enzymes not properly functioning) may affect BA metabolism. In addition, people with GI tract diseases may be poor metabolisers of BA's [25]. Under these circumstances the BA's enter the systematic circulation where they may promote release of adrenaline and noradrenaline, leading to many of the well documented problems (increased cardiac output), tachycardia, increased blood sugar and higher blood pressure.

In terms of toxicity via Oral absorption levels for putrescine, spermine, and spermidine are 2000, 600, and 600 ppm, respectively. The acute toxicity level for tyramine and cadaverine is greater than 2000 ppm. The no observed adverse effect level (NOAEL) is 2000 ppm for tyramine, putrescine, and cadaverine; 1000 ppm for spermidine; and 200 ppm for spermine [68].

4. Regulations policy

Without doubt, the toxicity of BAs is a very important parameter, however, determination of the exact toxicity threshold of BAs in an appropriate product of food is a difficult task, due to the fact



that their effect does not depend only on BAs presence alone, but is influenced by other compounds. In addition, toxic effect of BAs on organism is also dependent by on the specific efficiency of the detoxifying mechanisms in different individuals [16]. Therefore it can be stated that the BAs toxicity will depend on two factors associated with the food itself (quantitative and qualitative) and related to the consumer (individual susceptibility and state of health).

Due to the fact that there is still lack of information on toxic doses for each biogenic amine, not for all amines occurred in food legal limits are established. In fact, concentration level of histamine is regulated by law in some type of food product (fish food).

The European Food Safety Authority confirmed histamine and tyramine as the most toxic and particularly relevant for food safety [69]. Taking into account the data from food intoxication outbreaks a legal upper limit of 100 mg histamine·kg⁻¹ food and 2 mg L⁻¹ of ethanol has been suggested. In Brazil, the Regulation of Industrial and Sanitary Inspection of Animal Products does not mention the amine maximum level allowed in products of animal origin [67]. However, MERCOSUR Resolution and the Technical Regulation on the Identity and Quality of Fresh Fish (whole and gutted) establish a maximum level of 10 mg/100 g of histamine in the muscles of species of the Scombridae, Scomberesocidae, Clupeidae, Coryphaenidae and Pomatomidae families [67]. The maximum acceptable histamine levels in fish were also established in other countries. For example, the European Union has established regulations according to which histamine level should be below 100 mg kg⁻¹ in raw fish, and below 200 mg kg⁻¹ in salted fish, however, this is obligatory for species belonging to the Scombridae, Engraulidae, Coryfenidae, Pomatomidae and Clupeidae families [69]. In USA, the Food and Drug Administration, established a maximum limit of 5 mg histamine/100 g product at the port and 10 mg histamine/100 g product in pickled fish for species susceptible to form histamine [67]. The Nutritional codex of the Slovak Republic had determined the maximal tolerable limit for the following two BAs: histamine (20 mg kg⁻¹ in beer and 200 mg kg⁻¹ in fish and fish products) and tyramine (200 mg kg⁻¹ in cheese) [70]. A recommended upper limit of 100–200 mg kg⁻¹ for histamine in meat products has been proposed by the Netherlands Institute of Dairy Research and by the Czech Republic.

There are no established standards for cadaverine, putrescine or other BAs, only some recommendation are given, for instance, the recommended maximum level of tyramine has been proposed variously to be in the range of 100–800 mg kg⁻¹ of food. Value of 30 mg kg⁻¹ for β-phenylethylamine has been reported as toxic dose in food.

5. Presence of biogenic amines as an indicator of food products quality

Due to the fact that the level of concentration of BAs can increase (cadaverine, putrescine and tyramine), decrease (spermine and spermidine) or remain constant during the processing and storage of some food products (e.g. meat, meat products), their amounts and ratios have been proposed as an index of the hygienic conditions of raw material and/or manufacturing practices [16]. However, the usefulness of BAs as a quality index depend on the nature of the product what impact on the tendency to be more satisfactory in fresh meat as well as heat-treated products than in fermented products.

The Chemical Quality Index (CQI) was introduced in 1977 [71] for seafood and fish and was calculated using following equation:

$$CQI = \frac{cHIM + cPUT + cCAD}{1 + cSPD + cSPM}$$

where HIM – histamine, PUT – putrescine, CAD – cadaverine, SPD – spermidine, SPM – spermine and c – concentration [mg kg⁻¹].

Food freshness should be evaluated by considering an amine index, which includes all the BAs related to meat spoilage, and thus, considering the fact that tyramine increases during meat storage, this BA should also be included in a Biogenic Amine Index (BAI). This knowledge was used in BAI proposed in 1982. However, it needs to be remember, that not all spoilage or starter microorganisms can decarboxylate free amino acids. Even within the same species, not all strains develop the same decarboxylating capacity, so that a low concentration of BA does not always signal good microbiological quality [16]. Therefore, there is no simple way to establish a biogenic amine index that reliably predicts quality for products of this kind.

6. Analytical challenges and strategies of biogenic amines determination

Due to the fact that two reasons for importance of BAs determination in food exist which are potential toxicity and possibility of using them as food quality markers, the development of analytical methodologies for laboratory examining of BAs content is of high importance.

Undoubtedly, BAs identification remains one of the greatest challenges in food analysis. From the analytical point of view, bioactive amine detection in food is not simple due to reasons including strong polar character of the compounds, which results in a greater solubility in water rather than in the organic solvents frequently used; complexity of the matrix sample; variable concentration range which mainly is very low; the presence of potentially interfering compounds; absence of intrinsic properties of the compounds, which could enable their detection by usual physico-chemical methods as spectrophotometric, fluorometric or electrochemical methods; and the occurrence of several biogenic amines simultaneously. However, to solve these problems the developed analytical methods are generally based on amine extraction and derivation followed by separation and quantification. The extraction of amines represents the critical step of the process and it affects negatively the analytical recoveries, while derivatization also is not desirable process because it is additional step of analytical procedure, what brings further errors and losses in determined analyte concentration. Recently, extraction and derivatization processes are often performed simultaneously. After extraction (and derivatization if required), the determination of BAs is most commonly performed by means of chromatographic methods: high-performance liquid chromatography (HPLC), gas chromatography (GC) and capillary electrophoresis (CE).

6.1. Liquid chromatography

The most commonly popular technique for the determination of BAs in food products is the HPLC with Reversed-Phase separation, using C18 columns, however, other solutions are proposed. Table 1 summarizes the chromatographic methods used to quantify BAs in the food matrices, in the last decade.

Generally, the most common resolution to sample preparation prior to final separation is application of solvent extraction and derivatization process, not only to remove compounds that may interfere with the analysis but also to concentrate the analytes. In some cases, a simple treatment with polyvinylpyrrolidone (PVPP) to remove some phenolic compounds is used. Considering derivatization processes, pre-column or post-column modes are usually performed, since BAs do not have enough absorption in the UV–Vis or FLD wavelength ranges. This step is also applied to improve the separation in the RP columns, reducing the polarity

Table 1
Information on analytical methodologies based on gas and liquid chromatography developed for BAs determination in food samples.

Matrice	Analyte	Sample preparation	Derivatization	LOD/LOQ	Recovery	Separation technique	Detection	Ref.
Wine	TRYP, HIS, CAD, 2-PE, HEX, TYR, SPR	UA-DLLME	YES: BCEC-Cl	LOD: 1.1–7.8 ng/mL LOQ: 3.5–26.1 ng/mL	91.2–108.3%	HPLC (column: Hypersil C18)	FLD	[38]
	PUT, CAD, 2-PE, SPR, SPRE, HIS, TYR	Addition of PVPP	YES: dansyl chloride	LOD: 0.09–0.30 mg/L LOQ: 0.30–1.00 mg/L	>78%	HPLC (column: C ₁₈ YMC-Pack ODS-A)	UV	[84]
	HIS, TYR, 2-PE, SER, TRYP, OCT, DOP, CAD, PUT, AGM, SPR, SPRE	Solvent extraction	YES: OPA (post column derivatization)	LOD: 0.05–0.2 mg/L LOQ: 0.1–0.3 mg/L	93.56–103.28%	UHPLC (column: Acquity UPLC BEH C18)	FLD	[85]
	PROP, DMET, DET, MET, TRYP, CAD, SPR, 2-PE, TYR, PUT, HIS, BUT, HEX, IPENT, IBUT, SPRE, AGM	Dilution with water	YES: <i>p</i> -toluenesulfonyl chloride	LOD: 0.023–83 µg/dm ³ LOQ: 0.075–270 µg/dm ³	–	HPLC (column: Gemini C-18)	MS/MS	[86]
	BUT, CAD, DET, DMET, ET, MET, HIS, PROP, PUT, SPR, TRYP, TYR, 2-PE	DLLME	YES: IBCF	LOD: 1.1–4.1 µg/L LOQ: 3.3–12.3 µg/L	77–105%	GC (column: ZB-5MS, 0.25 µm)	MS	[24]
Beer	MET, DMET, ET, DET, PROP, IPROP, BUT, IBUT, AM, IAM, 2-MBUT, HEX, PYR, PIP, MOR, PUT, CAD, 2-PE, HIS, TYR	DLLME	YES: IBCF	LOD: 1.8–36.8 µg/L	85–111%	GC (column: HP-5MS, 0.25 µm)	MS	[87]
	TRYP, HIS, CAD, 2-PE, HEX, TYR, SPR	UA-DLLME	YES: BCEC-Cl	LOD: 1.1–7.8 ng/mL LOQ: 3.5–26.1 ng/mL	91.2–108.3%	HPLC (column: Hypersil C18)	FLD	[39]
	PUT, HIS, CAD, 2-PE, HEX, TYR, SPRE, SPR	Solvent extraction	YES: EAC	LOD: 0.27–0.69 ng/mL	94.18–104.02%	HPLC (column: Hypersil BDS C ₁₈)	FLD	[88]
	PROP, DMET, DET, MET, TRYP, CAD, SPR, 2-PE, TYR, PUT, HIS, BUT, HEX, IPENT, IBUT, SPRE, AGM	Dilution with water	YES: <i>p</i> -toluenesulfonyl chloride	LOD: 0.023–83 µg/dm ³ LOQ: 0.075–270 µg/dm ³	–	HPLC (column: Gemini C-18)	MS/MS	[86]
	TYR, MET, DMET, ET, IPROP, DET, IBUT, 2MBUT, PYR, IAM, MOR, PIP, AM, 2-PE, PUT, CAD, HIS	DLLME	YES: IBCF	LOD: 0.3–2.9 µg/L LOQ: 1.0–9.5 µg/L	72–113%	GC (column: DB-5MS, 0.18 µm)	MS	[59]
Cheese	TRYP, HIS, CAD, 2-PE, HEX, TYR, SPR	UA-DLLME	YES: BCEC-Cl	LOD: 1.1–7.8 ng/mL LOQ: 3.5–26.1 ng/mL	91.2–108.3%	HPLC (column: Hypersil C18)	FLD	[38]
	PUT, HIS, CAD, 2-PE, HEX, TYR, SPRE, SPR	Solvent extraction	YES: EAC	LOD: 0.27–0.69 ng/mL	94.98–101.38%	HPLC (column: Hypersil BDS C ₁₈)	FLD	[88]
	ETH, MET, AGM, HIS, DMET, ET, OCT, PYR, DOP, IPROP, PROP, TYR, PUT, BUT, CAD, TRYP, 2-PE, 3-METBUT, SPRE, SPR	Solvent extraction	YES: AQC	LOD: 4–162 mg/g	–	ULPC (column: Nova-Pak™ C ₁₈)	FLD	[3]
	HIS, TYR, 2-PE, SER, TRYP, OCT, DOP, CAD, PUT, AGM, SPR, SPRE	Solvent extraction	YES: OPA (post column derivatization)	LOD: 0.05–0.2 mg/L LOQ: 0.1–0.3 mg/L	79.10–103.00%	UHPLC (column: Acquity UPLC BEH C18)	FLD	[73]
	PUT, CAD, SPRE, SPR, 2-PE, HIS, TYR, TRYP	Solvent extraction	YES: dansyl chloride	LOD: 0.032–0.098 µg/L LOQ: 0.11–0.32 µg/L	–	UHPLC (column: Zorbax Eclipse XDB – C18 column)	UV	[89]
Yogurt	TRYP, HIS, CAD, 2-PE, HEX, TYR, SPR	UA-DLLME	YES: BCEC-Cl	LOD: 1.1–7.8 ng/mL LOQ: 3.5–26.1 ng/mL	91.2–108.3%	HPLC (column: Hypersil C18)	FLD	[38]
Sausage	TRYP, HIS, CAD, 2-PE, HEX, TYR, SPR	UA-DLLME	YES: BCEC-Cl	LOD: 1.1–7.8 ng/mL LOQ: 3.5–26.1 ng/mL	91.2–108.3%	HPLC (column: Hypersil C18)	FLD	[38]
	PUT, HIS, CAD, 2-PE, HEX, TYR, SPRE, SPR	Solvent extraction	YES: EAC	LOD: 0.27–0.69 ng/mL	96.37–101.30%	HPLC (column: Hypersil BDS C ₁₈)	FLD	[35]
	HIS, TYR, 2-PE, SER, TRYP, OCT, DOP, CAD, PUT, AGM, SPR, SPRE	Solvent extraction	YES: OPA (post column derivatization)	LOD: 0.05–0.2 mg/L LOQ: 0.1–0.3 mg/L	87.33–100.24	UHPLC (column: Acquity UPLC BEH C18)	FLD	[73]
Rice wine	TRYP, HIS, CAD, 2-PE, HEX, TYR, SPR	UA-DLLME	YES: BCEC-Cl	LOD: 1.1–7.8 ng/mL LOQ: 3.5–26.1 ng/mL	91.2–108.3%	HPLC (column: Hypersil C18)	FLD	[38]
Corn oil	HIS, 2-PE, PUT, TYR, CAD, TRYP, SPR, SPRE	Solvent extraction	YES: dansyl chloride	LOD: 0.01–0.10 mg/kg LOQ: 0.02–0.31	94.51–106.72%	HPLC (column: Nova-Pak C ₁₈ 4 µm)	UV–VIS	[90]
Fresh fish	MET, HIS, ET, TYR, PROP, TRYP, 2-PE, ISM, HEP, PUT, PUT, CAD	UA-LLE	YES: naphthalene-2,3-dicarboxaldehyde	LOD: 2.5 and 330 mg/kg	81.0–102.5%	HPLC (column: Inertsil ODS-3)	FLD	[85]
	PUT, HIS, CAD, 2-PE, HEX, TYR, SPRE, SPR	Solvent extraction	YES: EAC	LOD: 0.27–0.69 ng/mL	95.81–103.47%	HPLC (column: Hypersil BDS C ₁₈)	FLD	[88]
	HIS, TYR, 2-PE, SER, TRYP, OCT, DOP, CAD, PUT, AGM, SPR, SPRE	Solvent extraction	YES: OPA (post column derivatization)	LOD: 0.05–0.2 mg/L LOQ: 0.1–0.3 mg/L	83.08–98.81%	UHPLC (column: Acquity UPLC BEH C18)	FLD	[73]
	PUT, CAD, SPRE, SPR, 2-PE, HIS, TYR, TRYP	Solvent extraction	YES: dansyl chloride	LOD: 0.032–0.098 µg/L LOQ: 0.11–0.32 µg/L	–	UHPLC (column: Zorbax Eclipse XDB – C18 column)	UV	[88]

(continued on next page)

Table 1 (continued)

Matrice	Analyte	Sample preparation	Derivatization	LOD/LOQ	Recovery	Separation technique	Detection	Ref.
Canned fish	MET, HIS, ET, TYR, PROP, TRYP, 2-PE, ISM, HEP, PUT, PUT, CAD	UA-LLE	YES: naphthalene-2,3-dicarboxaldehyde	LOD: 2.5 and 330 mg/kg	81.0–102.5%	HPLC (column: Inertsil ODS-3)	FLD	[85]
		Solvent extraction	YES: dansyl chloride	LOD: 4.43–7.34 µg/L LOQ: 14.76–24.45 µg/L	90.9–106.3%	HPLC (column: Waters Spherisorb 5 µm ODS2)	UV–VIS	[91]
	HIS, TYR, PUT, CAD	DI-SPME	YES: IBCF	LOD: 2.98–45.3 µg/kg LOQ: .83–149 µg/kg	78.9–110%	GC (column: RXI-5MS, 0.25 µm)	MS	[75]
Shrimp sauce	TRYP, PUT, CAD, HIS, TYR, SPRE	HF-LPME	YES: dansyl chloride	LOD: 0.0075–0.030 µg/mL LOQ: 0.03–0.10 µg/mL	88.6–103%	HPLC (column: Waters Spherisorb 5 µm ODS2)	UV	[35]
		Solvent extraction	YES: dansyl chloride	LOD: 4.43–7.34 µg/L LOQ: 14.76–24.45 µg/L	90.9–106.3%	HPLC (column: Waters Spherisorb 5 µm ODS2)	UV–VIS	[91]
Tomato ketchup	TRYP, PUT, CAD, HIS, TYR, SPRE	HF-LPME	YES: dansyl chloride	LOD: 0.0075–0.030 µg/mL LOQ: 0.03–0.10 µg/mL	86.7–104%	HPLC (column: Waters Spherisorb 5 µm ODS2)	UV	[35]
		Solvent extraction	YES: dansyl chloride	LOD: 4.43–7.34 µg/L LOQ: 14.76–24.45 µg/L	90.9–106.3%	HPLC (column: Waters Spherisorb 5 µm ODS2)	UV–VIS	[91]
Meat	TYR, PUT, CAD, SPR, SPRE	Solvent extraction	YES: benzoyl chloride	LOD: 0.03–1.25 µg/L LOQ: 0.15–5.00 µg/L	64.40–112.22%	HPLC (column: Tracer Extrasil ODS2)	UV–VIS	[92]
		Solvent extraction	YES: dansyl chloride	LOD: 0.032–0.098 µg/L LOQ: 0.11–0.32 µg/L	–	UHPLC (column: Zorbax Eclipse XDB – C18 column)	UV	[89]
	HIS, 2-PE, PUT, TYR, CAD, TRYP, SPR, SPRE	Solvent extraction	YES: dansyl chloride	LOD: 0.01–0.10 mg/kg LOQ: 0.02–0.31	86.63–106.23	HPLC (column: Nova-Pak C ₁₈ 4 µm)	UV–VIS	[90]
Soy bean products	TRYP, PUT, CAD, HIS, TYR, SPRE	Solvent extraction	YES: dansyl chloride	LOD: 4.43–7.34 µg/L LOQ: 14.76–24.45 µg/L	90.9–106.3%	HPLC (column: Waters Spherisorb 5 µm ODS2)	UV–VIS	[91]
		Solvent extraction	YES: dansyl chloride	LOD: 4.43–7.34 µg/L LOQ: 14.76–24.45 µg/L	90.9–106.3%	HPLC (column: Waters Spherisorb 5 µm ODS2)	UV–VIS	[91]
Juice	HIS, 2-PE, PUT, TYR, CAD, TRYP, SPR, SPRE	Solvent extraction	YES: dansyl chloride	LOD: 0.01–0.10 mg/kg LOQ: 0.02–0.31	92.35–107.01%	HPLC (column: Nova-Pak C ₁₈ 4 µm)	UV–VIS	[90]
		Solvent extraction	YES: IBCF	LOD: 1.8–36.8 µg/L	92–112%	GC (column: HP-5MS, 0.25 µm)	MS	[87]
	MET, DMET, ET, DET, PROP, IPROP, BUT, IBUT, AM, IAM, 2-MBUT, HEX, PYR, PIP, MOR, PUT, CAD, 2-PE, HIS, TYR	Solvent extraction	YES: EAC	LOD: 0.27–0.69 ng/mL	93.74–103.40%	HPLC (column: Hypersil BDS C ₁₈)	FLD	[88]
Chocolate, vegetables, and fruits	TYR	Solvent extraction	NO	LOD: 10 ng/mL LOQ: 30 ng/mL	72.0–85.8%	HPLC (column: LiChrospher 100 RP-18e)	FLD-DAD	[93]
Fermented cow's and goat's milks	TYR, CAD, PUT, HIS, SPRE	Solvent extraction	YES: benzoyl chloride	LOD: 0.03–1.30 mg/L LOQ: 20–5.00 mg/L	91–107%	HPLC (column: Extrasil Tracer ODS2)	DAD	[47]
Peanut butter	HIS, 2-PE, PUT, TYR, CAD, TRYP, SPR, SPRE	Solvent extraction	YES: dansyl chloride	LOD: 0.01–0.10 mg/kg LOQ: 0.02–0.31	93.90–107.74%	HPLC (column: Nova-Pak C ₁₈ 4 µm)	UV–VIS	[90]
Mushrooms	PUT, CAD, SPRE, SPR, 2-PE, HIS, TYR, TRYP	Solvent extraction	YES: dansyl chloride	LOD: 0.032–0.098 µg/L LOQ: 0.11–0.32 µg/L	–	UHPLC (column: Zorbax Eclipse XDB – C18 column)	UV	[89]

AGM: agmatine; AM: amylamine; AQC: 6-aminoquinolyl-N-hydroxy-succinimidyl carbamate; BCEC-Cl: 2-(11H-benzo[a]carbazol-11-yl) ethyl carbonochloridate; BUT: butylamine; CAD: cadaverine; DET: diethylamine; DI-SPME: direct immersion solid phase microextraction; DLLME: dispersive liquid liquid microextraction; DMET: dimethylamine; DOP: dopamine; EAC: ethyl-acridine-sulfonyl chloride; ET: Ethylamine; ETH: ethanolamine; FLD: fluorescence detection; HEP: heptylamine; HEX: 1,6-hexamethylenediamine; HIS: histamine; HPLC: high-performance liquid chromatography; HS-SPME: head space solid phase microextraction; IAM: Isoamylamine; IBCF: isobutyl chloroformate; IBUT: isobutylamine; IPENT: isopentylamine; IPROP: isopropylamine; ISM: Isoamylamine; LLE: liquid-liquid extraction; 2-MBUT: 2-Methylbutylamine; MET: Methylamine; 3-METBUT: 3-methylbutylamine; MOR: morfoline; OCT: octopamine; 2-PE: 2-phenylethylamine; PIP: piperidine; PROP: propylamine; PUT: putrescine; PVPP: Polyvinylpyrrolidone; PYR: pyrrolidine; SER, serotonin; SPR: spermine; SPRE: spermidine; TRYP: Tryptamine; TYR: tyramine; UA-DLLME: ultrasound-assisted dispersive liquid–liquid microextraction; UA-LLE: ultrasound-assisted liquid–liquid microextraction; ULPC: ultra-performance liquid chromatography; UV: ultrafiolet.

of the original compounds. Several derivatization reagents have been reported, however, dansyl chloride (DNS-Cl) has been commonly applied in the last years, since that its derivatives can be detected using DAD, FLD and MS. Taking into account this three type of detection type, FLD revealed better sensitivity to detect dansylated amines. DNS-Cl is very often used as pre-column derivatizing reagent in the determination of BAs in food products, mainly because it produces stable derivatized compounds. From the other hand, the dansylation reaction is a time consuming process (10–60 min) and it requires the application of external temperature (40–70°C). However, a new method to perform the dansylation at room temperature, during 20 min, using ionic liquids as media for the derivatization of BAs in wines was introduced [72]. Another important reagent for derivatization of BAs is *ortho*-phthalaldehyde (OPA) compound. Although, OPA derivatives are less stable, the process can be performed at room temperature in a short time. OPA reagent can be used without using any preliminary separation or clean-up. Moreover, it can be used for post column derivatization [73].

6.2. Capillary electrophoresis

The second most commonly performed technique applied to the determination of biogenic amines in foods is capillary electrophoresis (CE), despite its low sensitivity when compared with other methods. CE has several advantages: it is simple, rapid, and reliable, making it a very useful tool for screening a large number of samples in a short period of time [26]. Moreover, CE is efficient, as well as inexpensive in terms of reagent consumption, and presents a lower reproducibility of migration times than LC. Therefore, the usefulness of this technique will depend on the expected BAs levels in food samples. Several approaches are used to solve the problem of BA determination in food samples using this technique: i) application of derivatization process, ii) application of selected buffer system without derivatization, iii) application of specific kits. Off-line pre-column derivatization is most commonly used, with the resulting derivatives being injected into the CE. Derivatization processes can also be coupled with some innovative on-line pretreatment procedures, contributing to enhanced sensitivity of these methods. Different type of derivatizing reagents are used, and these depend on the detection technique coupled to CE. Fluorometric methods are used owing to fluorescence of BAs at some pH range and reaction of BAs with suitable agents. This is achieved mainly by reaction with *o*-phthalaldehyde, dansylchloride or dabsylchloride and β -naphthol [26]. *o*-phthalaldehyde has the disadvantages that it reacts only with primary amines and the instability of its fluorescent derivatives, while dabsyl- and dansylchloride react with both primary and secondary amino groups and provide stable derivatives. Dabsyl derivatives are colored, so they absorb in the visible region, while dansyl derivatives combine a unique feature of being both fluorescent and detectable in the UV-region [74]. Indirect detection or electrochemical detection has also been used after CE separation of BAs [74]. In indirect methods, a previous step involving clean-up, such as added polyvinylpyrrolidone (PVP) followed by filtration, is usually performed. As a detection technique, UV, photometric detection and MS (or MS/MS) are usually applied. Information on the latest developments in determination of BAs in food samples by CE techniques are presented in Table 2.

6.3. Gas chromatography

Gas chromatography is not as commonly used as liquid chromatography or capillary electrophoresis for determination of BAs in food samples. However, some studies are reported, mainly with the application of mass spectrometer as a detection technique. Gas

chromatography has been specifically applied in fermented beverages by some authors in order to offer alternative methods to LC. Before analysis, derivatization of analytes need to be performed to change the properties of analytes, typically to increase volatile properties and to decrease polarities of BAs. Depending on the detection technique used for final determination, different type of derivatizing reagents can be applied, e.g. pentafluorobenzoyl chloride, however, the most popular are alkylated chloroformates. The fact that the derivatization reaction can be performed in an aqueous basic solution is one advantage of using these compounds. In addition, the derivatization process occurred at room temperature in a short time and can be coupled with extraction procedure. Moreover, the derivatives obtained are characterized by such properties which are useful for application of analytical procedures based on GC techniques at the mixture separation, detection, and quantitative determination stages.

Recently, several works focused on determination of BAs in food samples such as fish or alcoholic beverages by using isobutyl chloroformate coupled with solid phase microextraction (SPME) [75] or dispersive liquid-liquid microextraction (DLLME) [24,59]. Reported methods were efficient and highly reproducible, allowing the accurate identification and quantification of a higher number of biogenic amines. Furthermore, the derivatization procedure and the overall analysis time were faster than in some LC methods, reaching LODs of the same order.

6.4. Non-chromatographic approaches for determination of biogenic amines

Chromatographic techniques such as GC, LC or CE are among the most important instrumental methods for precise quantitative analysis of BAs. From the other site, these methods are often time-consuming as well as they require considerable skill [76]. In addition, a disadvantage of the HPLC, CE and GC-MS analyses in general is the long and tedious sample pretreatment. A further principal drawback of these methods is the requirement of (sometimes harmful) organic solvents of HPLC grade quality, whereby the cost for their purchase and disposal has to be taken into consideration. Enzymatic as well as immuno-enzymatic methods such as commercially available test kits designed to histamine determination can also be applied for routine analysis with semiquantitative and quantitative determination of biogenic amines.

Disposable screen-printed electrode biosensors with enzymes have been shown as a step forward to reduced sample pretreatment. Biosensors combines biological recognition through enzyme specificity with construction simplicity and they are a good and cheap alternative for the traditional methods of analytes determination. In biosensors used for BA determination, amine oxidases (AO) based electrodes, which catalyze the oxidative deamination of primary amines, diamines and substituted amines to produce aldehyde, ammonia and hydrogen peroxide, have been reported [77]. AO-sensors can base on platinum screen-printed electrodes (SPEs), using a glutaraldehyde – bovine serum albumin cross-linking immobilization procedure [78]. The procedure involves the measurement of the H₂O₂ produced by applying a potential higher than 600 mV. Monoamine oxidase (MAO)/horseradish peroxidase (HRP) and diamine oxidase (DAO)/horseradish peroxidase (HRP) based biosensors using screen-printed carbon electrodes for the determination of BAs in fish samples was also reported. The enzymes have been covalently immobilized onto the carbon working electrode, previously modified by an aryl diazonium salt, using hydroxysuccinimide and carbodiimide. The detection has been performed by measuring the cathodic current due to the reduction of the mediator hydroxymethylferrocene at a low potential. A linear response range from 0.2 up to 1.6 μ M and



Table 2
Information on analytical methodologies based on electrophoresis techniques developed for BAs determination in food samples.

Matrice	Analyte	Electrophoretic method	Sample preparation	Derivatization	Conditions	LOD/LOQ	Recovery	Detection	Ref.
Meat	TMA, PUT, CAD, SPR, TRYP, SPRE, 2-PE, TYR	Capillary zone electrophoresis	Homogenization, SPE	YES: dansyl chloride	Running buffer: 6 mM copper sulfate + 6 mM 18-crown-6-ether + 4 mM formic acid – pH 2.7 – in water	LOD: 0.2–0.6 µg/mL LOQ: 0.7–1.9 µg/m	–	Photometric detection	[74]
Beer	ETH, TRYP, TRPH	Microchip capillary electrophoresis	Degassed, Phosphate buffer	NO	Glass microchannel chips, model MC-BF4-001. Running buffer: 20 mM phosphate (pH 2.5) and a high voltage of 2.5 kV	LOD: 1.4–6.8 mg/L	–	Electrochemical detection	[94]
	HIS, TYR, PUT, CAD, SPR, SPRE, TRYP, 2-PE	Capillary electrophoresis	Degassed, filtration, addition of PVP	NO	PVA-coated silica capillary 700 mm, 50 µm. Background electrolyte: 0.5 M acetic acid (pH 2.5)	LOD: 1–2 µg/L LOQ: 3–8 µg/L	83–110%	MS/MS	[95]
		Capillary isotachopheresis	Degassed	NO	PTFE analytical capillary 160 mm, 300 µm. Leading electrolyte: 5 mM Ba(OH) ₂ + 15 mM valine + 1% hydroxyethylcellulose (pH 8.5). Terminating electrolyte: 0.02 M TRIS + 0.1 M HCl (pH 8.3).	LOD: 0.2–0.48 mg/L	90–101%	Conductometric detector	[96]
	Sausage	TMA, PUT, CAD, SPR, TRYP, SPRE, 2-PE, TYR	Capillary zone electrophoresis	Homogenization, SPE	YES: dansyl chloride	Running buffer: 6 mM copper sulfate + 6 mM 18-crown-6-ether + 4 mM formic acid – pH 2.7 – in water	LOD: 0.2–0.6 µg/mL LOQ: 0.7–1.9 µg/m	–	Photometric detection
Wine	HIS, TYR, PUT, CAD, SPR, SPRE, TRYP, 2-PE	Capillary electrophoresis	Filtration, addition of PVP	NO	PVA-coated silica capillary 700 mm, 50 µm. Background electrolyte: 0.5 M acetic acid (pH 2.5)	LOD: 1–2 µg/L LOQ: 3–8 µg/L	90–113%	MS/MS	[95]
		Capillary isotachopheresis	Filtration	NO	PTFE analytical capillary 160 mm, 300 µm; Leading electrolyte: 5 mM Ba(OH) ₂ + 15 mM valine + 1% hydroxyethylcellulose (pH 8.5). Terminating electrolyte: 0.02 M TRIS + 0.1 M HCl (pH 8.3).	LOD: 0.2–0.48 mg/L	90–101%	Conductometric detector	[96]
	HIS, TYR, PUT, CAD, SPRE, TRYP, 2-PE	MECK	Filtration	YES: FITC	Silica capillary 580 mm, 50 µm. Running buffer: Brij 35, borate buffer (pH 9.6)	LOD: 0.06–0.11 µg/L	93–104%	LIF	[97]
	HIS, TRYP, TYR, 2-PE	MECK	Direct analysis	NO	Fused silica capillary 645 mm, 50 µm. Electrophoresis buffer of 20 mM borate (pH 9.3) containing 30 mM SDS and 5% (v/v) methanol	–	–	UV	[98]
	2-PE, SPR, SPRE, CAD, PUT	MECK	Homogenization (6% trichloroacetic acid), filtration, dilution with water	YES: FBQCA	Running buffer: 25 mM pH 9.5 boric acid, 25 mM SDS, and 27% ACN. U: 22.5 kV	LOD: 0.4–12 nM	–	LIF	[99]
	Fish	2-PE, SPR, SPRE, CAD, PUT	MECK	Homogenization (6% trichloroacetic acid), filtration, dilution with water	YES: FBQCA	Running buffer: 25 mM pH 9.5 boric acid, 25 mM SDS, and 27% ACN. U: 22.5 kV	LOD: 0.4–12 nM	–	LIF
	TMA, PUT, CAD, SPR, TRYP, SPRE, 2-PE, TYR	Capillary zone electrophoresis	Homogenization, SPE	YES: dansyl chloride	Running buffer: 6 mM copper sulfate + 6 mM 18-crown-6-ether + 4 mM formic acid – pH 2.7 – in water	LOD: 0.2–0.6 µg/mL LOQ: 0.7–1.9 µg/m	–	Photometric detection	[74]
Soy sauce	2-PE, SPR, SPRE, CAD, PUT	MECK	Homogenization (6% trichloroacetic acid), filtration, dilution with water	YES: FBQCA	Running buffer: 25 mM pH 9.5 boric acid, 25 mM SDS, and 27% ACN. U: 22.5 kV	LOD: 0.4–12 nM	–	LIF	[99]
Oysters	HIS, TYR, 2-PE, PUT, SPR	Capillary electrophoresis	Homogenization (6% perchloric acid)	NO	Phosphate buffer at pH 7.00 in the detection cell, 5 mmol/L Ru(bpy) ₃ ²⁺ + and 50 mmol/L phosphate buffer at pH 7.00 as the separation buffer	LOD: 6.0 × 10 ⁻⁴ – 9.6 × 10 ⁻² µg/mL	92.5–104.2%	electrochemiluminescence	[100]

AGM: agmatine; AM: amylamine; BUT: butylamine; CAD: cadaverine; DET: diethylamine; DI-SPME: direct immersion solid phase microextraction; DLLME: dispersive liquid liquid microextraction; DMET: dimethylamine; DOP: dopamine; EAC: ethyl-acridine-sulfonyl chloride; ET: Ethylamine; ETH: ethanolamine; FBQCA: 3-(4-fluorobenzoyl)-2-quinolinecarboxaldehyde; FLD: fluorescence detection; FITC: Fluorescein isothiocyanate isomer; HEP: heptylamine; HEX: 1,6-hexamethylenediamine; HIS: histamine; HPLC: high-performance liquid chromatography; HS-SPME: head space solid phase microextraction; IAM: Isoamylamine; IBCF: isobutyl chloroformate; IBUT: isobutylamine; IPENT: isopentylamine; IPROP: isopropylamine; ISM: Isoamylamine; LIF: Laser-induced fluorescence detector; LLE: liquid-liquid extraction; 2-MBUT: 2-Methylbutylamine; MECK: micellar electrokinetic capillary chromatography; MET: Methylamine; 3-METBUT: 3-methylbutylamine; MOR: morfoline; OCT: octopamine; 2-PE: 2-phenylethylamine; PIP: piperidine; PROP: propylamine; PUT: putrescine; PVPP: Polyvinylpyrrolidone; PYR: pyrrolidine; SER, serotonin; SPR: spermine; SPRE: spermidine; TMA: trimethylamine; TRYP: Tryptamine; TRPH, tryptohane TYR: tyramine; UA-DLLME: ultrasound-assisted dispersive liquid–liquid microextraction; UA-LLE: ultrasound-assisted liquid–liquid microextraction; ULPC: ultra-performance liquid chromatography; UV: ultrafiolet.

from 0.4 to 2.4 μM of histamine was obtained for DAO/HRP and MAO/HRP based biosensors, respectively.

An interesting idea of application of polymer layers with different changes in absorption due to interaction with aromatic, aliphatic, and polyamines was reported [79]. Several colorimetric methods using test spots are also known and many times applied for routine analysis of food samples [80]. These can be semi-quantitatively evaluated by visual readout of the originated color in comparison to a reference color scale [76]. Colorimetric sensor arrays consisting either of several amine sensitive dyes [81] or organic liquid crystals [82] also have been applied for the determination of volatile amines in food and beverage samples.

Nowadays, an amperometric sensors are also many often used for BAs determination mainly in food products. An example of this type of biosensor composed with pea seedling AO and screen printed carbon electrode modified with MnO_2 for determining selected BAs was designed in 2007 [83]. The bio-component was immobilized with Nafion solution. The enzymatically produce hydrogen peroxide was determined to quantify BAs. The limits of detection were determined to be 0.3 μM for cadaverine and putrescine and 3.0 μM for tyramine and histamine. The sensor was used to quantify BAs in chicken meat samples. Enzymatic sensors reduce the time needed for analysis as well as offer a rapid screening method for industrial food quality testing. The use of disposable transducers, such as SPEs, boosts the intrinsic characteristics of enzymatic sensors. Screen-printing technology, which offers design flexibility, process automatization and good reproducibility in the transducers fabrication, had been shown as a method for mass production of biosensors at low cost [77].

In 2010, a new chromogenic and fluorogenic dry chemistry sensing spots based on filter paper containing an amine-reactive chromogenic probe and fluorescein as a green fluorescent (but amine insensitive) reference dye incorporated in a hydrogel matrix was presented [76]. These test spots can quantitate BAs upon dipping into the sample. The test spots were evaluated with six different biogenic amines at concentrations between 0.01 and 10 mM using the RGB readout option of a digital camera. This sensor spots represent an attractive alternative to existing schemes for sensing biogenic amines. Its digital read-out makes it more robust, and the use of conventional cameras goes along current trends toward simplified methods for absorption-based and emission-based detection schemes. In addition, the method also may enable high-throughput analysis and in-field examinations and does not require sophisticated instrumentation or trained personnel.

6.5. Green assessment of selected analytical procedures applied for determination of BAs in wine and fish samples

Green Analytical Chemistry (GAC) is an ideology concerned with the development of analytical procedures that minimize consumption of hazardous reagents and solvents as well as maximize safety for operators and the environment. Recently, there have been significant developments in methodological and technological tools to prevent and reduce the deleterious effects of analytical activities. And although, there are numerous examples of analytical procedures reported in the literature that are claimed to be green by their authors, these statements are not supported by any evidence, in the form of applied greenness metrics, and comparisons with previously developed analytical or standard procedures. Therefore, calculations that give an answer for question whether an analytical procedure can be regarded green, should be performed utilizing the tools that serve such assessment.

In this section several analytical procedures used for biogenic amines determination in wine [24,86,87,98] and fish samples [75,85,91,99] by application of different type of methods (GC, LC

and CE), that are mentioned in this paper are assessed in respect to the green character. To evaluate these selected protocols, Analytical Eco-Scale was used.

Considering the penalty points (PPs) calculated for each procedure used for wine analysis (Table 3), it can be concluded, that Procedure 4 (CE-DAD) can be assigned as green (Score: 85). However, this is not quantitative analysis. Also Procedure 3 gives satisfied results (Score: 73). The worst procedure in term of "green" profile is Procedure 1 which apply DLLME extraction type coupled with derivatization process (Score: 62), and although Procedure 2 (Score: 69) is based on the same techniques, it consume much less reagents.

Considering the PPs calculated for each procedure used for fish analysis, it can be stated that Procedure 8 (CE-LIF) can be assigned as green (Score: 77). Procedures: 6 and 7 are similar taking into account its "green" character, while Procedure 5 (Score: 48) is far from being green.

Without a doubt, this Analytical Eco-Scale is a good semi-quantitative tool for laboratory practice and educational purposes. It is simple and fast to use, has well-defined criteria of evaluation and can be applied to any known and new methodologies.

7. Conclusions

To protect the health of consumers, there is a requirement for more stringent regulations and more diligent monitoring of foods by regulators, vendors and producers. New green analytical extraction procedures are needed for monitoring of BAs in the food matrices and should be fast, inexpensive and easily adopted within analytical laboratories worldwide.

Considering the quality control as well as research requirements, a more precise identification of analytes is needed, hence the requirement for sensitive methods to detect slight changes in BAs profile.

The quantitation of BAs can be highly problematic due to several factors. Biogenic amines are small, polar compounds and these issues can make the separation challenging when using reverse phase chromatography. Thus, the assortment of chromatographic column is very important as some specific reverse phase columns can separate selected classes of BAs, and moreover, can add additional cost to whole analysis because are more expensive. Due to the fact that biogenic amines have highly similar structures to the compounds in the same sub-group as well similar chemical and physical properties to other compounds presented in food samples. This brings problem in providing effective separation by isocratic chromatography which is required for certain quantitative methods and is problematic when quantitating a number of different BAs simultaneously.

Additional problem in quantitation of biogenic amines is that they occur at very low concentration level (sub-ng/mL range), what means that a highly sensitive method is needed to accurately quantitate the analytes. This issue is even more problematic when working with low volume of the samples which means the absolute quantities of BAs in the total sample are low. Moreover, the method usually require time consuming and expensive sample preparation before analysis in order to achieve the high level of sensitivity required to quantitate the low concentration in food samples.

The food matrix in which BAs are commonly examined is often complex. Even aqueous matrixes such as wine, beer, dairy fermented products are problematic because usually contain interfering compounds such as polyphenols, lipids, proteins (depending on the sample). Hence, the sample preparation and extraction of these compounds is time consuming. It also means that these compounds need efficient separation and retention away from interfering compounds which can cause further problems with



Table 3
Calculated PPs for evaluated analytical procedures for biogenic amines determination in wine (Procedures 1–4) and fish samples (Procedures 5–8).

Wine analysis: PROCEDURE 1 [87]		Wine analysis: PROCEDURE 2 [24]		Wine analysis: PROCEDURE 3 [86]		Wine analysis: PROCEDURE 4 [98]	
Reagents	PPs	Reagents	PPs	Reagents	PPs	Reagents	PPs
NaOH	1	Pyridine	1	Water	0	Methanol	6
Phosphate buffer 0.5 M	0	Internal standard	4	Formic acid (0.1%)	2	Internal standard	4
Internal standard	4	HCl	3	Acetonitrile	8	Borate buffer (20 mM)	0
HCl	4	Chloroform	2	Borate buffer (0.5 M)	0	Sodium dodecyl sulfate	0
Sodium azide	1	Isobutyl chloroformate	8	Tosyl chloride	8		
Toluene	6	MeOH	6				
Isobutyl chloroformate	8						
MeOH	6						
	Σ 30		Σ 24		Σ 18		Σ 10
Instruments	PPs	Instruments	PPs	Instruments	PPs	Instruments	PPs
Transport	1	Transport	1	Transport	1	Transport	1
GC-MS	2	GC-MS	2	HPLC-MS/MS	4	CE-DAD	2
Occupational hazard	0	Occupational hazard	0	Occupational hazard	2	Waste	2
Centrifugation	1	Waste	1	Waste	2		
Temperature storage	1						
Waste	3						
	Σ 8		Σ 4		Σ 9		Σ 5
Total PPs: 38		Total PPs: 31		Total PPs: 27		Total PPs: 15	
Score: 62		Score: 69		Score: 73		Score: 85	
Fish analysis: PROCEDURE 5 [85]		Fish analysis: PROCEDURE 6 [91]		Fish analysis: PROCEDURE 7 [75]		Fish analysis: PROCEDURE 8 [99]	
Reagents	PPs	Reagents	PPs	Reagents	PPs	Reagents	PPs
Methanol	8	HCl (0.1 M)	3	n-hexane	8	HCl	3
Internal standard	4	NaHCO ₃	0	Internal standard	4	Water	0
Borate buffer	0	NaOH (2 M)	1	Isobutyl chloroformate	8	NaOH (1 M)	1
Acetonitrile	8	Dansyl chloride	8	Trichloroacetic acid (5%)	2	Sodium dodecyl sulfate	0
Acetone	4	Acetone	4	Iso-octane	4	Boric acid	1
HCl	3	Water	0	NaCl (25%)	0	Acetonitrile	4
Trichloroacetic acid	2	Acetonitrile	8			FBQCA	4
Hexane	8	Glutamic acid	4			Methanol	4
NaOH (1 M)	1					Trichloroacetic acid	2
Potassium cyanide	2						
	Σ 40		Σ 28		Σ 26		Σ 19
Instruments	PPs	Instruments	PPs	Instruments	PPs	Instruments	PPs
Transport	1	Transport	1	Transport	1	CE-LIF	2
HPLC-FLD	2	HPLC-UV-VIS	2	GC-MS	2	Homogenization	1
Occupational hazard	1	Occupational hazard	1	Occupational hazard	0	Occupational hazard	0
Centrifugation	1	Centrifugation	1	Centrifugation	1	Waste	1
Temperature storage	1	Temperature storage	1	Temperature storage	1		
Sonification	1	Waste	3	Homogenization	1		
Waste	5			Waste	3		
	Σ 12		Σ 9		Σ 9		Σ 4
Total PPs: 52		Total PPs: 37		Total PPs: 35		Total PPs: 23	
Score: 48		Score: 63		Score: 65		Score: 77	

detection. Moreover, many of BAs do not strong absorbance when using UV detection, thus, derivatization step is required when using gas chromatography as a means of separation.

Due to increased sensitivity and the specific structural information that MS or MS/MS detectors can provide, they are the most adequate detection technique for low concentrations of metabolites. However, matrix effects remains one of the main problematic issues when quantifying BAs in food samples which are composed of complex matrices. Therefore, the sample preparation step has been commonly used to minimize these matrix interferences. Isotopically labeled internal standards are also used to minimize the matrix effect, allowing the quantification of ultratrace quantities.

Reducing the time of analysis, being able to use a lower concentration of derivatizing agent and increasing sensitivity are the current challenges facing the BA research community. Thus, in the context of future regulations of BA in fermented products, fast and robust methods of analysis will be required.

Following previous considerations it should be helpful to actively promote the production of functional beverages and foods,

having a modified balance between amines and polyphenols, using different agricultural management practices and processing methods.

In order to assess human exposure and the risk associated with biogenic amines, there is a strong need to carry out comprehensive food surveys and related studies, such as gastrointestinal uptake studies, which are urgently required for a better understanding of the contribution of food pathway to consumer exposure to BAs.

Acknowledgments

The research was funded by the Polish Ministry of Science and Higher Education within the "Iuventus Plus" program in years 2015–2018, project no: IP2014 037573.

References

- [1] S. Ozcan, H.Z. Senyuva, Improved and simplified liquid chromatography/atmospheric pressure chemical ionization mass spectrometry method for the

- analysis of underivatized free amino acids in various foods, *J. Chromatogr. A* 1135 (2006) 179–185.
- [2] M.T. Kelly, A. Blaise, M. Larroque, Rapid automated high performance liquid chromatography method for simultaneous determination of amino acids and biogenic amines in wine, fruit and honey, *J. Chromatogr. A* 1217 (2010) 7385–7392.
- [3] H.K. Mayer, G. Fiechter, E. Fischer, A new ultra-pressure liquid chromatography method for the determination of biogenic amines in cheese, *J. Chromatogr. A* 1217 (2010) 3251–3257.
- [4] A. Jastrzębska, A. Piasta, E. Szyk, Application of ion chromatography for the determination of biogenic amines in food samples, *J. Anal. Chem.* 70 (2015) 1131–1138.
- [5] C.M. Mayr, P. Schieberle, Development of stable isotope dilution assays for the simultaneous quantitation of biogenic amines and polyamines in foods by LC-MS/MS, *J. Agric. Food Chem.* 60 (2012) 3026–3032.
- [6] V. Sirocchi, G. Caprioli, M. Ricciutielli, S. Vittori, G. Sagratini, Simultaneous determination of ten underivatized biogenic amines in meat by liquid chromatography-tandem mass spectrometry (HPLC-MS/MS), *J. Mass Spectr.* 49 (2014) 819–825.
- [7] E.J. Papavergou, I.N. Savvaidis, I.A. Ambrosiadis, Levels of biogenic amines in retail market fermented meat products, *Food Chem.* 135 (2012) 2750–2755.
- [8] H. Zhai, X. Yang, L. Li, G. Xia, J. Cen, H. Huang, S. Hao, Biogenic amines in commercial fish and fish products sold in southern China, *Food Control* 25 (2012) 303–308.
- [9] C. Palermo, M. Muscarella, D. Nardiello, M. Iammarino, D. Centonze, A multiresidual method based on ion-exchange chromatography with conductivity detection for the determination of biogenic amines in food and beverages, *Anal. Bioanal. Chem.* 405 (2013) 1015–1023.
- [10] R. Parchami, M. Kamalabadi, N. Alizadeh, Determination of biogenic amines in canned fish samples using head-space solid phase microextraction based on nanostructured polypyrrole fiber coupled to modified ionization region ion mobility spectrometry, *J. Chromatogr. A* 1481 (2017) 37–43.
- [11] Y. Fu, Z. Zhou, Y. Li, X. Lu, C. Zhao, G. Xu, High-sensitivity detection of biogenic amines with multiple reaction monitoring in fish based on benzoyl chloride derivatization, *J. Chromatogr. A* 1465 (2016) 30–37.
- [12] R. Romero-Gonzalez, M.I. Alarcon-Flores, J.L. Martinez Vidal, A. Garrido French, Simultaneous determination of four biogenic and three volatile amines in anchovy by ultra-high-performance liquid chromatography coupled to tandem mass spectrometry, *J. Agric. Food Chem.* 60 (2012) 5324–5329.
- [13] B. Bilgin, H. Genççelep, Determination of biogenic amines in fish products, *Food Sci. Biotech.* 24 (2015) 1907–1913.
- [14] Y. Hu, Z. Huang, J. Li, H. Yang, Concentrations of biogenic amines in fish, squid and octopus and their changes during storage, *Food Chem.* 135 (2012) 2604–2611.
- [15] J. Sun, H.X. Guo, D. Semin, J. Cheetham, Direct separation and detection of biogenic amines by ion-pair liquid chromatography with chemiluminescent nitrogen detector, *J. Chromatogr. A* 1218 (2011) 4689–4697.
- [16] J. Stadnik, Z.J. Dolatowski, Biogenic amines in meat and fermented meat products, *ACTA Scientiarum Polonorum Technologia Alimentaria* 9 (2010) 251–263.
- [17] F. Aflaki, V. Ghoulipour, N. Saemian, M. Salahinejad, A simple method for benzoyl chloride derivatization of biogenic amines for high performance liquid chromatography, *Anal. Meth.* 6 (2014) 1482.
- [18] Q. Sun, M. Lv, A. Hu, X. Yang, C.P. Yu, Seasonal variation in the occurrence and removal of pharmaceuticals and personal care products in a wastewater treatment plant in Xiamen, China, *J. Hazard. Mater.* 277 (2014) 69–75.
- [19] M. Coton, A. Romano, G. Spano, K. Ziegler, C. Vetrana, C. Desmarais, A. Lonvaud-Funel, P. Lucas, E. Coton, Occurrence of biogenic amine-forming lactic acid bacteria in wine and cider, *Food Microbiol.* 27 (2010) 1078–1085.
- [20] J. Pradenas, O. Galarce-Bustos, K. Henríquez-Aedo, R. Mundaca-Uribe, M. Aranda, Occurrence of biogenic amines in beers from Chilean market, *Food Control* 70 (2016) 138–144.
- [21] F. Gardini, Y. Ozogul, G. Suzzi, G. Tabanelli, F. Ozogul, Technological factors affecting biogenic amine content in foods: a review, *Front. Microbiol.* 7 (2016) 1218.
- [22] D.M. Linares, B. Del Rio, V. Ladero, N. Martinez, M. Fernandez, M.C. Martin, M.A. Alvarez, Factors influencing biogenic amines accumulation in dairy products, *Front. Microbiol.* 3 (2012) 180.
- [23] M. Saaid, B. Saad, I.A. Rahman, A.S. Ali, M.I. Saleh, Extraction of biogenic amines using sorbent materials containing immobilized crown ethers, *Talanta* 80 (2010) 1183–1190.
- [24] J. Płotka-Wasyłka, V. Simeonov, J. Namiesnik, An in situ derivatization – dispersive liquid-liquid microextraction combined with gas-chromatography – mass spectrometry for determining biogenic amines in home-made fermented alcoholic drinks, *J. Chromatogr. A* 1453 (2016) 10–18.
- [25] H. Dong, K. Xiao, Modified QuEChERS combined with ultra high performance liquid chromatography tandem mass spectrometry to determine seven biogenic amines in Chinese traditional condiment soy sauce, *Food Chem.* 229 (2017) 502–508.
- [26] J. Karović, Z. Kohajdová, Biogenic amines in food, *Chem. Pap.* 59 (2005) 70–79.
- [27] M. Giurfa, Associative learning: the instructive function of biogenic amines, *Curr. Biol.* 16 (2006) R892–R895.
- [28] A. Halász, A. Baráth, L. Simon-Sarkadi, W. Holzapfel, Biogenic amines and their production by microorganisms in food, *Trends Food Sci. Technol.* 5 (1994) 42–49.
- [29] M. Krížek, T. Pelikánová, Determination of seven biogenic amines in foods by micellar electrokinetic capillary chromatography, *J. Chromatogr.* 815 (1998) 243–250.
- [30] M.L. Latorre-Moratalla, T. Veciana-Nogues, S. Bover-Cid, M. Garriga, T. Aymerich, E. Zanardi, A. Ianieri, M.J. Fraqueza, L. Patarata, E.H. Drosinos, A. Laukova, R. Talon, M.C. Vidal-Carou, Biogenic amines in traditional fermented sausages produced in selected European countries, *Food Chem.* 107 (2008) 912–921.
- [31] D. Ansorena, M.C. Montel, M. Rokka, R. Talon, S. Eerola, A. Rizzo, M. Raemaekers, D. Demeyer, Analysis of biogenic amines in northern and southern European sausages and role of flora in amine production, *Meat Sci.* 61 (2002) 141–147.
- [32] P. Visciano, M. Schirone, R. Tofalo, G. Suzzi, Biogenic amines in raw and processed seafood, *Front. Microbiol.* 3 (2012) 1–10.
- [33] L. Gram, H.H. Huss, Microbiological spoilage of fish and fish products, *Int. J. Food Microbiol.* 33 (1996) 121–137.
- [34] K.B. Biji, C.N. Ravishankar, R. Venkateswarlu, C.O. Mohan, T.K.S. Gopal, Biogenic amines in seafood: a review, *J. Food Sci. Technol.* 53 (2016) 2210–2218.
- [35] M. Saaid, B. Saad, A.S. Ali, M.I. Saleh, C. Basheer, H.K. Lee, In situ derivatization hollow fibre liquid-phase microextraction for the determination of biogenic amines in food samples, *J. Chromatogr. A* 1216 (2009) 5165–5170.
- [36] F. Galgano, F. Favati, M. Bonadio, V. Lorusso, P. Romano, Role of biogenic amines as index of freshness in beef meat packed with different biopolymeric materials, *Food Res. Int.* 42 (2009) 1147–1152.
- [37] P. Paulsen, U. Hagen, F. Bauer, Changes in biogenic amine contents, non-protein nitrogen and crude protein during curing and thermal processing of *M. longissimus, pars lumborum* of pork, *Eur. Food Res. Technol.* 223 (2006) 603–608.
- [38] H. Wu, G. Li, S. Liu, Z. Ji, Q. Zhang, N. Hu, Y. Suo, J. You, Simultaneous determination of seven biogenic amines in foodstuff samples using one-step fluorescence labeling and dispersive liquid-liquid microextraction followed by HPLC-FLD and method optimization using response surface methodology, *Food Anal. Meth.* 8 (2014) 685–695.
- [39] C.A. Lázaro, C.A. Conte-Júnior, A.C. Canto, M.L.G. Monteiro, B. Costa-Lima, A.G.D. Cruz, E.T. Mársico, R.M. Franco, Biogenic amines as bacterial quality indicators in different poultry meat species, *LWT – Food Sci. Technol.* 60 (2015) 15–21.
- [40] S. Gerosa, J. Skoet, Milk availability: current production and demand and medium-term outlook, in: E. Muehlhoff, A. Bennett, D. McMahon (Editors), *Milk and Dairy Products in Human Nutrition, Food and Agricultural Organisation of the United Nations (FAO), Rome, Italy, 2013*, pp. 11–40.
- [41] J. Calzada, A. del Olmo, A. Picon, P. Gaya, M. Nuñez, Proteolysis and biogenic amine buildup in high-pressure treated ovine milk blue-veined cheese, *J. Dairy Sci.* 96 (2013) 4816–4829.
- [42] EFSA, Opinion of the Scientific Committee on a request from EFSA on the introduction of a Qualified Presumption of Safety (QPS) approach for assessment of selected microorganisms referred to EFSA, *EFSA J* 587 (2007) 1–16.
- [43] S. Andiç, H. Genççelep, Y. Tunçtürk, Ş. Köse, The effect of storage temperatures and packaging methods on properties of Motal cheese, *J. Dairy Sci.* 93 (2010) 849–859.
- [44] D. Restuccia, U.G. Spizzirri, F. Puoci, G. Cirillo, M. Curcio, O.I. Parisi, F. Iemma, N. Picci, A new method for the determination of biogenic amines in cheese by LC with evaporative light scattering detector, *Talanta* 85 (2011) 363–369.
- [45] N. Mascaro, R. Stocchi, M. Ricciutielli, N. Cammertoni, F. Renzi, S. Cecchini, A. Loschi, S. Rea, Biogenic amine content and chemical and physical features of Italian formaggio di Fossa, *Ital. J. Food Saf.* 1 (2010) 49–53.
- [46] L. Bunkova, G. Adamcova, K. Hudcova, H. Velichova, V. Pachlova, E. Lorencova, F. Bunka, Monitoring of biogenic amines in cheeses manufactured at small-scale farms and in fermented dairy products in the Czech Republic, *Food Chem.* 141 (2013) 548–551.
- [47] M.P. Costa, C.F. Balthazar, B.L. Rodrigues, C.A. Lazaro, A.C. Silva, A.G. Cruz, C.A. Conte Junior, Determination of biogenic amines by high-performance liquid chromatography (HPLC-DAD) in probiotic cow's and goat's fermented milks and acceptance, *Food Sci. Nutr.* 3 (2015) 172–178.
- [48] D. Restuccia, U.G. Spizzirri, O.I. Parisi, G. Cirillo, N. Picci, Brewing effect on levels of biogenic amines in different coffee samples as determined by LC-UV, *Food Chem.* 175 (2015) 143–150.
- [49] Ö. Özdeştan, Evaluation of bioactive amine and mineral levels in Turkish coffee, *Food Res. Int.* 61 (2014) 167–175.
- [50] E.C. Dias, R.G.F.A. Pereira, F.M. Borém, E. Mendes, R.R. de Lima, J.O. Fernandes, S. Casal, Biogenic amine profile in unripe Arabica coffee beans processed according to dry and wet methods, *J. Agric. Food Chem.* 60 (2012) 4120–4125.
- [51] R. Preti, M.L. Antonelli, R. Bernacchia, G. Vinci, Fast determination of biogenic amines in beverages by a core-shell particle column, *Food Chem.* 187 (2015) 555–562.
- [52] Z.-F. Liu, Y.-X. Wei, J.-J. Zhang, D.-H. Liu, Y.-Q. Hu, X.-Q. Ye, Changes in biogenic amines during the conventional production of stinky tofu, *Int. J. Food Sci. Technol.* 46 (2011) 687–694.

- [53] B.Y. Byun, X. Bai, J.-H. Mah, Occurrence of biogenic amines in Doubanjiang and Tofu, *Food Sci. Biotech.* 22 (2013) 55–62.
- [54] N. Toro-Funes, J. Bosch-Fuste, M.L. Latorre-Moratalla, M.T. Veciana-Nogues, M.C. Vidal-Carou, Biologically active amines in fermented and non-fermented commercial soybean products from the Spanish market, *Food Chem.* 173 (2015) 1119–1124.
- [55] R.-F. Guan, Z.-F. Liu, J.-J. Zhang, Y.-X. Wei, S. Wahab, D.-H. Liu, X.-Q. Ye, Investigation of biogenic amines in sufu (furu): a Chinese traditional fermented soybean food product, *Food Control* 31 (2013) 345–352.
- [56] J. Yang, X. Ding, Y. Qin, Y. Zeng, Safety assessment of the biogenic amines in fermented soya beans and fermented bean curd, *J. Agric. Food Chem.* 62 (2014) 7947–7954.
- [57] B. Kim, B.Y. Byun, J.H. Mah, Biogenic amine formation and bacterial contribution in Natto products, *Food Chem.* 135 (2012) 2005–2011.
- [58] S. Shukla, H.K. Park, J.K. Kim, M. Kim, Determination of biogenic amines in Korean traditional fermented soybean paste (Doenjang), *Food Chem. Toxicol.* 48 (2010) 1191–1195.
- [59] C. Almeida, J.O. Fernandes, S.C. Cunha, A novel dispersive liquid-liquid microextraction (DLLME) gas chromatography-mass spectrometry (GC-MS) method for the determination of eighteen biogenic amines in beer, *Food Control* 25 (2012) 380–388.
- [60] R.M. Ramos, I.M. Valente, J.A. Rodrigues, Analysis of biogenic amines in wines by salting-out assisted liquid-liquid extraction and high-performance liquid chromatography with fluorimetric detection, *Talanta* 124 (2014) 146–151.
- [61] M. Martuscelli, G. Arfelli, A.C. Manetta, G. Suzzi, Biogenic amines content as a measure of the quality of wines of Abruzzo (Italy), *Food Chem.* 140 (2013) 590–597.
- [62] B. Bach, S. Le Quere, P. Vuchot, M. Grinbaum, L. Barnavon, Validation of a method for the analysis of biogenic amines: histamine instability during wine sample storage, *Anal. Chim. Acta* 732 (2012) 114–119.
- [63] P. Comuzzo, D. Rauhut, M. Werner, C. Lagazio, R. Zironi, A survey on wines from organic viticulture from different European countries, *Food Control* 34 (2013) 274–282.
- [64] Y.Q. Wang, D.Q. Ye, B.Q. Zhu, G.F. Wu, C.Q. Duan, Rapid HPLC analysis of amino acids and biogenic amines in wines during fermentation and evaluation of matrix effect, *Food Chem.* 163 (2014) 6–15.
- [65] J.L. Ordonez, R.M. Callejon, M.L. Morales, M.C. Garcia-Parrilla, A survey of biogenic amines in vinegars, *Food Chem.* 141 (2013) 2713–2719.
- [66] M.K. Kim, J.H. Mah, H.J. Hwang, Biogenic amine formation and bacterial contribution in fish, squid and shellfish, *Food Chem.* 116 (2009) 87–95.
- [67] M.B.R. Rodriguez, C. Carneiro, M. Feijó, C. Júnior, S. Mano, Bioactive amines: aspects of quality and safety in food, *Food Nutr. Sci.* 5 (2014) 138–146.
- [68] H.P. Til, H.E. Falke, M.K. Prinsen, M.I. Willems, Acute and subacute toxicity of tyramine, spermidine, spermine, putrescine and cadaverine in rats, *Food Chem. Toxicol.* 35 (1997) 337–348.
- [69] Official Journal of the European Union, COMMISSION REGULATION (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs, L 338/1-L 338/26.
- [70] A. Halász, A. Baráth, W.Z. Holzappel, The influence of starter culture selection on sauerkraut fermentation, *Lebensm. Unters. Forsch.* 208 (1999) 434–438.
- [71] J.L. Mietz, E. Karmas, Chemical quality index of canned tuna as determined by highpressure liquid chromatography, *J. Food Sci.* 42 (1997) 155–158.
- [72] H.-L. Jiang, L.-Y. Ying, S.-C. Zhou, Chromatographic determination of biogenic amines in wines after treatment with ionic liquids as novel media, *J. Sep. Sci.* 34 (2011) 1055–1062.
- [73] M.L. Latorre-Moratalla, J. Bosch-Fustea, T. Lavizzaria, S. Bover-Cidb, M.T. Veciana-Nogués, M.C. Vidal-Carou, Validation of an ultra-high pressure liquid chromatographic method for the determination of biologically active amines in food, *J. Chromatogr. A* 1216 (2009) 7715–7720.
- [74] J. Ruiz-Jimenez, M.D. Luque de Castro, Pervaporation as interface between solid samples and capillary electrophoresis. Determination of biogenic amines in food, *J. Chromatogr. A* 1110 (2006) 245–253.
- [75] J. Huang, N. Gan, F. Lv, Y. Cao, C. Ou, H. Tang, Environmentally friendly solid-phase microextraction coupled with gas chromatography and mass spectrometry for the determination of biogenic amines in fish samples, *J. Sep. Sci.* 39 (2016) 4384–4390.
- [76] M.-S. Steiner, R.J. Meier, A. Duerkop, O.S. Wolfbeis, Chromogenic sensing of biogenic amines using a chameleon probe and the red-green-blue readout of digital camera images, *Anal. Chem.* 82 (2010) 8402–8405.
- [77] M.A. Alonso-Lomillo, O. Domínguez-Renedo, P. Matos, M.J. Arcos-Martínez, Disposable biosensors for determination of biogenic amines, *Anal. Chim. Acta* 665 (2010) 26–31.
- [78] W. Henao-Escobar, O. Domínguez-Renedo, M.A. Alonso-Lomillo, M.J. Arcos-Martínez, Simultaneous determination of cadaverine and putrescine using a disposable monoamine oxidase based biosensor, *Talanta* 117 (2013) 405–411.
- [79] T.L. Nelson, L. Tran, T.G. Ingallinera, M.S. Maynor, J.J. Lavigne, Multi-layered analyses using directed partitioning to identify and discriminate between biogenic amines, *Analyst* 132 (2007) 1024–1030.
- [80] S.B. Patange, M.K. Mukundan, K. Ashok Kumar, A simple and rapid method for colorimetric determination of histamine in fish flesh, *Food Control* 16 (2005) 465–472.
- [81] Z. Tang, J. Yang, J. Yu, B. Cui, A colorimetric sensor for qualitative discrimination and quantitative detection of volatile amines, *Sensors* 10 (2010) 6463–6476.
- [82] L. Sutarlie, H. Quin, K.-L. Yang, Polymer stabilized cholesteric liquid crystal arrays for detecting vaporous amines, *Analyst* 135 (2010) 1691–1696.
- [83] D. Telsnig, K. Kalcher, A. Leitner, A. Ortner, Design of an amperometric biosensor for the determination of biogenic amines using screen printed carbon working electrodes, *Electroanalysis* 25 (2013) 47–50.
- [84] A. Pineda, J. Carrasco, C. Peña-Farfal, K. Henríquez-Aedo, M. Aranda, Preliminary evaluation of biogenic amines content in Chilean young varietal wines by HPLC, *Food Control* 23 (2012) 251–257.
- [85] A. Zotou, M. Notou, Enhancing fluorescence lc analysis of biogenic amines in fish tissues by precolumn derivatization with naphthalene-2,3-dicarboxaldehyde, *Food Anal. Meth.* 6 (2012) 89–99.
- [86] K. Nalazek-Rudnicka, A. Wasik, Development and validation of an LC-MS/MS method for the determination of biogenic amines in wines and beers, *Monatsh Chem.* 148 (2017) 1685–1696.
- [87] S.C. Cunha, M.A. Faria, J.O. Fernandes, Gas chromatography-mass spectrometry assessment of amines in Port wine and grape juice after fast chloroformate extraction/derivatization, *J. Agric. Food Chem.* 59 (2011) 8742–8753.
- [88] G. Li, L. Dong, A. Wang, W. Wang, N. Hu, J. You, Simultaneous determination of biogenic amines and estrogens in foodstuff by an improved HPLC method combining with fluorescence labeling, *LWT – Food Sci. Technol.* 55 (2014) 355–361.
- [89] E. Dađáková, M. Křížek, T. Pelikánová, Determination of biogenic amines in foods using ultra-performance liquid chromatography (UPLC), *Food Chem.* 116 (2009) 365–370.
- [90] H. Yoon, J.H. Park, A. Choi, H.-J. Hwang, J.-H. Mah, Validation of an HPLC analytical method for determination of biogenic amines in agricultural products and monitoring of biogenic amines in Korean fermented agricultural products, *Toxicol. Res.* 31 (2015) 299–305.
- [91] M. Saaid, B. Saad, N.H. Hashim, A.S. Ali, M.I. Saleh, Determination of biogenic amines in selected Malaysian food, *Food Chem.* 113 (2009) 1356–1362.
- [92] C.A. Lázaro, C.A. Conte-Júnior, F.L. Cunha, E.T. Mársico, S.B. Mano, R.M. Franco, Validation of an HPLC methodology for the identification and quantification of biogenic amines in chicken meat, *Food Anal. Meth.* 6 (2013) 1024–1032.
- [93] I. Baranowska, J. Płonka, Simultaneous determination of biogenic amines and methylxanthines in foodstuff—sample preparation with HPLC-DAD-FL analysis, *Food Anal. Meth.* 8 (2015) 963–972.
- [94] N. Dossi, R. Toniolo, A. Pizzariello, S. Susmel, G. Bontempelli, A modified electrode for the electrochemical detection of biogenic amines and their amino acid precursors separated by microchip capillary electrophoresis, *Electrophoresis* 32 (2011) 906–912.
- [95] D. Daniel, V.B. Dos Santos, D.T. Vidal, C.L. do Lago, Determination of biogenic amines in beer and wine by capillary electrophoresis-tandem mass spectrometry, *J. Chromatogr. A* 1416 (2015) 121–128.
- [96] A. Jastrzębska, A. Piasta, E. Szyk, Simultaneous determination of selected biogenic amines in alcoholic beverage samples by isotachophoretic and chromatographic methods, *Food Addit. Contam. A* 31 (2014) 83–92.
- [97] S. Uzaşci, S. Baskan, F. Bedia Erim, Biogenic amines in wines and pomegranate molasses—a non-ionic micellar electrokinetic chromatography Assay with laser-induced fluorescence detection, *Food Anal. Meth.* 5 (2012) 104–108.
- [98] J. Płotka-Wasyłka, J. Namieśnik, E. Kłodzińska, Determination of biogenic amines in wine using micellar electrokinetic chromatography, *J. Res. Anal.* 3 (2017) 62–66.
- [99] N. Zhang, H. Wang, Z.-X. Zhang, Y.-H. Deng, H.-S. Zhang, Sensitive determination of biogenic amines by capillary electrophoresis with a new fluorogenic reagent 3-(4-fluorobenzoyl)-2-quinolinecarboxaldehyde, *Talanta* 76 (2008) 791–797.
- [100] D. An, Z. Chen, J. Zheng, S. Chen, L. Wang, Z. Huang, L. Weng, Determination of biogenic amines in oysters by capillary electrophoresis coupled with electrochemiluminescence, *Food Chem.* 168 (2015) 1–6.