



## Note

Tight bounds on the complexity of semi-equitable coloring of cubic and subcubic graphs<sup>☆</sup>Hanna Furmańczyk<sup>a,\*</sup>, Marek Kubale<sup>b</sup><sup>a</sup> Institute of Informatics, University of Gdańsk, Wita Stwosza 57, 80-308 Gdańsk, Poland<sup>b</sup> Department of Algorithms and System Modelling, Gdańsk University of Technology, Narutowicza 11/12, 80-233 Gdańsk, Poland

## ARTICLE INFO

## Article history:

Received 21 June 2016

Received in revised form 16 November 2017

Accepted 5 December 2017

Available online 28 December 2017

## Keywords:

Semi-equitable coloring

Equitable coloring

Cubic graphs

Subcubic graphs

## ABSTRACT

A  $k$ -coloring of a graph  $G = (V, E)$  is called semi-equitable if there exists a partition of its vertex set into independent subsets  $V_1, \dots, V_k$  in such a way that  $|V_1| \notin \{\lceil |V|/k \rceil, \lfloor |V|/k \rfloor\}$  and  $\| |V_i| - |V_j| \| \leq 1$  for each  $i, j = 2, \dots, k$ . The color class  $V_1$  is called non-equitable. In this note we consider the complexity of semi-equitable  $k$ -coloring,  $k \geq 4$ , of the vertices of a cubic or subcubic graph  $G$ . In particular, we show that, given a  $n$ -vertex subcubic graph  $G$  and constants  $\epsilon > 0$ ,  $k \geq 4$ , it is NP-complete to obtain a semi-equitable  $k$ -coloring of  $G$  whose non-equitable color class is of size  $s$  if  $s \geq n/3 + \epsilon n$ , and it is polynomially solvable if  $s \leq n/3$ .

© 2017 Elsevier B.V. All rights reserved.

## 1. Introduction

All graphs considered in this paper are finite, loopless, and without multiple edges. We refer the reader to [11] for terminology in graph theory. We say that a graph  $G = (V, E)$  is *equitably  $k$ -colorable* if and only if its vertex set can be partitioned into independent sets  $V_1, \dots, V_k \subset V$  such that  $|V_i| - |V_j| \in \{-1, 0, 1\}$  for all  $i, j = 1, \dots, k$ . The smallest  $k$  for which  $G$  admits such a coloring is called the *equitable chromatic number* of  $G$  and denoted by  $\chi_=(G)$ . A graph  $G$  on  $n$  vertices has a *semi-equitable coloring* if there exists a partition of its vertices into independent sets  $V_1, \dots, V_k \subset V$  such that one of these subsets, say  $V_1$ , is of size  $s \notin \{\lfloor \frac{n}{k} \rfloor, \lceil \frac{n}{k} \rceil\}$ , and the remaining subgraph  $G - V_1$  is equitably  $(k - 1)$ -colorable. In what follows, such a color class  $V_1$  will be called *non-equitable*. These two models of graph coloring are motivated by applications in multiprocessor scheduling of unit-execution time jobs [7,6].

In the following we will say that graph  $G$  has a  $(V_1, \dots, V_k)$ -coloring to express explicitly the partition of  $V$  into  $k$  independent sets. If, however, only cardinalities of color classes are important, we will use the notation of  $[|V_1|, \dots, |V_k|]$ -coloring. For a given coloring, we call the difference  $\max\{|V_i| - |V_j| : i, j \in \{1, \dots, k\}\}$  its *color width*. Thus, a coloring of a graph is equitable if and only if the color width does not exceed 1.

We mention the following two theorems on equitable graph coloring. First, Hajnal and Szemerédi [9] proved.

**Theorem 1.1** ([9]). *If  $G$  is a graph with maximum degree  $\Delta(G)$ ,  $\Delta(G) \leq k$ , then  $G$  has an equitable  $(k + 1)$ -coloring.*

This theorem implies that every cubic graph, i.e. a regular graph of degree 3, has an equitable  $k$ -coloring for every  $k \geq 4$ . Kierstead et al. [10] gave a simple algorithm for obtaining such a coloring in  $O(n^2)$  time. Secondly, Chen et al. [1] proved.

<sup>☆</sup> Project has been partially supported by Narodowe Centrum Nauki under contract DEC-2011/02/A/ST6/00201.

\* Corresponding author.

E-mail addresses: [hanna@inf.ug.edu.pl](mailto:hanna@inf.ug.edu.pl) (H. Furmańczyk), [kubale@eti.pg.edu.pl](mailto:kubale@eti.pg.edu.pl) (M. Kubale).

**Theorem 1.2** ([1]). *If  $G$  is a connected 3-chromatic cubic graph, then there exists an equitable 3-coloring of  $G$ .*

Actually, they proved that  $\chi(G) = \chi_=(G)$  for any connected cubic graph  $G$ . The proof starts from any proper 3-coloring of a connected cubic graph different from  $K_4$  and  $K_{3,3}$ , and it relies on successive decreasing of the color width of this coloring by one or by two, step by step, until the coloring is equitable. Moreover, Chen and Yen in [2] extended this result to disconnected subcubic graphs, where by a *subcubic graph* we mean a graph  $G = (V, E)$  with  $\deg(v) \leq 3$  for all  $v \in V$ .

**Theorem 1.3** ([2]). *A subcubic graph  $G$  with  $\chi(G) \leq 3$  is equitably 3-colorable if and only if exactly one of the following statements holds.*

1. *No components or at least two components of  $G$  are isomorphic to  $K_{3,3}$ .*
2. *Only one component of  $G$  is isomorphic to  $K_{3,3}$  and  $\alpha(G - K_{3,3}) > \frac{|V(G - K_{3,3})|}{3} > 0$ .*

By the above we immediately have the following corollary.

**Corollary 1.4.** *If  $G$  is a subcubic graph including neither  $K_{3,3}$  nor  $K_4$  as a component, then it admits an equitable 3-coloring.  $\square$*

The problem of semi-equitable 3-coloring of connected cubic graphs was introduced in [4]. We have shown that every cubic graph with  $t$  independent vertices has equitable 3-coloring for  $t \in \{\lceil n/3 \rceil, \lfloor n/3 \rfloor\}$  and semi-equitable 3-coloring for  $t \geq 2n/5$ . In this note we extend those results to an arbitrary number  $k \geq 4$  of colors and to, possibly disconnected, subcubic graphs. In contrast to equitable coloring not all cubic/subcubic graphs have a semi-equitable coloring (see  $K_4$  for example). Therefore, in the following we assume that all graphs under consideration have such a coloring. We will denote by  $N(v)$  the (open) neighborhood of the vertex  $v \in V$ , that is the set  $\{u \in V : \{u, v\} \in E\}$ . Let  $G_1 \cup G_2$  denote the union of graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  with disjoint vertex sets  $V_1$  and  $V_2$  and edge sets  $E_1$  and  $E_2$ , i.e. the graph  $G$  with  $V = V_1 \cup V_2$  and  $E = E_1 \cup E_2$ .

**Theorem 1.5** ([2]). *If two graphs  $G_1$  and  $G_2$  with disjoint vertex sets are both equitably  $k$ -colorable, then  $G_1 \cup G_2$  is also equitably  $k$ -colorable.*

The rest of the paper is organized as follows. In Section 2 we prove that, in contrast to equitable coloring, the problem of semi-equitable coloring becomes NP-complete for each  $k \geq 3$ . More precisely, we show that computing a semi-equitable  $k$ -coloring of a subcubic graph whose maximum color class is of size at least  $(1/3 + \epsilon)n$  for any  $\epsilon \in (0, 1/6)$  is NP-complete, if one exists. In Section 3 we show how to obtain in  $O(n^2)$  time a semi-equitable  $k$ -coloring of a subcubic graph with non-equitable color class of size at most  $n/3$ . Because we are interested in an algorithmic approach, in Appendix we reprove Corollary 1.4 by giving an appropriate algorithm resulting from a slight modification of Chen et al.'s proof [1]. The computational complexity of the whole equalizing procedure is  $O(n^2)$ .

## 2. NP-completeness of the problem

In this section we present one of the main results of the paper. We are interested in the computational complexity of deciding whether a subcubic graph  $G$  has a semi-equitable  $k$ -coloring ( $k \geq 4$ ) with non-equitable color class of size  $s$ . In [5] we proved that the problem of deciding whether a cubic graph has a coloring of type  $[4n/10, 3n/10, 3n/10]$  is NP-complete. In the following we strengthen and generalize this result to semi-equitable  $k$ -colorings ( $k \geq 4$ ) and subcubic graphs. We consider graphs not including  $K_4$  as a component. We say that  $G$  is a  $m$ -divisible graph if  $|V(G)|$  is divisible by  $m$ .

Let us define the following decision problem.

$(\frac{1}{3} + \epsilon)$ -Semi-Equitable Coloring of a Cubic  $m$ -Divisible graph (SECCD).

Instance: A  $m$ -divisible cubic graph  $G$ , an integer  $k \geq 4$ , and  $\epsilon > 0$ .

Question: Does  $G$  have a semi-equitable  $k$ -coloring whose non-equitable color class is of size at least  $(\frac{1}{3} + \epsilon)|V|$ ?

We want to prove that SECCD is NP-complete. The following lemma states a strong relationship between our problem and the Independent Set Problem.

**Lemma 2.1.** *Let  $G$  be a cubic or subcubic graph,  $k, s \in \mathbb{Z}^+$ , and  $k \geq 4$ . Then  $G$  has a semi-equitable  $k$ -coloring with non-equitable color class of size  $s$  if and only if  $G$  has an independent set of size  $s$ .*

**Proof.** If  $G$  has a semi-equitable  $k$ -coloring,  $k \geq 4$ , with non-equitable color class of size  $s$  then there must exist an independent set of size  $s$  in  $G$ .

Conversely, if there is an independent set of size  $s$  in  $G$ , it forms a non-equitable color class, say  $V_1$ . If  $k \geq 5$ , the existence of an equitable  $(k - 1)$ -coloring of the remaining subcubic graph  $G - V_1$  follows from Theorem 1.1. Let  $k = 4$ . If  $G - V_1$  fulfills the condition from Theorem 1.3, then we have an equitable 3-coloring of  $G - V_1$ . Let us assume that  $G - V_1$  is not equitably 3-colorable. This means that there is only one component of  $G - V_1$  isomorphic to  $K_{3,3}$  and  $\alpha(G - V_1 - K_{3,3}) \leq \frac{|V(G) - s - 6}{3}$  due to Theorem 1.3. Then we try to exchange one of vertices from  $V_1$  to another one in the copy of  $K_{3,3}$  in  $G - V_1$ . If we succeed, there would be no component isomorphic to  $K_{3,3}$  in  $G - V_1$ . Otherwise we conclude that exactly  $s$  vertices from  $V_1$  belong



to  $s$  subgraphs isomorphic to  $K_{3,3}$  in  $G$  and the subgraph  $G - V_1$  can be expressed as  $sK_{2,3} \cup K_{3,3} \cup H$ , where  $H$  is a subgraph (possibly empty) of  $G - V_1$  that is free from  $K_{3,3}$ . If  $V(H)$  includes at least one vertex, we exchange one vertex from  $V_1$  with any vertex of  $H$ . After this exchange a new graph  $G - V_1$  includes  $2K_{3,3}$  as a subgraph and such a graph is equitably 3-colorable due to [Theorem 1.3](#). If  $|V(H)| = 0$ , then we have  $G - V_1 = sK_{2,3} \cup K_{3,3}$  and the independence number of  $G - V_1 - K_{3,3}$  is equal to  $3s$  while  $|V(G - V_1 - K_{3,3})| = 5s$ . This means that the condition from [Theorem 1.3](#) is fulfilled, as  $3s > 5s/3$ , and the graph  $G - V_1$  is equitably 3-colorable.  $\square$

Now, we consider the following subproblem of the well known Maximum Independent Set (MIS) problem restricted to cubic graphs.

*Maximum Independent Set in a Cubic  $m$ -Divisible graph (MISCD)*

Instance: A  $m$ -divisible cubic graph  $G$ ,  $l \in \mathbb{Z}^+$ .

Question: Does  $G$  have an independent set of size at least  $l$ ?

**Lemma 2.2.** *Problem MISCD is NP-complete.*

**Proof.** Note that the MIS problem is NP-complete [8] even if  $G$  is 3-regular. We show that it remains so if  $m|n$  (MISCD). Let us denote by  $Cub_{t,t}$  any cubic bipartite graph on  $2t$  vertices such that  $n + 2t$  is divisible by  $m$ ,  $3 \leq t \leq m/2 + 2$ . We remark that a bipartite cubic graph has a perfect matching, hence by König's theorem [11] the size of a maximum independent set in such a graph on  $2t$  vertices is  $t$ . The statement holds because if we consider the graph  $G \cup Cub_{t,t}$ , the number of vertices in the new graph is divisible by  $m$  and, moreover,  $G$  has an independent set of size at least  $l$  if and only if  $G \cup Cub_{t,t}$  has an independent set of size at least  $l + t$ .  $\square$

We will see that MISCD remains NP-complete on the subset of instances where  $l = (\frac{1}{3} + \epsilon)|V|$ ,  $\epsilon > 0$ . Let us formally define the new decision problem.

*( $\frac{1}{3} + \epsilon$ )-Independent Set in a Cubic 6-Divisible graph (ISCD)*

Instance: A 6-divisible cubic graph  $G$ ,  $\epsilon > 0$ .

Question: Does  $G$  have an independent set of size at least  $(\frac{1}{3} + \epsilon)|V|$ ?

**Lemma 2.3.** *Problem ISCD is NP-complete.*

**Proof.** Let  $G$ ,  $l$ , and  $\epsilon$  be an instance of the MISCD problem with  $m = 6$ . Let  $n_G = |V(G)|$  and let  $p = n_G \lceil 1/\epsilon \rceil$ . Obviously,  $1/p < \epsilon$ . We will reduce the question about existence in  $G$  an independent set of size at least  $l$  to the question whether the subsequently defined cubic graph  $H$  has an independent set of size at least  $(1/3 + 1/p)n_H$ , where  $n_H = |V(H)|$ . Let  $q = \lceil (1/3 + 1/p)n_G - l \rceil$ .

If  $l \geq (1/3 + 1/p)n_G$ ,  $H$  is the union of  $G$  and  $pq/6$  copies of the  $P$  graph, where  $P$  is a 6-vertex prism (2 triangles joined by a 3-matching). This results in increasing the order of the graph by  $pq$ , i.e.  $n_H = n_G + pq$ , which is divisible by  $n_G$ . Each of these copies of  $P$  provides exactly 2 new vertices in any maximum independent set. Thus, if  $G$  has an independent set on at least  $l$  vertices then there exists in  $H$  an independent set of size at least

$$\begin{aligned} l + pq/3 &= (l - q) + (q + pq/3) = (1/3 + 1/p)n_G + (q + pq/3) \\ &= (1/3 + 1/p)n_G + pq(1/3 + 1/p) = (1/3 + 1/p)n_H, \end{aligned}$$

and vice versa.

If  $l < (1/3 + 1/p)n_G$ ,  $H$  is the union of  $q$  copies of  $p/6 - 2$  prisms and two graphs  $K_{3,3}$ . This time each such subgraph  $(p/6 - 2)P \cup 2K_{3,3}$  on  $p$  vertices provides  $(p/6 - 2)2 + 6 = p/3 + 2$  new independent vertices. Thus, if  $G$  has an independent set of size at least  $l$  then there exists in  $H$  an independent set of size

$$l + q(p/3 + 2) = (l + q) + q(p/3 + 1) = (1/3 + 1/p)n_G + (q + pq/3) = (1/3 + 1/p)n_H,$$

and vice versa.  $\square$

As a direct consequence of [Lemmas 2.1](#) and [2.3](#) we can conclude

**Theorem 2.4.** *For any fixed  $\epsilon \in (0, 1/6)$  problem SECCD is NP-complete.  $\square$*

### 3. Semi-equitable $k$ -coloring of subcubic graphs

In [Section 2](#) we have proved that for any constants  $k \geq 4$ ,  $\epsilon \in (0, 1/6)$  and any subcubic graph  $G$ , the problem of deciding if  $G$  has a  $(1/3 + \epsilon)$ -semi-equitable  $k$ -coloring is NP-complete. It turns out that if we reduce the size of non-equitable color class slightly, namely to  $\lceil n/3 \rceil$ , then the problem becomes polynomially solvable for any  $k \geq 4$ .

**Theorem 3.1.** *Given a  $n$ -vertex cubic or subcubic graph  $G$  not including  $K_4$  neither  $K_{3,3}$ , a constant  $k \geq 4$ , and an integer  $s \leq \lceil n/3 \rceil$ , finding a semi-equitable  $k$ -coloring of  $G$  with non-equitable color class of size  $s$  is solvable in  $O(n^2)$  time.*

**Table 1**

The computational complexity of semi-equitable  $k$ -coloring of cubic graphs.

$k$	$s \leq \lceil \frac{n}{3} \rceil$	$\lceil \frac{n}{3} \rceil < s < \frac{n}{2}$	$s = \frac{n}{2}$	$\frac{n}{2} < s$
3	$O(n)^a / -^b$	NPC	$O(n)^a / -^b$	$-^b$
$\geq 4$	$O(n^2)$	NPC	$O(n)^a / -^b$	$-^b$

<sup>a</sup> The linear solution concerns bipartite cubic graphs only.

<sup>b</sup> The corresponding solution may not exist.

**Table 2**

The computational complexity of semi-equitable  $k$ -coloring of subcubic graphs.

$k$	$s \leq \lceil \frac{n}{3} \rceil$	$\lceil \frac{n}{3} \rceil < s \leq \lfloor \frac{3n}{4} \rfloor$	$\lfloor \frac{3n}{4} \rfloor < s$
3	$-^a$	NPC	$-^b$
$\geq 4$	$O(n^2)$	NPC	$-^b$

<sup>a</sup> The corresponding solution may not exist.

<sup>b</sup> The corresponding solution does not exist in the case of connected graphs.

**Proof.** First, we have to determine an independent vertex set  $I$  of size  $s$ ,  $s \leq \lceil n/3 \rceil$ . This is an easy task, if we apply the following greedy approach. First, we find a vertex  $v$  of minimum degree and delete it from the graph together with its neighborhood  $N(v)$ . Note that, in all steps except the first, at most 3 vertices are deleted. Then, we repeat this step until  $s$  independent vertices are found. If it is not the case and one more independent vertex is needed, we can apply an equalizing 3-coloring procedure given in the proof of [Corollary 1.4](#) (see [Appendix](#)). It is easy to see that  $s$  must satisfy  $s \leq \lceil n/3 \rceil$  and this bound is tight. Next, we color graph  $G - I$  equitably with  $k - 1$  colors. If  $k \geq 5$  then  $(k - 1)$ -coloring of  $G - I$  is guaranteed by [Theorem 1.1](#). Such a coloring can be obtained in  $O(n^2)$  time. If  $k = 4$ ,  $G - I$  can be properly colored with 3 colors, as a subcubic graph different from  $K_4$ . Since  $G - I$  is also different from  $K_{3,3}$ , we can apply the procedure from the proof of [Corollary 1.4](#) for equalizing a given 3-coloring of  $G - I$ . Since this procedure is executed at most twice, the complexity of  $O(n^2)$  follows.  $\square$

One may ask about a semi-equitable 3-coloring of a subcubic graph  $G$ . The problem was discussed in [\[4\]](#), where we proved

**Theorem 3.2** ([\[4\]](#)). *If an  $n$ -vertex cubic graph  $G$  has an independent set  $I$  of size  $|I| \geq 2n/5$ , then it has a semi-equitable coloring of type  $[|I|, \lceil \frac{n-|I|}{2} \rceil, \lfloor \frac{n-|I|}{2} \rfloor]$ .*  $\square$

Moreover, we noticed that a cubic graph has an independent set of size  $|I| \geq 2n/5$  almost surely. This is so because Frieze and Suen [\[3\]](#) proved that for random cubic graphs  $G$  their independence number  $\alpha(G)$  fulfills the inequality  $\alpha(G) \geq 4.32n/10 - \epsilon n$  for any  $\epsilon > 0$  almost surely. Thus a random cubic graph is very likely to have an independent set of size  $s \geq 2n/5$  and the probability of this fact increases with  $n$ .

[Tables 1](#) and [2](#) gather the computational complexity status for semi-equitable  $k$ -colorings with non-equitable class of size  $s$  of  $n$ -vertex cubic and subcubic graphs, respectively.

**Acknowledgments**

The authors deeply thank the anonymous referee for her/his careful reading of the manuscript and her/his many insightful comments and suggestions. Moreover, the authors thank Professor Adrian Kosowski for taking great care in reading our manuscript and making several useful suggestions improving the presentation.

**Appendix. The proof of [Corollary 1.4](#)**

First, we recall some notations used by Chen et al. [\[1\]](#). Given  $A$  and  $B$  disjoint subsets of  $V(G)$ , let  $Ak_c B$  denote the set  $\{x \in A : x \text{ is adjacent to exactly } k \text{ vertices of } B \text{ in } G\}$ , while  $Adeg_c B$  denotes the set of all vertices in  $A$  having all its neighbors in set  $B$ , namely  $Adeg_c B = \{x \in A : N(x) \subset B\}$ . When it is clear we will use  $Ak B$  and  $A \deg B$ . Given a coloring of  $G$ , the notation  $A \Leftrightarrow B$  means that we exchange the color of vertices in  $A$  into the color of vertices in  $B$  and vice versa. A one-way arrow  $A \Leftarrow B$  means that we change the color of  $B$  into the color of  $A$ . We write  $A \Leftarrow x$  when  $B = \{x\}$ . By  $G(X, Y)$  we will denote a bipartite graph whose vertices can be divided into two disjoint sets  $X$  and  $Y$  such that every edge connects a vertex in  $X$  to one in  $Y$ .

**Lemma A.3** ([\[1\]](#)). *Let  $G(X, Y)$  be a connected bipartite subcubic graph such that  $|X| \geq |Y|$ . Then  $|X| - |Y| \leq |Y3X| + 1$ .*  $\square$

This lemma can be extended to the following one.

**Lemma A.4.** *Let  $G(X, Y, Z)$  be a tripartite subcubic graph and let  $H(X', Y')$ ,  $|X'| \geq |Y'|$ , be its connected bipartite subgraph such that  $X' \subset X$  and  $Y' \subset Y$ , and let  $t = |Y \deg_H X \cap Y'|$ . Then  $|X'| - |Y'| \leq t + 1$ .*

**Proof.** Let  $e$  be the number of edges in  $H$ . Since  $H$  is connected, we have  $e \geq |X'| + |Y'| - 1$ . On the other hand, the number of edges can be bounded from above by  $e \leq 3t + 2(|Y'| - t) = 2|Y'| + t$ . We get the thesis by combining these two inequalities.  $\square$

**Proof of Corollary 1.4.** We start with  $(A, B, C)$  a proper 3-coloring of  $G$  with  $|A| \geq |B| \geq |C|$ . If the coloring is not equitable we will decrease the color width of the coloring by 1 or 2. We repeat the color width decreasing procedure until the obtained coloring is equitable.

Let us assume that the 3-coloring is not equitable, this means that  $|A| - |C| \geq 2$ . We may assume that there is no isolated vertex in  $G$ , since if we have an equitably  $k$ -colored graph  $G$  and we add one isolated vertex, then we can always color the isolated vertex in such a way that the whole graph is equitably  $k$ -colored. Thus, we have  $1 \leq \deg(x) \leq 3$  for each  $x \in V(G)$ . Now, we consider the following steps (cases).

1. If  $A \deg B \neq \emptyset$ , then: Choose  $x \in A \deg B$  and do  $C \leftarrow x$ . The color width of the new coloring  $(A - \{x\}, B, C \cup \{x\})$  decreases in one or two units.  
Henceforth, in the following we assume  $A \deg B = \emptyset$ .
2. If  $C3A = \emptyset$  then: Consider the bipartite subgraph  $G(A, C)$  induced by  $A \cup C$ . Find a connected component  $G(A', C')$  of  $G(A, C)$  such that  $|A'| = |C'| + 1$  and do  $A' \leftrightarrow C'$ .  
Let us prove that there exists such a connected component  $G(A', C')$  of  $G(A, C)$ . In fact, since  $C3A = \emptyset$ , there is a component  $G(A', C')$  such that  $|A'| = |C'| + 1$ , by Lemma A.3. Then the color width of the new coloring decreases in one or two units.  
Henceforth, in the following we are assuming that  $C3A \neq \emptyset$ .
3. If  $|C \deg B| \geq |C3A|$  then: Consider the bipartite subgraph  $G(A, C)$  induced by  $A \cup C$ . Find a connected component  $G(A', C')$  of  $G(A, C)$  such that  $|A'| \geq |C'| + 1$ , choose  $S \subset (C \deg B) - C'$  such that  $|S| = |A'| - |C'| - 1$  and do  $A' \leftrightarrow C' \cup S$ .  
Let us prove that there exists such a connected component  $G(A', C')$  of  $G(A, C)$  and such a subset  $S$ . In fact, since  $|A| \geq |C| + 2$ , there must exist a connected component  $G(A', C')$  of  $G(A, C)$  for which  $|A'| \geq |C'| + 1$ . From the assumption of this case we have  $|C'3A'| \leq |C \deg B|$ . Moreover, we have  $|A'| - |C'| = t'$ , where  $t' \leq |C \deg B| + 1$ , by Lemma A.4, and the subset  $S \subset C \deg B - C'$  such that  $|S| = t' - 1$  may be chosen. Then the color width of the new coloring decreases by one unit.  
Henceforth, in the following we assume  $|C \deg B| \leq |C3A| - 1$ .
4. If  $B \deg A = \emptyset$  then: If  $|B| = |C|$  do  $B \leftrightarrow C$ . Otherwise consider the bipartite subgraph  $G(B, C)$  induced by  $B \cup C$ . Find a connected component  $G(B', C')$  of  $G(B, C)$  such that  $|B'| \geq |C'| + 1$ , choose  $S \subset (C \deg A) - C'$  such that  $|S| = |B'| - |C'|$  and do  $B' \leftrightarrow C' \cup S$ .  
Let us prove that there exists such a connected component  $G(B', C')$  of  $G(B, C)$  and such a subset  $S$ . Again, there must exist a connected component  $G(B', C')$  of  $G(B, C)$  for which  $|B'| \geq |C'| + 1$ . Since  $|C' \deg B'| \leq |C \deg B|$ , we have  $|B'| - |C'| = t$ , where  $t \leq |C \deg B| + 1$ . Therefore, a subset  $S \subseteq C \deg A - C'$  such that  $|S| = t$  may be chosen.  
Note that the new coloring does not decrease the color width but verifies  $B \deg A \neq \emptyset$ .  
Henceforth, in the following we assume  $B \deg A \neq \emptyset$ .
5. If  $A \deg C \neq \emptyset$  then: If  $|A| = |B|$  do  $C \leftarrow x$  for any  $x \in B \deg A$ , otherwise choose  $x \in B \deg A$  and  $y \in A \deg C$  and do  $B \leftarrow y$  and  $C \leftarrow x$ .  
Of course, all these operations are possible and the color width of the new coloring decreases.  
Henceforth, in the following we assume  $A \deg C = \emptyset$ .

From now on, we apply the following steps of the procedure on a one chosen connected component  $G(A', B')$  of  $G(A, B)$  such that  $|A'| \geq |B'| + 1$ .

6. If  $B' \deg A' = \emptyset$  then: Do  $C \leftarrow x$  and  $A' \leftrightarrow B'$  for any  $x \in B \deg A$ .  
Let us prove the correctness of this step. Since  $B' \deg A' = \emptyset$  then we have  $|A'| = |B'| + 1$ , by Lemma A.4. Thus, the color width of the new coloring decreases in one or two units. An example of such a situation is given in Fig. 1. Edges of a subgraph  $G(A', B')$  are depicted by dotted line.  
Henceforth, in the following we may assume that  $B' \deg A' \neq \emptyset$ .
7. If there exists some  $x \in B' \deg A'$  such that one of its neighbors, say  $y$ , satisfies  $y \in A'1B'$ , then: Do  $B \leftarrow y$  and  $C \leftarrow x$ .  
Of course, such operations are possible and the color width of the new coloring decreases in one or two units.  
Thus, since  $A \deg B = \emptyset$  we may assume that for every  $y \in N(x)$  we have  $y \in A'2B'$ .
8. If  $|B' \deg A'| \geq 2$  and for every  $x_1, x_2 \in B' \deg A'$  it holds  $N(x_1) \cap N(x_2) \neq \emptyset$ , then: Choose any  $w \in C \deg A$  and  $u \in N(x_1) \cap N(x_2)$ .  
First, we will show that we may assume  $N(x_1) \neq N(x_2)$ . Indeed, if  $\deg(x_1) = \deg(x_2) = 1$  then we have  $N(x_1) = N(x_2)$ , and therefore  $|A'| = |N(x_1)| = 1$  and  $B' = \{x_1, x_2\}$ , which contradicts  $|A'| > |B'|$ . Similarly, if  $\deg(x_1) = \deg(x_2) = 2$  then we also have a contradiction with  $|A'| > |B'|$ . So let  $\deg(x_1) = \deg(x_2) = 3$  and  $N(x_1) = N(x_2)$ . First, we do an exchange  $B \leftarrow z$  and  $C \leftarrow x_2$  for any  $z \in C3A$ . Then we have  $N(z) \neq N(x_1) = N(x_2)$ , since otherwise  $N(x_1) = N(x_2) = N(z)$  would force  $G = K_{3,3}$ , which is excluded by the assumption of the theorem. Therefore, we have  $N(x_1) \neq N(x_2)$ , as claimed.




Fig. 1. An example for the proof of Corollary 1.4 Case 6. A graph with its (a) [5, 4, 3]-coloring and (b) [4, 4, 4]-coloring;  $B = B_1 \cup B_2$ .




Fig. 2. An example of the last step in the proof of Corollary 1.4.

- If  $u \notin N(w)$  then: Do  $B \leftarrow u, B \leftarrow w$  and  $C \leftarrow \{x_1, x_2\}$ .  
The color width is decreased because the cardinality of color class  $A$  was decreased by 1, the number of vertices colored with 2 remains unchanged, while the cardinality of color class  $C$  was increased by 1.
- If  $u \in N(w)$  and  $B' \deg A' \setminus \{x_1, x_2\} \neq \emptyset$ , then: Choose  $x_3 \in B' \deg A' \setminus \{x_1, x_2\}$  and do  $B \leftarrow w, C \leftarrow u$ , and  $C \leftarrow x_3$ .  
This step is correct because  $u$  is not adjacent to  $x_3$ . Moreover, the number of vertices with 1 is decreased by 1 (vertex  $u \in A$  was recolored) while the number of vertices in class  $C$  was increased by 1.

9. Then, we can assume that for every  $\{x_1, x_2\} \subset B' \deg A'$  we have that  $N(x_1) \cap N(x_2) \neq \emptyset$  or  $B' \deg A' = \{x_1, x_2\}$  and for any  $w \in C \deg A$  and any  $u \in N(x_1) \cap N(x_2)$  we have  $u \in N(w)$ . Note, that it concerns also the case when  $|B' \deg A'| = 1$ , i.e.  $x_1 = x_2$ . Then:

- delete  $B' \deg A'$  and all those vertices which are adjacent to the both vertices of  $B' \deg A'$  from  $G(A', B')$ .  
Since we have assumed earlier that  $x \in B' \deg A'$  and  $y \in N(x)$  imply  $y \in A'2B'$  for any  $x$  and  $y$ , so each of the remaining vertices will have degree 1 or 2. Hence the resulting graph  $G'(A', B')$  decomposes into maximal paths.
- There must exist a path with initial and terminal vertices in the same partitions, otherwise  $|B'| = |A'|$ , which is a contradiction. Without loss of generality, let  $u_0, v_1, u_2, v_3, \dots, u_{2m}$  be a maximal path of  $G'(A', B')$  such that  $u_0$  is adjacent to  $x_1 \in B' \deg A'$  and  $u_{2m} \in A'$ .
  - If  $u_{2m} \notin N(x_2)$  for some  $x_2 \in B' \deg A'$  and  $x_2 \neq x_1$ , then: Do  $C \leftarrow x_1$  and  $A'' \leftrightarrow B''$ , where  $A'' = \{u_0, u_2, \dots, u_{2m}\} \subseteq A'$  and  $B'' = \{v_1, v_3, \dots, v_{2m-1}\} \subseteq B'$ .
  - If  $u_0 \in N(x_1) - N(x_2)$  and  $u_{2m} \in N(x_2) - N(x_1)$  for distinct  $x_1, x_2 \in B' \deg A'$ , then:

- \* If  $w \in N(v)$  for some  $v \in A''$ , then: Do  $B \leftarrow w, C \leftarrow \{v, x_2\}$ .  
In this case  $v$  is not adjacent to at least one of  $x_1$  and  $x_2$ , say  $x_2$  and the step is correct.
- \* If  $w \in N(v)$  for each  $v \in A''$ , then: Do  $B \leftarrow w, C \leftarrow \{x_1, x_2\}, A'' \leftrightarrow B''$ .  $w$  is independent in  $A''$ , thus the exchange is correct.

An example of such a situation from Case 9 is given in Fig. 2. Here  $G(A', B')$  consists of  $A' = \{u, a_1, \dots, a_5\}$ ,  $B' = \{x_1, x_2, b_1, \dots, b_3\}$  and dashed and dotted edges. Graph  $G'(A', B')$  consists of vertices  $A'' = \{a_1, \dots, a_5\}$ ,  $B'' = \{b_1, \dots, b_3\}$ , and edges drawn with a dashed line. We choose as the maximal path mentioned above—path  $(a_1, b_2, a_4, b_1, a_3)$ . We rename these vertices as  $(u_0, v_1, u_2, v_3, u_4)$ . Here  $u_0 \in N(x_1) \setminus N(x_2)$  and  $u_{2m} = u_4 \in N(x_2) \setminus N(x_1)$ . Since  $w \in N(u_0)$ , we do:  $B \leftarrow w, C \leftarrow \{u_0, x_2\}$ —the final result is given in Fig. 2(b).

We finally conclude that the color width has been decreased in all the cases of non-equitable 3-coloring of subcubic graph and therefore the proof is complete.  $\square$

It is easy to see that a single step of decreasing the color width can be done in linear time. Since such a decreasing procedure must be applied at most  $n/3 - 1$  times, we have  $O(n^2)$  as the computational complexity of the whole equalizing procedure.

Of course, if a graph is disconnected, each connected component may be colored and equalized separately. Finally, we may merge the equitable colorings of components into the equitable 3-coloring of the whole graph.

## References

- [1] B.L. Chen, K.W. Lih, P.L. Wu, Equitable coloring and the maximum degree, *European J. Combin.* 15 (1994) 443–447.
- [2] B.L. Chen, C.H. Yen, Equitable  $\Delta$ -coloring of graphs, *Discrete Math.* 312 (2012) 1512–1517.
- [3] A. Frieze, S. Suen, On the independence number of random cubic graphs, *Random Graphs and Structures* 5 (1994) 649–664.
- [4] H. Furmańczyk, M. Kubale, Equitable and semi-equitable coloring of cubic graphs and its application in batch scheduling, *Arch. Control Sci.* 25 (2015) 109–116.
- [5] H. Furmańczyk, M. Kubale, Equitable coloring of corona products of cubic graphs is harder than ordinary coloring, *Ars Math. Contemp.* 10 (2016) 333–347.
- [6] H. Furmańczyk, M. Kubale, Scheduling of unit-length jobs with bipartite incompatibility graphs on four uniform machines, *Bull. Pol. Acad. Sci. Tech. Sci.* 65 (2017) 29–34.
- [7] H. Furmańczyk, M. Kubale, Scheduling of unit-length jobs with cubic incompatibility graphs on three uniform machines, *Discrete Appl. Math.* 234 (2018) 210–217.
- [8] M.R. Garey, D.S. Johnson, L. Stockmeyer, Some simplified NP-complete problems, *Theoret. Comput. Sci.* 1 (1976) 237–267.
- [9] A. Hajnal, E. Szemerédi, Proof of a conjecture of Erdős, in: *Combinatorial Theory and Its Applications, II*, in: *Colloq. Math. Soc. János Bolyai*, vol. 4, North-Holland, Amsterdam, 1970, pp. 601–623.
- [10] H.A. Kierstead, A.V. Kostochka, M. Mydlarz, E. Szemerédi, A fast algorithm for equitable coloring, *Combinatorica* 30 (2010) 217–224.
- [11] D.B. West, *Introduction To Graph Theory*, Prentice Hall, 2001.