

# 1 Svalbard reindeer as an indicator of ecosystem changes in the Arctic terrestrial 2 ecosystem

3 Aneta Dorota Pacyna<sup>1</sup>, Katarzyna Kozirowska<sup>2</sup>, Stanisław Chmiel<sup>3</sup>, Jan  
4 Mazerski<sup>4</sup>, Żaneta Polkowska<sup>1\*</sup>

5

6 <sup>1</sup>Gdańsk University of Technology, Faculty of Chemistry, Department of Analytical Chemistry, 11/12 Narutowicza  
7 st, Gdańsk 80-233, Poland

8 <sup>2</sup>Institute of Oceanology Polish Academy of Sciences, ul. Powstańców Warszawy 55, Sopot, Poland

9 <sup>3</sup>Department of Hydrology, Faculty of Earth Sciences and Spatial Management, Maria Curie-Skłodowska  
10 University, Kraśnicka Ave. 2 cd, 20-718 Lublin, Poland

11 <sup>4</sup>Gdańsk University of Technology, Faculty of Chemistry, Department of Pharmaceutical Technology and  
12 Biochemistry, 11/12 Narutowicza st, Gdańsk 80-233, Poland

13 \*corresponding author [zanpolko@pg.edu.pl](mailto:zanpolko@pg.edu.pl)

14

## 15 Abstract

16 Over the years, noticeable effort has been directed towards contaminant determination in  
17 multiple biotic samples collected from the inhabitants of the Arctic. Little consideration has  
18 been given to polar herbivores, however, especially those from the European parts of the  
19 Arctic. To provide a broader perspective, we aimed to decipher trace element concentration  
20 in hairs of the key species in the Arctic, namely the Svalbard reindeer (*Rangifer tarandus*  
21 *platyrhynchus*), and to recognise whether diet variations could correspond with forward  
22 exposure. The effect of habitat and diet was investigated using the ratios of stable isotopes of

23 carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ), and previous literature studies on vegetation from the areas  
24 of interest. Analysis was performed for eighteen elements in total, both toxic and essential.  
25 Metals were present in a decreasing order Fe>Zn>Ba>Cu>Pb>Cr>Ni>V>Ga  
26 =La>Rb>As>Li>Co>Hg>Cd>Cs>Be. Similarity in trends in the studied subpopulations was  
27 observed for many metals. A significant log-linear correlation was observed for most of the  
28 elements, excluding nitrogen and carbon isotopes signature. Extremely high iron levels were  
29 determined in some of the samples, suggesting past iron overload. Zinc, in contrast to the  
30 remaining metals, did not correlate well with any other element. Mercury was determined at  
31 very low levels, in accordance with previous literature regarding its concentrations in moss  
32 and lichen species in Svalbard. The analysis of stable isotopes showed a high variation in  
33 nitrogen isotopes signatures. Further research is required to properly evaluate the potential  
34 health risks and ecological implications of elevated exposure.

35

36 **Capsule:** *Keratinised tissues can be a valuable source of information in ecotoxicological studies*  
37 *in the case of polar herbivores.*

38

39 **Key Words:** *Rangifer tarandus platyrhynchus*, hair, essential elements, toxic metals, stable  
40 isotopes, tundra

41

## 42 **1. Introduction**

43 Constant pollutant emission is undeniably a serious problem and it is considered a huge threat  
44 to ecosystem stability. Anthropogenic activities undoubtedly have significant ecological  
45 consequences worldwide. The Arctic is an invaluable source of information on the global-scale

46 impact due to long-range contaminant transport (Davis, 1996; Halbach et al. 2017). The  
47 accumulation of trace elements, particularly heavy metals, and the resulting enrichment in  
48 higher trophic levels, raise questions about its impact on native fauna. Due to its unique  
49 geographical location, the Svalbard Archipelago has become a significant recipient of  
50 pollutants emitted in the Northern Hemisphere. Natural sources of heavy metal emissions  
51 include volcanic activities, biogenic sources, soil-derived dusts, and sea salt aerosols. It is  
52 anthropogenic emissions, however, that are assumed to account for the observed heavy metal  
53 levels in the Arctic to the greatest degree (AMAP, 2005; Halbach et al. 2017). With only several  
54 local sources of pollution (such as mining activities, airport, ship traffic), most contaminants  
55 including heavy metals are atmospherically transported long-range from mid- and low-  
56 latitudes (Bard, 1999).

57 A growing amount of evidence arose in the recent years concerning the deposition of  
58 pollutants in polar, particularly marine biota (e.g. Burger et al. 2007). Physiological and  
59 ecological factors affecting the bioaccumulation process vary between terrestrial and aquatic  
60 ecosystems (van den Brink et al. 2015). Terrestrial species are often weakly investigated and  
61 yet crucial parts of any polar ecosystem. Reindeers are a key component of the Arctic  
62 terrestrial ecosystem (Duffy et al. 2005). Because they are a part of a simple food chain, the  
63 species is ideal for monitoring changes in the terrestrial trophic network (Elkin and Bethke,  
64 1995).

65 In this paper, we investigate the usefulness of molten fur collected from a broadly distributed  
66 resident of the European part of the Arctic, namely - the smallest reindeer subspecies  
67 (*Rangifer tarandus platyrhynchus*). This large herbivore, endemic to Svalbard, can be found in  
68 the majority of non-glaciated areas of the island. The Svalbard reindeer has certain



69 adaptations to the polar environment, including relatively short legs and thick fur with  
70 colouring and thickness varying between the seasons (Cuyler and Øritsland, 2002). Its total  
71 population size is estimated for 10,000 animals (npolar.no). Monitoring studies conducted in  
72 Brøggerhalvøya, Reindalen, Adventdalen, and Edgeøya suggest high annual fluctuations  
73 (mosj.no; Reimers, 2012) primarily caused by variations in climate condition (such as snow  
74 depth and rain-on-snow events), and partially by competition for food resources.

75 The primary function of the fur of the Svalbard reindeer is body insulation from cold and wind  
76 (Cuyler and Øritsland, 2002). In cervids, the coat is replaced annually. New fur develops from  
77 late spring/early summer to late fall. The trace element composition of fully grown hairs  
78 largely reflects summer and fall deposition (Drucker et al. 2010). Reindeer hairs develop a  
79 hollow, air-filled, stiff, close-packed structure with a primary heat transfer function. It also  
80 undergoes seasonal changes. Summer and winter fur of adults and calves is characterised by  
81 different properties such as hair length, density, and colour (Cuyler and Øritsland, 2002).

82 The Svalbard reindeer is the only large grazing mammal in the European High Arctic (Hayashi  
83 et al. 2014). It is exposed to contaminants particularly through its diet, composed of different  
84 types of vegetation, including lichen and moss (Robillard et al. 2002). Terrestrial plants receive  
85 metals sprayed from seawater (if they grow within the distance of sea spray influence), by dry  
86 and wet deposition, and from melting glaciers as trapped particles are released from ice (Xie  
87 et al. 2006; Samecka-Cymerman et al. 2011). Birds can also be an additional vector for  
88 contaminant transport (Savinov et al. 2003), as well as reindeer guano (van der Val et al. 2004).  
89 The Svalbard subpopulation eats almost all types of vegetation available. During the growing  
90 season, selection for plant quantity rather than quality is observed (Van der Wal et al. 2000).

91 Plants show variable stable isotope ratios ( $^{13}\text{C}/^{12}\text{C}$  and  $^{15}\text{N}/^{14}\text{N}$ ) depending on their physiology  
92 and environmental conditions, e.g. temperature, light intensity, air humidity, or precipitation  
93 (Drucker et al. 2010). Stable isotopes are incorporated into growing hair from diet, and can be  
94 used to assess spatial and temporal variation in diet components, to characterise the trophic  
95 niche (Boecklen et al. 2011), unravel the migration path (Hobson and Wassenaar, 2008), or  
96 determine habitat selection (Newsome et al. 2009). The ecology of the animal can be  
97 therefore investigated based on stable isotope analysis, as their abundance in tissues reflects  
98 that in the diet (Drucker et al. 2010).

99 The available data on exposure assessment in polar herbivores is still limited, particularly to  
100 the Alaskan and Canadian populations. Also studies concerning stable isotope analysis in  
101 reindeer tissues are scarce. To fill this gap in knowledge, the present study focused on the  
102 investigation on 18 trace elements (Fe, Zn, Ba, Cu, Pb, Cr, Ni, V, Ga, La, Rb, As, Li, Co, Hg, Cd,  
103 Cs, Be), and nitrogen and carbon stable isotopic composition in hairs collected in the summer  
104 season from reindeer herds. The Svalbard reindeer is a sedentary species, migrating only in  
105 the case of significantly reduced food resources (Hansen et al. 2010b). It is therefore  
106 vulnerable to any changes in local foraging conditions. Hairs can be used as a long-range  
107 record of contaminants deposition as they accumulate elements continuously by bounding  
108 them to sulphur-rich hair proteins during the hair growth period (Duffy et al. 2005).

109 The primary objective of this paper is to provide new background data on the levels of metals  
110 in reindeer fur, and a comparison between two subpopulations living in distant areas in order  
111 to establish the pollution level and determine variations in nitrogen ( $\delta^{15}\text{N}$ ) and carbon ( $\delta^{13}\text{C}$ )  
112 stable isotope composition.

## 113 **2. Materials and Methods**

## 114 2.1 Study area and sampling

115 Fur samples were collected in two consecutive summer seasons: in August 2015 from  
116 Longyearbyen region (N78° E015°, n=11) and in September 2016 from the Fuglebekken  
117 catchment in the vicinity of the Polish Polar Station in Hornsund (N77° E015°, n=16) (Fig.1).  
118 Samples were collected from the ground, after a herd moved to a new place. To avoid  
119 pseudoreplication, only freshly molten fur was collected (one sample per at least 4 m<sup>2</sup>  
120 distance). We assumed that samples were from separate individuals. All samples were  
121 individually packed in clean zip bags, and stored at a temperature of 4°C prior to analysis. Long,  
122 straight, white on entire length (except darker tip) guard hairs were collected. Mean  
123 temperature during the period of sample collection amounted to 2.9°C in August 2015  
124 (Longyearbyen) and 3.9°C in September 2016 (Hornsund) (yr.no). Sample weight varied from  
125 16 to 80 mg for samples collected from Longyearbyen, and from 9 to 100 mg for samples  
126 collected from the Hornsund area.

127

128 **Fig. 1 Study area with main coordinates, A – Longyearbyen area, B – Hornsund area [map**  
129 **source: [toposvalbard.npolar.no](http://toposvalbard.npolar.no)]; Svalbard reindeer (*Rangifer tarandus platyrhynchus*)**

130 The Svalbard reindeer, unlike other reindeer subspecies, is highly stationary. It is reluctant to  
131 migrate beyond its territory range mostly established by natural barriers (thin sea ice, glaciers,  
132 steep mountains) (Hansen et al. 2010b). Genetic differences between populations might occur  
133 even at distances <50 km<sup>2</sup> (Côté et al. 2002). Therefore, the studied herds are most likely from  
134 completely separate populations. Predation is almost non-existing, with the exception of local  
135 hunting and occasional evidence of polar bear hunting attempts (Hansen et al. 2011).

## 136 2.2 Analytical methods

137 18 trace elements and nitrogen and carbon stable isotopes composition were analysed. The  
138 basic course of the analytical procedure involves removal of external contamination and then  
139 elemental analysis preceded by acid mineralization in microwave emitter (trace elements  
140 except for mercury), thermal vaporization (mercury) and high temperature oxidation ( $\delta^{13}\text{C}$  and  
141  $\delta^{15}\text{N}$ ).

#### 142 **2.2.1. Trace elements (except for mercury)**

143 First, each hair strand was separated manually from the collected sample with clean tweezers  
144 to separate from any parts of moss collected with the fur ball. To remove the adherent  
145 external contamination such as dust and loosely bound particulate matter, each pooled  
146 sample from one individual was cleaned by vigorous shaking at least 2 times in double  
147 deionised water for 15 min in an automatic shaker, and then air-dried for 24 hours. Only white  
148 hairs were used, and all visible dust particles were washed out. Next, dry hairs were  
149 homogenised by cutting into small parts, weighed to the nearest 0.1 mg, and placed in a clean  
150 teflon vessel with 65%  $\text{HNO}_3$  (Merck, 99% purity). Digestion was carried out using a high-  
151 pressure microwave emitter (Microwave Digestion System, Anton Paar). The temperature was  
152 increased from room temperature to 90°C (app. 6-8°C/min). Such conditions were maintained  
153 for 25 min. After that, temperature was gradually cooled down. Subsequently mineralised  
154 samples were diluted with deionised water into 25 ml in clean plastic flasks. To ensure quality  
155 control, blank samples were run with every batch. The metals were determined by means of  
156 a quadrupole spectrometer ICP-MS Xseries2 by Thermo with inductively-coupled plasma. For  
157 the purpose of reduction of isobaric and polyatomic interferences, a collision/reaction cell was  
158 used with the application of a mix of helium and hydrogen gases, and the kinetic energy  
159 discrimination function (KED).



160 The accuracy of the analyses was verified by means of certified material Standard Reference  
161 Material NIST 1643e Trace Elements in Water and Analytical Reference Material EnviroMAT  
162 ES-H-2 CRM SCP SCIENCE. The retrieval of the elements water ranged from 87% to 109%.

163 The determination was performed at the Department of Hydrology, Faculty of Earth Sciences  
164 and Spatial Management, Marie Curie-Skłodowska University in Lublin.

165 **Tab.1 Detail information about analytical instrumentation (Supplementary Material)**

166 **2.2.2. Mercury analysis**

167 External contamination was washed out using the same procedure as for other trace  
168 elements. The pooled dry sample was cut into smaller pieces using sterilised stainless scissors,  
169 weighed (to the nearest 0.01 mg), and analysed by the thermal vaporisation atomic absorption  
170 method (MA-3000 Nippon Instruments Corporation). The samples were heat decomposed in  
171 a ceramic boat, first heated to 180°C for 120 s, and then to 850°C also for 120 s. The mercury  
172 collector collects the atomised mercury gas in a form of gold amalgam, condensing and  
173 purifying the mercury. After heat decomposition, the mercury collection tube was heated to  
174 650°C to liberate the mercury gas. Absorbance at a wavelength of 253.7 nm was then  
175 measured. Oxygen flow amounted to 0.4 L/min. Total mercury concentration was determined  
176 in triplicates, and based on them the variation coefficient was calculated. Quality control  
177 included blank samples every 5-6 subsamples run. The median of the coefficient of variation  
178 between replicates was equal to 10.0 (7.91-13.95) in samples collected from Longyearbyen,  
179 and 3.65 (1.64-8.98) in samples from Hornsund. Reference materials MODAS-4 Cormorant  
180 Tissue (M-3 CornTis), MODAS-3 Herring Tissue (M-3 HerTis), MODAS-5 Cod Tissue (M-5  
181 CodTis) were used to determine analytical accuracy, and to perform method and quality



182 control. Recovery of reference materials measured on three replicates of each RM varied from  
183 94 to 100%.

### 184 **2.2.3. Stable isotopes**

185 The analyses of carbon and nitrogen stable isotopes ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) were done in an Elemental  
186 Analyser Flash EA 1112 Series combined with an Isotopic Ratio Mass Spectrometer IRMS Delta  
187 V Advantage (Thermo Electron Corp., Germany). Details of these measurements are described  
188 by Kuliński et al. (2014). In short, the samples were dried, homogenised, and weighed into  
189 silver capsules (about 1 mg). This sample weight guarantees C and N loads significantly higher  
190 than those given by the limit of quantification (C = 20  $\mu\text{g}$ , N = 20  $\mu\text{g}$ ). Next, samples were  
191 oxidised in 1020°C in presence of  $\text{Cr}_2\text{O}_3$  and  $\text{Co}_3\text{O}_4$ . After catalytic oxidation, gases including  
192  $\text{CO}_2$ ,  $\text{NO}_x$  and  $\text{H}_2\text{O}$ , were transported to the second reactor, where  $\text{NO}_x$  was reduced to  $\text{N}_2$  on  
193 the metallic Cu (650°C). Subsequently, the analysis products were dried with  $\text{Mg}(\text{ClO}_4)_2$  and  
194 separated on GC (45°C). The separated gases ( $\text{CO}_2$  and  $\text{N}_2$ ) were transported to the IRMS. The  
195 isotopic composition of carbon and nitrogen was calculated using laboratory working pure  
196 reference gases ( $\text{CO}_2$  and  $\text{N}_2$ ) calibrated against IAEA standards: CO-8 and USGS40 for  $\delta^{13}\text{C}$  and  
197 N-1 and USGS40 for  $\delta^{15}\text{N}$ . Results of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  were given in the conventional delta  
198 notation, i.e., versus PDB for  $\delta^{13}\text{C}$  and versus air for  $\delta^{15}\text{N}$  as parts per thousand (‰) according  
199 to the following equation:

$$\delta X (\text{‰}) = \left[ \frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right] \times 1000$$

200  
201  
202 where: X is the stable isotope ratio of  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N}$ ; R is the ratio of  $^{13}\text{C}/^{12}\text{C}$  or  $^{15}\text{N}/^{14}\text{N}$ .  
203 The measurement precision was better than 0.20‰ for  $\delta^{13}\text{C}$  and 0.18‰ for  $\delta^{15}\text{N}$   
204 (n = 5).

### 205 **2.3 Quality assurance/quality control (QA/QC)**

206 To ensure high quality of results, the obtained data were subject to strict quality control  
207 procedures. All the analytical equipment was carefully washed before analysis. Based on  
208 duplicate and triplicate samples, the variance coefficient of metal concentration was  
209 calculated. If the coefficient >15%, samples were excluded from the analyses, assuming  
210 unreliable estimation of metal concentration. Background contamination was present in  
211 metal method blanks prepared after mineralisation, therefore blank correction was  
212 performed for all elements. Blank correction involved subtracting the total amount of analyte  
213 detected in the method blank from the total amount of analyte detected in the hair samples.  
214 Negative numbers and numbers below the limit of detection were reported as half of the limit  
215 of detection for statistical analysis. The obtained results were also corrected for sample  
216 weights and method dilution factor, and are reported as  $\mu\text{g/g dw}$ . All reagents were of the  
217 highest purity. Ultrapure water was produced from a Mili-Q Gradient A10 (Milipore, France).  
218 ICP-MS equipment calibration employed the multi-element standard by Inorganic Ventures  
219 ANALITYK - CCS-1, CCS-4, CCS-6. The optimised and validated methods showed good linearity  
220 ( $R^2 > 0.999$ ) over a wide range with low limits of detection. Both the method limit of detection  
221 (LOD) and the limit of quantitation (LOQ) were calculated based on the standard deviation of  
222 the response (s), and the slope of the calibration curve (b) according to the following formulas:  
223  $\text{LOD} = 3.3(s/b)$ ,  $\text{LOQ} = 10(s/b)$  (LOD/LOQ - Li, Fe, V, Cr, Ni, As, Rb, Ba, Pb 0.1/0.3 ppb; Be, Co,  
224 Ga, Cs, Cd, La 0.01/0.03 ppb; Cu, Zn 0.5/1.5 ppb). For mercury the method limit of detection  
225 and quantification was equal to 0.54 and 1.62 ppb, respectively.

226 Due to the fact that metals are bound to the keratin structure with variable affinity, removal  
227 efficiencies differ significantly among compounds when stronger solvents such as acetone are

228 used. Therefore, only double deionised water was used as a washing agent. Some part of  
229 surface contamination might not have been removed. Because it is difficult to distinguish  
230 between internal and external exposure, it can be assumed that hairs provide information of  
231 integral exposure.

232

## 233 **2.4 Statistical methods**

234

235 Data were log-transformed to meet the assumptions of normality, and consequently  
236 parametric tests were performed. A T-difference test of means was performed for trace  
237 metals and stable isotopes. A Pearson's correlation test was performed to investigate the  
238 relationships between metals and continuous explanatory variables (hair  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$   
239 values). High correlation values between the primary values of the metals in the analysed  
240 samples justify the principal component analysis. Two main components have been  
241 designated for interpretation, accounting for 81.79% of the cases. However, the analysis  
242 provides no meaningful information for the interpretation of data analysis. Therefore, data  
243 clustering was performed to provide an insight into the data structure. Clustering was done  
244 by the nearest neighbour's method, adopting tangent distance as a measure of distance.

245

## 246 **3. Results**

247 Median, mean, and standard error, log transformed mean, and t-difference test of means are  
248 presented in Table 2. For compiled samples, correlation coefficients are mostly high, many are  
249 close to one (Table 4). The correlation of variables with regard to the sampling site was also  
250 tested. In the majority of cases, stronger correlations between metals were observed in  
251 samples from the Longyearbyen area, compared to the Hornsund samples. For zinc,

252 correlations with other metals were notably lower (the highest occurs with gallium content:  
253  $R^2_{\text{tot}} = 0,54$ ). Those coefficients were used to measure similarity of variables by data clustering  
254 (Fig.2). As a result, two groups were obtained: zinc as an isolated element, and other elements  
255 forming a single cluster. After further division, we obtained a five-elemental cluster  
256 (containing V, Fe, Li, Cs, and La), a three-elemental cluster (As, Ga and Ba), and the remaining  
257 elements as isolated items. High variation was observed for nitrogen isotope composition. T-  
258 difference test of means ( $p < 0.05$ ) for nitrogen isotopes ( $\delta^{15}\text{N}$ ) was equal to -5.16, and for  
259 carbon ( $\delta^{13}\text{C}$ ) to -3.12. Three individuals from the Longyearbyen area showed elevated  
260 contents of all the measured elements, with extremely high levels of iron, chromium, nickel,  
261 and lead. The average value of nitrogen isotope  $\delta^{15}\text{N}$  for those outliers was equal to 6.95 [‰].  
262 Outliers were not excluded from statistical analysis.

263

264 **Table 2. Trace element concentration in reindeer fur samples collected from two separate**  
265 **populations ( $\mu\text{g/g dw}$ )**

266 **Table 3. Nitrogen and carbon stable isotopes concentration in Svalbard reindeer hairs**

267 **Table 4. Pearson correlation values indicating correlation between the various trace**  
268 **elements measured (n=26)**

269 **Fig. 2. Hierarchical dendrogram for clustering chemical elements. Lines indicate distance**  
270 **0.27 and 0.5, respectively. From the left: 1-V, 2-Fe, 3-Li, 4-Cs, 5-La, 6,5-Rb, 8-As, 9-Ga, 10-Ba,**  
271 **11,5-Be, 13-Ni, 14,5-Cr, 16-Co, 17,5-Pb, 19-Cu, 20,5-Cd, 22,5-Zn**

272

273 **4. Discussion**

274 This study reports the levels of essential and toxic elements and stable isotope composition in  
275 Svalbard reindeer hair samples collected from herds living in distant parts of the island.  
276 Keratinised tissues such as hairs, fur, or feathers can be collected non-lethally, and have been  
277 successfully used for stable isotopes and heavy metal analysis for many years (Duffy et al.  
278 2005; Burger et al. 2007; Sergiel et al. 2017). Hair tissue has several advantages in practical  
279 use. Owing to its stability, samples can be stored for a long time, they are relatively  
280 metabolically inactive (Duffy et al. 2005), and elements are accumulated over extended  
281 periods of time. Therefore, the exposure assessment covers several weeks or months. Molten  
282 hairs can be collected without direct contact, avoiding difficulties related to capturing a free-  
283 living individual. However, because factors such as specimen age and gender are often  
284 unknown, this mode of sampling also limits the possibility of result interpretation.

285 Svalbard reindeers consume various plants, including vascular plants, bryophytes, and lichens,  
286 all determined to accumulate high levels of essential and heavy elements (Jóźwik, 1990,  
287 Samecka-Cymerman et al. 2011, Garty, 2001). Their levels found in polar plant species can be  
288 elevated due to natural processes (such as volcano eruptions, rock weathering) or  
289 atmospheric deposition, mainly from long distance transboundary transport from lower  
290 latitudes (Grodzińska and Grodzik, 1991). Sea aerosol can be an additional source of elements  
291 such as lead, mercury, and cesium (Kłos et al. 2017).

292 Spatial and temporal heterogeneity in diet components might be responsible for significant  
293 seasonal differences in contaminant distribution across studies (Robillard et al. 2002). In our  
294 study, the majority of elements showed a strong positive correlation with multi-element  
295 totals, excluding zinc. High variability in trace element composition was observed even above  
296 an order of magnitude within samples of reindeer from one location. This is probably related  
297 to differences in age (herds were composed from both young and older individuals), gender,

298 and food preference. Due to lack of previous studies regarding trace elements in reindeer  
299 hairs, our data can be used as a reference for future investigations in the Svalbard Archipelago  
300 concerning reindeer and closely related species.

#### 301 **4.1 Accumulation route**

302 Vegetation covers only 6-7% of the area of Svalbard. The growing season lasts approximately  
303 90 days (Kłos et al. 2015). Because of the short grazing season, the Svalbard reindeer must  
304 restore its body reserves after winter, and accumulate fat at this time (Staaland, 1984). The  
305 plant species-specific physiology, age, and sampling location will correspond with forward  
306 exposure. Lower trace element levels are observed in vascular plants as compared to mosses  
307 and lichens (Wojtuń et al. 2013). This may be related to their higher morphological similarity,  
308 and more selective accumulation process (Chiarenzelli et al. 2001). Due to the lack of root  
309 system, slow growth rate, longevity, vast surface area, and lack of well-developed cuticle,  
310 plants such as lichens and bryophytes are prone to accumulating a varied cocktail of toxic  
311 compounds from the atmosphere (Robillard et al. 2002; Gamberg et al. 2005, Samecka-  
312 Cymerman et al. 2011). Essential elements such as copper and zinc, necessary for plant  
313 growth, can also be accumulated beyond physiological demands (Samecka-Cymerman et al.  
314 2011, Józwik, 1990). For instance, for zinc, enhanced exposure in lichens is above 500 µg/g,  
315 cadmium can be tolerated between 1 and 30 µg/g, and copper between 1 and 50 µg/g  
316 (Nieboer et al. 1978).

317 The accumulation route can be passive by water transpiration passage (e.g. Cu in lichens),  
318 active (e.g. zinc), and metabolic (e.g. manganese), or a mix of those factors (Józwik 1990).  
319 Mosses are evidenced to accumulate notably high levels of Cd, Co, Cr, Cu, Fe, Mn, and Zn,  
320 even higher than lichens (Wojtuń et al. 2013). Particularly moss species such as *Aulacomium*  
321 *palustre*, *A. turgidum*, *Hylocomium splendens*, *Sanionia uncinata*, and *Tortula ruralis* are  
322 suspected to be very good heavy metal accumulators in Svalbard (Grodzińska and Grodzik,  
323 1991).

324

## 325 **4.2 Toxic elements**

326 Mercury is a global pollutant that enters the Arctic terrestrial ecosystem mainly through rock  
327 weathering and long-range atmospheric deposition (Gamberg et al. 2015). During spring,  
328 atmospheric Hg(0) is oxidised into Hg(II), and deposited in the snow, ice, or ocean surfaces  
329 from where can be partly reemitted or further retained, transformed, and transported  
330 (Schroeder et al. 1998; Halbach et al. 2017). In addition to snow and ice, soil is believed to be  
331 a major land mercury reservoir in the Arctic (Gamberg et al. 2015). Our study shows low  
332 mercury contents in both studied subpopulations. Elevated mercury level is indeed usually  
333 found in marine biota, in contrast to terrestrial mammals, especially herbivores with a short  
334 food chain.

335 To the best of our knowledge, no studies are available regarding contaminant deposition in  
336 the hair of the Svalbard reindeer subspecies. Duffy et al. (2005) conducted a study on mercury  
337 levels in the hair of the Alaskan reindeer population, indicating low exposure (mean total  
338 mercury for free ranging individuals was equal to 0.055 µg/g). Mercury was also a major  
339 research interest in Lokken et al. (2009) pilot study performed on lichen and the Alaskan  
340 caribou population (mean hair levels varied from 0.0146 to 0.0834 µg/g). In the present study,  
341 the highest level was found in the Longyearbyen population. It does not exceed 0.160 µg/g  
342 (median equal to 0.112 and 0.060 µg/g).

343 Mercury and cadmium previously showed a clear pattern of accumulation towards higher  
344 trophic levels in the terrestrial ecosystem (Dietz et al. 2000). Cadmium binds to the low  
345 molecular weight sulphur-rich proteins, and accumulates mostly in kidneys (Chan et al. 2001).  
346 It also may significantly increase with age (Danielsson and Frank, 2009). In our study, however,  
347 age differences were not analysed, and hair bounding capacities are different than in internal  
348 tissues. Literature studies on both areas showed low cadmium exposure in vegetation (Wojtuń



349 et al. 2013; Samecka-Cymerman et al. 2011; Węgrzyn et al. 2013; Kłos et al. 2015), and as  
350 expected we found low levels in reindeer hair. To our best knowledge, no study has been  
351 published concerning cadmium accumulation in Svalbard mammal herbivores, therefore no  
352 comparison is possible.

353 On the other hand, high lead levels were found in the majority of samples, suggesting an  
354 accumulation path by vegetation. High levels of lead were also previously found in Greenland  
355 soils. However, it does not tend to accumulate towards higher trophic levels, as reindeers had  
356 lower lead levels than lichens (summarized in Dietz et al. 2000 based on Greenlandic studies  
357 of the AMAP programme). Notice that only reindeer internal tissues were used. In Svalbard  
358 area, levels of lead in vegetation is highly variable. Threshold values for lead in lichens are  
359 from 5 to 100  $\mu\text{g/g}$  and 15  $\mu\text{g/g}$  is a boundary for enhanced exposure (Nieboer and Richardson,  
360 1981). In hairs, lead is accumulated both externally and internally over a long period of time,  
361 until molting. It is possible that apart from internal contamination accumulated by foraging on  
362 high-lead level food sources, part of external contamination was not washed out during the  
363 cleaning procedure.

364  
365 **Fig. 3a,b. Plot of average Cd, Pb and Hg and Fe, Zn, Ba and Cu concentration in Longyearbyen**  
366 **(dark colors) and Hornsund (light colors) reindeer hair samples. Values are log transformed.**  
367 **The horizontal lines represent medians, the boxes – upper and lower (25-75% quartiles) and**  
368 **whiskers – minimum and maximum values**

### 370 4.3 Other elements

371 The studied samples showed particular patterns such as high intra-individual variations in the  
372 level of several compounds (iron, chromium, zinc etc.). All the analysed elements occur in

373 broad concentration ranges. Relatively high levels of mean nickel in the Longyearbyen  
374 subpopulation, before also observed in the population of moss *Hylocomium splendens*, could  
375 be associated with past mining activities in the area (Kłos et al. 2015). The main source of  
376 nickel in Longyearbyen is most likely rock waste derived from mining activities and aviation  
377 emissions, although discharges transported long-range from the Kola Peninsula are also  
378 suspected (Kłos et al. 2017). Iron was significantly elevated in some of the samples from the  
379 Longyearbyen area, with the highest level at 14640 µg/g dw. Other two samples were also  
380 above 5000 µg/g dw of iron. The effect of spontaneous iron overload was previously described  
381 in liver tissues of Svalbard reindeer (Borch-Iohnsen and Nilssen, 1987; Borch-Iohnsen and  
382 Thorstensen, 2009). It was caused by high uptake of dietary iron consumed with iron-rich  
383 forage plants (Borch-Iohnsen and Thorstensen, 2009). In Svalbard reindeers, spontaneous  
384 seasonal iron overload with massive siderosis is considered natural, and occurs mostly in  
385 winter when available vegetation is of poorer quality (Borch-Iohnsen and Thorstensen, 2009).  
386 It is possible that when reindeers' nutritional conditions improved after winter (Borch-Iohnsen  
387 and Thorstensen, 2009), accessory iron was redistributed from the liver to hairs. If that is the  
388 case, hairs can be used to reveal past iron overload. All other elements were also significantly  
389 elevated in those individuals, suggesting some health implications (with examples presented  
390 in Table 5, Supplementary material.). Mercury was not analysed in those samples. Levels of  
391 iron in samples from the Hornsund area were lower, not exceeding 5000 µg/g. In two cases,  
392 more than 1100 µg/g of iron was detected.

393

394 **Table 5. Outliers with significantly elevated element levels µg/g dw (Supplementary**  
395 **Material)**

396

397 Because reindeer subspecies *Rangifer tarandus platyrhynchus* lives exclusively in the Svalbard  
398 Archipelago, the nominative species was expected to receive more attention. Studies on  
399 Canada and Greenland caribou and reindeer populations mostly concerned internal tissues  
400 (Elkin and Bethke, 1995, Robillard et al. 2002, Larter and Nagy, 2000, Aastrup et al. 2000).  
401 Medvedev (1995) reported cadmium and lead levels in the bone, teeth, and antlers of forest  
402 reindeer (*Rangifer tarandus fennica*) from north-west Russia. The highest mean levels of  
403 cadmium and lead were found in the bone tissue ( $2.1\pm 1.1$  and  $41.6\pm 23.7$   $\mu\text{g/g dw}$ ,  
404 respectively). The levels did not depend on sex or age of individuals. Heavy metal levels were  
405 also reported for North Norway population in samples collected from semi-domesticated  
406 reindeer. Cadmium, lead, arsenic, nickel, and vanadium were determined in the muscle, liver,  
407 tallow, and bone marrow tissues, with the highest level of all the elements in the liver (except  
408 nickel) (Ali Hassan et al. 2012). A reliable comparison between those studies is not possible,  
409 however, because the relationship between deposition of compounds in hairs and internal  
410 tissues is not always clear. Svalbard is an Arctic semi-desert compared to other places  
411 inhabited by reindeers, with low precipitation and humidity, cold winter temperatures, and  
412 high wind speed, resulting in different feeding behaviour and patch choice (Lindner, 2002).  
413 The Svalbard reindeer also differs from other reindeer subspecies in its anatomy and  
414 physiology (Lindner, 2002).

415

#### 416 **4.4 Stable isotopes of carbon and nitrogen**

417

418 Stable isotopes (SI) of nitrogen ( $\delta^{15}\text{N}$ ) and carbon ( $\delta^{13}\text{C}$ ) are increasingly employed as an  
419 indispensable tool in ecological studies (Sergiel et al. 2017). The main sources of nitrogen in  
420 the Arctic include atmospheric discharge of  $\text{NO}_x$ ,  $\text{NH}_x$ , primary  $\text{N}_2$ -fixation from the

421 atmosphere, and bird guano (Skrzypek et al. 2015). In nitrogen-limited terrestrial ecosystems  
422 such as Arctic tundra, soil microbes are recognised to function as main nitrogen pools,  
423 competing for nitrogen with plants (Bardgett et al. 2007). Plant growth is limited by nitrogen  
424 availability. Consequently, the capacity for carbon sequestration is also restricted (Skrzypek et  
425 al. 2015). Arctic tundra contains a significant percent of the global soil carbon reserve. Its  
426 storage is controlled by factors such as e.g. temperature, vegetation type, soil hydrology, or  
427 shifts in vegetation state. The latter can be induced by herbivores (Van der Val, 2006; Speed  
428 et al. 2010).

429 Forage patch choice by reindeers and nitrogen content in plants are largely influenced by the  
430 timing of snowmelt (van der Wal et al. 2000). In Svalbard, seasonal variability of plant and soil  
431 nitrogen pools are mostly controlled by changes in temperature and soil moisture over the  
432 growing season. Such changes, however, are markedly lower than in the other seasonally cold  
433 ecosystems (Bardgett et al. 2007). Also Arctic tundra has a high capacity to retain nitrogen  
434 transported after extreme events, with non-vascular plants acting as a short-term sink, and  
435 vascular plants as a long-term reservoir (Choudhary et al. 2016). Our results indicate high  
436 variability in the  $^{15}\text{N}:$  $^{14}\text{N}$  ratio, suggesting that reindeers consume vegetation with different  
437  $^{15}\text{N}$  values. In the Fuglebekken catchment (Hornsund), high loads of nutrients are deposited  
438 by large bird colonies such as little auk (*Alle alle*). This influx impacts soil fertility and  
439 subsequently plant productivity and structure (Skrzypek et al. 2015). As a result, the available  
440 vegetation differs in protein, sugar composition, and digestibility (Staaland, 1984). Bird guano  
441 and additional N-sources from colonies, such as carcasses, dead chicks, and eggshells,  
442 constitute a huge N-load compared to other sources (Skrzypek et al. 2015). It could account  
443 for significant differences between the two subpopulations.

444 Moss tundra serves as an important sink for carbon sequestration (Nakatsubo et al. 2015).  
445 Here, relatively low variability was observed for stable carbon isotopes. No significant  
446 correlation was observed between C and N values and metal concentration, apart from zinc.

447  
448 No previous studies are available concerning stable isotope analysis in the keratinised tissues  
449 of the Svalbard reindeer. Mosbacher (et al. 2016) showed high inter- and intra-annual  
450 seasonality in the diet of the Greenland muskox (*Ovibos moschatus*) by the application of  
451 sequential data on nitrogen stable isotopes derived from guard hairs. Drucker (et al. 2010)  
452 studied the dietary references and habitat use of moose (*Alces alces*) and caribou (*Rangifer*  
453 *tarandus*) in plucked hair samples from Canada populations. The dietary strategies of those  
454 species differ in spite of the same habitat range. Differences in stable isotope abundance were  
455 significantly linked to the species' dietary specialisation (Drucker et al. 2010).

456  
457

458 **Fig. 4. Plot of average nitrogen (blue) and carbon (green) stable isotope composition in**  
459 **Longyearbyen (L) and Hornsund (H) reindeer hair samples. The horizontal lines represent**  
460 **medians, the boxes – upper and lower (25-75% quartiles) and whiskers – minimum and**  
461 **maximum values excluding outliers**

462  
463 The long-term variation in weather conditions may impact vegetation quality, consequently  
464 affecting the ungulates' nutritional profile and foraging conditions. Lower snow layer  
465 hardening in winter leads to changes in snow-pack properties, including ground icing, resulting  
466 in snowpack with impenetrable vegetation underneath (Hansen et al. 2011, Loe et al. 2016).  
467 Food availability can also be restricted by overgrazing (Węgrzyn et al. 2016). Therefore, some

468 populations are more likely to expand their foraging area, or alternatively use less preferred  
469 food sources such as goose droppings (van der Wal and Loonen, 1998) or marine algae  
470 (Hansen and Aanes, 2012). Because many factors are responsible for seasonal availability of  
471 various food sources, and Svalbard reindeers tend to forage for plant quantity rather than  
472 quality (Van der Wal et al. 2000), a complex study program concerning trace element levels in  
473 vegetation may help assess their future potential exposure.

474

## 475 **5. Conclusion**

476 The Svalbard reindeer is one of the least studied subspecies amongst family *Rangifer*. In this  
477 paper, we present to the best of our knowledge the first communication concerning trace  
478 element concentration in hairs of two separate subpopulations. Better knowledge of the  
479 potential impacts of metal on the terrestrial ecosystem is needed in polar mammal  
480 populations, especially to identify levels related to health dysfunction. In the present study,  
481 mercury is indicated as an insignificant thread in terrestrial ecosystem, although levels of lead,  
482 chromium, and nickel were noticeably elevated in some of the samples. Because hairs are a  
483 dead tissue accumulating elements over long period of time, reindeer may use it in a  
484 detoxification process for instance for depositing past iron overload.

485 Future climate changes will induce higher pressure on all terrestrial species. Rising  
486 temperatures, more frequent extreme weather events, heavy rain-on-snow events, and  
487 variations in seasonal precipitation patterns may cause negative implications for herbivores  
488 (Hansen and Aanes, 2012). In spite of their remarkable abilities to locate food beneath the  
489 snow-pack, severe icy conditions may induce changes in reindeer behaviour, including range  
490 expansion to mountainous terrain (Hansen et al. 2010), and eating marine algae (Hansen and  
491 Aanes, 2012) resulting in potential changes in the foraging profile and contaminant

492 accumulation. The research presented so far provides evidence that keratinised tissues can be  
493 a valuable source of information in ecotoxicological studies. Monitoring studies should involve  
494 not only marine species, but concurrently more terrestrial key species as an important part of  
495 the trophic network.

496

#### 497 **Acknowledgements**

498 Authors would like to thank Sara Lehmann-Konera and Katarzyna Kozak for sample collection  
499 in 2015. Katarzyna Koziarowska's participation in the study was supported by the Centre for  
500 Polar Studies, KNOW – Leading National Research Centre, Sosnowiec, Poland.

#### 501 **6. References:**

502 AMAP (2005) AMAP Assessment 2002: Heavy Metals in the Arctic. Arctic Monitoring and Assessment  
503 Programme (AMAP)

504 Aastrup P, Riget F, Dietz R, Asmund G. (2000) Lead, zinc, cadmium, mercury, selenium and copper in  
505 Greenland caribou and reindeer (*Rangifer tarandus*) Sci Total Environ 245, 149-159

506 Ali Hassan A, Rylander Ch, Brustad M, Sandanger TM. (2012) Level of selected toxic elements in meat,  
507 liver, tallow and bone marrow of young semidomesticated reindeer (*Rangifer tarandus tarandus* L.)  
508 from Northern Norway, Int J Circumpolar Health, 71, 18187

509 Bard SM. (1999) Global transport of anthropogenic contaminants and the consequences for the Arctic  
510 marine ecosystem. Mar Pollut Bull 38, 356–379

511 Bardgett RD, van der Wal R, Jónsdóttir IS, Quirk H, Dutton S. (2007) Temporal variability in plant and  
512 soil nitrogen pools in a high-Arctic ecosystem, Soil Biol Biochem 39, 2129–2137



- 513 Boecklen WJ, Yarnes CT, Cook BA, James AC. (2011) On the use of stable isotopes in trophic ecology,  
514 Annu Rev Ecol Evol Syst 42, 411–440
- 515 Borch-Iohnsen B, Nilssen KJ. (1987) Seasonal iron overload in Svalbard reindeer liver, J Nutr 117, 2072-  
516 2078
- 517 Borch-Iohnsen B, Thorstensen K. (2009) Iron Distribution in the Liver and Duodenum during Seasonal  
518 Iron Overload in Svalbard Reindeer, J Comp Path 141, 27-40
- 519 Burger J, Gochfeld M, Sullivan K, Irons D. (2007) Mercury, arsenic, cadmium, chromium lead, and  
520 selenium in feathers of pigeon guillemots (*Cepphus columba*) from Prince William Sound and the  
521 Aleutian Islands of Alaska, Sci Total Environ 387, 175–184
- 522 Chan HM, Kim C, Leggee D. (2001) Cadmium in caribou (*Rangifer tarandus*) kidneys : speciation, effects  
523 of preparation and toxicokinetics. Food Addit Contam 18, 607–614
- 524 Chiarenzelli J, Aspler L, Dunn C, Cousens B, Ozarko D, Powis K. (2001) Multi-element and rare earth  
525 element composition of lichens, mosses, and vascular plants from the Central Barrenlands, Nunavut,  
526 Canada, Appl Geochem 16, 245-270
- 527 Choudhary S, Blaud A, Osborn AM, Press MC, Phoenix GK. (2016) Nitrogen accumulation and  
528 partitioning in a High Arctic tundra ecosystem from extreme atmospheric N deposition events, Sci Total  
529 Environ 554-555, 303-310
- 530 Côté SD, Dallas JF, Marshall F, Irvine RJ, Langvatn R, Albon SD. (2002) Microsatellite DNA evidence for  
531 genetic drift and philopatry in Svalbard reindeer. Mol Ecol 11, 1923-1930
- 532 Cuyler Ch, Øritsland NA. (2002) Do seasonal changes in Svalbard reindeer fur have relevance for heat  
533 transfer?, Rangifer 22, 133-142
- 534 Danielsson R, Frank A. (2009) Cadmium in moose kidney and liver – age and gender dependency, and  
535 standardization for environmental monitoring, Environ Monit Assess 157, 73–88



- 536 Davis N. (1996) The Arctic wasteland: a perspective on Arctic pollution, *Polar Rec* 32, 237–248
- 537 Dietz R, Riget F, Cleemann M, Aarkrog A, Johansen P, Hansen JC. (2000) Comparison of contaminants  
538 from different trophic levels and ecosystems, *Sci Total Environ* 245, 221-231
- 539 Duffy LK, Duffy RS, Finstad G, Gerlach C. (2005) A note on mercury levels in the hair of Alaskan reindeer,  
540 *Sci Total Environ* 339, 273– 276
- 541 Drucker DG, Hobson KA, Ouellet J-P, Courtois R. (2010) Influence of forage preferences and habitat  
542 use on 13C and 15N abundance in wild caribou (*Rangifer tarandus caribou*) and moose (*Alces alces*)  
543 from Canada, *Isotopes Environ Health Stud*, 46, 107–121
- 544 Elkin BT, Bethke RW. (1995) Environmental contaminants in caribou in the Northwest Territories,  
545 Canada, *Sci Total Environ* 160/161, 307-321
- 546 Fuglei E, Øritsland NA, Prestrud P. (2003) Local variation in arctic fox abundance on Svalbard, Norway,  
547 *Polar Biol* 26, 93–98
- 548 Gamberg M, Braune B, Davey E, Elkin B, Hoekstra PF, Kennedy D, Macdonald C, Muir D, Nirwal A,  
549 Wayland M, Zeeb B. (2005) Spatial and temporal trends of contaminants in terrestrial biota from the  
550 Canadian Arctic, *Sci Total Environ* 351–352, 148– 164
- 551 Gamberg M, Chételat J, Poulain AJ, Zdanowicz C, Zheng J. (2015) Mercury in the Canadian Arctic  
552 terrestrial environment: an update. *Sci Total Environ* 509-510, 28-40
- 553 Garty J. (2001) *Biomonitoring Atmospheric Heavy Metals with Lichens: Theory and Application*, CRC  
554 *Crit Rev Plant Sci* 20, 309-371
- 555 Grodzińska K, Godzik B. (1991) Heavy metals and sulphur in mosses from Southern Spitsbergen, *Polar*  
556 *Res* 9, 133–140



- 557 Gustine DD, Barboza PS, Lawler JP, Arthur SM, Shults BS, Persons K, Adams LG. (2011) Characteristics  
558 of foraging sites and protein status in wintering muskoxen: insights from isotopes of nitrogen, *Oikos*  
559 120, 1546–1556
- 560 Halbach K, Mikkelsen Ø, Berg T, Steinnes E. (2017) The presence of mercury and other trace metals in  
561 surface soils in the Norwegian Arctic, *Chemosphere* 188, 567-574
- 562 Hansen BB, Aanes R, Sæther B.-E. (2010a) Feeding-crater selection by High-Arctic reindeer facing ice-  
563 blocked pastures, *Can J Zool* 88, 170-177
- 564 Hansen BB, Aanes R, Sæther B.-E. (2010b) Partial seasonal migration in high-arctic Svalbard reindeer  
565 (*Rangifer tarandus platyrhynchus*), *Can J Zool* 88, 1202-1209
- 566 Hansen BB, Aanes R, Herfindal I, Kohler J, Sæther B.-E. (2011) Climate, icing, and wild arctic reindeer:  
567 past relationships and future prospects, *Ecology* 92, 1917–1923
- 568 Hansen BB, Aanes R. (2012) Kelp and seaweed feeding by High-Arctic wild reindeer under extreme  
569 winter conditions, *Polar Res* 31, 17258
- 570 Hobson KA, Wassenaar LI, eds. (2008) Tracking animal migration using stable isotopes. Handbook of  
571 Terrestrial Ecology Series, Academic Press/Elsevier, Amsterdam, p. 188
- 572 Jóźwik Z. (1990) Heavy metals in tundra plants of Bellsund area, Spitzbergen, *Pol Polar Res* 11, 3-4, 401-  
573 409
- 574 Kłos A, Bochenek Z, Bjerke JW, Zagajewski B, Ziółkowski D, Ziembik Z, Rajfur M, Dołhańczuk-Śródka A,  
575 Tømmervik H, Krems P, Jerz D, Zielińska M. (2015) The use of mosses in biomonitoring of selected areas  
576 in Poland and Spitzbergen in the years from 1975 to 2014, *Ecol Chem Eng S* 22, 201-218
- 577 Kłos A, Ziembik Z, Rajfur M, , Dołhańczuk-Śródka A, Bochenek Z, Bjerke JW, Tømmervik H, Zagajewski  
578 B, Ziółkowski D, Jerz D, Zielińska M, Krems P, Godyń P. (2017) The origin of heavy metals and

579 radionuclides accumulated in the soil and biota samples collected in Svalbard, near Longyearbyen, Ecol  
580 Chem Eng S 24, 223-238

581 Kuliński K, Kędra M, Legeżyńska J, Głuchowska M, Zaborska A. (2014) Particulate organic matter sinks  
582 and sources in high Arctic fjord, J Mar Syst, 139, 27-37

583 Larter NC, Nagy JA. (2000) A comparison of heavy metal levels in the kidneys of High Arctic and  
584 mainland caribou populations in the Northwest Territories of Canada, Sci Total Environ 246, 109-119

585 Lindner E. (2002) Use of vegetation types by Svalbard reindeer from arctic winter to spring, Polar Rec  
586 39, 245–247

587 Lokken JA, Finstad GL, Dunlap KL, Duffy LK. (2009) Mercury in lichens and reindeer hair from Alaska:  
588 2005–2007 pilot survey, Polar Rec 45, 368–374

589 Loe LE, Hansen BB, Stien A, Albon SD, Bischof R, Carlsson A, Irvine RJ, Meland M, Rivrud IM, Ropstad E,  
590 Veiberg V, Myrsterud A. (2016) Behavioral buffering of extreme weather events in a high-Arctic  
591 herbivore, Ecosphere 7(6):e01374. 10.1002/ecs2.13

592 Medvedev N. (1995) Concentrations of cadmium, lead and sulphur in tissues of wild, forest reindeer  
593 from north-west Russia, Environ Pollut 90, 1-5

594 Mosbacher JB, Michelsen A, Stelvig M, Hendrichsen DK, Schmidt NM. (2016) Show Me Your Rump Hair  
595 and I Will Tell You What You Ate – The Dietary History of Muskoxen (*Ovibos moschatus*) Revealed by  
596 Sequential Stable Isotope Analysis of Guard Hairs, PLoS ONE 11(4): e0152874

597 Mosj.no, [WWW Document]. URL [http://www.mosj.no/en/fauna/terrestrial/svalbard-reindeer-](http://www.mosj.no/en/fauna/terrestrial/svalbard-reindeer-population.html)  
598 [population.html](http://www.mosj.no/en/fauna/terrestrial/svalbard-reindeer-population.html) [available 20.12.17]

599 Nakatsubo T, Uchida M, Sasaki A, Kondo M, Yoshitake S, Kanda H. (2015) Carbon accumulation rate of  
600 peatland in the High Arctic, Svalbard: Implications for carbon sequestration, Polar Sci 9, 267-275

601 Newsome SD, Tinker MT, Monson DH, Oftedal O, Ralls K, Fogel ML, Estes JA. (2009) Using stable  
602 isotopes to investigate individual diet specialization in California sea otters (*Enhydra lutris nereis*),  
603 Ecology 90, 961–974

604 Nieboer E, Richardson DHS, Tomassini FD. (1978) Mineral uptake and release by lichens: an overview,  
605 The Bryologist 81, 226–246

606 Nieboer E, Richardson DHS. (1981) Lichens as monitors of atmospheric deposition. In S. J. Eisenreich  
607 (Ed.), Atmospheric pollutants in natural waters (pp. 339–388). Ann Arbor MI: Ann Arbor Science  
608 Publishers

609 npolar.no, [WWW Document]. URL <http://cruise-handbook.npolar.no/en/svalbard/wildlife.html>  
610 [available 20.12.17]

611 Reimer E. (2012) Svalbard reindeer population size and trends in four sub-areas of Edgeøya, Polar Res  
612 31, 11089, DOI: 10.3402/polar.v31i0.11089

613 Robillard S, Beauchamp G, Paillard G, Belanger D. (2002) Levels of Cadmium, Lead, Mercury and  
614 <sup>137</sup>Caesium in Caribou (*Rangifer tarandus*) Tissues from Northern Québec, Arctic 55, 1-9

615 Savinov VM, Gabrielsen GW, Savinova TN. (2003) Cadmium, zinc, copper, arsenic, selenium and  
616 mercury in seabirds from the Barents Sea: levels, inter-specific and geographical differences, Sci Total  
617 Environ 306, 133–158

618 Samecka-Cymerman A, Wojtuń B, Kolon K, Kempers AJ. (2011) *Sanionia uncinata* (Hedw.) loeske as  
619 bioindicator of metal pollution in polar regions, Polar Biol 34, 381–388

620 Schroeder WH, Anlauf KG, Barrie La, Lu JY, Steffen A. (1998) Arctic springtime depletion of mercury,  
621 Nature 394, 331-332

622 Sergiel A, Hobson KA, Janz DM, Cattet M, Selva N, Kapronczai L, Gryba C, Zedrosser A. (2017)  
623 Compatibility of preparatory procedures for the analysis of cortisol concentrations and stable isotope

624 ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) ratios: a test on brown bear hair. *Conserv Physiol* 5(1): cox021;  
625 doi:10.1093/conphys/cox021

626 Speed JDM, Woodin SJ, Tømmervik , van der Wal R. (2010) Extrapolating herbivore-induced carbon  
627 loss across an arctic landscape, *Polar Biol* 33, 789–797

628 Staaland H. (1984) On the quality of Svalbard reindeer pasture in the summer and autumn, *Rangifer* 4,  
629 16-23

630 Toposvalbard.npolar.no, [WWW Document]. URL <http://toposvalbard.npolar.no/> [available 20.12.17]

631 Van der Wal R, Loonen MJJE. (1998) Goose droppings as food for reindeer, *Can J Zool* 76, 1117-1122

632 Van der Wal R, Madan N, van Lieshout S, Dormann C, Langvatn R, Albon SD. (2000) Trading forage  
633 quality for quantity? Plant phenology and patch choice by Svalbard reindeer, *Oecologia* 123, 108–115

634 Van der Wal R, Bardgett RD, Harrison RD, Stien A. (2004) Vertebrate herbivores and ecosystem control:  
635 cascading effects of faeces on tundra ecosystem, *Ecography* 27, 242-252

636 Van der Wal R. (2006) Do herbivores cause habitat degradation or vegetation state transition?  
637 Evidence from the tundra, *Oikos* 114, 177–186

638 Węgrzyn M, Lisowska M, Nicia P. (2013) The value of the terricolous lichen *Cetrariella delisei* in the  
639 biomonitoring of heavy–metal levels in Svalbard, *Pol Polar Res* 34, 375–382

640 Węgrzyn M, Wietrzyk P, Lisowska M, Klimek B, Nicia P. (2016) What influences heavy metals  
641 accumulation in arctic lichen *Cetrariella delisei* in Svalbard?, *Polar Sci* 10, 532-540

642 Xie Z, Sun L, Blum JD, Huang Y, He W. (2006) Summertime aerosol chemical components in the marine  
643 boundary layer of the Arctic Ocean, *J Geophys Res* 111:D10309

644 yr.no, n.d. Weather statistics for Longyearbyen (Svalbard) - yr.no [WWW Document]. URL  
645 <http://www.yr.no/place/Norway/Svalbard/Longyearbyen/statistics> [available 20.12.17]



646

647

648 **Fig. 1 Study area with main coordinates, A-Longyearbyen area, B- Hornsund area [map**  
 649 **source: toposvalbard.npolar.no]; Svalbard reindeer (*Rangifer tarandus platyrhynchus*)**

650

651

652


653

654

655

656

657



658

659 Fig. 2 Hierarchical dendrogram for clustering the chemical elements. Lines indicate distance  
 660 0.27 and 0.5, respectively. From the left: 1-V, 2-Fe, 3-Li, 4-Cs, 5-La, 6,5-Rb, 8-As, 9-Ga, 10-Ba,  
 661 11,5-Be, 13-Ni, 14,5-Cr, 16-Co, 17,5-Pb, 19-Cu, 20,5-Cd, 22,5-Zn

662

663

664

665

666

667


668

669


670

671

672



673




674

675 Fig. 3a,b. Plot of average Cd, Pb and Hg and Fe, Zn, Ba and Cu concentration in Longyearbyen  
 676 (dark colors) and Hornsund (light colors) reindeer hair samples. Values are log transformed.  
 677 The horizontal lines represent medians, the boxes – upper and lower (25-75% quartiles) and  
 678 whiskers – minimum and maximum values

679





680

681

682 Fig. 4. Plot of average nitrogen (blue) and carbon (green) stable isotope composition in  
 683 Longyearbyen (L) and Hornsund (H) reindeer hair samples. The horizontal lines represent  
 684 medians, the boxes – upper and lower (25-75% quartiles) and whiskers – minimum and  
 685 maximum values excluding outliers

686

687

688

689

690

691

692

693

ICP-MS parameter and accessories	Value
Radio frequency power generator	1400 V
Gas type	Argon
Plasma gas flow rate	12 L/min
Auxiliary gas flow rate	0.7 L/min
Nebulization gas flow rate	0.9 L/min
Torch Option	Standard one-piece quartz torch with PlasmaScreenPlus
Nebulizer	Standard glass concentric
Spray chamber	Quartz impact bead
Cones	Xt
Internal Standard	<sup>6</sup> Li, Sc, Y, In, Tb, Bi
Sample Uptake Rate (mL/min)	0.4 approx.
Sampling depth	98 mm
Collision Cell Gas flow (7 % H <sub>2</sub> in He)	5.5 mL/min
Number of replicates	3

695

Mercury analyzer specification	
Detectors	Photo tubes (Reference-background; Absorption cell 1; Absorption cell 2)
Wave length	253.7nm
Maximum measurement range	70,000ng

Measuring time	Approx. 5 minutes
Maximum decomposition temp.	Up to 1,000°C
Combustion tube	Quartz (Filled with catalyst)
Gas	Oxygen (>90% purity), 0.1~0.29MPa
Sample boat	Ceramic (standard supply)

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713 Tab. 2 Trace element concentration in reindeer fur samples collected from two separate  
 714 populations ( $\mu\text{g/g dw}$ )

Element	Longyearbyen (n=11)			Hornsund (n=16)			t – difference test of means ( $p < 0.05$ )
	Median	Mean $\pm$ standard error (CI95%)	Log transformed mean	Median	Mean $\pm$ standard error (CI95%)	Log transformed mean	
Li	0.43	4.36 $\pm$ 2.18	-0.04	0.51	0.49 $\pm$ 0.08	-0.49	2.15
Be	0.01	0.09 $\pm$ 0.05	-1.11	0.02	0.025 $\pm$ 0.004	-1.80	1.79
V	0.83	3.05 $\pm$ 1.20	0.14	0.73	0.94 $\pm$ 0.21	-0.24	2.08
Cr	0.89	2.82 $\pm$ 1.17	0.08	2.24	3.28 $\pm$ 0.81	0.06	-0.34
Co	0.13	1.31 $\pm$ 0.65	-0.48	0.15	0.34 $\pm$ 0.11	-0.97	1.76
Ni	0.89	3.81 $\pm$ 1.72	0.13	1.26	1.90 $\pm$ 0.54	-0.05	1.23
Ga	0.37	0.97 $\pm$ 0.38	-0.32	0.81	1.00 $\pm$ 0.20	-0.12	-0.07
As	0.54	1.06 $\pm$ 0.39	-0.21	0.65	0.74 $\pm$ 0.13	-0.24	0.91
Rb	0.62	3.12 $\pm$ 1.42	-0.01	0.66	0.76 $\pm$ 0.10	-0.19	2.02
Cd	0.05	0.30 $\pm$ 0.23	-1.08	0.11	0.17 $\pm$ 0.04	-1.01	0.68
Cs	0.09	0.73 $\pm$ 0.40	-0.83	0.03	0.04 $\pm$ 0.01	-1.61	2.08
La	0.32	2.22 $\pm$ 1.08	-0.18	0.72	0.79 $\pm$ 0.14	-0.34	1.59
Pb	1.68	5.14 $\pm$ 2.19	0.37	1.96	3.20 $\pm$ 0.82	0.29	0.95
Hg	0.13*	0.34 $\pm$ 0.23*	0.29*	0.06*	0.06 $\pm$ 0.01*	-1.17*	<sup>L</sup>
Fe	602	3300 $\pm$ 1550	3.03	494	530 $\pm$ 97	2.54	2.17
Zn	65.9	90.6 $\pm$ 24.8	1.82	141	154 $\pm$ 16	2.15	-2.23
Cu	13.2	19.95 $\pm$ 4.63	1.19	15.2	18.45 $\pm$ 3.04	1.18	0.28
Ba	12.5	27.50 $\pm$ 8.85	1.24	26.3	26.50 $\pm$ 3.73	1.33	0.11

715 \*Longyearbyen (n=4), Hornsund (n=5), <sup>L</sup>- low sample size

716  
717  
718  
719  
720  
721  
722  
723

724 **Tab. 3 Nitrogen and carbon stable isotopes concentration in Svalbard reindeer hairs**

	Longyearbyen (n=10)		Hornsund (n=22)	
	$\delta^{15}\text{N}$ [‰]	$\delta^{13}\text{C}$ [‰]	$\delta^{15}\text{N}$ [‰]	$\delta^{13}\text{C}$ [‰]
Arythmetic Mean	<b>6.73</b>	<b>-26.19</b>	<b>10.96</b>	<b>-25.47</b>
SD	<b>1.40</b>	<b>0.24</b>	<b>2.01</b>	<b>0.76</b>
Median	<b>7.41</b>	<b>-26.22</b>	<b>10.66</b>	<b>-25.17</b>
Min	<b>3.73</b>	<b>-26.48</b>	<b>7.49</b>	<b>-26.67</b>
Max	<b>8.00</b>	<b>-25.82</b>	<b>14.04</b>	<b>-24.02</b>

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742 **Tab. 4 Pearson correlation values indicating correlation between the various trace elements**  
 743 **measured (n=26)**

Variable	15N	13C	Li	Be	V	Cr	Fe	Co	Ni	Cu	Zn	Ga	As	Rb	Cd	Cs	Ba	La	Pb
<b>15N</b>	1.00	-0.06	-0.16	-0.25	-0.21	-0.02	-0.26	-0.16	-0.03	-0.02	0.64	0.24	0.04	-0.10	0.20	-0.36	0.17	-0.04	0.01
<b>13C</b>	1.00	0.08	0.25	0.08	-0.10	0.09	0.07	-0.09	-0.10	-0.32	-0.19	-0.06	-0.07	-0.09	0.12	-0.10	-0.01	-0.07	
<b>Li</b>	1.00	0.95	0.97	0.77	0.97	0.77	0.89	0.79	0.44	0.79	0.87	0.92	0.75	0.94	0.81	0.94	0.94		
<b>Be</b>	1.00	0.87	0.74	0.86	0.52	0.68	0.59	0.49	0.71	0.79	0.83	0.57	0.90	0.78	0.86	0.56			
<b>V</b>	1.00	0.82	0.98	0.76	0.88	0.80	0.39	0.78	0.88	0.87	0.74	0.92	0.82	0.95	0.81				
<b>Cr</b>	1.00	0.80	0.48	0.80	0.71	0.52	0.76	0.74	0.65	0.76	0.70	0.82	0.88	0.64					
<b>Fe</b>	1.00	0.73	0.88	0.78	0.38	0.75	0.86	0.91	0.71	0.97	0.81	0.95	0.78						
<b>Co</b>	1.00	0.73	0.71	0.33	0.65	0.72	0.72	0.66	0.67	0.61	0.64	0.89							
<b>Ni</b>	1.00	0.85	0.55	0.80	0.80	0.80	0.85	0.79	0.81	0.89	0.88								
<b>Cu</b>	1.00	0.50	0.79	0.80	0.75	0.82	0.67	0.78	0.82	0.86									
<b>Zn</b>	1.00	0.74	0.60	0.53	0.57	0.27	0.67	0.55	0.52										
<b>Ga</b>	1.00	0.94	0.84	0.81	0.64	0.97	0.84	0.83											
<b>As</b>	1.00	0.91	0.74	0.78	0.94	0.87	0.82												
<b>Rb</b>	1.00	0.66	0.89	0.85	0.87	0.80													
<b>Cd</b>	1.00	0.63	0.83	0.79	0.82														
<b>Cs</b>	1.00	0.71	0.87	0.69															
<b>Ba</b>	1.00	0.88	0.79																
<b>La</b>	1.00	0.78																	
<b>Pb</b>	1.00																		

744

745

746

747

748

749

750

751

752

753

754

755 Tab.5 Outliers with significantly elevated element levels  $\mu\text{g/g dw}$  (Supplementary material)

No	$\delta^{15}\text{N}$ [‰]	$\delta^{13}\text{C}$ [‰]	Iron	Chromium	Cobalt	Barium	Nickel	Lead	Arsenic
1	5.88	-25.92	14640	11.1	6.31	69.1	15.5	20.6	4.22
2	7.56	-26.32	11450	8.27	4.36	91.7	13.9	18.0	2.61
3	7.41	-25.82	5810	6.34	2.47	49.4	6.79	7.35	1.61

756