

RESEARCH

Open Access



Bernstein-type theorem for ϕ -Laplacian

Jakub Maksymiuk^{1*}, Jakub Ciesielski¹ and Maciej Starostka¹

*Correspondence: jakub.maksymiuk@pg.edu.pl
¹Faculty of Applied Physics and Mathematics, Gdańsk University of Technology, Gdańsk, Poland

Abstract

In this paper we obtain a solution to the second-order boundary value problem of the form $\frac{d}{dt}\Phi'(\dot{u}) = f(t, u, \dot{u})$, $t \in [0, 1]$, $u: \mathbb{R} \rightarrow \mathbb{R}$ with Sturm–Liouville boundary conditions, where $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ is a strictly convex, differentiable function and $f: [0, 1] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and satisfies a suitable growth condition. Our result is based on a priori bounds for the solution and homotopical invariance of the Leray–Schauder degree.

Keywords: Φ -Laplacian; Boundary value problem; Fixed point; A priori bounds; Leray–Schauder degree

1 Introduction

In this paper we study the existence of solutions to the boundary value problems (BVPs)

$$\frac{d}{dt}\Phi'(\dot{u}) = f(t, u, \dot{u}), \quad t \in [0, 1] \tag{P}$$

$$-\alpha u(0) + \beta \dot{u}(0) = A, \quad au(1) + b\dot{u}(1) = B, \tag{BC}$$

where Φ' is an increasing homeomorphism, the scalar function f is continuous, $\alpha, a > 0$ and $\beta, b \geq 0$.

The solvability of various second-order two-point BVPs with p - or Φ -Laplacian has been discussed extensively in the literature, see the recent works [1–9] for results, methods, and references.

In 1912, Bernstein [10] proved that the BVP

$$u'' = f(t, u, u') \tag{1}$$

$$u(0) = A, \quad u(1) = B, \tag{2}$$

has a unique C^2 -solution if $f(t, u, v)$ is continuous, has continuous partial derivatives f_u and f_v on $[0, 1] \times \mathbb{R}^2$, there is a constant $K > 0$ such that

$$f_u(t, u, v) \geq K \quad \text{on } [0, 1] \times \mathbb{R}^2 \tag{3}$$

and

$$|f(t, u, v)| \leq A(t, u)v^2 + B(t, u) \quad \text{on } [0, 1] \times \mathbb{R}^2,$$

where A, B are functions bounded on each compact subset of $[0, 1] \times \mathbb{R}$.

In 1978, Granas et al. [11] proved similar results for (1) with either Dirichlet, Neumann, or periodic boundary conditions. The authors have established the existence of solutions to the considered problems by replacing (3) with the following assumption: There is a constant $M > 0$ such that

$$uf(t, u, 0) > 0 \quad \text{for } t \in [0, 1] \text{ and } |u| > M.$$

The uniqueness of the solution to (1.4), (BC) follows from the assumption that the partial derivatives f_u and f_v exist, are bounded, and $f_u \geq 0$ on $[0, 1] \times \mathbb{R}^2$.

In 1983, Baxley [12] proved Bernstein-type theorems for boundary value problems for (1) with nonlinear boundary conditions. In 1988, Frigon and O'Regan [13] established existence results of this type for (1), (2) and (1), (BC).

The aim of this paper is to give Bernstein-type existence theorems for BVPs with Φ -Laplacian. Throughout this paper we assume that $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ satisfies the following conditions:

- (Φ_1) Φ is strictly convex, differentiable and $\Phi(x)/|x| \rightarrow \infty$ as $|x| \rightarrow \infty$;
- (Φ_2) $\Phi(0) = \Phi'(0) = 0$;
- (Φ_3) $(\Phi')^{-1}$ is continuously differentiable;
- (Φ_4) there exists a constant $K_\Phi > 1$ such that

$$K_\Phi \Phi(x) \leq \Phi'(x)x \quad \text{for all } x \in \mathbb{R}.$$

Assumption (Φ_1) guarantees that Φ' is an increasing homeomorphism and so $(\Phi')^{-1}$ exists. Note that $\Phi(x) = \frac{1}{p_1}|x|^{p_1} + \dots + \frac{1}{p_n}|x|^{p_n}$, $1 < p_i \leq 2$, is in the considered class of functions, and if $n = 1$, then the differential operator on the left-hand side of the equation is a p -Laplacian. A more general form of Φ is provided by an N-function satisfying the ∇_2 -condition (see [14]).

We assume also that $f: [0, 1] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and satisfies the following:

- (f_1) There exists a constant $M > 0$ such that

$$xf(t, x, 0) > 0 \quad \text{for } |x| > M,$$

- (f_2) There exist positive functions S, T bounded on bounded sets such that

$$|f(t, x, v)| \leq S(t, x)(\Phi'(v) \cdot v - \Phi(v)) + T(t, x).$$

Now, we can state our main result.

Main Theorem *Suppose that Φ and f satisfy (Φ_1)–(Φ_4) and (f_1), (f_2), respectively. Then problem (P), (BC) has at least one solution in $C^2([0, 1], \mathbb{R})$.*

To establish the validity of the above result, we apply the Leray–Schauder degree theory on a suitable constructed map. To define its domain, we use a priori bounds.

To prove the existence, we use topological methods. This approach has already been used by many authors. In [11] and [13] the authors considered the case of a Laplace operator with various boundary conditions. Generalizations to the p -Laplacian and to the

operator defined by an arbitrary increasing homeomorphism were developed in [3] and [5], respectively. The main idea in the paper [11] was to use the topological transversality theorem. This is a fixed point type theorem (see [15]). We decided to use an approach via Leray–Schauder degree theory instead, since it is essentially equivalent but the degree theory is familiar to a broader audience.

However, in [3] and [5] authors subject the equation to very specific boundary conditions, namely $u(0) = A, u(1) = B$. In order to show the existence for general Sturm–Liouville conditions, more effort has to be put in as can be seen below.

2 Auxiliary results

Lemma 2.1 *Let X be a metric space, and let $G: X \times \mathbb{R} \rightarrow \mathbb{R}$ be continuous. Suppose that*

- (1) *for every $v \in X$, function $g_v: \mathbb{R} \rightarrow \mathbb{R}$, defined by $g_v(c) = G(v, c)$, is an increasing homeomorphism;*
- (2) *if $\{v_n\}$ is bounded and $b_n \rightarrow \pm\infty$, then $G(v_n, b_n) \rightarrow \pm\infty$.*

Then, for each fixed constant $C \in \mathbb{R}$, the function $c: X \rightarrow \mathbb{R}$ defined by $G(v, c(v)) = C$ is continuous.

Proof Suppose that function c is not continuous, i.e., there exist $\epsilon > 0$ and a sequence v_n converging to some v_0 such that $|c(v_n) - c(v_0)| > \epsilon$. By the definition of c , $G(v_n, c(v_n)) = C$. In particular, both v_n and $G(v_n, c(v_n))$ are bounded. This, together with (2), implies that $c(v_n)$ is bounded. Take a subsequence $c(v_{n_k})$ which converges to some c' . Note that $c' \neq c(v_0)$ because $|c(v_n) - c(v_0)| > \epsilon$. By the continuity of G , we have $G(v_{n_k}, c(v_{n_k})) \rightarrow G(v_0, c')$. But $G(v_{n_k}, c(v_{n_k})) = C$ and $G(v_0, c') \neq G(v_0, c(v_0)) = C$ by (1). A contradiction. \square

If g_v is differentiable and g'_v is positive, then the conclusion follows from implicit function theorem. However, in the problem that we consider, g'_v is only non-negative.

Remark 2.2 Note that this trivializes in [3, 5]. For boundary conditions considered therein c_1 and c_2 are constants independent of v . We cannot proceed in such a way here.

Now introduce the map $\hat{K}: C^0([0, 1]) \times \mathbb{R} \times \mathbb{R} \rightarrow C^1([0, 1])$ defined by

$$\hat{K}(v, c_1, c_2)(t) = c_1 + \int_0^t (\Phi')^{-1} \left(\int_0^\tau v(s) ds + c_2 \right) d\tau,$$

and by $C^1_{BC}([0, 1])$ denote the set of the functions in $C^1([0, 1])$ which satisfies (BC).

For every v , we would like to choose c_1 and c_2 in such a way that $u = \hat{K}(v, c_1, c_2)$ is an element of C^1_{BC} . Moreover, we need that c_1 and c_2 depend continuously on v .

Lemma 2.3 *Let (Φ_1) and (Φ_3) hold. Then, for every fixed $v \in C^0([0, 1])$, there exists a unique pair of constants $c_1(v), c_2(v)$ such that $\hat{K}(v, c_1(v), c_2(v)) \in C^1_{BC}([0, 1])$. Moreover, the functions $c_1, c_2: C^0([0, 1]) \rightarrow \mathbb{R}$ are continuous.*

Proof Put $u = K(v, c_1, c_2)$. Then

$$u(0) = c_1, \quad u(1) = c_1 + \int_0^1 (\Phi')^{-1} \left(\int_0^\tau v(s) ds + c_2 \right) d\tau$$

and

$$\dot{u}(0) = (\Phi')^{-1}(c_2), \quad \dot{u}(1) = (\Phi')^{-1}\left(\int_0^1 v(s) ds + c_2\right).$$

Clearly, u will satisfy the boundary conditions (BC) if c_1 and c_2 are such that $-\alpha c_1 + \beta(\Phi')^{-1}(c_2) = A$ and

$$a\left[c_1 + \int_0^t (\Phi')^{-1}\left(\int_0^\tau v(s) ds + c_2\right) d\tau\right] + b(\Phi')^{-1}\left(\int_0^1 v(s) ds + c_2\right) = B,$$

from where we get

$$\begin{aligned} & a\left[-\frac{A}{\alpha} + \frac{\beta}{\alpha}(\Phi')^{-1}(c_2) + \int_0^1 (\Phi')^{-1}\left(\int_0^\tau v(s) ds + c_2\right) d\tau\right] \\ & + b(\Phi')^{-1}\left(\int_0^1 v(s) ds + c_2\right) = B. \end{aligned} \tag{4}$$

Since $(\Phi')^{-1}$ is increasing, the function

$$\begin{aligned} G(v, c) = & a\left[-\frac{A}{\alpha} + \frac{\beta}{\alpha}(\Phi')^{-1}(c) + \int_0^1 (\Phi')^{-1}\left(\int_0^\tau v(s) ds + c\right) d\tau\right] + \\ & + b(\Phi')^{-1}\left(\int_0^1 v(s) ds + c\right) \end{aligned}$$

is increasing with respect to c . We can apply Lemma 2.1 for $C = B$ to conclude that (4) defines a unique constant c_2 depending continuously on v , and so c_1 is also unique and depends continuously on v . □

Now, for $\lambda \in [0, 1]$, consider the family of differential equations

$$\frac{d}{dt} \Phi'(\dot{u}) = \lambda f(t, u, \dot{u}), \quad t \in [0, 1]. \tag{P_\lambda}$$

Note that if u is a C^1 solution to problem (P_λ) , then $u \in C^2$. Indeed, \dot{u} reads

$$\dot{u}(t) = (\Phi')^{-1}\left(\int_0^t \lambda f(\tau, u, \dot{u}) d\tau + c\right),$$

and by assumption (Φ_3) and the continuity of f , it is continuously differentiable.

The next lemma is a variant of [13, Theorem 3.3].

Lemma 2.4 *Assume that (Φ_1) – (Φ_3) and (f_1) hold. Let $u \in C^1([0, 1])$ be a solution to (P_λ) , (BC) for $\lambda \in [0, 1]$. If $|u|$ achieves its maximum at $t_0 \in (0, 1)$, then*

$$|u(t)| \leq M \quad \text{for } t \in [0, 1].$$

Proof Suppose on the contrary that $|u|$ achieves its maximum at $t_0 \in (0, 1)$. We can assume that $u(t_0) > M$. In the case $u(t_0) \leq -M$ the proof is similar. It is clear that $\dot{u}(t_0) = 0$. For

$t \in [0, 1]$ we have

$$\int_{t_0}^t (t - \sigma)u(\sigma) \frac{d}{d\tau} \Phi'(\dot{u})(\tau) \Big|_{\tau=\sigma} d\sigma = t \int_{t_0}^t u \frac{d}{d\tau} \Phi'(\dot{u}) d\sigma - \int_{t_0}^t \sigma u \frac{d}{d\tau} \Phi'(\dot{u}) d\sigma.$$

Since $\Phi'(\dot{u}(t_0)) = \Phi'(0) = 0$,

$$\int_{t_0}^t u \frac{d}{d\tau} \Phi'(\dot{u}) d\sigma = u\Phi'(\dot{u}) \Big|_{t_0}^t - \int_{t_0}^t \dot{u}\Phi'(\dot{u}) d\sigma = u(t)\Phi'(\dot{u}(t)) - \int_{t_0}^t \dot{u}\Phi'(\dot{u}) d\sigma$$

and

$$\begin{aligned} \int_{t_0}^t \sigma u \frac{d}{d\tau} \Phi'(\dot{u}) d\sigma &= \sigma u\Phi'(\dot{u}) \Big|_{t_0}^t - \int_{t_0}^t (u + \sigma \dot{u})\Phi'(\dot{u}) d\sigma \\ &= tu(t)\Phi'(\dot{u}(t)) - \int_{t_0}^t u\Phi'(\dot{u}) d\sigma - \int_{t_0}^t \sigma \dot{u}\Phi'(\dot{u}) d\sigma. \end{aligned}$$

Combining the above, we get

$$\int_{t_0}^t (t - \sigma)u(\sigma) \frac{d}{d\tau} \Phi'(\dot{u})(\tau) \Big|_{\tau=\sigma} d\sigma = \int_{t_0}^t u\Phi'(\dot{u}) d\sigma + \int_{t_0}^t (\sigma - t)\dot{u}\Phi'(\dot{u}) d\sigma.$$

Hence, using (P_λ) , we have

$$\int_{t_0}^t (t - \sigma)(\lambda u(\sigma)f(\sigma, u(\sigma), \dot{u}(\sigma)) + \dot{u}(\sigma)\Phi'(\dot{u}(\sigma))) d\sigma = \int_{t_0}^t u(\sigma)\Phi'(\dot{u}(\sigma)) d\sigma.$$

Note that, for $0 < \lambda \leq 1$, $xf(t, x, 0) > 0$, $|x| > M$ implies $\lambda xf(t, x, 0) > 0$, $|x| > M$. Thus, by assumption (f_1) , $\lambda u(t_0)f(t_0, u(t_0), 0) > 0$. The continuity of f , u , and \dot{u} implies that there exists a neighborhood $N \subset (0, 1)$ of t_0 such that

$$\lambda u(t)f(t, u(t), \dot{u}(t)) > 0 \quad \text{for } t \in N.$$

Since $u \in C^1$ and achieves its maximum at t_0 , there exist t_0^- and t_0^+ such that

- $u(t) > M$ for $t \in (t_0^-, t_0^+)$,
- $\dot{u}(t) \geq 0$ on $(t_0^-, t_0]$,
- $\dot{u}(t) \leq 0$ on $[t_0, t_0^+)$.

Hence $\Phi'(\dot{u}(t)) \geq 0$ for $t \in (t_0^-, t_0]$ and $\Phi'(\dot{u}(t)) \leq 0$ for $t \in [t_0, t_0^+)$, since Φ' is increasing.

This implies that

$$\int_{t_0}^t (t - \sigma)\dot{u}(\sigma)\Phi'(\dot{u}(\sigma)) d\sigma \geq 0 \quad \text{for } t \in (t_0^-, t_0^+)$$

and

$$\int_{t_0}^t u(\sigma)\Phi'(\dot{u}(\sigma)) d\sigma \leq 0 \quad \text{for } t \in (t_0^-, t_0^+).$$

It follows that for t close to t_0

$$0 < \int_{t_0}^t (t - \sigma)(\lambda u(\sigma)f(\sigma, u(\sigma), \dot{u}(\sigma)) + \dot{u}(\sigma)\Phi'(\dot{u}(\sigma))) d\sigma = \int_{t_0}^t u(\sigma)\Phi'(\dot{u}(\sigma)) d\sigma \leq 0,$$

a contradiction. Thus $u(t_0) \leq M$. □

Lemma 2.5 Assume that (Φ_1) – (Φ_3) and (f_1) hold. Let $u \in C^1([0, 1])$ be a solution to (P_λ) , (BC) for $\lambda \in [0, 1]$. There exists a constant $M_0 > 0$ independent of λ and u such that

$$|u(t)| \leq M_0 \quad \text{for } t \in [0, 1].$$

Proof For $\lambda = 0$, problem (P_λ) has a unique linear solution, so there is a constant $C > 0$ such that $|u(t)| \leq C$ for $t \in [0, 1]$. Let $0 < \lambda \leq 1$. If $|u|$ achieves its maximum at $t = 0$, then $u(0)\dot{u}(0) \leq 0$. The boundary conditions give

$$u(0)(A + \alpha u(0)) = \beta u(0)\dot{u}(0) \leq 0,$$

and consequently $|u(0)| \leq |A/\alpha|$. Similarly, $|u(1)| \leq |B/a|$. If the maximum is at any $t_0 \in (0, 1)$, then by Lemma 2.4 we get $|u(t)| \leq M$. As a result, for $\lambda \in [0, 1]$, we have

$$|u(t)| \leq M_0 = \max\{M, |A/\alpha|, |B/a|\} \quad \text{for } t \in [0, 1]. \quad \square$$

Now we provide bounds for \dot{u} . The proof of the following theorem is based on [13].

Lemma 2.6 Assume that (Φ_1) – (Φ_4) , (f_1) , and (f_2) hold. Let $u \in C^1([0, 1])$ be a solution to (P_λ) , (BC) for $\lambda \in [0, 1]$. There exists a constant $M_1 > 0$, independent of λ and u , such that

$$|\dot{u}(t)| \leq M_1 \quad \text{for } t \in [0, 1].$$

Proof Since we have obtained a priori bounds $|u(t)| \leq M_0$, it is easy to observe that there exists a constant $C \geq 0$ independent of λ and u such that

$$|\dot{u}(t_0)| \leq C$$

for some $t_0 \in [0, 1]$. The point t_0 belongs to an interval $[\mu, \nu] \subset [0, 1]$ such that the sign of $\dot{u}(t)$ does not change in $[\mu, \nu]$ and $\dot{u}(\mu) = \dot{u}(t_0)$ and/or $\dot{u}(\nu) = \dot{u}(t_0)$.

Assume that $\dot{u}(\mu) = \dot{u}(t_0)$ and $\dot{u}(t) \geq 0$ for every $t \in [\mu, \nu]$. The other cases are treated similarly and the same bound is obtained.

Denote by S_0, T_0 the upper bounds of S and T , respectively, on $[0, 1] \times [-M_0, M_0]$. Since

$$|\lambda f(t, u, \dot{u})| \leq S_0(\Phi'(\dot{u})\dot{u} - \Phi(\dot{u})) + T_0,$$

we have

$$\int_\mu^t \frac{S_0 \dot{u} \left| \frac{d}{dt} \Phi'(\dot{u}) \right|}{S_0(\Phi'(\dot{u})\dot{u} - \Phi(\dot{u})) + T_0} dt \leq S_0 \int_\mu^t \dot{u} dt \leq 2S_0M_0.$$

For $\mu \leq \tau \leq t$, we have

$$\begin{aligned} & (\Phi'(\dot{u}(\tau))\dot{u}(\tau) - \Phi(\dot{u}(\tau))) - (\Phi'(\dot{u}(\mu))\dot{u}(\mu) - \Phi(\dot{u}(\mu))) \\ &= \int_\mu^t \frac{d}{dt} (\Phi'(\dot{u}(\tau))\dot{u}(\tau) - \Phi(\dot{u}(\tau))) \Big|_{t=\sigma} dt = \int_\mu^\tau \dot{u} \frac{d}{dt} \Phi'(\dot{u}) d\sigma. \end{aligned}$$

Note that since Φ is a convex differentiable function and $\Phi(0) = 0$, we have $\Phi(x) \leq \Phi'(x)x$ for every $x \in \mathbb{R}$. Thus, $0 \leq S_0(\Phi'(\dot{u}(\mu))\dot{u}(\mu) - \Phi(\dot{u}(\mu)))$. On the other hand, there exists $C_0 \geq 0$ such that $S_0(\Phi'(\dot{u}(\mu))\dot{u}(\mu) - \Phi(\dot{u}(\mu))) + T_0 \leq C_0$. Hence,

$$0 \leq S_0(\Phi'(\dot{u}(\tau))\dot{u}(\tau) - \Phi(\dot{u}(\tau))) + T_0 \leq S_0 \int_{\mu}^{\tau} \dot{u} \left| \frac{d}{dt} \Phi'(\dot{u}) \right| d\sigma + C_0 + T_0.$$

Set $g(\tau) = S_0 \int_{\mu}^{\tau} \dot{u} \left| \frac{d}{dt} \Phi'(\dot{u}) \right| d\sigma + C_0$, then integration by substitution yields

$$\log\left(\frac{g(t) + T_0}{C_0 + T_0}\right) = \int_{C_0}^{g(t)} \frac{1}{x + T_0} dx = \int_{\mu}^t \frac{S_0 \dot{u} \frac{d}{dt} \Phi'(\dot{u})}{g(\tau) + T_0} d\tau \leq 2S_0M_0.$$

Thus

$$g(t) \leq (T_0 + C_0)e^{2S_0M_0} - T_0$$

and by (Φ_4)

$$(k_{\Phi} - 1)\Phi(\dot{u}(t)) \leq \Phi'(\dot{u}(t))\dot{u}(t) - \Phi(\dot{u}(t)) \leq \frac{1}{S_0}((T_0 + C_0)e^{2S_0M_0} - T_0).$$

The last inequality gives $|\dot{u}(t)| \leq M_1$ for all $t \in [0, 1]$. □

3 Proof of the main theorem

Introduce the map $N: C^1_{BC}([0, 1]) \rightarrow C^0([0, 1])$ defined by

$$N(u)(t) = f(t, u, \dot{u}),$$

and for $\lambda \in [0, 1]$ consider the composition $\widehat{K} \circ \lambda N$, where the map $\widehat{K}: C([0, 1]) \rightarrow C^1([0, 1])$ is well defined by Lemma 2.3. Moreover, by the Arzela–Ascoli theorem, \widehat{K} is compact. Since N is continuous, the composition $\widehat{K} \circ \lambda N$ is also compact.

The fixed points of $\widehat{K} \circ \lambda N$ are of interest to us. Instead of looking for fixed points of $\widehat{K} \circ \lambda N$, one can look for zeros of $Id - \widehat{K} \circ \lambda N$. For this we will use the Leray–Schauder degree and its homotopical invariance. Consider the homotopy $H: [0, 1] \times C^1_{BC}([0, 1]) \rightarrow C^1_{BC}([0, 1])$ given by

$$H(\lambda, u) = (Id - \widehat{K}(\lambda N))(u).$$

Observe first that $H(0, l) = 0$, where $l = \widehat{K}(0)$ is unique. Thus, if $\overline{B}_r(0)$ is a closed ball with center 0 and radius r with the property $l \in B_r(0)$, then

$$\deg(H(0, u), B_r(0)) = 1.$$

It is well known that if r is such that

$$H(\lambda, u) \neq 0 \quad \text{for } \lambda \in [0, 1] \text{ and } u \in \partial B_r(0), \tag{5}$$

then also

$$\text{deg}(H(1, u), B_r(0)) = 1.$$

It is not hard to check that $u \in C^1_{BC}([0, 1])$ is a zero of $Id - \widehat{K} \circ \lambda N$ if and only if u is a solution to BVP (P_λ) , (BC). Thus, each zero $u \in C^1_{BC}([0, 1])$ of $Id - \widehat{K} \circ \lambda N$ satisfies the bound

$$\|u\|_{C^1([0,1])} < K,$$

where $K = \max\{M_0, M_1\} + 1$, where M_0 and M_1 are the constants from Lemmas 2.5 and 2.6. Clearly, (5) holds for $r = K$ and so, since in particular $l \in B_K(0)$, we have

$$\text{deg}(H(1, u), B_K(0)) = 1.$$

This means that $Id - \widehat{K} \circ \lambda N$ has at least one zero $u_0 \in B_K(0)$, which is a $C^1([0, 1])$ -solution to BVP of family (P_λ) arisen when $\lambda = 1$, that is, u_0 is a $C^1([0, 1])$ -solution to (P), (BC). However, as a solution of (P), u_0 is such that, for some constant c , we have

$$u'_0(t) = (\Phi')^{-1} \left(\int_0^t f(\tau, u_0(\tau), u'_0(\tau)) d\tau + c \right),$$

from where, keeping in mind (Φ_3) and the continuity of f , we get $u_0 \in C^2([0, 1])$.

4 Examples

Example 4.1 Let $\Phi : \mathbb{R} \rightarrow \mathbb{R}$, $\Phi(x) = \frac{1}{p}|x|^p$, $1 < p \leq 2$. It is easy to see that the function Φ satisfies assumptions (Φ_1) – (Φ_3) . Moreover, since $\frac{1}{p}|x|^p \leq x^2|x|^{p-2}$, one can take $K_\Phi = p$ in (Φ_4) .

Define the function $f : \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ by the formula

$$f(t, x, v) = \frac{(x^3 - x)(1 + |v|^{\frac{p+1}{2}})}{1 + t^2}.$$

One can easily check that

$$xf(t, x, 0) = \frac{x^4 - x^2}{1 + t^2} > 0$$

for $|x| > 1$. Since $\Phi'(v)v - \Phi(v) = \frac{1-p}{p}|v|^p$ and

$$|f(t, x, v)| \leq \frac{p|x^3 - x|}{(p-1)(1+t^2)} \cdot \frac{1-p}{p}|v|^p + \frac{|x^3 - x|}{1+t^2},$$

assumption (f_2) is satisfied with $S(t, x) = \frac{p|x^3 - x|}{(p-1)(1+t^2)}$ and $T(t, x) = \frac{|x^3 - x|}{1+t^2}$.

Assume that Φ satisfies assumptions (Φ_1) – (Φ_4) and that the functions $S, T : \mathbb{R} \times \mathbb{R} \rightarrow (0, \infty)$ are continuous and such that $xT(t, x) > 0$ for $|x| > M$. Then the function

$$f(t, x, v) = S(t, x)(\Phi'(v)v - \Phi(v)) + T(t, x)$$

satisfies our assumptions.

Example 4.2 Let $\Phi(x) = \sum_{i=1}^n \frac{1}{p_i} |x|^{p_i}$, $1 < p_i \leq 2$, for $i = 1, 2, \dots, n$. Then Φ satisfies assumptions (Φ_1) – (Φ_3) . Assumption (Φ_4) is satisfied with $K_\Phi = \min\{p_1, \dots, p_n\}$.

Example 4.3 Let $\Phi(x) = \frac{1}{p} |x|^p \log(1 + x^2)$. The function Φ satisfies all assumptions. In particular, as $\Phi'(x) = x|x|^{p-2} \log(1 + x^2) + \frac{1}{p} |x|^p \frac{2x}{1+x^2}$ and

$$p \frac{1}{p} |x|^p \log(1 + x^2) \leq x^2 |x|^{p-2} \log(1 + x^2) + \frac{1}{p} \frac{2x}{1 + x^2},$$

we can take $K_\Phi = p$. One can also consider functions of the form $\Phi(x) = \frac{1}{p} |x|^p \log^r(1 + |x|^s)$ for suitable choice of p, r, s .

Acknowledgements

The authors would like to thank Professor Andrzej Granas for suggesting this problem during the seminar in Gdańsk in 2016.

Funding

M. Starostka was partially supported by Grants Beethoven2 and Preludium9 of the National Science Centre, Poland, no. 2016/23/G/ST1/04081 and no. 2015/17/N/ST1/02527.

Abbreviations

Not applicable.

Availability of data and materials

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The three authors contributed equally in this paper. All authors read and approved the final manuscript.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 8 May 2018 Accepted: 21 December 2018 Published online: 07 January 2019

References

- Bereanu, C., Mawhin, J.: Nonhomogeneous boundary value problems for some nonlinear equations with singular ϕ -Laplacian. *J. Math. Anal. Appl.* **352**(1), 218–233 (2009)
- Cabada, A., Dimitrov, N.D.: Existence results for singular φ -Laplacian problems in presence of lower and upper solutions. *Anal. Appl.* **13**(2), 135–148 (2015)
- Kelevedjiev, P.S., Tersian, S.A.: The barrier strip technique for a boundary value problem with p -Laplacian. *Electron. J. Differ. Equ.* **2013**, 28 (2013)
- Kelevedjiev, P., Bojerikov, S.: On the solvability of a boundary value problem for p -Laplacian differential equations. *Electron. J. Qual. Theory Differ. Equ.* **2017**, 9 (2017)
- Ma, R.Z.L., Liu, R.: Existence results for nonlinear problems with ϕ -Laplacian. *Electron. J. Qual. Theory Differ. Equ.* **2015**, 22 (2015)
- Mawhin, J.: Homotopy and nonlinear boundary value problems involving singular ϕ -Laplacians. *J. Fixed Point Theory Appl.* **13**(1), 25–35 (2013)
- Rachůnková, I., Staněk, S., Tvrđý, M.: Chap. 7. Singularities and Laplacians in boundary value problems for nonlinear ordinary differential equations. In: *Handbook of Differential Equations: Ordinary Differential Equations*, vol. 3, pp. 607–723. North-Holland, Amsterdam (2006)
- Staněk, S.: Positive and dead-core solutions of two-point singular boundary value problems with φ -Laplacian. *Adv. Differ. Equ.* **2010**, 17 (2010)
- Wang, X., Liu, Q., Qian, D.: Existence and multiplicity results for some nonlinear problems with singular ϕ -Laplacian via a geometric approach. *Bound. Value Probl.* **2016**, 27 (2016)
- Bernstein, S.: Sur les équations du calcul des variations. *Ann. Sci. Éc. Norm. Supér. (3)* **29**, 431–485 (1912)
- Granás, A., Guenther, R., Lee, J.: On a theorem of S. Bernstein. *Pac. J. Math.* **74**(1), 67–82 (1978)
- Baxley, J.V.: Nonlinear second order boundary value problems: an existence-uniqueness theorem of S. N. Bernstein. In: *Ordinary Differential Equations and Operators*, to F. V. Atkinson, Proc. Symp., Dundee/Scotl. 1982. Lect. Notes Math., vol. 1032, pp. 9–16 (1983)
- Frigon, M., O'Regan, D.: On a generalization of a theorem of S. Bernstein. *Ann. Pol. Math.* **48**, 297–306 (1988)
- Krasnoselskiĭ, M.A., Rutickiĭ, J.B.: *Convex Functions and Orlicz Spaces*. Noordhoff, Groningen (1961)
- Granás, A., Dugundji, J.: *Fixed Point Theory*. Springer, New York (2010)

