

PAPER • OPEN ACCESS

## Understanding the formation of metastable furan dication in collisions with ions

To cite this article: E Erdmann *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 132002

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

## Understanding the formation of metastable furan dication in collisions with ions

E Erdmann<sup>1\*</sup>, S Indrajith<sup>2</sup>, J Chiarinelli<sup>3,4</sup>, S Díaz-Tendero<sup>5</sup>, N F Aguirre<sup>5,6</sup>, M Alcamí<sup>5</sup>,  
A Domaracka<sup>2</sup>, P Rousseau<sup>2</sup>, B A Huber<sup>2</sup>, P Bolognesi<sup>3</sup>, L Avaldi<sup>3</sup> and M Łabuda<sup>1</sup>

<sup>1</sup>Gdańsk University of Technology, Narutowicza 11/12, 80-233 Gdańsk, Poland

<sup>2</sup>Normandie Université, ENSICAEN, UNICAEN, CEA, CNRS, CIMAP, 14000 Caen, France

<sup>3</sup>CNR - Istituto di Struttura della Materia, Area della Ricerca di Roma 1, Monterotondo Scalo, Italy

<sup>4</sup>Dipartimento di Scienze, Università di Roma Tre, 00146 Roma, Italy

<sup>5</sup>Departamento de Química, Módulo 13, Universidad Autónoma de Madrid, 28049, Madrid, Spain

<sup>6</sup>Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

**Synopsis** This work relies on complementary theoretical and experimental studies of the processes induced by ion-furan collisions. Results of the Molecular Dynamics simulations and exploration of the energy profiles combined with coincidence mass spectrometry provide complete picture of the fragmentation of furan dication.


The main aim of this work is to investigate and understand the fragmentation processes of a furan dication induced by ion collisions with complementary theoretical methodology and experimental technique. The furan molecule has been chosen as a target due to its model structure that can be seen as an elementary analogue of deoxyribose, fundamental component of the DNA backbone chain. Depending on the charge state, the furan molecule is expected to dissociate in various ways [1].

Our theoretical approach consists of *ab initio* Molecular Dynamics describing the evolution of the system following deposition of the energy, and exploration of the Potential Energy Surface (PES) studied at high level of theory. Therefore, the dynamical and energetical considerations of the studied process are expected to create a complete picture of the fragmentation mechanism.

Results of the dynamics simulations lead to the observation of four different processes: isomerization, skeleton fragmentation, H<sup>+</sup>/H<sub>2</sub><sup>+</sup> loss and H/H<sub>2</sub> loss. Total occurrence of the most abundant channels varies depending on the energy deposited to the system (from 5 to 30 eV). More insight into three- and four-body fragmentation processes is obtained by investigation of the order of events occurring as the trajectory progresses. Therefore, average time scales of sequential events leading to the major channels are also estimated.

Calculated fragmentation pathways, presenting minima and transition states, indicate stability of intermediate structures leading to chan-

nels with many fragments. Investigation of the PES and MD suggests that fragmentation of the furan dication follows a dissociation path with one channel clearly distinguished as most probable and producing fragments of formyl cation (CHO<sup>+</sup>) and cyclopropenyl cation (C<sub>3</sub>H<sub>3</sub><sup>+</sup>).



**Figure 1.** Snapshots of the dynamical simulation.

In order to support theoretical results, coincidence time-of-flight (TOF) mass spectrometry has been employed to study the interaction of 46 keV O<sup>6+</sup> ions with neutral furan. Measured relative intensities of the coincident ion pairs are in good agreement with calculated energy barriers of the lowest-energy pathways. Geometrical parameters of the measured correlation islands in the fragment-fragment map indicate the mechanism of a specific dissociation channel. Long tail in the TOF mass spectrum points out to slow charge separation associated with the production of a metastable furan dication. This feature can be explained by the properties of the calculated PES.

**Acknowledgments:** This work has been done within the XLIC COST Action and CNRS-PICS BIFACE project.

### References

- [1] Erdmann E *et al* 2018 *J. Phys. Chem. A.* **122** 4153

\* E-mail: [ewa.erdmann@pg.edu.pl](mailto:ewa.erdmann@pg.edu.pl)

