

© 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Colloidal transport of lipid digesta i n human and porcine small intestinal mucus

Adam Macierzanka ^{a,b,c,*}, Olivia Ménard ^d, Didier Dupont ^d, Krzysztof Gutkowski ^e, Robert Staroń ^e,
Lukasz Krupa ^e

^a Gdańsk University of Technology, Faculty of Chemistry, Department of Colloid and Lipid Sciences, Gabriela Narutowicza 11/12, 80-322 Gdańsk, Poland

^b Riddet Institute, Massey University, Private Bag 11 222, Palmerston North 4442, New Zealand

^c Institute of Food Research, Norwich Research Park, Colney Lane, Norwich, NR4 7UA, United Kingdom.

^d STLO, INRAE, Institut Agro, 65 Rue de St. Briec, 35042 Rennes, France

^e Teaching Hospital No 1, Department of Gastroenterology and Hepatology with Internal Disease Unit, Chopina 2, 35-055 Rzeszów, Poland

*Corresponding author. E-mail address: adam.macierzanka@pg.edu.pl (A. Macierzanka).

Abbreviations:

CLSM, confocal laser scanning microscopy; D, diffusion coefficient; DG, diglyceride; FFA, free fatty acid; GI, gastrointestinal; MG, monoglyceride; TG, triglyceride.

34 **Abstract**

35

36 Small intestinal mucus transport of food-derived particulates has not been extensively studied, despite mucus
37 being a barrier nutrients need to cross before absorption. We used complex dispersions of digesta obtained
38 from simulated, dynamic gastrointestinal digestion of yogurt to examine the penetrability of human and
39 porcine mucus to the particles formed of lipolysis products. Quantitative, time-lapse confocal microscopy
40 revealed a sieve-like behaviour of the pig jejunal and ileal mucus. The digesta diffusivity decreased significantly
41 over the first 30 min of mucus penetration, and then remained constant at ca. $5 \times 10^{-12} \text{ m}^2 \text{ s}^{-1}$ (approx. 70%
42 decrease from initial values). A non-significantly different penetrability was recorded for the ileal mucus of
43 adult humans. The digesta diffusion rates in neonatal, jejunal mucus of 2 week old piglets were 5–8 times
44 higher than in the three different types of adult mucus. This is the first report that validates the mucus of fully-
45 grown pigs as a human-relevant substitute for mucus permeation studies of nutrients/bio-actives and/or
46 complex colloidal dispersions (e.g., post-digestion food particulates, orally-administrated delivery systems).

47

48

49

50 **Keywords**

51 Nutrient transport; Dynamic digestion model; Mucus barrier; Lipolysis; Proteolysis; Particle diffusion.

52

53 1. Introduction

54

55 The small intestine is the part of the alimentary tract where most of food digestion and nutrient
56 absorption take place. In order to get absorbed by the body, any nutrients released from food during digestion
57 in the intestinal lumen need to penetrate through the mucus layer that separates the underlying epithelium
58 from luminal contents (Johansson, Sjövall, & Hansson, 2013). The mucus layer is a selective barrier. It allows
59 the passage of molecules (e.g., water-soluble nutrients, bioactives, pharmaceuticals, etc.) and small particles
60 (e.g., lipid/bile salt mixed micelles, etc.), but prevents the epithelium from direct exposure to luminal
61 microorganisms, including pathogens (Cone, 2009). It also lubricates the mucosal epithelium, which is crucial in
62 protecting the tissue from abrasions that can be caused by the peristaltic movement of luminal contents.

63 The small intestinal mucus is a viscoelastic hydrogel made from a number of components, amongst
64 which the MUC2 mucin glycoprotein is the major gel-forming macromolecule responsible for the porous
65 structure and rheological properties of the mucus (Corfield, Carroll, Myerscough, & Probert, 2001), (Round et
66 al., 2012). Several studies have also pointed out at the importance of extracellular DNA in maintaining the
67 microstructural organisation and high viscosity of the mucus as well as its barrier properties with regard to
68 penetrability to particles and bacteria (Macierzanka et al., 2014), (Lock et al., 2020).

69 Despite the importance of small intestinal mucus in regulating the access of nutrients to the
70 epithelium, there is a limited number of studies looking at the penetrability of the mucus layer by digested
71 food. So far, most of the research in this area has either focused on monitoring the local diffusivity of
72 molecules (e.g., polyphenols, alginates) inside the mucus matrix (Gonzales et al., 2015), (Mackie et al., 2016), or
73 on characterising the mucus structure by a combination of qualitative and quantitative microscopy methods,
4 often with the use of synthetic particles simulating digesta. In the latter case, one of regularly used methods is
5 multiple-particle tracking (MPT), which relies on simultaneous real-time monitoring of the displacement of
6 hundreds or thousands of individual particles (S. K. Lai, Wang, Wirtz, & Hanes, 2009). Evaluation of the
7 diffusivity of identical-in-size nano- or micro-particles allows for probing the permeability, microrheology, and

78 heterogeneity in microstructural organisation of mucus (S. K. Lai, Wang, Hida, Cone, & Hanes, 2010),
79 (Macierzanka, Mackie, & Krupa, 2019), (Lock et al., 2020). This method has been used to determine the role of
80 the size and surface chemistry of particles on their transport in mucus (Yildiz, McKelvey, Marsac, & Carrier,
81 2015), (Maisel, Ensign, Reddy, Cone, & Hanes, 2015). For instance, it was shown that a high negative charge of
82 the surface of polystyrene beads, caused by adsorption of anionic intestinal surfactants – bile salts (BS),
83 significantly enhanced the ability of the beads to diffuse in the small intestinal mucus (Macierzanka et al.,
84 2011). It was suggested this was due to the BS adsorption preventing mucoadhesion of the beads to the
85 negatively-charged mucus matrix. MPT has also been used to assess the intestinal mucus diffusion of post-
86 digestion emulsion droplets (i.e., the droplets obtained after *in vitro* gastro-duodenal proteolysis of protein-
87 stabilised oil-in-water emulsions, (Macierzanka et al., 2011), (Macierzanka et al., 2012)). However, these
88 measurements were only conducted using size-narrowed monomodal fractions of droplets with well-defined
89 diameters. Whilst monitoring the diffusion of spherical particles with known, uniform sizes does allow for
90 calculating microrheological parameters of the mucus, the application of such particles can be of limited
91 physiological relevance, as gastrointestinal (GI) digestion of food emulsions usually yields a broad distribution
92 of particles in the gut lumen – from nano-size mixed micelles of bile salts and the lipolysis products to larger,
93 micro-size droplets of partially digested lipids (Armand et al., 1999). Under physiological conditions of the small
94 intestine, it is these complex dispersions of food-derived particles that interact simultaneously with the mucus
95 layer. Therefore, studies looking at the intestinal bioaccessibility and transport rates of nutrients should use
96 such physiologically relevant dispersions.

97 Another potential limitation of most scientific studies that have been conducted over the recent years
98 in this area is that they used animal mucus, usually obtained from mice, rats or pigs, to simulate a difficult-to-
9 obtain human mucus. Even for porcine mucus, its human-relevance cannot be assumed without scientific
0 validation, despite the human GI physiology being historically considered most similar to that of pigs (Zhang,
1 Widmer, & Tzipori, 2013). This might be especially true for studying interactions with digesta produced from

102 real foods, where colloidal complexity of such dispersions can highlight even small differences in
103 microstructural organisation and penetrability of the human mucus relative to mucus of animal origin.

104 This study had more than one goal. Firstly, we used digesta samples obtained from semi-liquid food
105 emulsion (natural yogurt was used), following different digestion times, to look at their ability to penetrate the
106 small intestinal mucus and whether the transport characteristics change in time. By investigating this, we also
107 intended to give an example of how *in vitro* digestion models can be complemented with simulating the
108 passive transport of nutrients/particles through the mucus layer under intestinal conditions. Secondly, we
109 wanted to compare the permeability of porcine and human small intestinal mucus to complex digesta
110 produced from real food. This aimed to examine the hypothesis of a similar barrier function of the porcine and
111 human mucus, and, if confirmed, would validate the use of porcine mucus for future investigations on the
112 mucus transport of post-digestion food particulates under simulated human small intestinal conditions.

113

114 **2. Materials and methods**

115

116 **2.1. Human small intestinal mucus**

117 The collection and studies of the human small intestinal mucus were approved by the ethics committee
118 of the Regional Medical Chamber in Rzeszów, Poland (certificate no. 4/B/2015). All methods were planned and
119 conducted in accordance with the ethical principles outlined in the Declaration of Helsinki. Aspiration of mucus
120 from the terminal ileum was done during diagnostic colonoscopy performed at the Teaching Hospital No 1 in
121 Rzeszów. The procedure duration was extended for a maximum of 5 min to obtain mucus samples. A typical
122 indication for colonoscopy was either occasional lower-GI bleeding, persistent unexplained abdominal pain, or
123 screening and surveillance of colorectal polyps, and 51 individuals (31 men and 20 women), aged 34–67 years,
124 were initially included in the study. Any evidence of inflammatory changes to the mucosa of the colon and/or
125 the terminal ileum confirmed during the procedure disqualified a subject from the study. All individuals who
126 agreed to take part in the study were clearly informed about the procedure and instructed by a clinician
127 regarding the bowel preparation. Informed consent was provided by all participants prior to examination.
128 Personal information of the volunteers was de-identified.

129 All participants included in the study followed a low residue diet protocol for 5 days prior to
130 colonoscopy in order to improve bowel preparation. Individuals listed for colonoscopy received bowel
131 preparation with 4L of Fortrans® (Ipsen Pharma) aqueous solution (1 sachet dissolved in 1L of water). One
132 sachet of Fortrans® powder contained Macrogol 4000 (64 g), anhydrous sodium sulphate (5.7 g), sodium
133 bicarbonate (1.68 g), sodium chloride (1.46 g), and potassium chloride (0.75 g). The timing of bowel
134 preparation prior to colonoscopy has a very significant impact on the preparation quality. It has been reported
5 that the optimal time for the colonoscopy procedure after completion of bowel preparation is 3-4 h, and
6 should be less than 8 h after completion of the preparation (Seo et al., 2012). This concept has led to a split-
7 dose preparation, which requires taking a portion of the bowel preparation solution the night prior to
8 colonoscopy (50% of the total dose) and the remaining portion on the day of colonoscopy (Bucci et al., 2014).

139 The split-dose preparation was used in this study. The participants received 2 sachets of Fortrans® diluted in 2L
140 of water between 6:00–8:00 pm on the day before colonoscopy and the remaining 2L of solution between
141 5:00–6:00 am on the day of the procedure.

142 As a routine part of colonoscopy examination, the intubation of the terminal ileum is performed (Meral
143 et al., 2018). This part of the small bowel contains a relatively thin layer of the mucus which is distributed on
144 the mucosal surface of the bowel wall. We only performed aspiration of the mucus samples from the terminal
145 ileum in participants who had bowel preparation rated as excellent (i.e., rate 9 of the Boston Scale (E. J. Lai,
146 Calderwood, Doros, Fix, & Jacobson, 2009)). Once the terminal ileum was intubated with the colonoscope, the
147 mucosal layer was inspected and an optimal area for aspiration of mucus chosen (i.e., an area free of residual
148 liquid, etc.). At the same time, an assistant endoscopist inserted a disposable plastic catheter through the
149 biopsy channel of the colonoscope. While inserting the catheter, air was being constantly insufflated with a
150 syringe attached to one end of the catheter in order to prevent incidental aspiration of any material that might
151 have resided in the biopsy channel and would potentially lead to the contamination of a sample. Once the
152 catheter was in the lumen of the terminal ileum, the principal endoscopist was in charge of the control of the
153 colonoscope tip and the catheter. The assistant endoscopist performed aspiration of the mucus by applying a
154 gentle suction with a syringe. Mucus was only aspirated into the tip of the catheter in order to limit disturbing
155 of the sampled mucus, and from the mucosal surface of the terminal ileum located between 5 cm and 25 cm
156 from the ileocecal valve. Typically, no more than 0.5 mL mucus was aspirated from one subject over the limited
157 time that was allowed for mucus collection. Immediately after aspiration, the plastic catheter was removed and
158 the sample gently transferred into 0.5 mL plastic test tubes. The tubes were sealed and instantly immersed in
159 liquid nitrogen for snap freezing. Samples were stored at -80 °C prior to further examination.

0 Mucus samples collected from five individuals (three women and two men; 51-56 years old) were used
1 in the experiments described below in order to narrow down the age range of adult humans.

2

3

2.2. Porcine small intestinal mucus

164 The collection of porcine mucus, including the *ex vivo* handling of intestinal tissue, was done as
165 described previously (Macierzanka et al., 2014), (Macierzanka, Mackie, et al., 2019). In brief, mucus was gently
166 removed with a soft-rubber scraper from the mucosal surface of freshly excised and cleaned small intestines of
167 (i) 6–8-month old pigs (this being referred to as the ‘adult pig mucus’ or the ‘mucus from fully-grown pigs’
168 throughout the paper) and (ii) 2 week old piglets (this being referred to as the ‘piglet mucus’). Pig mucus was
169 collected from the most proximal, 1-meter-long jejunal segment and/or the most distal, 1-metre-long ileal
170 segment. The last 5 cm of the ileum before the ileocecal valve was discarded. Piglet mucus was only collected
171 from the most proximal jejunum (0.5–0.7-meter-long segment). Aliquots of collected mucus were immediately
172 transferred to 0.5 mL plastic screw-cap tubes for snap freezing in liquid nitrogen, and then stored at -80 °C. The
173 collection was carried out within 20 min from animal slaughter. Mucus was incubated for 10 min at RT before
174 use. As reported previously (Macierzanka et al., 2011), (Macierzanka, Mackie, et al., 2019), the freezing, storing
175 and thawing cause no significant differences in the macro- and microrheological properties of small intestinal
176 mucus.

177

178 **2.3. Yogurt *in vitro* dynamic digestion and characterisation of digesta samples**

179 A commercially available, natural yogurt (pH 4.2) was used. The yogurt was obtained from cow's milk
180 and contained 3.5 wt% fat, 4.5 wt% protein and 6.6 wt% carbohydrates, according to the information provided
181 by the producer (Danone). The yogurt (120 g) was put through the dynamic *in vitro* gastrointestinal digestion
182 simulator DIDGI® (Fig 1A, (Ménard et al., 2014)), mimicking the adult human gastric and small intestinal
183 conditions. The digestion was done in triplicate, following the procedure described before (Ménard, Famelart,
184 et al., 2018), with some modifications. The specific gastrointestinal digestion parameters applied have been
5 summarised in Table 1. The flow of enzyme secretions, prepared with either simulated gastric fluid (SGF) or
6 simulated intestinal fluid (SIF; Table 1), as well as the emptying rates and pH, were continuously monitored
7 using the STORM® software of the DIDGI® system (INRAE, France). Gastric and intestinal emptying rates
8 followed an exponential equation as described by (Elashoff, Reedy, & Meyer, 1982). The gastric and intestinal

189 half-times ($t_{1/2}$) of 70 min and 160 min, respectively, and β coefficient of 2 and 1.6, respectively, were applied
190 according to the values reported by Minekus et al. (Minekus, Marteau, Havenaar, & Huisintveld, 1995) for
191 yoghurt digestion in adults. A gastric acidification curve was obtained after modelling data from adult humans
192 (Malagelada, Longstreth, Summerskill, & Go, 1976). Evolutions in the volumes of gastric and small intestinal
193 contents, as well as in pH values, have been shown in Supplementary Figs. S1 and S2. In the gastric
194 compartment of digestion, porcine pepsin (Sigma-Aldrich, P6887) was used at 2000 U/mL of gastric content
195 (Minekus et al., 2014), and lipase A12 from *Aspergillus niger* (Amano) at 18 U/mL of gastric content. The
196 intestinal conditions were set up as reported by Minekus et al. (Minekus et al., 1995), using pancreatin from
197 porcine pancreas (Sigma-Aldrich, P7545) and porcine bile extract (Sigma-Aldrich, B8631). Intestinal pH was
198 maintained at 6.6.

199 The gastrointestinal digestion was carried out continuously for 4.5h at 37°C. Samples of digesta were
200 taken periodically from the gastric compartment (after 0.25, 0.5, 1, and 2h from the beginning of the digestion
201 procedure) and the intestinal compartment (after 1, 2, 3 and 4h from the beginning of the procedure) and
202 analysed using the methods described below. The enzymatic activity in digesta samples, that were withdrawn
203 for subsequent analysis of the proteolysis progress, was immediately inhibited by using protease inhibitors,
204 pepstatin A (0.72 mM) for gastric samples and phenylmethanesulfonyl fluoride (0.37 mg/mL; Sigma-Aldrich,
205 93482) for intestinal samples. The extent of proteolysis in digesta samples was assessed by SDS-PAGE using the
206 procedure described elsewhere (Ménard, Famelart, et al., 2018), (Böttger et al., 2019). The intestinal digesta
207 samples that were used in the mucus penetration experiments (section 2.4) were additionally treated with
208 Orlistat (Sigma-Aldrich, O4139) at the final concentration of 200 μ M. Lipolysis in the digesta samples
209 withdrawn for the purpose of analysing changes in the lipid profiles was inhibited by the addition of 0.1 M 4-
0 bromophenylboronic acid solution in methanol (Sigma-Aldrich, B75956; 50 μ L of solution per 1 mL of digesta).
1 A direct extraction of lipids from digesta was performed after sampling, and was based on the Folch method
2 (Ménard, Bourlieu, et al., 2018). Thin layer chromatography (TLC) was used to assess the overall lipid profiles of
3 digesta samples, according to the method described previously (Ménard, Bourlieu, et al., 2018). The TLC



214 allowed for monitoring the disappearance of triglycerides (TGs) present in the yogurt and the appearance of
215 lipolysis products during digestion.

216 The particle size distribution of yogurt, digesta samples, as well as some dispersions of materials used
217 in the digestion model (i.e., insoluble particles in the bile extract and pancreatin preparations) was obtained
218 from dynamic light scattering (Mastersizer 2000, Malvern Instruments, Malvern, UK) using the method
219 described before (Ménard, Bourlieu, et al., 2018). Samples were measured either in their original state or after
220 treatment with 0.5 % (w/v) sodium dodecyl sulfate (SDS) to evaluate the size distribution of deflocculated
221 particles. The zeta-potential (ζ) of the gastrointestinal digesta samples was measured using the method
222 described previously (Macierzanka et al., 2011), (Macierzanka et al., 2012). The gastrointestinal digesta
223 samples were used in experiments looking at the diffusion of partially digested lipids through human and
224 porcine small intestinal mucus.

225

226 ***2.4. Diffusion of gastrointestinal digesta in human and porcine small intestinal mucus***

227 The ability of the gastrointestinal digesta (i.e., the digesta samples withdrawn from the intestinal
228 compartment, after the yogurt had been exposed to the gastric followed by the small intestinal digestion for 1,
229 2, 3 and 4 h, Fig. 1A,B) to penetrate into the mucus layer was assessed using time-lapse confocal laser scanning
230 microscopy (CLSM). The mucus was stained for mucins with WGA-Oregon Green (Invitrogen, W6748; 1 mg/mL
231 in PBS buffer (Sigma-Aldrich, P4417), pH 7.4, containing 2% (w/v) sodium azide). The mucus was gently mixed
232 with the dye stock at a high mucus to stock ratio, 99: 1 (v/v), in order to minimise the dilution of mucus. This
233 produced the final dye concentration of 10 $\mu\text{g}/\text{mL}$ and sodium azide concentration of 0.02 % (w/v) in mucus.
234 Separately, the digesta was stained for lipids with Nile Red (Sigma-Aldrich, 72485) at a final dye concentration
5 of 8 $\mu\text{g}/\text{mL}$. The mucus was placed in a custom-made optical cell developed for this study. The cell consisted of
6 a chamber (6×6×1 mm) for mucus and an adjacent channel (luminal space) for introducing digesta (Fig. 2A).
7 After the chamber was filled with mucus to 60-80% of its volume, the cell was covered with a coverslip, and the
8 mucus boundary localised under the microscope. The cell with mucus was incubated on the microscope stage

239 at 37 ± 0.1 °C for 3 min. Subsequently, the liquid digesta (preconditioned at RT) was gently introduced through
240 the channel inlet to the luminal space (Fig. 2A; ca. 100 μ L of digesta was required), and finally the channel
241 sealed to prevent water evaporation during the experiment. This part of the procedure was done in less than
242 60 s, which was sufficient to bring the temperature of the introduced digesta to ca. 37 °C. The introduction of
243 digesta was done after the cell with mucus had been securely positioned on the microscope stage, in order to
244 start monitoring the penetration of mucus with digesta immediately after the two components were brought
245 into contact with each other. The transport progress was recorded at 37 ± 0.1 °C with a Leica TCS SP upright
246 confocal microscope (Leica Microsystems (UK) Ltd, UK). The penetration was followed for up to 90 min, and
247 assessed using the procedure adopted from previous study (Mackie et al., 2016). Briefly, linear fluorescent
248 intensity profiles of the Nile Red stained lipids in digesta were generated from the time-lapse images (500 \times 500
249 μ m, 1024 \times 1024 pixel, 30 s time intervals) using Image-Pro Analyzer 7.0 software (Media Cybernetics Inc., Silver
250 Spring, MD, USA). The diffusion coefficient was calculated from the fluorescence profiles using the following
251 equation, $F(x,t) = a \times \text{erfc}(x/(4Dt)^{1/2})$, where F is the fluorescence of digesta stained for lipids described as a
252 function of distance from the starting boundary x and time t . Here, a is an arbitrary scalar, erfc is the
253 complimentary error function and D is the diffusion coefficient of the fluorescent lipids.

254 All experiments were performed for mucus samples obtained from 3–5 individual pigs, piglets or
255 humans, and at least three times (i.e., for three separate mucus specimens) for each individual mucus source.
256 Results are presented as the mean \pm SD and/or distributions of data from measurements, for each condition
257 used. Statistical comparisons between two groups were made using a Student's t -test, and three (or more)
258 groups were evaluated using 1-way ANOVA (significance level, $\alpha = 0.05$).

259

260 **3. Results and discussion**

261

262 **3.1. Yogurt *in vitro* digesta**

263 The first step of the study was to obtain digesta from real food. Natural yogurt was selected as it is an
264 example of a semi-liquid dairy product that is consumed worldwide and contains substantial quantities of all
265 macronutrients. It was not the aim of this study to investigate in detail the gastrointestinal digestion of yogurt,
266 but rather to produce *in vitro* a selection of liquid digesta samples that could be used to investigate the
267 transport of partially digested food particulate matter through small intestinal mucus of various origins. The
268 dynamic *in vitro* model of human gastrointestinal digestion was chosen to accurately mimic the digestive
269 conditions in adult humans and to produce digesta.

270 Consequently, we have only carried out qualitative analyses of the proteolysis progress using SDS-PAGE
271 (Fig. 1C). The protein bands corresponding to milk caseins showed a progressive and substantial reduction in
272 intensity, which implied a rapid hydrolysis of the caseins by pepsin. After 0.5h of gastric digestion, there were
273 no casein bands observed. The pepsinolysis of whey proteins, β -lactoglobulin and α -lactalbumin, seemed to be
274 largely completed after 2h of the gastric digestion. The *in vitro* gastrointestinal proteolysis of yogurt has been
275 studied before. Ménard et al. (Ménard, Famelart, et al., 2018) reported on relatively quick gastric degradation
276 of caseins in several isocaloric yogurts that differed in the original viscosity (0.3 – 2.2 Pa s) and milk protein
277 content (3.1 – 8.1 wt%). In that study, the amount of caseins in the yogurts was reduced to only 4 wt% of the
278 original amount after 2h of the simulated dynamic gastric digestion with pepsin. The amounts of residual β -
279 lactoglobulin and α -lactalbumin at the end of the gastric digestion were more dependent on the original
280 protein concentration and yogurt viscosity. However, for conventional yogurt (i.e., with a total protein content
1 of 3.1 wt%; viscosity, 1.3 Pa s), β -lactoglobulin and α -lactalbumin were reduced to only 3.1 wt% and 6.4 wt% of
2 the original amounts, respectively. The digestion of both caseins and whey proteins was finalised in the
3 intestinal compartment, and resulted in complete disappearance of the protein SDS-PAGE bands after 3h of the
4 dynamic gastrointestinal digestion. A similar, rapid proteolysis of caseins and whey proteins was observed

285 under static gastrointestinal conditions for yogurts prepared with different casein:whey protein ratios (4.5:1,
286 2.8:1 and 1.5:1) and 3.70 – 3.75 % total protein contents (Rioux & Turgeon, 2012). For all types of yogurt, the
287 vast majority of the original amounts of caseins and whey proteins was hydrolysed during the gastric phase of
288 digestion. Overall, the finding of the two studies described above are in good agreement with our results
289 shown in Fig. 1C.

290 In the present study, we have also looked at changes in the lipid profile of the milk fat fraction of
291 yogurt (Fig. 1D). Unsurprisingly, the TLC showed triglycerides (TGs) as a major lipid class in the yogurt before
292 digestion. They were hydrolysed to some extent to free fatty acids (FFAs) and partial glycerides – di- and
293 monoglycerides (DGs, MGs) – during gastric digestion, and after 2h TGs coexisted with the lipolysis products.
294 The complete conversion of remaining TGs seemed to largely take place during the first 1h of the intestinal
295 lipolysis with pancreatin (Fig. 1D). The intestinal phase of digestion is where chyme is exposed to highly
296 surface-active bile salts (Macierzanka, Torcello-Gómez, Jungnickel, & Maldonado-Valderrama, 2019). These
297 physiological surfactants aid in effective intestinal lipolysis by emulsifying ingested lipids, and by solubilising
298 and removing the lipolysis products from the oil–water interface. All these processes can lead to the reduction
299 in the size of original TG droplets and in the formation of mixed micelles, composed of the lipolysis products,
300 bile salts, as well as other polar lipids delivered with bile or ingested food (e.g., phospholipids). The resulting
301 gastrointestinal digesta is usually a complex mixture of dispersed/solubilised and surface-active lipids, including
302 FFAs, DGs, MGs, bile salts and other lipids (e.g., phospholipids, cholesterol) (Armand et al., 1999), (Fatouros,
303 Walrand, Bergenstahl, & Müllertz, 2009). In our study, these types of lipids were observed in digesta produced
304 after 1–4h of the gastrointestinal digestion (Fig. 1D).

305 The ζ -potential values did not differ significantly ($P>.05$) between the four gastrointestinal digesta
6 samples (i.e., GI-1h, -49.8 ± 1.1 mV; GI-2h, -48.6 ± 2.1 mV; GI-3h, -49.0 ± 2.3 mV; GI-4h, -49.2 ± 1.8 mV), and
7 were comparable to the results obtained for other post-digestion emulsions that had been digested in the
8 presence of bile salts (Macierzanka et al., 2011), (Macierzanka et al., 2012). It was also noticed that the original
9 size of lipid droplets in yogurt was reduced due to lipolysis and interfacial activity of bile salts (Fig. 1E). This

310 behaviour is consistent with the results of other studies (Bonnaire et al., 2008), (Armand et al., 1999). However,
311 it was difficult to compare the mean particle size between the gastrointestinal digesta and the original yogurt,
312 as the digesta samples revealed multimodal size distributions (Fig. 1E). Treating samples with SDS before
313 measurement did not change the size profiles (data not shown), suggesting the polydispersity was not the
314 effect of particle flocculation. The particle size distributions of the digesta samples seemed to be obscured to
315 some extent by a particulate matter introduced to the intestine digestion mix with the pancreatin and the bile
316 salt extract. These insoluble particles were likely to be cellular debris remaining in the preparations after
317 extraction of pancreatic enzymes and bile salts from animal materials by the producer (Sigma-Aldrich). When
318 measured separately, the pancreatin and the bile salt extract showed an existence of insoluble particles
319 ranging in size from ca. 1 μm to $>1000 \mu\text{m}$, with most particles being well over 20 μm in diameter (Fig. 1E).
320 Those particles contributed to the volume size distributions of the digesta samples withdrawn during digestion.
321 However, the majority were much larger than the particles of digested lipids and did not dominate the
322 distribution profiles of digesta samples (Fig. 1E). This meant they could not greatly contribute to the overall
323 number of particles, the major fraction of which was represented by considerably smaller lipid particles. The
324 negligible contribution of the insoluble particulate matter that derived from pancreatin and bile extract was
325 confirmed by converting the volume distributions of the four gastrointestinal digesta samples to number
326 distributions (Fig. 1F). The gastrointestinal digesta was further used in the major part of the study looking at
327 the effect of the type of small intestinal mucus on its penetrability to the post-digestion lipid particles.

329 **3.2. Diffusion of gastrointestinal digesta in the small intestinal mucus**

330 1 **3.2.1. Sieve-like barrier properties of mucus**

2 We have developed a simple optical cell that allowed for convenient and reproducible monitoring of
3 interactions between mucus and colloidal dispersions. The cell was used to bring the liquid digesta samples
4 into contact with the small intestinal mucus without compromising the integrity of the mucus gel. This way, the

335 penetration of the mucus layer by the lipid particles undergoing Brownian motion in the digesta could be
336 monitored immediately after the digesta and mucus were brought together (Fig. 2A,B), mimicking the passage
337 of dispersed/solubilised lipids from the small intestinal lumen into the mucus overlaying the intestinal
338 epithelium. Fluorescent intensity profiles of stained lipids were generated from the time-lapse CLSM images
339 (Fig. 2C) and used to assess the diffusion of lipid digesta in the mucus layer as explained in section 2.4.

340 By applying this approach, we have initially evaluated an influence of the digestion time on the
341 diffusion rate of digesta in mucus. The mucus scraped from the jejunal segments of the small intestine in adult
342 pigs was chosen for experiments as this type of *ex vivo* mucus had previously been shown to produce relatively
343 little individual variability regarding rheological properties (Macierzanka et al., 2011). It has also been shown,
344 by multiple-particle tracking of 500 nm latex beads, that porcine mucus collected *ex vivo* maintains the same
345 microstructural characteristics and permeability to particles as the native mucus overlaying the jejunal mucosa,
346 even if the *ex vivo* mucus was subjected to freezing and thawing after collection (Macierzanka, Mackie, et al.,
347 2019).

348 Figure 3A shows the comparison of diffusion coefficients of the digesta samples obtained after 1, 2, 3
349 and 4h of gastrointestinal digestion. For each mucus penetration time, there was a subtle inverse correlation
350 between the mean diffusivity value and the digestion time. However, the correlations were not statistically
351 significant ($P > .05$) within the same penetration times, even during the first several minutes after the start of
352 mucus penetration when the effect was most noticeable (Fig. 3A). This might have been due to the relatively
353 small differences in the size distribution profiles between the four time-point digesta samples collected during
354 the course of digestion (Fig. 1F). Much more profound was a progressive decrease in the apparent diffusion of
355 digesta over the time of mucus penetration (Fig. 3A). This trend was consistent for all four types of
6 gastrointestinal digesta. A significant drop in diffusion coefficient was observed during the first 30 min after the
7 initial penetration of mucus, and then the diffusivity remained fairly constant until the end of experiment. The
8 ratios of diffusion coefficients from 5 min to 90 min after the start of penetration into the mucus (D_{90}/D_5)
9 showed consistently for each digesta that there was a nearly 70% reduction of the initial diffusivity after 90 min

360 (Fig. 3B). An explanation for this phenomenon may lie in the way the small intestinal mucus is organised at the
361 microscale. The gel-forming MUC2 mucin oligomers are able to form a porous network in the mucus, which can
362 limit movement only to particles not restricted by the mesh size. In murine jejunal mucus, an average pore size
363 of 200–220 nm was found (Bajka, Rigby, Cross, Macierzanka, & Mackie, 2015). However, the reported sizes
364 ranged broadly, from 75 nm to approx. 500 nm. Furthermore, the transport of particles can be affected by the
365 spatial organisation of mucin microscale domains in the mucus. A hierarchical model of colloidal transport
366 through the small intestinal mucus was postulated by Round and co-workers (Round et al., 2012). It involves an
367 existence of individual lamellae of MUC2 mucin network rather than a continuous three-dimensional network
368 of the mucin polymer. According to the model, the weak interactions between the lamellae allow for the
369 passage of particles larger than the pores in lamellae as those particles are able to diffuse along transient
370 channels between lamellae. Particles smaller than the pore size can traverse directly through the lamellae as
371 long as they are not sterically trapped. More recently, Meldrum et al. (Meldrum et al., 2018) reported on a
372 similar mechanism of a network assembly of MUC2 mucin in which viscoelastic microscale domains of mucin
373 oligomers were proposed to form via hydrogen bonding and Ca^{2+} -mediated links. Individual domains were then
374 assembled in a yield stress, gel-like fluid. In the previous studies on multiple-particle tracking in mucus
375 (Macierzanka et al., 2011), (Macierzanka et al., 2014), (Bajka et al., 2015), a broad diffusivity distribution was
376 observed for non-mucoadhesive, monodisperse nano- and micro-beads that were used to probe the
377 microviscosity of the porcine small intestinal mucus. Those results confirmed a high degree of mucus
378 heterogeneity and might also support the postulated hierarchical organisation of the mucus microstructure.

379 Taking into account these previous findings as well as the results of the present study (Fig. 3), the small
380 intestinal mucus may be considered to act as a sieve to a polydisperse digesta through the network of channels
1 and pores of various sizes. The concept of size filtering has been previously introduced for different types of
2 mucus (Lieleg, Vladescu, & Ribbeck, 2010); it assumes that particles smaller than mucus mesh size are allowed
3 to pass while larger particles are rejected. The contrasting concept of interaction filtering is based on the
4 strength of interaction between particles and the mucus polymer network, so the weakly interacting particles

385 are not retained in the mucus and allowed to diffuse. In this study, the gastrointestinal digesta was produced in
386 the presence of bile extract containing bile salts, which must have been responsible for the high negative
387 charge of the digesta dispersions (ζ -potential ca. -49 mV, see section 3.1.). The adsorption of anionic bile salts
388 has been shown previously to hugely enhance the diffusivity of particles in small intestinal mucus by
389 strengthening their negative surface charge and thus preventing mucoadhesion (Macierzanka et al., 2011),
390 (Macierzanka et al., 2012). Therefore, the size filtering mechanism seems to be the most likely explanation in
391 the present study for the sieve-like behaviour of the mucus confronted with the yogurt digesta. It is plausible
392 that the largest lipid particles in digesta were immobilised within a relatively short time after penetrating into
393 the mucus layer, whereas smaller particles were able to diffuse deeper into the mucus. The former was
394 suggested by an increase in red fluorescence in the boundary region of the mucus over the fluorescence
395 recorded for digesta in the luminal space (Fig. 2C). Consequently, it was only the smallest particles that were
396 likely to penetrate through the surface region of mucus clogged with the lipid droplets larger than the mucus
397 mesh size (i.e., the size of channels and pores), and continued to diffuse freely in the mucus at a constant rate
398 until the end of the penetration experiment (90 min, Fig. 3A). This may suggest that, under physiological
399 conditions of the gut, lipolysis continues in the intestinal mucus after partially digested fat droplets leave the
400 intestinal lumen and get entangled in the mucus. The diffusion through the mucus and towards the underlying
401 epithelium would then only continue after they become gradually reduced to smaller droplets and/or particles
402 of assembled lipolysis products due to enzymatic hydrolysis and assistance of the surface-active bile salts.

403 A similar clogging of intestinal mucus was observed after it was exposed to the soluble dietary fibre
404 sodium alginate (Mackie et al., 2016). As a result, the apparent diffusivity of that high molecular weight
405 polymer was reduced in the function of mucus penetration time, despite the lack of interaction between the
6 mucin and the alginate. The local diffusivity of lipid digesta inside the small intestinal mucus matrix has been
7 studied before (Mackie, Rigby, Harvey, & Bajka, 2016). However, our present report is the first time the sieve-
8 like behaviour of mucus has been shown to affect the transport rates of a complex, polydisperse mixture of
9 post-digestion lipid particles during the penetration into the mucus layer from a simulated luminal space.

410

411 3.2.2. Penetration of human and porcine mucus layer

412 Having analysed the diffusion of various digesta samples in the pig jejunal mucus, we looked at how the
413 permeation of digesta may be affected by the type of small intestinal mucus. The penetrability was compared
414 between mucus samples obtained from the jejunum of adult pigs, and the ileum of adult pigs and humans. We
415 also included jejunal mucus from 2 week old piglets in the study. This aimed to investigate the following,
416 individual comparisons of mucus types: (i) age-specific (i.e., piglet vs. adult pig; both for jejunal mucus), (ii)
417 intestinal location-specific (i.e., jejunal mucus vs. ileal mucus; both for adult pig), and (iii) species-specific (i.e.,
418 adult pig vs. adult human; both for ileal mucus). The yogurt digesta produced after 2h of gastrointestinal
419 digestion (GI-2h) was used for analysing penetrability of different types of mucus. As for the initial experiments
420 shown in Figs. 2 and 3, mucus samples were exposed to the digesta at 37°C, and diffusion of fluorescently-
421 labelled lipids into the mucus monitored in the function of time (Fig. 4).

422 The fluorescence profiles revealed a striking difference in penetration rates between the two jejunal
423 types of mucus analysed (Fig. 4A). The piglet mucus was significantly more permeable to the diffusing digesta
424 than the pig mucus, and after 5 min from the start of penetration the respective mean diffusion coefficients
425 were $6.56 \times 10^{-11} \text{ m}^2\text{s}^{-1}$ and $1.29 \times 10^{-11} \text{ m}^2\text{s}^{-1}$ ($P < .05$, Fig. 4B). This indicates the piglet mucus presents a less
426 effective barrier to passively diffusing lipid particles. The microstructural organisation of the pig and the piglet
427 jejunal mucus has been compared previously, using direct tracking of the movement of 500 nm latex beads in
428 mucus (Macierzanka et al., 2014). The two mucus secretions used in that study were pre-treated with DNase in
429 order to reduce the impact of extracellular DNA on the viscosity experienced by the probe particles. The study
430 revealed a substantial difference in the diffusivity of beads, which was largely due to contrasting ways the
1 mucin matrix was organised between the two mucus types. In the mucus of fully-grown pigs, the mucin
2 produced a coherent porous network, whereas in the piglet mucus the network was more heterogeneous and
3 fragmented. It consisted of large aggregates of mucin polymer surrounded by regions with lower local
4 concentrations of the glycoprotein. The most rapid diffusion of the tracer particles was observed in those

435 regions. According to the same report, the piglet mucus also contained approx. 40% less extracellular DNA than
436 the pig mucus. The differences in both the mucin organisation and the DNA concentration were suggested to
437 be caused by disparities in the distribution of mucin-producing goblet cells and in the epithelial cell turnover
438 during neonatal development compared to adulthood. Because of this, the mean microviscosity experienced
439 by the latex beads diffusing in the piglet mucus was roughly half the value recorded for the pig counterpart
440 (Macierzanka et al., 2014). The impact of developmental age on differences in chemical composition, structural
441 properties and permeability to particles of small intestinal mucus was studied recently in rats (Lock et al.,
442 2020). The gut immaturity was suggested to be responsible for significantly lower concentrations of mucin and
443 DNA in the ileal mucus of 5 day old rat pups relative to 21 day old rats. Particle tracking experiments, using 200
444 nm PEG-, carboxyl- and amine-modified polystyrene particles, showed the mucus from early developmental
445 age rats (5 day old) was substantially more permeable to diffusing particle compared to 21 day old rats. The
446 mucus of the former age group was also significantly more penetrable to flagellated bacteria (*E. coli*),
447 suggesting immaturity may contribute to enhanced exposure of the intestinal epithelium to microbes.

448 Since the structural barrier properties of mucus appear to develop during infancy, it is not surprising
449 that in the present study we observed a higher diffusivity of digesta in the 2 week old piglet mucus than in the
450 adult pig mucus. The broad size distribution of lipid particles in the digesta (Fig. 1E) does not allow for using
451 them to probe the microviscosity of mucus, as the diffusion of an individual particle greatly depends on its
452 diameter (S. K. Lai et al., 2010). However, results of the time-lapse measurements with digesta produced from
453 a real dairy product may provide new insights into how the permeability of jejunal mucus to lipids can change
454 postneonatally. Because of the relatively fast penetration of digesta into the piglet mucus, the experiment was
455 terminated after 50 min from the initial penetration of this type of mucus (Fig. 4B). Although the diffusivity of
6 digesta was reduced progressively over the first 30 min in the piglet mucus, the difference between the
7 diffusion coefficients after 5 min and 30 min was not significant ($P>.05$). This suggests the channels and pores
8 in the mucus matrix were not substantially clogged with large lipid particles, and allowed fast transport. This
9 also supports the previous findings of a loose and easy-to-penetrate structure of neonatal mucus (Macierzanka



460 et al., 2014), (Lock et al., 2020). The diffusivity in piglet mucus remained unchanged for the remaining 20 min.
461 This is in contrast to what was observed for the pig jejunal mucus. There was a significant drop in the rate of
462 digesta diffusion measured between 5 min and 30 min ($P < .05$) from the start of mucus penetration. The
463 diffusion coefficient remained constant after 30 min and until the end of experiment. The difference in
464 penetrability of the two types of jejunal mucus was further emphasised by comparing the ratios of diffusion
465 coefficients from 5 min to 50 min (D_{50}/D_5 , Fig. 4C). For the piglet mucus, the ratio was almost twice as high as
466 for the pig mucus, meaning that the apparent diffusivity of digesta was reduced far more substantially by the
467 pig mucus.

468 The above suggest that the transmucus transport of lipids in a neonatal small intestine might be much
469 faster than in a mature gut. It can potentially involve the arrival of lipid droplets/particles with a large spectrum
470 of sizes in close proximity to the epithelium, which may enhance the rate of intestinal absorption of lipolysis
471 products that are eventually released from hydrolysed fat droplets. This, however, can be limited by low
472 activity/concentration of pancreatic lipase in the neonatal small intestine, as reported for the human pre-term
473 and full-term infants relative to adults (Bourlieu et al., 2014). Further studies would be required to assess the
474 impact of mucus structure on the rate of intestinal lipid absorption.

475 Comparing the permeability of mucus from different segments of the small intestine has not attracted
476 much scientific attention so far despite the fact that the mucosal architecture varies longitudinally as the villi
477 length decreases from the duodenum to the ileum, which may affect the structural properties of the mucus
478 layer. In our study, the sieve-like barrier behaviour, characteristic of the pig jejunal mucus, was also observed
479 for the ileal mucus obtained from fully-grown pigs. The time evolution in the fluorescence intensity profiles of
480 digesta, and thus in the diffusivity observed in the ileal mucus mirrored that recorded for the jejunal mucus
1 (Fig. 4A,B). The D_{50}/D_5 ratios for both types of pig mucus were very similar, and showed that there was an
2 almost 70% reduction in the apparent diffusion of digesta after 50 min of mucus penetration regardless of the
3 original, anatomical location of the mucus (Fig. 4C). The ratios did not change significantly after extending the
4 penetration to 90 min (Supplementary Fig. S3). The reduction of the diffusion rate in the function of

485 penetration time, over the initial 50 min, meant that for both types of mucus some lipid particles could
486 penetrate as deep as 100–150 μm into the mucus layer during the first 5–10 min, whereas the penetration
487 depth could only increase to ca. 300 μm after 50 min (Fig. 4A). The fact that the diffusion coefficients were very
488 similar for both types of mucus (Fig. 4B) suggests that *in vivo* the transmucosal transport times might depend
489 on variations in the thickness of the mucus layer between the jejunum and the ileum, rather than on
490 differences in mucus microstructure. However, the effect of varying thickness on transport times might be
491 difficult to account for. To our knowledge, there are no studies looking at how these variations might look in
492 humans. In rats, the ileal mucus layer was reported to be significantly thicker than the duodenal and jejunal
493 mucus gel layers ($480 \pm 47 \mu\text{m}$, $170 \pm 38 \mu\text{m}$ and $123 \pm 4 \mu\text{m}$, respectively, (Atuma, Strugala, Allen, & Holm,
494 2001)). It was also observed that the mean mucus thickness can be reduced approx. 7-fold in the rat ileum and
495 even >20-fold in the jejunum after feeding relative to the animals fasted for 48h (Szentkuti & Lorenz, 1995).
496 However, in the mouse, no substantial difference was found between the thickness of jejunal and ileal mucus
497 (Ermund, Schütte, Johansson, Gustafsson, & Hansson, 2013). There were no significant differences observed
498 for the mean pore size of mucus and the size distribution profiles between the two intestinal segments either
499 (Bajka et al., 2015). The latter mouse study supports our finding of the very comparable penetrability of jejunal
500 and ileal mucus to digesta (Fig. 4B), and implies a similar lack of difference in microstructural organisation of
501 mucus in proximal and distal regions of the pig small intestine.

502 Importantly, we also looked at how the human mucus was penetrated by the lipid particles of digesta.
503 The results showed a very similar behaviour of the human ileal mucus and the pig jejunal and ileal mucus (Fig.
504 4B). Analogically to the pig mucus, the diffusion coefficient of digesta decreased significantly ($P < .05$) over the
505 first 30 min after bringing the digesta into contact with human mucus, and remained largely unchanged after
6 that time at ca. $5 \times 10^{-12} \text{m}^2\text{s}^{-1}$. There was no significant difference in how the diffusivity evolved in time
7 between the human mucus and the pig mucus, which has been shown by comparing the actual values of
8 diffusion coefficient (Fig. 4B) as well as the ratios of diffusivities at various stages of mucus penetration (Fig. 4C,
9 Supplementary Fig. S3). All these results imply a very similar microstructural organisation of the adult human

510 and pig small intestinal mucus with regard to penetrability to polydisperse colloidal particles. These data bring
511 about the proof for the human-relevance of adult pig small intestinal mucus in terms of selective barrier
512 properties to post-digestion luminal lipids.

513

514 **4. Conclusions**

515

516 We compared *ex vivo* the penetrability of several types of small intestinal mucus to digesta obtained
517 after simulated dynamic gastrointestinal digestion of yogurt. The overall apparent diffusivity of digesta in adult
518 pig mucus decreased significantly during the first 30 min of penetration, suggesting a sieve-like response of the
519 mucus matrix to the post-digestion lipid particles varying in size. The rate of digesta diffusion in the mucus of 2
520 week old piglets was several times faster. This suggests that neonatal mucus is a less effective barrier to
521 passively diffusing particles relative to adult mucus.

522 Most importantly, we showed that the evolution in penetrability of the adult human small intestinal
523 mucus to diffusing lipids was analogous to the penetration profiles of the mucus obtained from adult pigs. To
524 our knowledge, it is the first time such analogies have been reported for the transport of gastrointestinal
525 digesta. This validates the use of small intestinal mucus from fully-grown pigs as a human-relevant substitute in
526 studies focusing on the transmucous transports of molecules and complex colloidal dispersions (e.g.,
527 nutrients/bioactives, digested foods, orally-administrated drug delivery systems) under simulated conditions of
528 the adult human small intestine. The close resemblance in penetrability to diffusing entities of the adult pig
529 mucus and the adult human mucus suggests that some analogy might also be expected between the examined
530 piglet mucus and the mucus of human infants, which was not studied. However, confirming any such similarity
1 would certainly require further studies that, for obvious ethical reasons, might be problematic.

2

3 **Acknowledgements**



534 The work was supported by the Polish National Agency for Academic Exchange NAWA through a
535 Bekker Programme grant (PPN/BEK/2018/1/00073). The work was co-funded by the BBSRC, UK (research grant
536 BBS/E/F/00044424). A.M., O.M. and D.D. are participants of the INFOGEST international network on Food
537 Digestion. The funding sources had no role in study design; in the collection, analysis and interpretation of
538 data; in preparation of the manuscript; or in the decision to submit the article for publication.

539

540 **Supplementary Material**

541 Additional characterisation of methods and data.

542

543 **Declarations of interest**

544 None.

545

546 **References**

547

- 548 Armand, M., Pasquier, B., André, M., Borel, P., Senft, M., Peyrot, J., ... Lairon, D. (1999). Digestion and
549 absorption of 2 fat emulsions with different droplet sizes in the human digestive tract. *American Journal*
550 *of Clinical Nutrition*, 70(6), 1096–1106. <https://doi.org/10.1093/ajcn/70.6.1096>
- 551 Atuma, C., Strugala, V., Allen, A., & Holm, L. (2001). The adherent gastrointestinal mucus gel layer: thickness
552 and physical state in vivo. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, 280(5),
553 G922–G929. <https://doi.org/10.1152/ajpgi.2001.280.5.G922>
- 554 Bajka, B. H., Rigby, N. M., Cross, K. L., Macierzanka, A., & Mackie, A. R. (2015). The influence of small intestinal
555 mucus structure on particle transport ex vivo. *Colloids and Surfaces B: Biointerfaces*, 135, 73–80.
556 <https://doi.org/10.1016/j.colsurfb.2015.07.038>
- 557 Bonnaire, L., Sandra, S., Helgason, T., Decker, E. A., Weiss, J., & McClements, D. J. (2008). Influence of lipid
558 physical state on the in vitro digestibility of emulsified lipids. *Journal of Agricultural and Food Chemistry*,
559 56(10), 3791–3797. <https://doi.org/10.1021/jf800159e>
- 560 Böttger, F., Dupont, D., Marcinkowska, D., Bajka, B., Mackie, A., & Macierzanka, A. (2019). Which casein in
561 sodium caseinate is most resistant to in vitro digestion? Effect of emulsification and enzymatic
562 structuring. *Food Hydrocolloids*, 88, 114–118. <https://doi.org/10.1016/j.foodhyd.2018.09.042>
- 563 Bourlieu, C., Ménard, O., Bouzerzour, K., Mandalari, G., Macierzanka, A., Mackie, A. R., & Dupont, D. (2014).
4 Specificity of Infant Digestive Conditions: Some Clues for Developing Relevant In Vitro Models. *Critical*
5 *Reviews in Food Science and Nutrition*, 54(11), 1427–1457.
6 <https://doi.org/10.1080/10408398.2011.640757>
- 7 Bucci, C., Rotondano, G., Hassan, C., Rea, M., Bianco, M. A., Cipolletta, L., ... Marmo, R. (2014). Optimal bowel
8 cleansing for colonoscopy: split the dose! A series of meta-analyses of controlled studies. *Gastrointestinal*
9 *Endoscopy*, 80(4), 566-576.e2. <https://doi.org/10.1016/j.gie.2014.05.320>
- 0 Cone, R. A. (2009). Barrier properties of mucus. *Advanced Drug Delivery Reviews*, 61(2), 75–85.
1 <https://doi.org/10.1016/j.addr.2008.09.008>

- 572 Corfield, A. P., Carroll, D., Myerscough, N., & Probert, C. S. (2001). Mucins in the gastrointestinal tract in health
573 and disease. *Frontiers in Bioscience : A Journal and Virtual Library*, 6(1), D1321-57.
574 <https://doi.org/10.2741/corfield>
- 575 Elashoff, J. D., Reedy, T. J., & Meyer, J. H. (1982). Analysis of Gastric Emptying Data. *Gastroenterology*, 83(6),
576 1306–1312. [https://doi.org/10.1016/S0016-5085\(82\)80145-5](https://doi.org/10.1016/S0016-5085(82)80145-5)
- 577 Ermund, A., Schütte, A., Johansson, M. E. V., Gustafsson, J. K., & Hansson, G. C. (2013). Studies of mucus in
578 mouse stomach, small intestine, and colon. I. Gastrointestinal mucus layers have different properties
579 depending on location as well as over the Peyer's patches. *American Journal of Physiology-*
580 *Gastrointestinal and Liver Physiology*, 305(5), G341–G347. <https://doi.org/10.1152/ajpgi.00046.2013>
- 581 Fatouros, D. G., Walrand, I., Bergenstahl, B., & Müllertz, A. (2009). Colloidal structures in media simulating
582 intestinal fed state conditions with and without lipolysis products. *Pharmaceutical Research*, 26(2), 361–
583 374. <https://doi.org/10.1007/s11095-008-9750-9>
- 584 Gonzales, G. B., Smagghe, G., Mackie, A., Grootaert, C., Bajka, B., Rigby, N., ... Van Camp, J. (2015). Use of
585 metabolomics and fluorescence recovery after photobleaching to study the bioavailability and intestinal
586 mucus diffusion of polyphenols from cauliflower waste. *Journal of Functional Foods*, 16, 403–413.
587 <https://doi.org/10.1016/j.jff.2015.04.031>
- 588 Johansson, M. E. V., Sjövall, H., & Hansson, G. C. (2013). The gastrointestinal mucus system in health and
589 disease. *Nature Reviews Gastroenterology and Hepatology*, 10(6), 352–361.
590 <https://doi.org/10.1038/nrgastro.2013.35>
- 591 Lai, E. J., Calderwood, A. H., Doros, G., Fix, O. K., & Jacobson, B. C. (2009). The Boston bowel preparation scale:
592 a valid and reliable instrument for colonoscopy-oriented research. *Gastrointestinal Endoscopy*, 69(3
593 SUPPL.), 620–625. <https://doi.org/10.1016/j.gie.2008.05.057>
- 594 Lai, S. K., Wang, Y.-Y., Wirtz, D., & Hanes, J. (2009). Micro- and macrorheology of mucus. *Advanced Drug*
595 *Delivery Reviews*, 61(2), 86–100. <https://doi.org/10.1016/j.addr.2008.09.012>
- 596 Lai, S. K., Wang, Y. Y., Hida, K., Cone, R., & Hanes, J. (2010). Nanoparticles reveal that human cervicovaginal
597 mucus is riddled with pores larger than viruses. *Proceedings of the National Academy of Sciences of the*
598 *United States of America*, 107(2), 598–603. <https://doi.org/10.1073/pnas.0911748107>
- 599 Lieleg, O., Vladescu, I., & Ribbeck, K. (2010). Characterization of particle translocation through mucin hydrogels.
600 *Biophysical Journal*, 98(9), 1782–1789. <https://doi.org/10.1016/j.bpj.2010.01.012>
- 601 Lock, J. Y., Carlson, T. L., Yu, Y., Lu, J., Claud, E. C., & Carrier, R. L. (2020). Impact of Developmental Age,
602 Necrotizing Enterocolitis Associated Stress, and Oral Therapeutic Intervention on Mucus Barrier
603 Properties. *Scientific Reports*, 10(1), 6692. <https://doi.org/10.1038/s41598-020-63593-5>
- 604 Macierzanka, A., Böttger, F., Rigby, N. M., Lille, M., Poutanen, K., Mills, E. N. C., & Mackie, A. R. (2012).
605 Enzymatically structured emulsions in simulated gastrointestinal environment: Impact on interfacial
606 proteolysis and diffusion in intestinal mucus. *Langmuir*, 28(50), 17349–17362.
607 <https://doi.org/10.1021/la302194q>
- 608 Macierzanka, A., Mackie, A. R., Bajka, B. H., Rigby, N. M., Nau, F., & Dupont, D. (2014). Transport of Particles in
609 Intestinal Mucus under Simulated Infant and Adult Physiological Conditions: Impact of Mucus Structure
610 and Extracellular DNA. *PLoS ONE*, 9(4), e95274. <https://doi.org/10.1371/journal.pone.0095274>
- 611 Macierzanka, A., Mackie, A. R., & Krupa, L. (2019). Permeability of the small intestinal mucus for physiologically
612 relevant studies: Impact of mucus location and ex vivo treatment. *Scientific Reports*, 9(1), 17516.
613 <https://doi.org/10.1038/s41598-019-53933-5>
- 4 Macierzanka, A., Rigby, N. M., Corfield, A. P., Wellner, N., Böttger, F., Mills, E. N. C., & Mackie, A. R. (2011).
5 Adsorption of bile salts to particles allows penetration of intestinal mucus. *Soft Matter*, 7(18), 8077–8084.
6 <https://doi.org/10.1039/c1sm05888f>
- 7 Macierzanka, A., Torcello-Gómez, A., Jungnickel, C., & Maldonado-Valderrama, J. (2019). Bile salts in digestion
8 and transport of lipids. *Advances in Colloid and Interface Science*, 274, 102045.
9 <https://doi.org/10.1016/j.cis.2019.102045>
- 0 Mackie, A. R., Macierzanka, A., Aarak, K., Rigby, N. M., Parker, R., Channel, G. A., ... Bajka, B. H. (2016). Sodium
1 alginate decreases the permeability of intestinal mucus. *Food Hydrocolloids*, 52, 749–755.



- 622 <https://doi.org/10.1016/j.foodhyd.2015.08.004>
- 623 Mackie, A., Rigby, N., Harvey, P., & Bajka, B. (2016). Increasing dietary oat fibre decreases the permeability of
624 intestinal mucus. *Journal of Functional Foods*, 26, 418–427. <https://doi.org/10.1016/j.jff.2016.08.018>
- 625 Maisel, K., Ensign, L., Reddy, M., Cone, R., & Hanes, J. (2015). Effect of surface chemistry on nanoparticle
626 interaction with gastrointestinal mucus and distribution in the gastrointestinal tract following oral and
627 rectal administration in the mouse. *Journal of Controlled Release*, 197, 48–57.
628 <https://doi.org/10.1016/j.jconrel.2014.10.026>
- 629 Malagelada, J.-R., Longstreth, G. F., Summerskill, W. H. J., & Go, V. L. W. (1976). Measurement of Gastric
630 Functions During Digestion of Ordinary Solid Meals in Man. *Gastroenterology*, 70(2), 203–210.
631 [https://doi.org/10.1016/S0016-5085\(76\)80010-8](https://doi.org/10.1016/S0016-5085(76)80010-8)
- 632 Meldrum, O. W., Yakubov, G. E., Bonilla, M. R., Deshmukh, O., McGuckin, M. A., & Gidley, M. J. (2018). Mucin
633 gel assembly is controlled by a collective action of non-mucin proteins, disulfide bridges, Ca²⁺-mediated
634 links, and hydrogen bonding. *Scientific Reports*, 8(1), 5802. <https://doi.org/10.1038/s41598-018-24223-3>
- 635 Ménard, O., Bourlieu, C., De Oliveira, S. C., Dellarosa, N., Laghi, L., Carrière, F., ... Deglaire, A. (2018). A first step
636 towards a consensus static in vitro model for simulating full-term infant digestion. *Food Chemistry*,
637 240(March 2017), 338–345. <https://doi.org/10.1016/j.foodchem.2017.07.145>
- 638 Ménard, O., Cattenoz, T., Guillemin, H., Souchon, I., Deglaire, A., Dupont, D., & Picque, D. (2014). Validation of a
639 new in vitro dynamic system to simulate infant digestion. *Food Chemistry*, 145, 1039–1045.
640 <https://doi.org/10.1016/j.foodchem.2013.09.036>
- 641 Ménard, O., Famelart, M.-H., Deglaire, A., Le Gouar, Y., Guérin, S., Malbert, C.-H., & Dupont, D. (2018). Gastric
642 Emptying and Dynamic In Vitro Digestion of Drinkable Yogurts: Effect of Viscosity and Composition.
643 *Nutrients*, 10(9), 1308. <https://doi.org/10.3390/nu10091308>
- 644 Meral, M., Bengi, G., Kayahan, H., Akarsu, M., Soytürk, M., Topalak, Ö., ... Sağol, Ö. (2018). Is ileocecal valve
645 intubation essential for routine colonoscopic examination? *European Journal of Gastroenterology &*
646 *Hepatology*, 30(4), 432–437. <https://doi.org/10.1097/MEG.0000000000001065>
- 647 Minekus, M., Alminger, M., Alvito, P., Ballance, S., Bohn, T., Bourlieu, C., ... Brodkorb, A. (2014). A standardised
648 static in vitro digestion method suitable for food-an international consensus. *Food and Function*, 5(6),
649 1113–1124. <https://doi.org/10.1039/c3fo60702j>
- 650 Minekus, M., Marteau, P., Havenaar, R., & Huisintveld, J. H. J. (1995). A multicompartmental dynamic
651 computer-controlled model simulating the stomach and small intestine. *Atla-Alternatives to Laboratory*
652 *Animals*, 23(2), 197–209.
- 653 Rioux, L. E., & Turgeon, S. L. (2012). The ratio of casein to whey protein impacts yogurt digestion in vitro. *Food*
654 *Digestion*, 3(1–3), 25–35. <https://doi.org/10.1007/s13228-012-0023-z>
- 655 Round, A. N., Rigby, N. M., Garcia De La Torre, A., Macierzanka, A., Mills, E. N. C., & Mackie, A. R. (2012).
656 Lamellar structures of MUC2-rich mucin: A potential role in governing the barrier and lubricating
657 functions of intestinal mucus. *Biomacromolecules*, 13(10), 3253–3261.
658 <https://doi.org/10.1021/bm301024x>
- 659 Seo, E. H., Kim, T. O., Park, M. J., Joo, H. R., Heo, N. Y., Park, J., ... Moon, Y. S. (2012). Optimal preparation-to-
660 colonoscopy interval in split-dose PEG bowel preparation determines satisfactory bowel preparation
661 quality: An observational prospective study. *Gastrointestinal Endoscopy*, 75(3), 583–590.
662 <https://doi.org/10.1016/j.gie.2011.09.029>
- 663 Szentkuti, L., & Lorenz, K. (1995). The thickness of the mucus layer in different segments of the rat intestine.
4 *Histochemical Journal*, 27(6), 466–472. <https://doi.org/10.1007/bf00173712>
- 5 Yildiz, H. M., McKelvey, C. A., Marsac, P. J., & Carrier, R. L. (2015). Size selectivity of intestinal mucus to diffusing
6 particulates is dependent on surface chemistry and exposure to lipids. *Journal of Drug Targeting*, 23(7–8),
7 768–774. <https://doi.org/10.3109/1061186X.2015.1086359>
- 8 Zhang, Q., Widmer, G., & Tzipori, S. (2013). A pig model of the human gastrointestinal tract. *Gut Microbes*, 4(3),
9 193–200. <https://doi.org/10.4161/gmic.23867>
- 0

671 **Tables**

672

673 Table 1. Gastrointestinal parameters of *in vitro* dynamic digestion of yogurt


674

Gastric conditions (37°C)		
Simulated Gastric Fluid (SGF) (stock solution adjusted at pH 6.5)	Na ⁺	100 mmol/L
	Ca ²⁺	1 mmol/L
Yogurt	Ingested amount	120 g
Gastric pH (acidification curve) (using 1M HCl)	pH = 1.68 + 3.82 ^(-t/42) (t, time after ingestion in min)	
SGF + pepsin (porcine)	Pepsin	2000 U/mL of gastric content
	Flow rate	0.75 mL/min from 0 to 10 min
	Flow rate	0.25 mL/min from 10 min to the end of the gastric phase
SGF + lipase (fungal)	Lipase	18 U/mL of gastric content
	Flow rate	0.75 mL/min from 0 to 10 min
	Flow rate	0.25 mL/min from 10 min to the end of the gastric phase
Gastric emptying (Elashoff fitting)	t _{1/2}	70 min
	β	2
Intestinal conditions (37°C)		
Simulated Intestinal Fluid (SIF) (stock solution adjusted at pH 6.2)	Na ⁺	100 mmol/L
	Ca ²⁺	1 mmol/L
Intestinal pH (using 1M NaHCO ₃)	pH	6.6
SIF + bile extract (porcine)	Bile	4% from 0 to 30 min
	Bile	2% from 30 min to the end
	Flow rate	0.5 mL/min from 0 min to the end
SIF + pancreatin (porcine)	Pancreatin	7%
	Flow rate	0.25 mL/min from 0 min to the end
Intestinal emptying (Elashoff fitting)	t _{1/2}	160 min
	β	1.6

675


676





681
682
683 **Fig. 1. *In vitro* dynamic digestion of yogurt and characterisation of digesta samples.** (A) The DIDGI® dynamic *in vitro* gastrointestinal (GI) digestion system comprising interconnected gastric and small intestinal compartments. (B) The small intestinal compartment with GI digesta produced after 4h of yogurt digestion. (C) SDS-PAGE, (D) TLC, and (E,F) particle size distributions ((E) volume distribution and (F) number distribution) of digestion time-point samples. Samples withdrawn from the gastric and the small intestinal compartments are marked with 'G' and 'GI', respectively, followed by the time from the beginning of the digestion experiment. The graph (E) also shows the size distribution profiles for particles in the bile extract and pancreatin preparations used in digestion. The 'Control' (in C-E) refers to the yogurt before digestion. Abbreviations: CNs, caseins; β-LG, β-lactoglobulin; α-LA, α-lactalbumin; TG, triglyceride; DG, diglyceride; MG, monoglyceride; PL, phospholipid; Ch, cholesterol; FFA, free fatty acid.

Downloaded from mostwiedzy.pl
681
682
683
684
685
686
687
688
689
690
1
2
3



694

695

696

697

698

699

700

701


702

703

704

705

Fig. 2. The time-lapse CLSM experimental set-up for tracking the extent of digesta penetration into the small intestinal mucus. (A) Schematic representation of a custom-made optical cell used for monitoring the boundary between mucus and digesta (i.e., the area enclosed by the dashed line). (B) Representative CLSM images showing a layer of pig jejunal mucus stained with WGA-Oregon Green for mucin (green channel) and positioned in the optical cell. The mucus layer was exposed for up to 90 min to digesta (red channel) produced from yogurt after 3h of dynamic *in vitro* gastrointestinal digestion (GI-3h). The red fluorescence shows progressive penetration of the mucus layer by the digesta stained for lipids with Nile Red. (C) Evolution in the red fluorescence intensity profiles of digesta in the function of the distance of penetration into the mucus. The profiles were generated at different times over a period of 90 min after the initial penetration of the mucus layer.



706

707

708

709

710

711

712

713

714

715

716

Fig. 3. Diffusion of yogurt digesta in the pig jejunal mucus. The effect of gastrointestinal digestion time on the diffusivity of digesta in the mucus. (A) Evolution of the diffusion coefficients of digesta samples as measured from the fluorescence intensity profiles of lipids in digesta over a period of 90 min after the initial penetration of the mucus layer. The digesta samples were obtained after either 1, 2, 3 or 4h of dynamic *in vitro* gastrointestinal digestion of yogurt (GI-1h, GI-2h, GI-3h or GI-4h), stained fluorescently for lipids, and exposed *ex vivo* to the mucus (Fig. 2). Bars marked with identical letters within the same type of digesta are not significantly different at $\alpha = 0.05$. (B) The ratio of diffusion coefficients of the digesta at 5 min and 90 min (D_{90}/D_5) after initial penetration of the mucus layer. All measurements were conducted at 37 ± 0.1 °C. Mean \pm SD ($n = 3$).




Fig. 4. Comparison of the penetration rates of gastrointestinal yogurt digesta into the human and porcine small intestinal mucus. The digesta was obtained after 2h of dynamic *in vitro* gastrointestinal digestion of yogurt (GI-2h) and exposed *ex vivo* to the mucus collected from the human ileum, the pig ileum, the pig jejunum, and the piglet jejunum. (A) Representative fluorescence intensity profiles (normal to the mucus boundary) of digesta stained for lipids in the function of the distance of penetration into the mucus. The fluorescence profiles were recorded at 1, 5, 10, 30 and 50 min after the initial penetration of the mucus layer. Each fluorescence graph is accompanied by CLSM images showing the mucus boundary (MB; mucus is shown in green channel) and the extent of digesta penetration (red channel) into mucus after 10 min. The scale bars correspond to 100 μ m. (B) Evolution of the diffusion coefficient of the yogurt digesta in the four mucus types over a period of up to 90 min after the initial penetration of mucus, as measured from the fluorescence intensity profiles of lipids in digesta. Bars marked with identical letters within the same type of mucus are not significantly different at $\alpha = 0.05$. (C) The ratio of diffusion coefficients of the digesta at 5 min and 50 min (D_{50}/D_5) after initial penetration of the mucus layer. All measurements were conducted at 37 ± 0.1 °C. Mean \pm SD ($n = 3-5$); * $P < .05$; NS, not significant ($P > .05$).