



# Stability by linear approximation for time scale dynamical systems



Sergey Kryzhevich<sup>a,b,c,\*</sup>, Alexander Nazarov<sup>a,d</sup>

<sup>a</sup> Faculty of Mathematics and Mechanics, Saint-Petersburg State University, 28, Universitetsky pr., Peterhof, 198503, Saint-Petersburg, Russia

<sup>b</sup> School of Natural Sciences and Mathematics, The University of Texas at Dallas, 800 W. Campbell Road, 75080-3021, Richardson, TX, USA

<sup>c</sup> University of Nova Gorica, Vipavska, 13, SI-5000, Nova Gorica, Slovenia

<sup>d</sup> Saint-Petersburg Department of Steklov Institute, Fontanka, 27, 191203, Saint-Petersburg, Russia

## ARTICLE INFO

### Article history:

Received 11 December 2015

Available online 11 January 2017

Submitted by M.J. Schlosser

### Keywords:

Time scale system

Linearization

Lyapunov functions

Millionschikov rotations

Stability

## ABSTRACT

We study systems on time scales that are generalizations of classical differential or difference equations and appear in numerical methods. In this paper we consider linear systems and their small nonlinear perturbations. In terms of time scales and of eigenvalues of matrices we formulate conditions, sufficient for stability by linear approximation. For non-periodic time scales we use techniques of central upper Lyapunov exponents (a common tool of the theory of linear ODEs) to study stability of solutions. Also, time scale versions of the famous Chetaev's theorem on conditional instability are proved. In a nutshell, we have developed a completely new technique in order to demonstrate that methods of non-autonomous linear ODE theory may work for time-scale dynamics.

© 2017 Elsevier Inc. All rights reserved.

## 1. Introduction

We study dynamic equations on time scales i.e. on unbounded closed subsets of  $\mathbb{R}$ . The time scale approach first introduced by S. Hilger and his collaborators (see [1] and references therein) was intensively developing during last decades. The first advantage of such approach is the common language that fits both for flows and diffeomorphisms. On the other hand, there are many numerical methods that correspond to non-uniform steps. Especially, this is applicable for modeling non-smooth or strongly non-linear dynamical systems.

Consider a motion of a particle in two distinct media, e.g. water and air. Evidently, to model such system, it is not effective to use equidistant nodes. It is better to take more of them inside time periods, corresponding

\* Corresponding author at: University of Nova Gorica, Slovenia.

E-mail addresses: kryzhevicz@gmail.com, s.kryzhevich@spbu.ru (S. Kryzhevich).

to motions in water. This is a natural way to obtain a non-trivial time scale in a real life problem (see [29] and references therein). Another application of time scale analysis may be found for systems with delay [5].

In this sense it seems to be useful to generalize some results on stability theory well-known for ODEs for time scale case. Mainly, we consider a general linear system (autonomous or non-autonomous) and its small non-linear perturbation. For the continuous dynamics, there exists a well-developed theory on stability by first approximation. For autonomous case there are classical stability criteria related to eigenvalues of a matrix of coefficients for linear approximation (call it  $\mathcal{A}$ ).

For non-autonomous systems or for cases when eigenvalues do not give information on stability of the perturbed system, there might be two approaches. The first one is based on the theory of Lyapunov functions (see [21] and references therein for review on the time scale version of this method). The second one involves integral inequalities, particularly the Grönwall–Bellman inequality (see [4,6]). For ordinary differential equations there is a very powerful tool that allows to find stability of solutions via the so-called central upper Lyapunov exponents [10]. In this paper we combine all referred methods in order to study time scale dynamics.

Exponential stability for solutions of time-varying dynamic equations on a time scale have been investigated by many authors. We mention recent papers by Bohner and Martynyuk [7] (this article is also a good introduction to the theory of time scale systems), Du and Tien [15], Hoffacker and Tisdell [18] and Martynyuk [24]. We also refer to papers [16,19,20,23,26] where related problems are studied and new approaches have been introduced.

A “multidimensional” analog of time scales called discrete differential geometry is also studied, see [3] and references therein. In such problems, time scales may appear, for instance, as discretizations of geodesic flows.

However, the following problems were open by now.

1. For constant matrices  $\mathcal{A}$ , are there any stability criteria for perturbed time-scale systems?
2. Is there any analog of Chetaev’s theorem on instability by the first approximation for time scale systems?
3. Are there any sufficient conditions on stability by the first approximation, close to necessary ones?

One of principal difficulties in the theory of time scale systems is that, generally speaking, in “autonomous” case (i.e. when the right hand side of the system does not depend on  $t$ ) the system does not define a flow (a shift of a solution is not necessarily a solution, group property may be violated etc). Also, one must carefully check basic properties like smoothness of solutions that can be violated even for systems with smooth right hand sides.

In our paper we give positive answers to all mentioned questions. The main idea of our paper is very simple: methods of classical theory of linear non-autonomous differential equations are applicable for time scale systems. Here we notice that in time scale analysis there are two types of derivatives: the so-called  $\Delta$ - and  $\nabla$ -derivatives (see [7] for details). In this paper we study  $\Delta$ -derivatives only. However, it seems that the main ideas of our work can be easily transferred to equations with  $\nabla$ -derivatives.

We have two principal objectives.

First, we provide sufficient conditions on stability by the first approximation. We demonstrate that the obtained conditions are close to necessary ones. In our proofs, we use the techniques of central upper Lyapunov exponents. This approach seems to be novel for time scale analysis. We can use many related tools such as Millionschikov’s rotations to obtain instability.

Secondly, we prove an analog of Chetaev’s theorem on instability by first approximation. Specifics of time scales demands a novel, non-classical approach to proof since, generally speaking, we cannot use tools of the theory of autonomous systems, anymore.

The paper is organized as follows. In Section 2 we give a brief introduction to time scale analysis. In Section 3 we give a review of existing results on stability of time scale equations. In these two sections




Fig. 1. A time scale.

we mostly refer to results of [7], changing some notations. In Section 4 we introduce one of the main “characters” of our paper: central upper exponent. Using this concept and the Grönwall–Bellman lemma, we study stability of solutions of time scale dynamical systems. In Section 5 we demonstrate that a linear time scale system may be “embedded” into the ODE system. In this connection we introduce so-called syndetic time scales i.e. time scales that do not have arbitrarily big gaps. We demonstrate that there is a correspondence between linear time scale systems and linear systems of ordinary differential equations. We develop new technical tools that play a key role in proofs of following sections. In Section 6 we provide a time scale generalization of the classical Millionschikov’s result on attainability of the central upper exponent. The proof of Millionschikov’s result is given in Appendix. As a corollary, we deduce a time scale version of the condition on instability by the first approximation [22]. The Lyapunov approach is developed in Section 7. Similarly to what happens in ODEs theory, we construct Lyapunov functions for time scale systems as quadratic forms and thus relate stability by the first approximation with certain estimates on eigenvalues of the matrix of coefficients. We also give a time scale version of Chetaev’s instability theorem and a condition on instability by the first approximation.

## 2. Time scale analysis

We use following notation:  $B(r, x)$  is the Euclidean ball in  $\mathbb{R}^n$  with radius  $r$ , centered at  $x$ ;  $B_r = B(r, 0)$ ,  $\mathbf{M}_n$  is the space of  $n \times n$  complex matrices,  $|\cdot|$  stands for a vector norm in  $\mathbb{R}^n$  or  $\mathbb{C}^n$  and the corresponding operator norm.  $E_n$  is  $n \times n$  identity matrix.

**Definition 2.1.** A *time scale*  $\mathbb{T}$  is an unbounded closed subset of  $\mathbb{R}$  with the inherited metric. Without loss of generality we assume that  $0 \in \mathbb{T}$ .

We consider two spaces of matrix functions:  $\mathcal{M}_R$  that is a space of continuous functions  $\mathcal{A} : \mathbb{R} \rightarrow \mathbf{M}_n$  and  $\mathcal{M}_T$  that is the space of similarly defined functions  $\mathcal{A} : \mathbb{T} \rightarrow \mathbf{M}_n$ .

We set  $\mathbb{T}_a^+ = [a, \infty) \cap \mathbb{T}$ .

We introduce basic notions connected to the theory of time scales, which summarize the material from the recent book by Bohner and Peterson [8] (see also [9,17]).

**Definition 2.2.** Let  $t \in \mathbb{T}$ . We define the *forward jump operator*  $\sigma : \mathbb{T} \rightarrow \mathbb{T}$  by  $\sigma(t) := \inf\{s \in \mathbb{T}; s > t\}$ .

If  $\sigma(t) > t$ , we say that  $t$  is right-scattered, while if  $\sigma(t) = t$ , then  $t$  is called right-dense. Denote by  $\mathbb{S}$  the set of right-scattered points and by  $\mathbb{D}$  the set of right-dense points. Evidently,  $\mathbb{T} = \mathbb{S} \cup \mathbb{D}$  is a disjoint union. We always assume that  $\sup \mathbb{S} = +\infty$ .

The *graininess function*  $\mu : \mathbb{T} \rightarrow [0, \infty)$  is defined by  $\mu(t) := \sigma(t) - t$  (Fig. 1).

**Definition 2.3.** A function  $f : \mathbb{T} \rightarrow \mathbb{R}$  is called *rd-continuous* provided it is continuous at right-dense points in  $\mathbb{T}$  and finite left-sided limits exist at left-dense points in  $\mathbb{T}$ . Denote the class of rd-continuous functions by  $\mathcal{C}_{rd} = \mathcal{C}_{rd}(\mathbb{T}, \mathbb{R})$ . We use the similar notation for vector and matrix functions.

**Definition 2.4.** The function  $f : \mathbb{T} \rightarrow \mathbb{R}$  is called  $\Delta$ -*differentiable* at a point  $t \in \mathbb{T}$  if there exists  $\gamma \in \mathbb{R}$  such that for any  $\varepsilon > 0$  there exists a neighborhood  $W$  of  $t$  satisfying

$$|[f(\sigma(t)) - f(s)] - \gamma[\sigma(t) - s]| \leq \varepsilon|\sigma(t) - s|$$

for all  $s \in W$ . In this case we write  $f^\Delta(t) = \gamma$ . We use the similar notation for vector and matrix functions.

When  $\mathbb{T} = \mathbb{R}$ ,  $x^\Delta(t) = \dot{x}(t)$ . When  $\mathbb{T} = \mathbb{Z}$ ,  $x^\Delta(n)$  is the standard forward difference operator  $x(n + 1) - x(n)$ .

**Definition 2.5.** If  $F^\Delta(t) = f(t)$ ,  $t \in \mathbb{T}$ , then  $F$  is a  $\Delta$ -antiderivative of  $f$ , and the Cauchy  $\Delta$ -integral is given by

$$\int_\tau^s f(t)\Delta t = F(s) - F(\tau), \quad \text{for all } s, \tau \in \mathbb{T}.$$

**Definition 2.6.** A function  $p : \mathbb{T} \rightarrow \mathbb{R}$  is called *regressive* provided that  $1 + \mu(t)p(t) \neq 0$  for all  $t \in \mathbb{T}$  and *positively regressive* if  $1 + \mu(t)p(t) > 0$  for all  $t \in \mathbb{T}$ . The set of all regressive and rd-continuous functions is denoted by  $\mathcal{R} = \mathcal{R}(\mathbb{T}, \mathbb{R})$ . The set of all positively regressive and rd-continuous function is denoted by  $\mathcal{R}^+$ .

A matrix mapping  $\mathcal{A} : \mathbb{T} \rightarrow \mathbf{M}_n(\mathbb{R})$  is called *regressive* if for each  $t \in \mathbb{T}$  the  $n \times n$  matrix  $E_n + \mu(t)\mathcal{A}(t)$  is invertible, and *uniformly regressive* if in addition the matrix function  $(E_n + \mu(t)\mathcal{A}(t))^{-1}$  is bounded.

If  $\mathcal{A}$  is constant, it is uniformly regressive if and only if

$$\inf_{\mathbb{T}} |\lambda_k \mu(t) + 1| > 0, \quad k = 1, \dots, n \tag{2.1}$$

where  $\lambda_k$  are the eigenvalues of  $\mathcal{A}$ . Note that in this case solutions of the system

$$x^\Delta = \mathcal{A}x \tag{2.2}$$

are unique and have finite Lyapunov exponents.

To prove this statement, it suffices to reduce (2.2) to the normal form and thus reduce it to a set of linear first order equations.

**Definition 2.7.** For  $p \in \mathcal{R}$ , we define the *generalized exponential function*  $e_p(t, s)$  by

$$e_p(t, s) = \exp \left( \int_s^t \xi_{\mu(\tau)} p(\tau) \Delta \tau \right), \tag{2.3}$$

where  $\xi_h$  is the cylinder transformation given by formula

$$\xi_h(z) = \begin{cases} \log(1 + zh)/h & \text{if } h \neq 0; \\ z & \text{if } h = 0. \end{cases}$$

Note that the function  $x(t) = e_p(t, t_0)x_0$  is the unique solution of the Cauchy problem

$$x^\Delta(t) = p(t)x(t), \quad x(t_0) = x_0, \quad t_0 \in \mathbb{T},$$

see [7] for details.

**Remark 2.8.** If  $p \in \mathcal{R}$  then

$$1 + \int_a^t p(u)\Delta u \leq e_p(t, a) \leq \exp \left( \int_a^t p(u)\Delta u \right) \quad \forall t \in \mathbb{T}_a^+.$$

**Theorem 2.9** (Comparison Theorem, [8]).

Let  $t_0 \in \mathbb{T}$ ,  $x, f \in C_{rd}$  and  $p \in \mathcal{R}^+$ . Then

$$x^\Delta(t) \leq p(t)x(t) + f(t), \quad \text{for all } t \in \mathbb{T}_{t_0}^+$$

implies

$$x(t) \leq x(t_0)e_p(t, t_0) + \int_{t_0}^t e_p(t, \sigma(\tau))f(\tau)\Delta\tau, \quad \forall t \in \mathbb{T}_{t_0}^+.$$

**Definition 2.10.** Let the matrix function  $\mathcal{A}$  be regressive. The matrix function  $\Phi_{\mathcal{A}}(t, t_0)$  satisfying

$$\Phi_{\mathcal{A}}^\Delta(t, t_0) = \mathcal{A}(t)\Phi_{\mathcal{A}}(t, t_0), \quad \Phi_{\mathcal{A}}(t_0, t_0) = E_n, \quad t, t_0 \in \mathbb{T}, \quad t \geq t_0,$$

is called *matrix exponential function* or *fundamental matrix*.

**Theorem 2.11** ([17]). Suppose that  $n \times n$  matrix function  $\mathcal{A}$  on the time scale  $\mathbb{T}$  is regressive. Then

- (i) the matrix exponential function  $\Phi_{\mathcal{A}}(t, t_0)$  is uniquely defined for any  $t_0 \in \mathbb{T}$ ;
- (ii)  $\Phi_{\mathcal{A}}(t, r)\Phi_{\mathcal{A}}(r, s) = \Phi_{\mathcal{A}}(t, s)$  for  $r, s, t \in \mathbb{T}$ ,  $s \leq r \leq t$ ;
- (iii)  $\Phi_{\mathcal{A}}(\sigma(t), s) = (E_n + \mu(t)\mathcal{A}(t))\Phi_{\mathcal{A}}(t, s)$ ;
- (iv) If  $\mathbb{T} = \mathbb{R}$  and  $\mathcal{A}$  is constant, then  $\Phi_{\mathcal{A}}(t, s) = \exp(\mathcal{A}(t - s))$ ;
- (v) If  $\mathbb{T} = h\mathbb{Z}$  with  $h > 0$  and  $\mathcal{A}$  is constant, then  $\Phi(t, s) = (E_n + h\mathcal{A})^{\frac{t-s}{h}}$ .

### 3. Types of stability

Let us consider a linear system

$$x^\Delta = \mathcal{A}(t)x \tag{3.1}$$

and its nonlinear perturbation

$$x^\Delta = \mathcal{A}(t)x + f(t, x). \tag{3.2}$$

We always suppose that the matrix  $\mathcal{A}(t)$  is bounded on  $\mathbb{T}$ . When discuss systems (3.1) and (3.2), we denote by  $x(t, t_0, x_0)$  the solution of the system subject to the initial value  $x(t_0) = x_0$ .

**Definition 3.1.**

- a) System (3.1) is said to be *stable* if, for every  $t_0 \in \mathbb{T}$  and for every  $\varepsilon > 0$  there exists a  $\delta = \delta(\varepsilon, t_0) > 0$  such that

$$|x_0| < \delta \implies |x(t, t_0, x_0)| < \varepsilon, \quad \forall t \in \mathbb{T}_{t_0}^+. \tag{3.3}$$

- b) System (3.1) is said to be *uniformly stable* if for every  $\varepsilon > 0$  there exists a  $\delta = \delta(\varepsilon) > 0$  independent on initial point  $t_0$ , such that (3.3) is satisfied.
- c) System (3.1) is said to be *asymptotically stable* if it is stable and for any  $t_0 \in \mathbb{T}$  there exists a positive value  $c$  such that  $|x_0| < c$  implies  $x(t, t_0, x_0) \rightarrow 0$  as  $t \rightarrow +\infty$ .

**Theorem 3.2** (Choi et al. [13]; DaCunha [14]). *Let the matrix function  $\mathcal{A}$  be regressive. Linear system (3.1) is stable if and only if all its solutions are bounded on  $\mathbb{T}_0^+$ . It is uniformly stable if and only if there exists a positive constant  $\gamma$ , such that  $|\Phi_{\mathcal{A}}(t, t_0)| \leq \gamma$ , for all  $t_0 \in \mathbb{T}$ ,  $t \in T_{t_0}^+$ .*

Later on, when discuss systems (3.1) and (3.2) we always assume that the matrix function  $\mathcal{A}$  is regressive. When discuss the system (2.2) we always assume that  $\mathcal{A}$  is uniformly regressive i.e. inequality (2.1) is true unless the opposite statement is specified.

#### 4. Stability. Grönwall–Bellman approach

The principal objective of this section is to establish a condition on stability of a solution of a time scale system by first approximation. We use a tool well-known in the theory of linear systems. Namely, we introduce the so-called central upper exponents.

**Definition 4.1.** A function  $f : \mathbb{T}_0^+ \times B_r \rightarrow \mathbb{R}^n$  belongs to the class  $\mathcal{F}$  if it satisfies conditions

1.  $f(t, 0) = 0$  for any  $t \geq 0$ ;
2.  $\frac{\partial f}{\partial x}(t, 0) = 0$  for any  $t \geq 0$ ;
3. the Jacobi matrix  $\frac{\partial f}{\partial x}(t, x)$  is uniformly continuous at  $\mathbb{T}_0^+ \times B_r$ .

Observe that for any  $f \in \mathcal{F}$  and any  $\varepsilon > 0$  there exists  $r_1 > 0$  such that

$$|f(t, x)| \leq \varepsilon|x|, \quad \forall t \in \mathbb{T}_0^+, \quad x \in B_{r_1}. \quad (4.1)$$

We consider systems (3.1) and (3.2) where  $f \in \mathcal{F}$  for an  $r > 0$ .

**Definition 4.2.** An rd-continuous function  $u(t) : \mathbb{T}_0^+ \rightarrow \mathbb{R}$  is called *upper function* for system (3.1) if there exists a  $C > 0$  such that the fundamental matrix of (3.1) satisfies<sup>1</sup>

$$|\Phi_{\mathcal{A}}(t, s)| \leq C e_u(t, s), \quad \forall t, s \in \mathbb{T}_0^+, \quad t \geq s.$$

Let  $U$  be the set of all upper functions of (3.1). We call the value

$$\chi(\mathcal{A}) := \inf_{u \in U} \chi_u := \inf_{u \in U} \limsup_{t \rightarrow \infty} \frac{1}{t} \int_0^t u(\tau) \Delta \tau \quad (4.2)$$

central upper exponent for (3.1).

Observe that this exponent is not less than the greatest Lyapunov exponent for solutions of system (3.1). On the other hand, it is not greater than

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \int_0^t |\mathcal{A}(s)| \Delta s.$$

<sup>1</sup> See (2.3) for the definition of generalized exponent  $e_u(t, s)$ .

**Remark 4.3.** Central upper exponent may be greater than the greatest Lyapunov exponent. For linear systems of ODEs, corresponding example was given by Perron [27]. He controlled velocities of growth for solutions specifying the matrix of coefficients. Below in Example 4.6 we obtain the same effect for a system with a constant matrix and a special time scale controlling switches between discrete and continuous regimes.

**Theorem 4.4.** *If  $\chi(\mathcal{A}) < 0$ , there exists  $\varepsilon > 0$  such that for any  $f$  satisfying (4.1), the zero solution of system (3.2) is asymptotically stable.*

This statement is very close to Lemma 3.1 of [7]. To prove it we use the time scale version of the Grönwall–Bellman Lemma first given in [8].

**Lemma 4.5 (Grönwall–Bellman Inequality).** *Let  $t_0 \in \mathbb{T}$ ,  $x, g, p \in \mathcal{C}_{rd}$ ,  $p \geq 0$ . Then*

$$x(t) \leq g(t) + \int_{t_0}^t x(s)p(s) \Delta s \quad \text{for all } t \in \mathbb{T}_{t_0}^+$$

implies

$$x(t) \leq g(t) + \int_{t_0}^t e_p(t, \sigma(s))g(s)p(s) \Delta s \quad \text{for all } t \in \mathbb{T}_{t_0}^+.$$

**Proof of Theorem 4.4.** Set  $g(t, x) = f(t, x)\eta(|x|)$  where  $\eta \in C^\infty(\mathbb{R}^+ \rightarrow \mathbb{R})$  is a cut-off function such that  $\eta(s) = 1$  for  $s \in (0, r_1/2)$ ,  $\eta(s) = 0$  for  $s \geq r_1$ , and  $\eta'(s) \leq 0$  (here  $r_1 = r_1(\varepsilon)$  is a constant from (4.1)). Evidently, the zero solution of the system  $x^\Delta = \mathcal{A}(t)x + g(t, x)$  is asymptotically stable if and only if one of (3.2) is.

Hence, we may assume without loss of generality that inequality (4.1) is satisfied for all  $t \in \mathbb{T}_0^+$  and  $x \in \mathbb{R}^n$ . Then for any solution of (3.2) we have

$$|x(t)|^\Delta = \frac{x^\Delta(t) \cdot x(t)}{|x(t)|} \leq |\mathcal{A}(t)||x(t)| + \varepsilon.$$

By the Comparison Theorem, any solution of (3.2) can be spread to  $\mathbb{T}_0^+$ .

Next, we rewrite (3.2) in standard way as

$$x(t) = \Phi_{\mathcal{A}}(t, t_0)x(t_0) + \int_{t_0}^t \Phi_{\mathcal{A}}(t, s)f(s, x(s)) \Delta s.$$

By (4.1), for any solution  $x$  and for all  $t \in \mathbb{T}_{t_0}^+$  we have

$$|x(t)| \leq |\Phi_{\mathcal{A}}(t, t_0)||x(t_0)| + \varepsilon \int_{t_0}^t |\Phi_{\mathcal{A}}(t, s)||x(s)| \Delta s. \quad (4.3)$$

Fix an upper function  $u$  such that  $\chi_u \in (\chi(\mathcal{A}), 0)$  where  $\chi_u$  is defined by (4.2). It follows from (4.3) that

$$|x(t)| \leq C|x(t_0)|e_u(t, t_0) + C\varepsilon \int_{t_0}^t e_u(t, s)|x(s)| \Delta s.$$

Let  $v(t) = |x(t)|/e_u(t, t_0)$ . Then previous inequality can be rewritten as

$$v(t) \leq C|x(t_0)| + C\varepsilon \int_{t_0}^t v(s)\Delta s.$$

By the Grönwall–Bellman inequality, we have

$$v(t) \leq C|x(t_0)| + C^2\varepsilon|x(t_0)| \int_{t_0}^t e_{C\varepsilon}(t, \sigma(s)) \Delta s.$$

Taking into account [Remark 2.8](#), we obtain

$$|x(t)| \leq C|x(t_0)| \left( 1 + C\varepsilon \int_{t_0}^t \exp(C\varepsilon(t - \sigma(s))) \Delta s \right) e_u(t, t_0).$$

Evidently, if we choose  $\varepsilon$  is sufficiently small i.e.  $C\varepsilon < -\chi_u/2$ , then  $x(t)$  tends to zero exponentially. This completes the proof.  $\square$

Now we turn to the case of the constant matrix  $\mathcal{A}$  and, respectively, to system [\(2.2\)](#). Let  $\lambda_k, k = 1, \dots, n$ , be eigenvalues of the matrix  $\mathcal{A}$ . It is easy to see that the Lyapunov exponents are equal to<sup>2</sup>

$$\nu_k = \limsup_{t \rightarrow \infty} \frac{1}{t} \int_{t_0}^t \Re \xi_{\mu(t)}(\lambda_k) \Delta t = \limsup_{t \rightarrow \infty} \frac{1}{t} \log |e_{\lambda_k}(t, t_0)|.$$

To obtain this formula we proceed to the Jordan normal form in the corresponding system [\(2.2\)](#). Then we apply [Theorem 2.9](#) to estimate norms of solutions of obtained equations.

Evidently for any constant matrix  $\mathcal{A}$  and any time scale  $\mathbb{T}$  we have  $\chi(\mathcal{A}) \geq \max_k \nu_k$ . For  $\mathbb{T} = \mathbb{R}$  or  $\mathbb{T} = \mathbb{Z}$  the converse inequality is true, so  $\chi(\mathcal{A}) = \max_k \nu_k$ . However, in general this is not the case.

**Example 4.6.** Take values  $a \in (0, 1)$  such that

$$\begin{aligned} \nu_1(a) &:= -2a + \log 11(1 - a)/6 < 0; \\ \nu_2(a) &:= -0.5a + \log 2(1 - a)/6 < 0; \\ \chi(a) &:= -0.5a + \log 11(1 - a)/6 > 0 \end{aligned} \tag{4.4}$$

(observe that the set of such values  $a$  is open and non-empty, we can take  $a$  such that  $\chi(a) = 0$  and slightly decrease it). We can choose  $a$  rational, i.e.  $a = p/q, p, q \in \mathbb{N}$ .

Let  $\mathcal{A} = \text{diag}(-2, -0.5)$ . Notice that for  $\mathbb{T}_c = \mathbb{R}$  the system  $x^\Delta = \mathcal{A}x$  is “stable”. For the case  $\mathbb{T}_d = 6\mathbb{Z}$  the situation is opposite (the Lyapunov exponents are  $(\log 11)/6$  and  $(\log 2)/6$  respectively).

For all  $m, n \in \mathbb{N}$  consider the time scale

$$\mathbb{T}_{m,n} = \bigcup_{k \in \mathbb{Z}} [6qmk + 6qn, 6qmk + 6pm + 6qn] \cup 6\mathbb{Z}.$$

<sup>2</sup> See [Definition 2.7](#) for the formula of the transformation  $\xi_h$ .



This time scale is continuous for  $a$ -th part of the time and discrete for  $(1 - a)$ -th part. By (4.4), this implies  $\nu_{1,2} < 0$ .

Now we take an increasing sequence  $n_k \rightarrow \infty$  such that  $m_k := n_{k+1} - n_k \rightarrow \infty$  and  $n_{k+1}/n_k \rightarrow 1$ . Introduce the time scale  $\mathbb{T}$  by formula:

$$\mathbb{T} \cap [6n_jq, 6n_{j+1}q] = \mathbb{T}_{m_j, n_j} \cap [6n_jq, 6n_{j+1}q], \quad j \in \mathbb{N}.$$

For this time scale, Lyapunov exponents of the system  $x^\Delta = \mathcal{A}x$  are still equal to  $\nu_{1,2}$ . Meanwhile,  $\chi(\mathcal{A}) = \chi(a) > 0$  that follows from the definition of central upper exponent, see also (A.1).

### 5. Reduction to ordinary differential equations

The main objective of this section is to prove that a linear time scale system with a bounded and uniformly regressive matrix of coefficients can be “embedded” into a linear system of ordinary differential equations with a bounded matrix. Results of Lemmas 5.1 and 5.2 and Theorem 5.3 given below are applied in the next section. However, they are of independent interest.

**Lemma 5.1.** *For every linear regressive time scale system (3.1), there exists a linear system of ordinary differential equations*

$$\dot{x} = \tilde{\mathcal{A}}(t)x, \quad t \geq 0, \tag{5.1}$$

such that if  $\Phi_{\mathcal{A}}(t, s)$  is the fundamental matrix of system (3.1) and  $\tilde{\Phi}_{\mathcal{A}}(t, s)$  is one for system (5.1), then  $\tilde{\Phi}_{\mathcal{A}}|_{\mathbb{T}_0^+ \times \mathbb{T}_0^+} = \Phi_{\mathcal{A}}$ . If the matrix function  $\mathcal{A}(t)$  is bounded and uniformly regressive, the matrix function  $\tilde{\mathcal{A}}(t)$  is bounded.

**Proof.** We introduce the notation

$$[t]_{\mathbb{T}} := \sup(\mathbb{T} \cap (-\infty, t]).$$

We define  $\tilde{\mathcal{A}}(t)$  as follows:

$$\tilde{\mathcal{A}}(t) = \begin{cases} \mathcal{A}(t), & \text{if } t \in \mathbb{D} \cap \mathbb{T}_0^+; \\ \frac{1}{\mu([t]_{\mathbb{T}})} \text{Log}(E_n + \mu([t]_{\mathbb{T}})\mathcal{A}([t]_{\mathbb{T}})), & \text{if } t \in \mathbb{R}_+ \setminus \mathbb{D}. \end{cases} \tag{5.2}$$

Notice that  $\tilde{\mathcal{A}}(t)$  is not uniquely defined. In general, it cannot be selected real even for a real matrix  $\mathcal{A}(t)$ . However, we can take it rd-continuous on  $\mathbb{R}$  and constant on every connected subset of  $\mathbb{R} \setminus \mathbb{T}$ . Moreover, for any bounded and uniformly regressive matrix function  $\mathcal{A}(t)$  we can select branches of matrix logarithm so that  $\tilde{\mathcal{A}}(t)$  is bounded. In what follows we fix such branches and use the notion log rather than Log that is the multivalued matrix logarithm.

Further, define  $\hat{\Phi}(t)$  by the formula

$$\hat{\Phi}(t) = \begin{cases} \Phi_{\mathcal{A}}(t, 0), & \text{if } t \in \mathbb{T}_0^+; \\ \exp(\tilde{\mathcal{A}}(t)(t - [t]_{\mathbb{T}}))\Phi_{\mathcal{A}}([t]_{\mathbb{T}}, 0), & \text{if } t \in \mathbb{R}_+ \setminus \mathbb{T}. \end{cases}$$

Direct calculation shows that  $\tilde{\Phi}_{\mathcal{A}}(t, s) = \hat{\Phi}(t)\hat{\Phi}^{-1}(s)$  is the fundamental matrix for system (5.1).  $\square$

The following statement is evident.

**Lemma 5.2.** *Let  $\mathcal{A}(t)$  be a bounded, rd-continuous and uniformly regressive matrix function on  $\mathbb{T}_0^+$ , and let  $\tilde{\mathcal{A}}(t)$  be defined by (5.2). Then system (3.1) is stable (asymptotically stable) if and only if corresponding system (5.1) is.*

Given a bounded rd-continuous uniformly regressive matrix function  $\mathcal{A}(t)$ , we notice that for arbitrary rd-continuous matrix function  $\mathcal{B}(t)$  such that  $\|\mathcal{B}\|_{\mathbb{L}^\infty} \leq \delta$ ,  $(\mathcal{A} + \mathcal{B})(t)$  is also uniformly regressive provided that  $\delta > 0$  is sufficiently small. Thus, we can define the extension  $(\widetilde{\mathcal{A} + \mathcal{B}})(t)$  similarly to (5.2), and it is easy to see that the nonlinear mapping

$$\mathcal{L}[\mathcal{B}] \equiv \widehat{\mathcal{B}} := \widetilde{\mathcal{A} + \mathcal{B}} - \tilde{\mathcal{A}}$$

is Lipschitz.

Now let  $\tilde{\Phi}(t, s)$  be the fundamental matrix for the system

$$\dot{x} = (\tilde{\mathcal{A}}(t) + \widehat{\mathcal{B}}(t))x. \tag{5.3}$$

We define

$$\mathcal{B}(t) =: \tilde{\mathcal{L}}[\widehat{\mathcal{B}}] = \begin{cases} \widehat{\mathcal{B}}(t), & \text{if } t \in \mathbb{D} \cap \mathbb{T}_0^+; \\ \frac{\tilde{\Phi}(\sigma(t), t) - E_n}{\mu(t)} - \mathcal{A}(t), & \text{if } t \in \mathbb{S} \cap \mathbb{T}_0^+. \end{cases} \tag{5.4}$$

We are in the position to prove the main result of this section.

**Theorem 5.3.** *Suppose that  $\mathcal{A}(t)$  is a bounded, rd-continuous and uniformly regressive matrix function on  $\mathbb{T}_0^+$ , and let  $\tilde{\mathcal{A}}(t)$  be defined by (5.2). Let a continuous matrix function  $\widehat{\mathcal{B}}(t)$  on  $\mathbb{R}_+$  satisfy the following assumptions:*

$$\begin{aligned} & \|\widehat{\mathcal{B}}\|_{\mathbb{L}^\infty} \leq \delta; \\ & \widehat{\mathcal{B}}(t) = 0 \quad \text{for all } t \in \mathbb{R}, \quad \text{dist}(t, \mathbb{T}) > S, \end{aligned} \tag{5.5}$$

where  $S > 0$  is arbitrary given constant while  $\delta > 0$  is sufficiently small. Then formula (5.4) defines the rd-continuous matrix function  $\mathcal{B}(t)$  on  $\mathbb{T}_0^+$  such that  $(\mathcal{A} + \mathcal{B})(t)$  is uniformly regressive. Moreover, the nonlinear operator  $\tilde{\mathcal{L}}$  defined by (5.4) is Lipschitz left inverse to  $\mathcal{L}$ , and its Lipschitz constant depends only on  $\mathcal{A}$ ,  $S$  and  $\delta$  (in particular, it does not depend on the time scale  $\mathbb{T}$ ).

**Proof.** Fix a matrix  $\widehat{\mathcal{B}}_0$  subject to (5.5), such that  $\|\widehat{\mathcal{B}}_0\|_{\mathbb{L}^\infty} = 1$ , and define  $\Phi_\delta(t, s)$ ,  $\delta \in [0, 1]$ , as the Cauchy matrix for the system

$$\dot{x} = (\tilde{\mathcal{A}}(t) + \delta \widehat{\mathcal{B}}_0(t))x.$$

Let  $R_\delta := \tilde{\mathcal{L}}(\delta \widehat{\mathcal{B}}_0)$ . By (5.4),

$$\frac{\partial}{\partial \delta} R_\delta(t) = \begin{cases} \widehat{\mathcal{B}}_0(t), & \text{if } t \in \mathbb{D} \cap \mathbb{T}_0^+; \\ \frac{1}{\mu(t)} \frac{\partial \Phi_\delta(\sigma(t), t)}{\partial \delta}, & \text{if } t \in \mathbb{S} \cap \mathbb{T}_0^+. \end{cases}$$

It is easy to see that  $\partial\Phi_\delta(\sigma(t), t)/\partial\delta = U(\sigma(t))$ , where  $U(s)$  is the solution of the matrix initial value problem

$$\dot{U}(s) = (\tilde{\mathcal{A}}(s) + \delta\widehat{\mathcal{B}}_0(s))U(s) + \widehat{\mathcal{B}}_0(s)\Phi_\delta(s, t); \quad U(t) = 0.$$

Consequently,

$$\frac{\partial\Phi_\delta(\sigma(t), t)}{\partial\delta} = \int_t^{\sigma(t)} \Phi_\delta(\sigma(t), s)\widehat{\mathcal{B}}_0(s)\Phi_\delta(s, t) ds. \tag{5.6}$$

Without loss of generality we can assume that  $S \geq 1$ . If  $0 < \mu(t) \leq 2S$  then

$$\begin{aligned} \left| \frac{\partial\Phi_\delta(\sigma(t), t)}{\partial\delta} \right| &\leq \int_t^{\sigma(t)} |\Phi_\delta(\sigma(t), s)| |\widehat{\mathcal{B}}_0(s)| |\Phi_\delta(s, t)| ds \\ &\leq \int_t^{\sigma(t)} \exp((\|\tilde{\mathcal{A}}\|_{\mathbb{L}^\infty} + \delta)(\sigma(t) - s)) \exp((\|\tilde{\mathcal{A}}\|_{\mathbb{L}^\infty} + \delta)(s - t)) ds \\ &\leq \mu(t) \exp((\|\tilde{\mathcal{A}}\|_{\mathbb{L}^\infty} + \delta)\mu(t)), \end{aligned}$$

and thus

$$\left| \frac{\partial}{\partial\delta} R_\delta(t) \right| \leq \exp(2S(\|\tilde{\mathcal{A}}\|_{\mathbb{L}^\infty} + 1)) =: M.$$

Otherwise, for  $\mu(t) > 2S$ , from (5.6) we obtain the following estimate for the function  $w(\delta) = |\Phi_\delta(\sigma(t), t)|$ :

$$\begin{aligned} \frac{dw(\delta)}{d\delta} &\leq \left| \frac{\partial\Phi_\delta(\sigma(t), t)}{\partial\delta} \right| \leq \int_t^{\sigma(t)} |\Phi_\delta(\sigma(t), s)| |\widehat{\mathcal{B}}_0(s)| |\Phi_\delta(s, t)| ds \\ &\leq \int_t^{t+S} |\Phi_\delta(\sigma(t), t)| |\Phi_\delta(t, s)| |\widehat{\mathcal{B}}_0(s)| |\Phi_\delta(s, t)| ds \\ &\quad + \int_{\sigma(t)-S}^{\sigma(t)} |\Phi_\delta(\sigma(t), s)| |\widehat{\mathcal{B}}_0(s)| |\Phi_\delta(s, \sigma(t))| |\Phi_\delta(\sigma(t), t)| ds \\ &\leq 2S(\exp((\|\tilde{\mathcal{A}}\|_{\mathbb{L}^\infty} + \delta)S))^2 w(\delta) \leq 2SMw(\delta). \end{aligned}$$

Since  $w(0) = |E_n + \mu(t)\mathcal{A}(t)|$ , we obtain

$$\left| \frac{\partial}{\partial\delta} R_\delta(t) \right| \leq 2SM \exp(2\delta SM) \cdot \frac{|E_n + \mu(t)\mathcal{A}(t)|}{\mu(t)},$$

and the derivative of  $R_\delta$  is uniformly bounded for all  $t$ .

The identity  $\tilde{\mathcal{L}}[\mathcal{L}[\mathcal{B}]] \equiv \mathcal{B}$  is established by direct calculation. Finally, for sufficiently small  $\delta$  the matrix function  $(\mathcal{A} + \mathcal{B})(t)$  is evidently uniformly regressive.  $\square$

**Definition 5.4.** We say that the time scale  $\mathbb{T}$  is *syndetic*, if  $\limsup_{\mathbb{T} \ni t \rightarrow +\infty} \mu(t) < +\infty$ . This notion is similar to one used in Combinatorics and Number Theory.

**Corollary 5.5.** Let  $\mathbb{T}$  be syndetic time scale. Then we can select the value  $S$  so that the relation (5.5) is satisfied for all matrix functions  $\widehat{\mathcal{B}}$ .

**Remark 5.6.** For non-syndetic time scales operator  $\widetilde{\mathcal{L}}$  is not continuous without assumption (5.5). Namely, there exists a small matrix  $\widehat{\mathcal{B}}$  such that  $\widetilde{\mathcal{L}}[\widehat{\mathcal{B}}]$  is unbounded with respect to  $t$ .

## 6. Instability via Millionschikov's rotations

Now we prove two statements in a sense converse to [Theorem 4.4](#).

**Theorem 6.1.** Let the matrix  $\mathcal{A}(t)$  in (3.1) be bounded and uniformly regressive, and let  $\chi(\mathcal{A}) \geq 0$ . Suppose that the time scale  $\mathbb{T}$  is syndetic. Then for any  $\delta > 0$  there exists a matrix  $\mathcal{B}_\delta(t)$  such that

$$\|\mathcal{B}_\delta\|_{\mathbb{L}^\infty} \leq \delta, \quad (6.1)$$

and the system

$$x^\Delta = (\mathcal{A}(t) + \mathcal{B}_\delta(t))x \quad (6.2)$$

is unstable.

We reproduce the classical result on attainability of upper center exponents for systems of ordinary differential equations. In Appendix we provide a full proof of that result cause we need its details later on. We slightly modify the original proof in order to be able to apply [Theorem 5.3](#).

**Theorem 6.2** (Millionschikov [25]). Consider a system (5.1) and suppose that  $\|\widetilde{\mathcal{A}}\|_{\mathbb{L}^\infty} = a < \infty$ . Then for any  $\varepsilon > 0$  and  $\delta > 0$ , there exists a continuous matrix  $\widehat{\mathcal{B}}(t)$  such that

$$\|\widehat{\mathcal{B}}\|_{\mathbb{L}^\infty} \leq (2a + 1)\delta \quad (6.3)$$

and the greatest Lyapunov exponent of system (5.3) is greater than  $\chi(\widetilde{\mathcal{A}}) - \varepsilon$ , where  $\chi(\widetilde{\mathcal{A}})$  is the central upper exponent of system (5.1).

**Proof of Theorem 6.1.** 1. First, we assume  $\chi(\mathcal{A}) > 0$ .

We embed the system (3.1) to a linear system of ODEs (5.1) and observe the evident fact that  $\chi(\widetilde{\mathcal{A}}) \geq \chi(\mathcal{A})$ . Denote  $a := \|\widetilde{\mathcal{A}}\|_{\mathbb{L}^\infty}$ .

By [Theorem 6.2](#) and [Corollary A.1](#), for any  $\delta_1 > 0$  there exists a continuous perturbation  $\widehat{\mathcal{B}}(t)$  such that  $\|\widehat{\mathcal{B}}\|_{\mathbb{L}^\infty} \leq (2a + 1)\delta_1$ , and the greatest Lyapunov exponent of ODE system (5.3) is greater than  $\chi(\mathcal{A})/2$ . Since the time scale is syndetic, the reduction of (5.3) to  $\mathbb{T}$  does also have a positive Lyapunov exponent.

Denote  $\mathcal{B}_\delta = \widetilde{\mathcal{L}}[\widehat{\mathcal{B}}]$ . By [Theorem 5.3](#) (with regard to [Corollary 5.5](#)), there exists a  $K > 0$  such that  $\|\mathcal{B}_\delta\|_{\mathbb{L}^\infty} < K(2a + 1)\delta_1$  for small values of  $\delta_1$ .

To finish the proof it suffices to take  $\delta_1 \leq \frac{\delta}{K(2a+1)}$ .

2. Now we study the case  $\chi(\mathcal{A}) = 0$ . First of all, observe that the transformation  $y = \exp(\varepsilon t)x$  transfers a system  $\dot{x} = P(t)x$  to  $\dot{y} = (P(t) + \varepsilon E_n)y$ .

We begin with the same procedure as in part 1 and construct a perturbation  $\widehat{\mathcal{B}}'(t)$  such that  $\|\widehat{\mathcal{B}}'\|_{\mathbb{L}^\infty} \leq \frac{\delta}{3K(2a+1)}$ , and the greatest Lyapunov exponent of the system

$$\dot{x} = (\widetilde{\mathcal{A}}(t) + \widehat{\mathcal{B}}'(t))x$$

is greater than  $-\frac{\delta}{3K(2a+1)}$ .

Now it is easy to see that the matrix  $\mathcal{B}_\delta = \widetilde{\mathcal{L}}[\widehat{\mathcal{B}}' + \frac{2\delta}{3K(2a+1)}E_n]$  satisfies (6.1) and provides unstable system (6.2).  $\square$

For non-syndetic time scales the similar result is true under the positivity assumption for the upper central exponent.

**Theorem 6.3.** *Let the matrix  $\mathcal{A}(t)$  in (3.1) be bounded and uniformly regressive, and let  $\chi(\mathcal{A}) > 0$ . Then for any  $\delta > 0$  there exists a matrix  $\mathcal{B}_\delta(t)$  satisfying (6.1) and such that system (6.2) is unstable.*

**Proof.** Notice that direct repetition of the proof of Theorem 6.1 does not work since the assumption (5.5) for the perturbation  $\widehat{\mathcal{B}}(t)$  constructed in Theorem 6.2 may be violated. So, we need to modify the Millionschikov method.

As in the proof of Theorem 6.1, we embed system (3.1) to linear system of ODEs (5.1) and denote  $a := \|\widetilde{\mathcal{A}}\|_{\mathbb{L}^\infty}$ . Recall that

$$\widetilde{\mathcal{A}}(t) = \frac{1}{\mu([t]_{\mathbb{T}})} \log(E_n + \mu([t]_{\mathbb{T}})\mathcal{A}([t]_{\mathbb{T}})), \quad t \in \mathbb{R}_+ \setminus \mathbb{D}.$$

We choose  $S_1(\varepsilon, a)$  such that  $\log(1 + sa)/s < \varepsilon/4$  provided  $s > S_1$ .

Now we follow the proof of Theorem 6.2 (see Appendix) up to definition of segments  $Q_j$  (Fig. 4). Without loss of generality we assume that  $T_0 \geq S_1 \max(1, 8a/\varepsilon)$ .

We start with the segment  $[T, 2T]$  where  $T$  is defined in Step 2 of the proof of Millionschikov’s theorem, see (A.6).

There may be a segment  $[\tau_j, \tau_j + 1]$  and a segment  $[\tau_{j+1}, \tau_{j+1} + 1]$  where the perturbation from Theorem 6.2 is non-zero. On these segments two steps of rotation from  $x_0(t)$  to  $x_1(t)$  are performed.

Observe that

$$[\tau_j, \tau_{j+1}] \cap \mathbb{T} \neq \emptyset. \tag{6.4}$$

Otherwise, the segment  $[\tau_j, \tau_{j+1}]$  completely belongs to a “gap” of the time scale of length greater than  $S_1$  and, consequently, inequality (A.8) fails for  $i = j$ .

We introduce time instants  $\tau'_j$  and  $\tau'_{j+1}$  as follows.

$$\tau'_j = \begin{cases} \tau_j, & \text{if } \text{dist}(\tau_j, \mathbb{T}) \leq S_1; \\ \sigma([\tau_j]_{\mathbb{T}}), & \text{if } \text{dist}(\tau_j, \mathbb{T}) > S_1; \end{cases}$$

$$\tau'_{j+1} = \begin{cases} \tau_{j+1}, & \text{if } \text{dist}(\tau_{j+1}, \mathbb{T}) \leq S_1; \\ [\tau_{j+1}]_{\mathbb{T}}, & \text{if } \text{dist}(\tau_{j+1}, \mathbb{T}) > S_1. \end{cases}$$

Notice that  $\tau_{j+1} \geq \tau'_{j+1} \geq \tau'_j \geq \tau_j$  by virtue of (6.4).

By definition of  $S_1$ , the following analog of inequality (A.8) is satisfied in any case:

$$\frac{|x_1(\tau'_{j+1})|}{|x_1(\tau'_j)|} : \frac{|x_0(\tau'_{j+1})|}{|x_0(\tau'_j)|} \geq \exp\left(\frac{\varepsilon T_0}{2} - \frac{\varepsilon T_0}{4}\right) = \exp\left(\frac{\varepsilon T_0}{4}\right). \tag{6.5}$$

This implies  $\tau'_j + 1 < \tau'_{j+1}$  (recall that  $aS_1 \leq \varepsilon T_0/8$ ). Consequently,

$$[\tau'_j, \tau'_j + 1] \cap [\tau'_{j+1}, \tau'_{j+1} + 1] = \emptyset.$$

Observe that inequality (6.5) is still enough to imply item **C** of the Step 3 in the proof of Theorem 6.2 (see also the footnote to Eq. (A.3)).

So, the proof of Theorem 6.2 still passes with  $\tau_j$  and  $\tau_{j+1}$  replaced with  $\tau'_j$  and  $\tau'_{j+1}$ . On the interval  $[T, 2T]$  the perturbation  $\widehat{\mathcal{B}}(t)$  is non-zero on segments  $[\tau'_j, \tau'_j + 1]$  and  $[\tau'_{j+1}, \tau'_{j+1} + 1]$  only. The similar statement is true for all other segments  $[kT, (k + 1)T]$ .

Thus, for any  $\delta_1 > 0$  and  $\varepsilon > 0$  we have constructed a continuous perturbation  $\widehat{\mathcal{B}}(t)$  such that  $\|\widehat{\mathcal{B}}\|_{\mathbb{L}^\infty} \leq (2a + 1)\delta_1$ , the greatest Lyapunov exponent of ODE system (5.3) is greater than  $\chi(\mathcal{A}) - \varepsilon$ , and the inequality (5.5) holds with  $S = S_1 + 1$ .

Denote  $\mathcal{B}_\delta = \widetilde{\mathcal{L}}[\widehat{\mathcal{B}}]$ . By Theorem 5.3, there exists a  $K(\varepsilon, a) > 0$  such that  $\|\mathcal{B}_\delta\|_{\mathbb{L}^\infty} < K(2a + 1)\delta_1$  for small values of  $\delta_1$ .

Now we put  $\varepsilon = \chi(\mathcal{A})/2$  and claim that the greatest Lyapunov exponent of the time scale system (6.2) is not less than  $\chi(\mathcal{A})/2$ . Indeed, let  $\Phi(t)$  be a fundamental matrix of system (5.3). By construction, there exists an unbounded sequence  $t_k \in \mathbb{R}$  such that  $|\Phi(t_k)| > \exp(\chi(\mathcal{A})t_k/2)$ .

Denote  $s_k = [t_k]_{\mathbb{T}}$  and notice that if  $t_k - s_k > S_1$  then  $\mu(s_k) > S_1$  and therefore  $|\widetilde{\mathcal{A}}(t)| \leq \varepsilon/4$  for  $t \in [s_k, t_k]$ . This gives

$$\begin{aligned} |\Phi(s_k)| &\geq |\Phi(t_k)| \min\{\exp(-\varepsilon(t_k - s_k)/4), \exp(-aS_1)\} \\ &\geq \exp(\chi(\mathcal{A})t_k/2 - \varepsilon(t_k - s_k)/4) \exp(-aS_1) \\ &\geq \exp(-aS_1) \exp(\chi(\mathcal{A})s_k/2), \end{aligned}$$

and the claim follows.

To finish the proof it suffices to take  $\delta_1 \leq \frac{\delta}{K(2a+1)}$ .  $\square$

The next statement is a generalization of the result of [22] for time scales.

**Theorem 6.4.** *Let the matrix  $\mathcal{A}(t)$  in (3.1) be bounded and uniformly regressive, and let  $\chi(\mathcal{A}) > 0$ . Then there exists a continuous map  $f : \mathbb{T}_0^+ \times B_1 \rightarrow \mathbb{R}^n$  such that<sup>3</sup>  $f \in \mathcal{F}$  and the solution  $x(t) \equiv 0$  of the corresponding system (3.2) is unstable. If the time scale  $\mathbb{T}$  is syndetic, the same is true provided  $\chi(\mathcal{A}) \geq 0$ .*

**Proof.** By Theorems 6.1 and 6.3, there exist continuous perturbation matrices  $\mathcal{B}_\ell(t)$ ,  $\ell \in \mathbb{N}$ , such that

- 1)  $|\mathcal{B}_\ell(t)| \leq 2^{-\ell}$  for all  $\ell \in \mathbb{N}$ ,  $t \in \mathbb{T}$ ;
- 2) for every  $\ell \in \mathbb{N}$  the system

$$x^\Delta = (\mathcal{A}(t) + \mathcal{B}_\ell(t))x \tag{6.6}$$

has a solution with positive Lyapunov exponent.

Fix an unbounded solution  $x_1(t)$  of system (6.6) for  $\ell = 1$  such that  $|x_1(0)| \leq 1$ . Select  $T_1 > 0$  so that  $|x_1(T_1)| \geq 2$ ,  $|x_1(t)| < 2$  while  $0 \leq t < T_1$ . Then we construct an unbounded solution  $x_2(t)$  of system (6.6) for  $\ell = 2$  such that  $|x_2(t)| < |x_1(t)|/2$  for  $0 \leq t \leq T_1$ . Given  $x_2(t)$  we select the first time instant  $T_2$  such that  $|x_2(T_2)| \geq 2$ . Then we construct  $x_3(t)$  and  $T_3$  and so on (Fig. 2).

<sup>3</sup> See Definition 4.1.




Fig. 2. Solutions of the perturbed system ( $\mathbb{T} = \mathbb{R}$ ).

Now we construct a map  $f : [0, \infty) \times B_1 \rightarrow \mathbb{R}^n$  such that the system (3.2) coincides with (6.6) in some neighborhood of the graph of  $x_\ell$  on  $[0, T_\ell]$  for all  $\ell \in \mathbb{N}$ . This implies that all  $x_\ell(t)$  are solutions of (3.2) on  $[0, T_\ell]$ . Since  $|x_\ell(0)| \rightarrow 0$  as  $\ell \rightarrow \infty$  and  $|x_\ell(T_\ell)| \geq 2$ , the zero solution of (3.2) is unstable.

We set

$$\Psi_\ell(t, x) = \begin{cases} \mathcal{B}_\ell(t)\phi\left(\frac{|x-x_\ell(t)|}{\epsilon_\ell(t)}\right), & \text{if } |x - x_\ell(t)| \leq \epsilon_\ell(t), \quad t < T_\ell; \\ 0, & \text{otherwise,} \end{cases}$$

where  $\phi(s)$  is a smooth cut-off function equal to one for  $s \leq 1/2$  and to zero for  $s \geq 1$ , while  $\epsilon_\ell(t) = |x_\ell(t)|/4$ . Evidently, all  $\Psi_\ell(t, x)$  are continuous and continuously differentiable with respect to  $x$ .

We define

$$f(t, x) = \sum_{\ell=1}^{\infty} \Psi_\ell(t, x) \cdot x.$$

Notice that for any  $k < \ell$  and  $t < T_k$  we have  $|x_\ell(t)| < \frac{|x_{\ell-1}(t)|}{2} < \dots < \frac{|x_k(t)|}{2^{\ell-k}}$  and therefore  $|x_\ell(t) - x_k(t)| > |\epsilon_\ell(t)| + |\epsilon_k(t)|$ . Thus, balls  $B(x_k(t), \epsilon_k(t))$  and  $B(x_\ell(t), \epsilon_\ell(t))$  are pairwise disjoint, and the map  $f$  satisfies the above assumption. Moreover, evidently  $f(t, 0) \equiv 0$ . So, we should only examine differentiated series

$$\sum_{\ell=1}^{\infty} \Psi_\ell(t, x) + \sum_{\ell=1}^{\infty} \nabla_x \Psi_\ell(t, x) \cdot x. \tag{6.7}$$

Since  $|\Psi_\ell(t, x)| \leq |\mathcal{B}_\ell(t)| < 1/2^\ell$ , the first series in (6.7) uniformly converges on  $\mathbb{T}_0^+ \times B_1$ . Further, for  $(t, x)$  in support of  $\Psi_\ell$  we have

$$\begin{aligned} |\nabla_x \Psi_\ell(t, x) \cdot x| &\leq |\mathcal{B}_\ell(t)| \frac{\max_s |\phi'(s)|}{\epsilon_\ell(t)} |x| \\ &\leq \frac{\max_s |\phi'(s)|}{2^\ell} \frac{|x_\ell(t)| + |x - x_\ell(t)|}{\epsilon_\ell(t)} \leq \frac{5 \max_s |\phi'(s)|}{2^\ell}, \end{aligned}$$

and the second series in (6.7) also uniformly converges. Therefore, the sum of the series (6.7) equals  $\frac{\partial f}{\partial x}(t, x)$  and is uniformly continuous on  $\mathbb{T}_0^+ \times B_1$ . The equality  $\frac{\partial f}{\partial x}(t, 0) \equiv 0$  is evident since supports of  $\Psi_\ell$  do not intersect  $t$  axis. This implies  $f \in \mathcal{F}$ .  $\square$

**Remark 6.5.** Recall that in Example 4.6 we construct a linear system with constant matrix on a syndetic time scale which has negative Lyapunov exponents but positive central upper exponent. Theorem 6.1 implies that this (asymptotically stable and even exponentially stable) system becomes unstable under arbitrarily

small linear perturbation. [Theorem 6.4](#) shows that a nonlinear system with exponentially stable first approximation can be unstable. Such examples can be found, for instance, in [\[28\]](#), but for non-regressive time scale systems.

## 7. Stability and instability by first approximation

First of all, we recall the time scale version of the Lyapunov theorem on asymptotic stability by first approximation, proved in [\[8\]](#), see also [\[2\]](#).

**Definition 7.1.** Let  $r > 0$ . We say that a continuous function  $V(t, x) : \mathbb{T}_0^+ \times B_r \rightarrow \mathbb{R}$  is a *strict Lyapunov function* for a time scale system

$$x^\Delta = F(t, x), \quad t \in \mathbb{T}_0^+, \quad x \in \mathbb{R}^n; \quad F(t, 0) \equiv 0, \quad (7.1)$$

if the following conditions are fulfilled for some  $a > 0$  and for all  $t \in \mathbb{T}_a^+$ ,  $x \in B_r$ :

1.  $V(t, x) \geq w_+(x)$ , and  $V(t, 0) \equiv 0$ ;
2. the trajectory  $\Delta$ -derivative of  $V$  satisfies  $V^\Delta(t, x) \leq -w_-(x)$ .

Here  $w_\pm(x) : B_r \rightarrow \mathbb{R}$  are positive definite functions.

**Remark 7.2.** Note that condition 2 means

$$\begin{aligned} \frac{\partial V}{\partial t}(t, x) + \frac{\partial V}{\partial x}(t, x) \cdot F(t, x) &\leq -w_-(x) \quad \forall t \in \mathbb{T}_a^+ \cap \mathbb{D}; \\ V(\sigma(t), x + \mu(t)F(t, x)) - V(t, x) &\leq -\mu(t)w_-(x) \quad \forall t \in \mathbb{T}_a^+ \cap \mathbb{S}. \end{aligned}$$

**Theorem 7.3** (*Lyapunov's Theorem, [8]*). *If there is a strict Lyapunov function for system (7.1), then the zero solution of this system is asymptotically stable.*

**Definition 7.4.** A constant matrix  $\mathcal{A}$  is called *strongly stable* with respect to  $\mathbb{T}$  if its eigenvalues  $\lambda_k$ ,  $k = 1, \dots, n$ , satisfy inequality  $\limsup_{\mathbb{S} \ni t \rightarrow +\infty} \frac{1}{\mu(t)} (|1 + \mu(t)\lambda_k|^2 - 1) < 0$ .

**Remark 7.5.** It is easy to see that if  $\mathcal{A}$  is strongly stable then the following is true:

1.  $\Re(\lambda_k) < 0$ ,  $k = 1, \dots, n$ ;
2. time scale  $\mathbb{T}$  is syndetic.

**Theorem 7.6.** *Suppose that the matrix  $\mathcal{A}$  is strongly stable. Then there exists  $\varepsilon > 0$  such that for any  $r > 0$  and any  $f : \mathbb{T}_0^+ \times B_r \rightarrow \mathbb{R}^n$  satisfying condition (4.1), the solution  $x = 0$  of the system*

$$x^\Delta = \mathcal{A}x + f(t, x) \quad (7.2)$$

*is asymptotically stable.*

**Proof.** We show that, under the assumptions of theorem, there is a positive definite matrix  $\mathcal{B}$  such that the quadratic form  $V(x) = x^T \mathcal{B}x$  is a strict Lyapunov function for the system (7.2).



Making a non-degenerate transformation  $x = \mathcal{S}y$ , we can reduce the first approximation system (2.2) to the Jordan form

$$y^\Delta = \mathcal{J}y, \tag{7.3}$$

where  $\mathcal{J} = \mathcal{S}^{-1}\mathcal{A}\mathcal{S} = \text{diag}(\mathcal{J}_1, \dots, \mathcal{J}_k)$  while for any  $m = 1, \dots, k$

$$\mathcal{J}_m = \begin{pmatrix} \lambda_m & 0 & 0 & \dots & 0 & 0 \\ \delta & \lambda_m & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & \delta & \lambda_m \end{pmatrix} =: \lambda_m E + \delta I. \tag{7.4}$$

A parameter  $\delta > 0$  may be selected arbitrarily small.

System (7.2) takes form

$$y^\Delta = \mathcal{J}y + g(t, y),$$

where  $g(t, y) = \mathcal{S}^{-1}f(t, \mathcal{S}y)$ . The assumption (4.1) implies

$$|g(t, y)| \leq |\mathcal{S}| \cdot |\mathcal{S}^{-1}| \cdot \varepsilon|y| \leq C(\delta) \cdot \varepsilon|y|.$$

First, we construct the desired quadratic form for the system (7.3). It suffices to consider the system

$$z^\Delta = \mathcal{J}_m z. \tag{7.5}$$

We set  $V(z) = |z|^2$  for (7.5). Direct calculation of trajectory  $\Delta$ -derivative gives

$$V^\Delta = \begin{cases} \frac{|1 + \mu(t)\lambda_m|^2 - 1}{\mu(t)}|z|^2 + 2\delta\Re((1 + \mu(t)\lambda_m)z \cdot Iz) + \delta^2\mu(t)|Iz|^2, & t \in \mathbb{S}; \\ 2\Re(\lambda_m)|z|^2 + 2\delta\Re(z \cdot Iz), & t \in \mathbb{D}. \end{cases}$$

Taking into account Remark 7.5, we obtain  $V^\Delta \leq -\varkappa|z|^2$  with some  $\varkappa > 0$ , if  $\delta$  is sufficiently small and  $t \in \mathbb{T}_a^+$  for  $a$  sufficiently large.

For nonlinear system (7.2) we set  $V(y) = |y|^2$  and observe that  $V^\Delta \leq -\frac{\varkappa}{2}|y|^2$  if  $\varepsilon$  is sufficiently small.  $\square$

**Corollary 7.7.** *If a matrix  $\mathcal{A}$  is strongly stable with respect to  $\mathbb{T}$  then the central upper exponent of the system (2.2) is negative.*

**Proof.** By Theorem 7.6, there exists a transformation  $x = \mathcal{S}y$  such that the trajectory derivative of the Lyapunov function  $V(y) = |y|^2$  satisfies  $V^\Delta \leq -\varkappa|y|^2$  with some  $\varkappa > 0$ . This means that for any solution  $\varphi(t)$  of linear system (2.2) we have

$$|\varphi(t)| \leq Ce_{-\varkappa/2}(t, s)|\varphi(s)|, \quad t > s, \quad t, s \in \mathbb{T},$$

where  $C$  is a positive constant depending on the transformation matrix  $\mathcal{S}$ .

So,  $u(t) \equiv -\varkappa/2$  is an upper function for system (2.2) and thus  $\chi(\mathcal{A}) < 0$ .  $\square$

Now we prove an analog of the famous Chetaev theorem on instability by first approximation (see [12] for the classical theorem and [11] for the “discrete” one) for time scale systems.




Fig. 3. Selection of  $x_0$ .

**Definition 7.8.** Let  $r > 0$ . We say that a continuous function  $V(t, x) : \mathbb{T}_0^+ \times B_r \rightarrow \mathbb{R}$  is a *Chetaev function* for the system (7.1) if the following conditions are fulfilled for some  $a > 0$  and for all  $t \in \mathbb{T}_a^+$ :

1.  $0 \in \partial\Omega_t$ , where  $\Omega_t = \{x \in B_r : V(t, x) > 0\}$ ;
2.  $V$  is continuous at the origin uniformly with respect to  $t$ ;
3. the trajectory  $\Delta$ -derivative of  $V$  satisfies  $V^\Delta(t, x) \geq w(x)$ , where  $w(x) : \Omega_t \rightarrow \mathbb{R}$  is a positive definite function (compare with condition 2 in Definition 7.1).

**Theorem 7.9.** *If there is a Chetaev function for system (7.1), then the zero solution of this system is unstable.*

**Proof.** Let  $t_0 \in \mathbb{T}_a^+$ , and let  $x_0 \in \Omega_{t_0}$ . Denote by  $\hat{x}(t)$  the solution of (7.1) corresponding to initial conditions  $x(t_0) = x_0$  (Fig. 3). By condition 3, the function  $V(t, \hat{x}(t))$  increases while  $\hat{x}(t) \in B_r$ . Moreover, the set  $\{(t, x) : x \in \Omega_t, V(t, x) \geq V(t_0, x_0)\}$  is uniformly separated from zero by the condition 2, and therefore  $(V(t, \hat{x}(t)))^\Delta \geq b > 0$ . This means that  $\hat{x}(t)$  leaves the ball  $B_r$ , since otherwise  $V(t, \hat{x}(t))$  is unbounded. Since  $x_0$  can be chosen arbitrarily close to zero by condition 1, the zero solution is unstable.  $\square$

**Definition 7.10.** A constant matrix  $\mathcal{A}$  is called *strongly unstable* with respect to  $\mathbb{T}$  if we can split its eigenvalues into two sets (the second one may be empty)

$$\lambda_k^{(1)}, \quad k = 1, \dots, \ell; \quad \lambda_j^{(2)}, \quad j = 1, \dots, n - \ell; \quad 1 \leq \ell \leq n,$$

such that the following inequalities are satisfied:

1.  $\liminf_{\mathbb{S} \ni t \rightarrow +\infty} \frac{1}{\mu(t)} (|1 + \mu(t)\lambda_k^{(1)}|^2 - 1) > 0, \quad k = 1, \dots, \ell;$
2.  $\liminf_{\mathbb{S} \ni t \rightarrow +\infty} \frac{1}{\mu(t)} (|1 + \mu(t)\lambda_k^{(1)}|^2 - |1 + \mu(t)\lambda_j^{(2)}|^2) > 0, \quad k = 1, \dots, \ell, \quad j = 1, \dots, n - \ell;$
3. if  $\sup \mathbb{D} = +\infty$ , we assume in addition that  $\Re(\lambda_k^{(1)}) > 0, \Re(\lambda_k^{(1)}) > \Re(\lambda_j^{(2)})$  for all  $k = 1, \dots, \ell, j = 1, \dots, n - \ell$ .

**Remark 7.11.** Given a time scale  $\mathbb{T}$ , strongly stable and strongly unstable matrices form two non-intersecting classes. For time-invariant systems of ordinary differential equations ( $\mathbb{T} = \mathbb{R}$ ) these matrices satisfy the assumptions  $\max_k \Re \lambda_k < 0$  (Hurwitz matrices) and  $\max_k \Re \lambda_k > 0$ , respectively.

**Theorem 7.12.** Let  $\mathbb{T}$  be a syndetic time scale. Suppose that the matrix  $\mathcal{A}$  is strongly unstable. Then there exists  $\varepsilon > 0$  such that for any  $r > 0$  and any  $f : \mathbb{T}_0^+ \times B_r \rightarrow \mathbb{R}^n$  satisfying condition (4.1), the solution  $x = 0$  of the system (7.2) is unstable.

**Proof.** We show that, under the assumptions of theorem, there is a matrix  $\mathcal{B}$  such that the quadratic form  $V(x) = x^T \mathcal{B}x$  is a Chetaev function for the system (7.2).

As in the proof of Theorem 7.6, we can reduce the first approximation system (2.2) to the Jordan form

$$(y^{(1)})^\Delta = \mathcal{J}^{(1)}y^{(1)}; \quad (y^{(2)})^\Delta = \mathcal{J}^{(2)}y^{(2)}, \tag{7.6}$$

where  $y^{(1)} \in \mathbb{R}^\ell$ ,  $y^{(2)} \in \mathbb{R}^{n-\ell}$ , and, similarly to (7.4),

$$\mathcal{J}^{(1)} = \text{diag}(\lambda_1^{(1)}E + \delta I, \dots, \lambda_m^{(1)}E + \delta I); \quad \mathcal{J}^{(2)} = \text{diag}(\lambda_1^{(2)}E + \delta I, \dots, \lambda_l^{(2)}E + \delta I).$$

A parameter  $\delta > 0$  may be selected arbitrarily small.

We set  $V(y) = |y^{(1)}|^2 - |y^{(2)}|^2$  for (7.6). Direct calculation of trajectory  $\Delta$ -derivative gives

$$V^\Delta = \begin{cases} \sum_{i=1}^{\ell} \frac{|1 + \mu(t)\lambda_i^{(1)}|^2 - 1}{\mu(t)} (y_i^{(1)})^2 - \sum_{i=1}^{n-\ell} \frac{|1 + \mu(t)\lambda_i^{(2)}|^2 - 1}{\mu(t)} (y_i^{(2)})^2 + O(\delta|y|^2), & t \in \mathbb{S}; \\ 2 \sum_{i=1}^{\ell} \Re(\lambda_i^{(1)})(y_i^{(1)})^2 - 2 \sum_{i=1}^{n-\ell} \Re(\lambda_i^{(2)})(y_i^{(2)})^2 + O(\delta|y|^2), & t \in \mathbb{D}. \end{cases}$$

By assumptions 1–3, we conclude that  $V > 0$  implies  $V^\Delta \geq \varkappa|y|^2$  with some  $\varkappa > 0$ , if  $\delta$  is sufficiently small and  $t \in \mathbb{T}_a^+$  for  $a$  sufficiently large.

For nonlinear system (7.2), similarly to Theorem 7.6, we obtain  $V^\Delta \geq \frac{\varkappa}{2}|y|^2$  if  $\varepsilon$  is sufficiently small.  $\square$

**Acknowledgments**

The first author was partially supported by RFBR grants 14-01-00202 and 15-01-03797-a, by Saint Petersburg State University under Thematic Plan 6.38.223.2014, and by the Fulbright Foundation (Fulbright Visiting Scholar Program, grantee id 68150272). The second author was supported by RFBR grant 15-01-07650 and by Saint Petersburg State University grant 6.38.670.2013.

Authors are grateful to Prof. Martin Bohner, Prof. Anatoly Martynyuk and Prof. Andrejs Reinfelds for their attention to our research and for their precious advices and remarks. We also thank Prof. Vitaly Slyn’ko who provided us with reference [28], and anonymous referee for useful comments.

**Appendix A. Proof of Millionschikov’s theorem (Theorem 6.2)**

We use the following relation (see [10, page 116, (8.8)]):

$$\chi(\tilde{\mathcal{A}}) = \lim_{T \rightarrow \infty} \limsup_{k \rightarrow \infty} \frac{1}{kT} \sum_{i=0}^{k-1} \log |\Phi((i+1)T, iT)|, \tag{A.1}$$

where  $\Phi$  is the Cauchy matrix for the system (5.1).

We start with the main idea of the proof. Consider  $T > 0$  so that the value

$$\limsup_{k \rightarrow \infty} \frac{1}{kT} \sum_{i=0}^{k-1} \log |\Phi((i+1)T, iT)|$$

is close to  $\chi(\tilde{\mathcal{A}})$ , see (A.1). Let  $x_i$  ( $i = 0, 1, 2, \dots$ ) be a unit vector such that

$$|\Phi((i + 1)T, iT)x_i| = |\Phi((i + 1)T, iT)|, \tag{A.2}$$

and put  $x_i(t) = \Phi(t, iT)x_i$ .

It is  $x_0(t)$  that has the fastest growth among solutions of (5.1) on  $[0, T]$ . Without loss of generality, we may say that on  $[T, 2T]$ , the solution  $x_1(t)$  increases faster than  $x_0(t)$ .

We perturb system (5.1) in the following way. First of all, we rotate the solution  $x_0(t)$  in the plane  $\langle x_0(t), x_1(t) \rangle$  by an angle  $\delta > 0$ . Thus we obtain a function  $y_0(t)$ . This rotation can be done on a time segment of length  $\ll T$ . Then, for greater values of  $t$ , we set perturbation zero. Since  $x_1(t)$  increases faster than  $x_0(t)$ , the angle between vectors  $y_0(t)$  and  $x_1(t)$  becomes less than  $\delta$ . This happens on a time period of length  $\ll T$ . Then we perturb system (5.1) so that  $y_0(t)$  becomes parallel to  $x_1(t)$ . Then we set the perturbation equal to zero up to  $t = 2T$ .

Similarly, we consider segment  $[2T, 3T]$  and later ones. Finally, we obtain a solution  $y_0(t)$  of the perturbed system that has Lyapunov exponent, close to  $\chi(\tilde{\mathcal{A}})$ .

Now we proceed to the detailed proof.

**Step 1.** Given  $\varepsilon > 0$  and  $\delta > 0$ , we fix a  $T_0 > 1$  so that<sup>4</sup>

$$\exp(\varepsilon T_0/4) \cdot \sin^2 \delta \geq 1. \tag{A.3}$$

Let triangles  $\triangle ABC$  and  $\triangle A_1 B_1 C_1$  be such that

$$\frac{B_1 C_1}{A_1 C_1} : \frac{BC}{AC} \geq \exp(\varepsilon T_0/4); \quad \sphericalangle A = \delta. \tag{A.4}$$

Then (A.3), (A.4) and Sine Theorem imply that

$$\sin \sphericalangle B_1 \leq \frac{\sin \sphericalangle B_1}{\sin \sphericalangle A_1} = \frac{A_1 C_1}{B_1 C_1} \leq \exp(-\varepsilon T_0/4) \cdot \frac{1}{\sin \delta} \leq \sin \delta. \tag{A.5}$$

Since  $A_1 C_1 / B_1 C_1 \leq 1$  we have  $\sphericalangle B_1 \leq \sphericalangle A_1$ , and, consequently,  $\sphericalangle B_1 \leq \pi/2$ . Therefore, (A.5) implies  $\sphericalangle B_1 \leq \delta$ .

**Step 2.** Fix  $T > 0$  such that  $m = T/T_0 \in \mathbb{N}$ ,  $(2a + (2a + 1)\delta)/m < \varepsilon/8$  and

$$\limsup_{k \rightarrow \infty} \frac{1}{kT} \sum_{i=0}^{k-1} \log |\Phi((i + 1)T, iT)| > \chi(\tilde{\mathcal{A}}) - \frac{\varepsilon}{4}. \tag{A.6}$$

**Step 3.** Take a unit vector  $x_i$  ( $i = 0, 1, 2, \dots$ ) such that (A.2) is satisfied. Let

$$x_i(t) = \Phi(t, iT)x_i \tag{A.7}$$

be solutions of (5.1).

<sup>4</sup> It is sufficient to take  $\varepsilon T_0/2$  instead of  $\varepsilon T_0/4$  in formulae (A.3)–(A.5). In fact, we need such selection of  $T_0$  in Theorem 6.3 where we reproduce a part of the proof of Millionschikov’s theorem.




Fig. 4. Segments  $Q_l$ .

Set  $\widehat{\mathcal{B}}(t) = 0$  for  $0 \leq t \leq T$ . Suppose that

$$\frac{|x_1(2T)|}{|x_1(T)|} : \frac{|x_0(2T)|}{|x_0(T)|} > \exp(3\varepsilon T/4)$$

(if this is wrong, we set  $\widehat{\mathcal{B}}(t) = 0$  for  $T < t \leq 2T$ ).

Divide the segment  $[T, 2T]$  to  $m$  segments of length  $T_0$ :

$$Q_l = [\tau_l, \tau_{l+1}] = [T + (l - 1)T_0, T + lT_0] \quad (l = 1, 2, \dots, m).$$

Let  $Q_j$  be the first of segments  $Q_1, Q_2, \dots, Q_{m-1}$  where

$$\frac{|x_1(\tau_{l+1})|}{|x_1(\tau_l)|} : \frac{|x_0(\tau_{l+1})|}{|x_0(\tau_l)|} \geq \exp(\varepsilon T_0/2). \tag{A.8}$$

So,  $\tau_j < \tau_{j+1} < \tau_{j+2}$  are ends of segments  $Q_j$  and  $Q_{j+1}$  (Fig. 4).

Note that the number of values of  $l$  for which (A.8) is satisfied is not less than 2. Indeed, otherwise

$$\frac{|x_1(2T)|}{|x_1(T)|} : \frac{|x_0(2T)|}{|x_0(T)|} \leq \exp((m - 1)\varepsilon T_0/2) \cdot \frac{|x_1(\tau_{l+1})|}{|x_1(\tau_l)|} : \frac{|x_0(\tau_{l+1})|}{|x_0(\tau_l)|}.$$

Since for any nonzero solution  $x(t)$  of (5.1) we have

$$\exp(-aT_0) \leq \frac{|x(\tau_{l+1})|}{|x(\tau_l)|} \leq \exp(aT_0)$$

(we recall that  $a = \|\widetilde{\mathcal{A}}\|_{\mathbb{L}^\infty}$ ), this implies

$$\frac{|x_1(2T)|}{|x_1(T)|} : \frac{|x_0(2T)|}{|x_0(T)|} \leq \exp(\varepsilon T/2) \cdot \exp(2aT_0) \leq \exp(5\varepsilon T/8),$$

a contradiction. Therefore,  $\tau_{j+1} < 2T$ .

Define the perturbation  $\widehat{\mathcal{B}}(t)$  for  $T < t \leq 2T$  in the following way.

- A. If  $t \notin [\tau_j, \tau_j + 1] \cup [\tau_{j+1}, \tau_{j+1} + 1]$  we set  $\widehat{\mathcal{B}}(t) = 0$ .
- B. For  $t \in [\tau_j, \tau_j + 1]$  we set

$$\widehat{\mathcal{B}}(t) = U_\delta^{-1}(t)\widetilde{\mathcal{A}}(t)U_\delta(t) - U_\delta^{-1}(t)\dot{U}_\delta(t) - \widetilde{\mathcal{A}}(t), \tag{A.9}$$

where  $U_\delta(t)$  is an orthogonal matrix such that

$$U_\delta(\tau_j) = E_n, \quad |\dot{U}_\delta(t)| \leq \delta. \tag{A.10}$$




Fig. 5. Millionschikov's rotations.

Namely, we define  $U_\delta(t)$  as a rotation in the plane  $\langle x_0(\tau_j + 1), x_1(\tau_j + 1) \rangle$  in the direction from  $x_0(\tau_j + 1)$  to  $x_1(\tau_j + 1)$  with the speed not greater then  $\delta$ . From (A.9), (A.10) and orthogonality of  $U_\delta$  we deduce inequality (6.3).

By construction, there exist  $\alpha_1 \geq 0, \alpha_2 > 0$  such that

$$U_\delta^{-1}(\tau_j + 1)x_0(\tau_j + 1) = \alpha_1 x_0(\tau_j + 1) + \alpha_2 x_1(\tau_j + 1), \tag{A.11}$$

and

$$\angle(x_0(\tau_j + 1), y_0(\tau_j + 1)) = \delta. \tag{A.12}$$

C. Due to relations (A.11), (A.12), (A.8) and to statements of Step 1, we have

$$\angle(\alpha_1 x_0(\tau_{j+1} + 1) + \alpha_2 x_1(\tau_{j+1} + 1), x_1(\tau_{j+1} + 1)) \leq \delta$$

(here  $\alpha_1$  and  $\alpha_2$  are defined by (A.11)).

For  $t \in [\tau_{j+1}, \tau_{j+1} + 1]$  we take  $\widehat{\mathcal{B}}(t)$  that satisfies (A.9) and (A.10) (with  $\tau_j$  replaced by  $\tau_{j+1}$ ). Instead of inequalities (A.11) and (A.12) we demand that

$$U_\delta^{-1}(\tau_{j+1} + 1)(\alpha_1 x_0(\tau_{j+1} + 1) + \alpha_2 x_1(\tau_{j+1} + 1)) = \beta x_1(\tau_{j+1} + 1)$$

for some  $\beta > 0$  (Fig. 5).

Observe that since  $x_0(t)$  is a solution of (5.1), the function

$$y_0(t) = \begin{cases} x_0(t) & \text{if } T \leq t \leq \tau_j \\ U_\delta^{-1}(t)x_0(t) & \text{if } \tau_j \leq t \leq \tau_j + 1, \\ \alpha_1 x_0(t) + \alpha_2 x_1(t) & \text{if } \tau_j + 1 \leq t \leq \tau_{j+1}, \\ U_\delta^{-1}(t)(\alpha_1 x_0(t) + \alpha_2 x_1(t)) & \text{if } \tau_{j+1} \leq t \leq \tau_{j+1} + 1, \\ \beta x_1(\tau_{j+1} + 1) & \text{if } \tau_{j+1} + 1 \leq t \leq 2T \end{cases}$$

is a solution of system (5.3) with constructed matrix  $\widehat{\mathcal{B}}(t)$ .

**Step 4.** We construct the perturbation  $\widehat{\mathcal{B}}(t)$  on segments  $[iT, (i + 1)T]$ ,  $i = 2, 3, \dots$ , basing on solution  $x_i(t)$  similarly to what we have done above.

**Step 5.** Consider the constructed solution  $y_0(t)$  of the system (5.3). We claim that  $y_0(t)$  has the Lyapunov exponent greater than  $\chi(\widehat{\mathcal{A}}) - \varepsilon$ . Indeed, due to (A.2), (A.6) and (A.7) it suffices to prove that for any  $i = 0, 1, 2, \dots$

$$\frac{|y_0((i + 1)T)|}{|y_0(iT)|} \geq \frac{|x_i((i + 1)T)|}{|x_i(iT)|} \exp\left(-\frac{3\varepsilon T}{4}\right).$$

It follows from construction of  $y_0(t)$  that for any fixed  $i$  the number of indices  $l$  such that inequality

$$\frac{|y_0(iT + (l + 1)T_0)|}{|y_0(iT + lT_0)|} \geq \frac{|x_i(iT + (l + 1)T_0)|}{|x_i(iT + lT_0)|} \exp\left(-\frac{\varepsilon T_0}{2}\right) \tag{A.13}$$

is not fulfilled, does not exceed 2 (for  $i = 1$  this might be only segments  $Q_j$  and  $Q_{j+1}$ , see Fig. 4). If (A.13) is not satisfied, we use the inequality

$$\frac{|y_0(iT + (l + 1)T_0)|}{|y_0(iT + lT_0)|} \geq \frac{|x_i(iT + (l + 1)T_0)|}{|x_i(iT + lT_0)|} \exp(-(2a + (2a + 1)\delta)T_0). \tag{A.14}$$

Multiplying inequalities (A.13) and (A.14) corresponding to  $l = 0, 1, \dots, m - 1$ , we obtain

$$\begin{aligned} \frac{|y_0((i + 1)T)|}{|y_0(iT)|} &> \frac{|x_i((i + 1)T)|}{|x_i(iT)|} \exp\left(-\left(\frac{\varepsilon}{2} + \frac{2(2a + (2a + 1)\delta)T_0}{T}\right)T\right) \\ &\geq \frac{|x_i((i + 1)T)|}{|x_i(iT)|} \exp\left(-\frac{3\varepsilon T}{4}\right) \end{aligned}$$

(the last inequality holds by the choice of  $T$  in **Step 2**). This completes the proof.  $\square$

**Corollary A.1.** *The perturbation  $\widehat{\mathcal{B}}(t)$  may be taken continuous.*

**Proof.** It follows from the proof that  $\widehat{\mathcal{B}}(t)$  is piecewise continuous i.e. has finitely many discontinuity points on bounded subsets of  $\mathbb{R}$ . So, we may construct a continuous matrix  $\widehat{\mathcal{B}}_1(t)$  such that  $|\widehat{\mathcal{B}}_1(t)| \leq (2a + 1)\delta$ , and

$$M = \{t : \widehat{\mathcal{B}}(t) \neq \widehat{\mathcal{B}}_1(t)\} = \bigcup_{j=1}^{\infty} \Delta_j,$$

where the length of intervals  $\Delta_j$  can be chosen arbitrarily small.

Consider the system

$$\dot{x} = (\widetilde{\mathcal{A}}(t) + \widehat{\mathcal{B}}_1(t))x. \tag{A.15}$$

Let  $\Psi(t)$  and  $\Xi(t)$  be fundamental matrices of (A.15) and (5.3) respectively, so that  $\Psi(0) = \Xi(0) = E_n$ . Then

$$\Xi(t) = \Psi(t) + \Psi(t) \int_0^t \Psi^{-1}(\tau)(\widehat{\mathcal{B}}(\tau) - \widehat{\mathcal{B}}_1(\tau))\Xi(\tau) d\tau,$$

and thus

$$|\Xi(t)| \leq |\Psi(t)| + |\Psi(t)| \int_0^t |\Psi^{-1}(\tau)| |\widehat{\mathcal{B}}(\tau) - \widehat{\mathcal{B}}_1(\tau)| |\Xi(\tau)| d\tau. \tag{A.16}$$

Denote  $u(t) = |\Xi(t)|/|\Psi(t)|$ ,  $v(t) = |\Psi^{-1}(t)||\Psi(t)||\widehat{\mathcal{B}}(t) - \widehat{\mathcal{B}}_1(t)|$ . Dividing both parts of (A.16) by  $|\Psi(t)|$ , we obtain

$$u(t) \leq 1 + \int_0^t u(\tau)v(\tau) d\tau,$$

which implies by the Grönwall–Bellman lemma

$$\frac{|\Xi(t)|}{|\Psi(t)|} = u(t) \leq \exp \left( \int_0^t v(\tau) d\tau \right) \leq \exp \left( (2a+1)\delta \sum_{j=1}^{\infty} |\Delta_j| \cdot \sup_{s \in \Delta_j} \exp(2(a+(2a+1)\delta)s) \right).$$

The last expression can be made arbitrarily close to 1, and the statement follows.  $\square$

## References

- [1] B. Aulbach, S. Hilger, Linear dynamic processes with inhomogeneous time scale, in: In Nonlinear Dynamics and Quantum Dynamical Systems, Gaussig, 1990, in: Math. Res., vol. 59, Akademie Verlag, Berlin, 1990, pp. 9–20.
- [2] A. BenAbdallah, M. Dlala, M.A. Hammami, A new Lyapunov function for stability of time-varying nonlinear perturbed systems, *Systems Control Lett.* 56 (2007) 179–187.
- [3] A.I. Bobenko, Yu.B. Suris, *Discrete Differential Geometry. Integrable Structure*, Graduate Studies in Mathematics, vol. 98, AMS, 2008, xxiv+404 pp.
- [4] S. Bodine, D.A. Lutz, Exponential functions on time scales: their asymptotic behavior and calculation, *Dynam. Systems Appl.* 12 (2003) 23–43.
- [5] M. Bohner, Some oscillation criteria for first order delay dynamic equations, *Far East J. Appl. Math.* 18 (3) (2005) 289–304.
- [6] M. Bohner, D.A. Lutz, Asymptotic behavior of dynamic equations on time scales, *J. Difference Equ. Appl.* 7 (1) (2001) 21–50.
- [7] M. Bohner, A.A. Martynyuk, Elements of stability theory of A.M. Liapunov for dynamic equations on time scales, *Nonlinear Dyn. Syst. Theory* 7 (3) (2007) 225–251.
- [8] M. Bohner, A. Peterson, *Dynamic Equations on Time Scales. An Introduction with Applications*, Birkhäuser Boston Inc., Boston, MA, 2001.
- [9] M. Bohner, A. Peterson, *Advances in Dynamic Equations on Time Scales*, Birkhäuser Boston Inc., Boston, MA, 2003.
- [10] B.F. Bylov, R.E. Vinograd, D.M. Grobman, V.V. Nemytskii, *Teoriya pokazatelei Lyapunova i ee prilozheniya k voprosam ustoichivosti* (Theory of Lyapunov Exponents and Its Application to Problems of Stability), Nauka, Moscow, 1966 (in Russian), 576 pp.
- [11] C. Carcamo, C. Vidal, The Chetaev theorem for ordinary difference equations, *Proyecciones* 31 (4) (2012) 391–402.
- [12] N.G. Chetaev, *The Stability of Motion*, Pergamon Press, New York, 1961.
- [13] S.K. Choi, D.M. Im, N. Koo, Stability of linear dynamic systems on time scales, *Adv. Difference Equ.* 2008 (2008) 670203.
- [14] J.J. DaCunha, Stability for time varying linear dynamic systems on time scales, *J. Comput. Appl. Math.* 176 (2) (2005) 381–410.
- [15] N.H. Du, L.H. Tien, On the exponential stability of dynamic equations on time scales, *J. Math. Anal. Appl.* 331 (2007) 1159–1174.
- [16] T. Gard, J. Hoffacker, Asymptotic behavior of natural growth on time scales, *Dynam. Systems Appl.* 12 (1–2) (2003) 131–148.
- [17] G.Sh. Guseinov, Integration on time scales, *J. Math. Anal. Appl.* 285 (2003) 107–127.
- [18] J. Hoffacker, C.C. Tisdell, Stability and instability for dynamic equations on time scales, *Comput. Math. Appl.* 49 (9–10) (2005) 1327–1334.
- [19] G. Hovhannisyan, Asymptotic stability for dynamic equations on time scales, *Adv. Difference Equ.* 2006 (2006) 18157.
- [20] G. Hovhannisyan, Asymptotic stability for 2x2 linear dynamic systems on time scales, *Int. J. Difference Equ.* 2 (1) (2007) 105–121.
- [21] P.E. Kloeden, A. Zmorzynska, Lyapunov functions for linear nonautonomous dynamical equations on time scales, *Adv. Difference Equ.* 2006 (2006) 69106.
- [22] S.G. Kryzhevich, The relation between central exponents of linear systems of ordinary differential equations and stability of the excited systems, *Differ. Equ.* 36 (10) (2000) 1430–1431.
- [23] W.N. Li, Some Pachpatte type inequalities on time scales, *Comput. Math. Appl.* 57 (2009) 275–282.
- [24] A.A. Martynyuk, On the exponential stability of a dynamical system on a time scale, *Dokl. Akad. Nauk* 421 (2008) 312–317.
- [25] V.M. Millionschikov, A proof of attainability for central exponents of linear systems, *Sib. Math. J.* 10 (1) (1969) 99–104.
- [26] D.B. Pachpatte, Explicit estimates on integral inequalities with time scale, *J. Inequal. Pure Appl. Math.* 7 (4) (2006) 143.
- [27] O. Perron, Über Stabilität und Asymptotisches Verhalten der Integrale von Differentialgleichungssystemen, *Math. Z.* 29 (1928) 129–160 (in German).
- [28] C. Pötzsche, S. Siegmund, F. Wirth, A spectral characterization of exponential stability for linear time-invariant systems on time scales, *Discrete Contin. Dyn. Syst.* 9 (2003) 1223–1241.
- [29] A. Reinfelds, L. Sermone, Stability of impulsive differential systems, *Abstr. Appl. Anal.* 2013 (2013) 253647.

