

# Relativistic Bennett-Brassard cryptographic scheme, relativistic errors, and how to correct them

Marek Czachor\* and Marcin Wilczewski

*Katedra Fizyki Teoretycznej i Metod Matematycznych, Politechnika Gdańska, ul. Narutowicza 11/12, 80-952 Gdańsk, Poland*

(Received 12 March 2003; published 17 July 2003)

The Bennett-Brassard cryptographic scheme needs two bases, at least one of them linearly polarized. The problem is that linear polarization, formulated in terms of helicities, is not a relativistically covariant notion: State which is linearly polarized in one reference frame becomes depolarized in another one. We show that a relativistically moving receiver of information should define linear polarization with respect to projection of Pauli-Lubanski's vector in a principal null direction of the Lorentz transformation which defines the motion, and not with respect to the helicity basis. Such qubits do not depolarize.

DOI: 10.1103/PhysRevA.68.010302

PACS number(s): 03.67.Dd, 03.30.+p

In nonrelativistic quantum mechanics a generic state of a free particle with spin takes the form where spin and momentum degrees of freedom are nonentangled, i.e.,

$$\begin{pmatrix} \psi_0(\mathbf{p}) \\ \psi_1(\mathbf{p}) \end{pmatrix} = \begin{pmatrix} \psi_0 \\ \psi_1 \end{pmatrix} \psi(\mathbf{p}). \quad (1)$$

This is the reason why it is possible to base the concept of a nonrelativistic qubit on a two-dimensional Hilbert space. In particular, observables associated with spin are always of form  $A \otimes \mathbf{1}$ , where  $\mathbf{1} = \int d^3p |\mathbf{p}\rangle \langle \mathbf{p}|$  is the identity in momentum space and  $A$  stands for a spin operator. Formula  $\text{Tr}_p(A \otimes \mathbf{1}) = \text{Tr}_p A$  allows one to define states of qubits in terms of  $2 \times 2$  reduced density matrices.

In relativistic quantum mechanics, a generic state satisfies

$$\begin{pmatrix} \psi_0(\mathbf{p}) \\ \psi_1(\mathbf{p}) \end{pmatrix} \neq \begin{pmatrix} \psi_0 \\ \psi_1 \end{pmatrix} \psi(\mathbf{p}). \quad (2)$$

The origin of this property is very deeply rooted in the structure of unitary representations of the Poincaré group. A qubit which takes form (1) in one reference frame will be seen in form (2) by another observer. A Poincaré transformation necessarily involves multiplication by  $\mathbf{p}$ -dependent  $SU(2)$  matrices, a fact making form (1) noncovariant. Definitions of qubits in terms of reduced density matrices with traced-out momenta are no longer justified. This is why quantum information theory based on such a formal notion of qubit [1–4] is in danger of internal physical inconsistency.

Constructing nonzero-spin unitary representations of the Poincaré group, we always encounter certain spinor structure. The simplest representation corresponds to mass  $m$  and spin  $1/2$ . Whenever we write the state in form (2), we *implicitly* choose a “spin-quantization axis” and spin is here associated with the second Casimir invariant of the group,  $W^a W_a$ , where  $W_a$  is the Pauli-Lubanski (PL) vector.

The most popular choice of quantization axis corresponds to a timelike direction  $t^a = (1, 0, 0, 0)$ . The resulting spin operator  $t^a W_a$  is proportional to the helicity (in order to obtain directly the helicity, one should choose  $t^a = (1/|\mathbf{p}|, 0, 0, 0)$ ). In application to quantum cryptography we need several differ-

ent yes-no observables, and helicity eigenstates are not sufficient. Natural candidates for such yes-no observables are projectors on linear combinations of opposite helicities, i.e., linear polarizations. The problem with linear polarizations defined in terms of helicities is that different momentum components undergo different  $SU(2)$  transformations. In the photon case, the  $SU(2)$  transformations are diagonal and multiply opposite helicities by phase factors whose phases are opposite and momentum dependent: A wave packet which is linearly polarized in one reference frame becomes a combination of different linear polarizations in another reference frame and, hence, depolarized (cf. Sec. 2.5 in Ref. [5]). An exception occurs for massless wave packets consisting exclusively of parallel momenta since the Wigner phase factor is independent of  $|\mathbf{p}|$ . Below, we shall see that geometrically the effect is rooted in noninvariance of  $t^a = (1, 0, 0, 0)$  under nontrivial Lorentz boosts.

Different quantization axes lead to different yes-no observables. Taking  $t^a = (0, \mathbf{t})$  we arrive at observables equivalent to spins defined via relativistic center of mass [6] or, in the Dirac case, to the so-called even part of the Dirac spin. The even part is obtained from Dirac's  $\Sigma$  by sandwiching it between projectors on signs of energy. Physically, one eliminates in this way the *Spinbewegung* oscillations [7]. The applications of such spins to the relativistic Einstein-Podolsky-Rosen (EPR) problem were given by one of us many years ago (even part of Dirac's spin in Ref. [8], relativistic center of mass, the PL vector, and even spin in Refs. [9,10]). Quite recently, the review [11] discusses in the same context Dirac's and rest-frame spins, however, the link of Dirac's  $\Sigma$  to the results of Ref. [9] was here overlooked.

During the past year, various relativistic aspects of EPR correlations were discussed in a series of detailed works. The intriguing paper [12] starts with a definition of spin in terms of a generator of rotations but taken from a representation of the Poincaré group involving a nonstandard clock synchronization. This seems to be the first work where details related to spatial localization of measurements were taken into account, and the conclusion is that EPR correlations might, in principle, reveal a preferred reference frame. An approach starting *ab initio* from the level of quantum electrodynamics can be found in Ref. [13]; as opposed to the approach advocated in Refs. [1,11] the momentum degrees of freedom are taken into account in measurements of spin and the loophole

\*Email address: mczachor@sunrise.pg.gda.pl

of the argument given in Ref. [1] is not present. The roles of appropriate choices of spin measurements in EPR experiments were discussed in Refs. [14,15]. The authors stress differences of their analysis with the one given in Ref. [9], but the main conclusion (the degree of violation of the Bell inequality decreases with increasing velocities of the observers) remains unchanged. Similar conclusions can be found in Ref. [16]. Finally, a completely new direction of theoretical investigation was initiated by the work in Ref. [17], and extended in Refs. [18,19]. The problem is what happens if the observers move noninertially. One expects here a host of new quantum-field-theoretic phenomena related to inequivalence of vacua in accelerated frames. The situation awaits a detailed review, especially in the context of possible experiments.

In the present paper, we concentrate on the choice of qubits appropriate for cryptographic problems involving relativistically moving observers. At the level of first quantization we do not experience the subtleties related to the Unruh effect [17–19] and can, in principle, also discuss noninertial motions. Quantization in curved spaces and accelerated systems is still an open problem [20], so we prefer to concentrate on purely kinematic phenomena which can be naturally treated at the level of representations of the Poincaré group.

We define qubits by vectors from the mass- $m$  spin-1/2 unitary representation of (covering space of) the Poincaré group. In momentum space the qubits are given by pairs of functions  $(f_0(p), f_1(p))$  and  $p^2 = m^2$ . It is essential that whenever one writes pair  $(f_0(p), f_1(p))$ , one implicitly chooses a basis of states corresponding to a projection of the PL vector in some direction  $t^a \in \mathbb{R}^4$  in Minkowski space. This choice is implicitly present in the transformation properties of the qubit [21].

PL vector  $W_a = P^b * S_{ab}$  is a tensor operator [22] and under the action of the Poincaré group its projection in direction  $t^a$  gets transformed by

$$\mathcal{U}_{y,\Lambda}^{-1} t^a W_a \mathcal{U}_{y,\Lambda} = t^a \Lambda_a^b W_b, \quad (3)$$

where  $\mathcal{U}_{y,\Lambda}$  is a unitary representation. One can say that a moving particle experiences measurements of spins in Lorentz-modified directions  $t^a \Lambda_a^b$ . Of particular interest are directions  $t^a$  satisfying the eigenvalue condition  $t^a \Lambda_a^b \sim t^b$  since they lead to Lorentz-invariant yes-no observables. Eigenvectors of Lorentz transformations are known to be given by null vectors ( $t^2 = t^a t_a = 0$ ) and any  $\Lambda \in \text{SL}(2, \mathbb{C})$  possesses at least one and at most two, such eigendirections [principal null directions (PNDs)] [23].

Accordingly, it is very natural to contemplate projections of the PL vector in null directions instead of the usual timelike or spacelike ones. Moreover, the projection of the PL vector in momentum direction vanishes,  $P^a W_a = 0$ , and therefore we obtain a kind of gauge freedom: For any parameter  $\theta$ , observables  $t^a W_a$  and  $(t^a + \theta P^a) W_a$  are identical. An appropriate choice of  $\theta$  and a  $P$ -dependent  $t^a$  will allow us to work with invariant yes-no observables which are equivalent to projections of PL vector in directions perpendicular to the four-momentum. In what follows, we shall describe a simple procedure leading to such invariant yes-no observables.

A formalism which almost ideally suits the purposes of relativistic quantum information theory is the two-spinor calculus, especially in the form developed by Penrose [23]. The unitary representations of the Poincaré group can be translated into a two-spinor language by means of ‘‘Bargmann-Wigner (BW) spinors’’ [24]. One exploits here the special role played in two-spinor formalism by null directions. Four-momentum  $p_a$  is split into a linear combination of two null directions  $\pi_a$  and  $\omega_a$ , defined by

$$p_a = \pi_a + (m^2/2)\omega_a = \pi_A \bar{\pi}_{A'} + (m^2/2)\omega_A \bar{\omega}_{A'}, \quad (4)$$

where  $\pi_A, \omega_A$  is a field of spin-frames, i.e.,  $\omega_A \pi^A = 1$ . If  $\nu_A$  is any  $p$ -independent spinor, the spin frame may be taken as

$$\omega^A(p) = \frac{\nu^A}{\sqrt{p^{BB'} \nu_B \bar{\nu}_{B'}}} = \omega^A(\nu, p), \quad (5)$$

$$\pi^A(p) = \frac{p^{AA'} \bar{\nu}_{A'}}{\sqrt{p^{BB'} \nu_B \bar{\nu}_{B'}}} = \pi^A(\nu, p). \quad (6)$$

One can directly verify that the spin frame satisfies Eq. (4) and

$$\Lambda \pi_A(\nu, p) = \Lambda_A^B \pi_B(\nu, \Lambda^{-1} p) = \pi_A(\Lambda \nu, p), \quad (7)$$

$$\Lambda \omega_A(\nu, p) = \Lambda_A^B \omega_B(\nu, \Lambda^{-1} p) = \omega_A(\Lambda \nu, p). \quad (8)$$

The latter formulas will be crucial for our analysis of relativistic qubits. The simplest unitary representation is characterized by mass  $m$  and spin-1/2 and its BW-spinor form reads

$$\mathcal{U}_{y,\Lambda} f_A(p) = e^{iy p} U(\Lambda, p)_A^B f_B(\Lambda^{-1} p), \quad (9)$$

where  $U(\Lambda, p)_A^B \in \text{SU}(2)$  and  $p^2 = m^2$ . The BW-spinor indices are written in the calligraphic font to distinguish them from the  $\text{SL}(2, \mathbb{C})$  ones. The matrix

$$U(\Lambda, p)_A^B = \begin{pmatrix} \omega_A(p) \Lambda \pi^A(p) & -\frac{m}{\sqrt{2}} \omega_A(p) \Lambda \omega^A(p) \\ \frac{m}{\sqrt{2}} \bar{\omega}_{A'}(p) \Lambda \omega^{A'}(p) & \bar{\omega}_{A'}(p) \Lambda \pi^{A'}(p) \end{pmatrix} \quad (10)$$

is responsible for changes of the ‘‘polarization’’ whereas  $f_B(\Lambda^{-1} p)$  introduces Doppler shifts.

One can show [24] that the amplitudes  $f_0(p)$  and  $f_1(p)$  play, for Dirac electrons, the roles of momentum-space wave functions associated with eigenvectors of the projection of the PL vector in null direction  $t^a = \omega^a$ , with eigenvalues  $-1/2$  and  $+1/2$ , respectively. Choosing  $\theta = -m^{-2}$  we find that the projection  $\omega^a W_a$  is an observable identical to the projection of  $W_a$  in direction  $\omega^a - m^{-2} p^a$  which is spacelike and orthogonal to the four-momentum.

At the level of BW spinors we do not have to make any reference to the Dirac equation but can directly compute the

generators, the PL vector, and its projection in any direction. In particular, in momentum space the projection in the null direction turns out to be

$$\omega^a(p)W_a(p)_{A^B} = \frac{1}{2} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = -\frac{1}{2}\sigma_3, \quad (11)$$

which agrees with the fact that amplitudes  $f_0(p)$  and  $f_1(p)$  correspond, respectively, to eigenvalues  $-1/2$  and  $+1/2$  at the level of the Dirac equation.

The spin operator we have introduced through projection of the PL vector on the null quantization axis  $\omega^a(p)$ , has led to the standard-looking spin. So what have we gained with respect to the earlier works where  $\sigma_3$  was taken for granted as the correct operator associated with relativistic qubits? The gain is that we have arrived at  $\sigma_3$  by means of a systematic procedure and have the relativistic transformation properties of qubits under control. Recall that in addition to Eq. (11) we have representation (9) where matrix  $U(\Lambda, p)_{A^B}$  does not, in general, commute with  $\sigma_3$ .

Notice, however, that the actual problem we will need to solve in practical quantum communication is how to correct the errors which are due to a relativistic and perhaps noninertial motion  $s \mapsto \Lambda(s)$  of an observer. The problem can be reduced to an appropriate choice of quantization axis that defines the qubit.

A PND associated with  $\Lambda \in \text{SL}(2, C)$  is the flagpole direction of an eigenspinor of  $\Lambda$ , i.e.,

$$\Lambda_A^B \nu_B = \lambda \nu_A, \quad (12)$$

where  $\lambda = |\lambda|e^{i\varphi}$  is in general, complex. Classification of PNDs of  $\text{SL}(2, C)$  transformations can be found in Ref. [23]. Inserting Eq. (12) into Eqs. (5)–(8), we find

$$\Lambda \pi_A(\nu, p) = e^{-i\varphi} \pi_A(\nu, p), \quad \Lambda \omega_A(\nu, p) = e^{i\varphi} \omega_A(\nu, p)$$

and

$$U(\Lambda, p)_{A^B} = \begin{pmatrix} e^{-i\varphi} & 0 \\ 0 & e^{i\varphi} \end{pmatrix}, \quad (13)$$

where  $\varphi$  is momentum independent. The independence of momentum is important since transformation  $s \mapsto \Lambda(s)$  affects all the momentum components in the same way. An arbitrary linearly polarized state is now transformed as follows:

$$\begin{pmatrix} f_0(p) \\ f_1(p) \end{pmatrix} \mapsto \begin{pmatrix} \mathcal{U}_{y,\Lambda} f_0(p) \\ \mathcal{U}_{y,\Lambda} f_1(p) \end{pmatrix} = e^{iyp} \begin{pmatrix} e^{-i\varphi} f_0(\Lambda^{-1}p) \\ e^{i\varphi} f_1(\Lambda^{-1}p) \end{pmatrix},$$

and a linear polarization goes into linear polarization, perhaps rotated by some angle. In particular, product form (1) is covariant. This was possible only because we replaced helicity qubits by qubits related to an invariant direction.

For quantum cryptographic protocols, such as Bennett-Brassard cryptographic scheme (BB84), it may be important to allow for motions that are characterized by two different

PNDs. As an example, consider a general accelerated motion (of Alice, say) in a  $z$ -direction. The relevant  $\text{SL}(2, C)$  transformation reads

$$\Lambda_A^B = \begin{pmatrix} w^{1/2} & 0 \\ 0 & w^{-1/2} \end{pmatrix},$$

$w = \sqrt{1+\beta}/\sqrt{1-\beta}$  and  $\beta = v(s)/c$ . The eigenvalues of  $\Lambda$  are real and there are two eigenvectors

$$\nu_A^{(-)} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \nu_A^{(+)} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (14)$$

In both cases, we find  $U(\Lambda, p) = \mathbf{1}$  (since  $\varphi = 0$ ) and

$$\begin{pmatrix} \mathcal{U}_{y,\Lambda} f_0(\nu^{(\pm)}, p) \\ \mathcal{U}_{y,\Lambda} f_1(\nu^{(\pm)}, p) \end{pmatrix} = e^{iyp} \begin{pmatrix} f_0(\nu^{(\pm)}, \Lambda^{-1}p) \\ f_1(\nu^{(\pm)}, \Lambda^{-1}p) \end{pmatrix}. \quad (15)$$

Amplitudes  $f_A(\nu^{(-)}, p)$  and  $f_A(\nu^{(+)}, p)$  represent wave functions associated with projections of the PL vector in invariant directions  $\omega^a(\nu^{(\pm)}, p) = (p^0 \mp p^3)^{-1} (1, 0, 0, \pm 1)$ . The  $p$ -dependent denominators come from the denominator in  $\omega_a(\nu^{(\pm)}, p) = \nu_a^{(\pm)}/(p \nu^{(\pm)})$  and could be also skipped since the null directions of  $\nu_a^{(\pm)}$  and  $\omega_a(\nu^{(\pm)}, p)$  are identical. The yes-no observables are defined by normalization of eigenvalues to  $\pm 1$ . This is similar to the problem of choosing  $t^a$  associated with the helicity.

Formula (15) illustrates the role of appropriate choices of quantization axes. No rotations of qubits are involved, the relativistic corrections are reduced to the Doppler shifts, and form (1) will be preserved.

The associated yes-no observables are given in both cases by  $\sigma_3$  but, of course, two different bases are involved. For future references, we give here the general form of the  $\text{SU}(2)$  transformation that maps qubits associated with a null  $\omega^a$  direction into those associated with the most general direction  $t^a$  (null, timelike, or spacelike, and in general  $p$ -dependent). Let

$$\begin{pmatrix} \Omega(t, p)_A \\ \Omega(t, p)_{A'} \end{pmatrix} = -\{8\lambda(t, p)[\lambda(t, p) + tp - m^2(t\omega)]\}^{-1/2} \times \begin{pmatrix} [2\lambda(t, p) + tp] \pi_A - \pi_C t^C_{X'} p_A^{X'} + 3\frac{m^2}{2} t_{AX'} \bar{\omega}^{X'} \\ \frac{m}{\sqrt{2}} \{ [2\lambda(t, p) - tp] \bar{\omega}_{A'} - \bar{\omega}_C t^C_{X'} p_{A'}^X - 3t_{XA'} \pi^X \} \end{pmatrix}.$$

Then  $f_A(t, p) = \mathcal{W}(p)_{A^B} f_B(\omega, p)$ , where

$$\mathcal{W}(p)_{A^B} = \begin{pmatrix} \bar{\omega}(p)^{A'} \bar{\Omega}(t, p)_{A'} & \omega(p)^A \bar{\Omega}(t, p)_A \\ -\bar{\omega}(p)^{A'} \Omega(t, p)_{A'} & \omega(p)^A \Omega(t, p)_A \end{pmatrix}, \quad (16)$$

$\bar{\Omega}(t,p)_A = \overline{\Omega(t,p)_{A'}}$ ,  $\bar{\Omega}(t,p)_{A'} = \overline{\Omega(t,p)_A}$ , and  $\lambda(t,p) = \sqrt{(tp)^2 - m^2 t^2}$ . Notice that in momentum space, the eigenvalues of the projection of the PL vector in a direction  $t^a$  are given by  $\pm \frac{1}{2}\lambda(t,p)$ .

Now, set  $\omega^A(p) = \omega^A(\nu^{(-)}, p)$ ,  $\pi^A(p) = \pi^A(\nu^{(-)}, p)$ , and  $t^a(p) = \omega^a(\nu^{(+)}, p)$ , where the invariant spinors are given by Eq. (14). We find

$$f_{\mathcal{A}}(\nu^{(+)}, p) = \mathcal{W}(p)_{\mathcal{A}}^{\mathcal{B}} f_{\mathcal{B}}(\nu^{(-)}, p), \quad (17)$$

$$\mathcal{W}(p)_{\mathcal{A}}^{\mathcal{B}} = \frac{1}{\sqrt{1+|\zeta|^2}} \begin{pmatrix} |\zeta| & e^{i\chi} \\ -e^{-i\chi} & |\zeta| \end{pmatrix} \in \text{SU}(2),$$

where  $\zeta = (p_1 + ip_2)/m = |\zeta|e^{i\chi}$  is invariant under Lorentz boosts along the third axis. The matrix in Eq. (17) itself is, therefore, also invariant under transformations that do not change the quantization axes.

As the next application let us consider the case where the  $\text{SL}(2, \mathbb{C})$  transformations do not commute with one another if taken at different points on curve  $s \mapsto \Lambda(s)$ , i.e.,  $[\Lambda(s), \Lambda(s')] \neq 0$ . A good illustration is a composition of the previously discussed boost with a null rotation, i.e.,

$$\Lambda(s)_A^B = \begin{pmatrix} w(s)^{1/2} & 0 \\ 0 & w(s)^{-1/2} \end{pmatrix} \begin{pmatrix} 1 & \alpha(s) \\ 0 & 1 \end{pmatrix}. \quad (18)$$

There is only one eigenvector, namely,  $\nu_A^{(-)}$ . The construction is unchanged but we must make sure the spin is projected on null axis  $\omega^a(\nu^{(-)}, p)$ . This presence of only one invariant direction is not a problem since one direction is enough to define linear polarizations.

Finally, let us consider the case of photons. The Poincaré transformation reads

$$\mathcal{U}_{y,\Lambda} f_{\mathcal{A}\mathcal{A}}(p) = e^{iyp} U(\Lambda, p)_{\mathcal{A}}^{\mathcal{B}} U(\Lambda, p)_{\mathcal{A}}^{\mathcal{B}} f_{\mathcal{B}\mathcal{B}}(\Lambda^{-1}p), \quad (19)$$

where  $U(\Lambda, p)_{\mathcal{A}}^{\mathcal{B}}$  is the zero-mass version of Eq. (10). There are only two degrees of freedom, represented by  $f_{00}(p)$  and  $f_{11}(p)$ .

Diagonal elements  $U(\Lambda, p)_0^0$  and  $U(\Lambda, p)_1^1$  are momentum-dependent phase factors. As before, an appropriate choice of  $\nu$  in spin frames reduces  $U(\Lambda, p)$  to Eq. (13), where  $\varphi$  is momentum independent. In the massless case there exists a topological restriction that does not occur for massive fields: For a fixed  $\nu_A$ , there exists a momentum  $p_a$  parallel to null vector  $\nu_a = \nu_A \bar{\nu}_{A'}$ , where the spin frames are not defined (actually this happens for the direction given by the class of momenta parallel to  $p_a$ ). Usually this is not a difficulty since one can define the spin frames locally. In our problem, this is a difficulty that restricts the choice of  $\nu_a$  to null vectors that are not proportional to any four-momentum in the wave packet. The case where the wave packet does contain a vector parallel to  $\nu_a$  has to be treated with some care.

The noise due to relativistic helicity-basis depolarization can be reduced for massless fields if wave packets  $f_{\mathcal{A}\mathcal{A}}(p)$  consist of vectors parallel to some given direction. To what extent an approximation of a realistic signal by such a plane-wave-like front is acceptable depends on the geometry of the experiment. Qubits associated with PNDs do not impose any restriction on momentum-space wave packets.

We are grateful to J. Rembieliński for comments.

- [1] A. Peres, P.F. Scudo, and D.R. Terno, Phys. Rev. Lett. **88**, 230402 (2002).
- [2] A. Peres and D.R. Terno, Rev. Mod. Phys. (to be published), e-print quant-ph/0212023.
- [3] R.M. Gingrich and C. Adami, Phys. Rev. Lett. **89**, 270402 (2002).
- [4] A.J. Bergou, R.M. Gingrich, and C. Adami, e-print quant-ph/0302095.
- [5] S. Weinberg, *The Quantum Theory of Fields: Foundations* (Cambridge University Press, Cambridge, 1995).
- [6] G.N. Fleming, Phys. Rev. **137**, B188 (1965); **139**, B963 (1965).
- [7] R.F. Guertin and E. Guth, Phys. Rev. D **7**, 1057 (1973).
- [8] M. Czachor, University of Gdansk preprint, 1984 (unpublished).
- [9] M. Czachor, Phys. Rev. A **55**, 72 (1997).
- [10] M. Czachor, in Photonic Quantum Computing, Proceedings of AeroSense97, Orlando, FL, 1997, edited by S.P. Hotaling and A.R. Piritch [Proc. SPIE **3076**, 141 (1997)]; e-print quant-ph/0205187.
- [11] D.R. Terno, Phys. Rev. A **67**, 014102 (2003)
- [12] J. Rembieliński and K.A. Smoliński, Phys. Rev. A **66**, 052114 (2002).
- [13] J. Pachos and E. Solano, Quantum Inf. Comput. **3**, 115 (2003); e-print quant-ph/0203065.
- [14] H. Terashima and M. Ueda, Q. Inf. Comput. **3**, 224 (2003); e-print quant-ph/0204138.
- [15] H. Terashima and M. Ueda, Int. J. Quant. Inf. **1**, 93 (2003); e-print quant-ph/0211177.
- [16] D. Ahn, H. Lee, Y.H. Moon, and S.W. Hwang, Phys. Rev. A **67**, 012103 (2003).
- [17] P.M. Alsing and G.J. Milburn, Quantum Inf. Comput. **2**, 487 (2002).
- [18] X. Anderson, S.J. van Enk, and T. Rudolph, e-print quant-ph/0302091.
- [19] P.M. Alsing and G.J. Milburn, e-print quant-ph/0302179.
- [20] N.D. Birell and P.C.W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1982).
- [21] C.H. Bennett and G. Brassard, in *Proceedings of IEEE, International Conference on Computers, Systems and Signal Processing*, Bangalore, India (IEEE, New York, 1984), p. 175.
- [22] A.O. Barut and R. Rączka, *Theory of Group Representations and Applications* (Polish Scientific Publishers, Warszawa, 1980)
- [23] R. Penrose and W. Rindler, *Spinors and Space-Time* (Cambridge University Press, Cambridge, 1984), Vol. 1.
- [24] M. Czachor, in *Photon and Poincaré Group*, edited by V. V. Dvoeglazov (Nova, New York, 1999), p. 32; e-print hep-th/9701135.